

Interactive comment on “Synergistic effect of water-soluble species and relative humidity on morphological changes of aerosol particles in Beijing mega-city during severe pollution episodes” by Xiaole Pan et al.

Xiaole Pan et al.

panxiaole@mail.iap.ac.cn

Received and published: 8 November 2018

The authors appreciate the anonymous reviewer very much for providing insight comments on the manuscript, we would like to revise the context on the basis of suggestions and advises carefully. The specific replies are as follows:

Major comment #1, Line 143-145, page 5: Since the aerosol flow was further diluted by zero air (920 ccm, 38% RH), what was the RH of the flow after dilution? I assume that the POPC measured the depolarization ratios at $\sim 38\%$ RH since the dilution fac-

C1

tor by the zero air was >10 . This information is critical because the morphology and measured depolarization ratios of aerosol particles depend on RH. Please clarify it.

Reply: we agree with the comment that physical property of aerosol is critically depend on the humid condition, and mixing of sampling aerosol flow with dry diluting flow will change the morphology of particles. For the measurement of instruments, the exhausted air from measuring chamber was filtered and recycled to sever as diluting and sheath flow. Since the sampling flow rate was much lower than the sum of diluting and sheath flow air, relative humidity of the flow after dilution was the same as the dilution flow. As the reviewer mentioned, the original morphology of particle in the measuring chamber might be altered as a result of mixing with dry air. Besides, the temperature in the measuring chamber was $\sim 4^\circ\text{C}$ higher than the ambient temperature which also lead to decrease of relative humidity. Therefore, the depolarization ratio in this study refer to the value in the dry condition. We will clarify this point in the revised manuscript.

Major comment #2: Section 3.5: I consider this section as the most important part of this manuscript. However, the discussion is far too short and rather descriptive. A number of previous studies have investigated the hygroscopic properties of $\text{Ca}(\text{NO}_3)_2$ aerosol particles, and I would refer the authors to these papers (Gibson et al., 2006; Guo et al., 2018). I can understand depolarization ratio depended on nitrate fraction (as shown in Figure 8), but what was it also affected by RH? It is related to dependence of aerosol liquid water content on RH? This question is also related to my first major comment: did the author measured the depolarization ratio at ambient RH or at the RH after the aerosol flow was diluted by zero air?

Reply: As suggested by reviewer, the relationship between depolarization ratio of particles in coarse mode and the relative humidity is an important indicator for investigating the heterogeneous reaction on the surface of dust particles. During the observation period, the relative humidity in the measuring chamber was stable at $34.3 \pm 1.6\%$, indicating that physical properties of particles in measuring chamber would have been altered, in particular at high ambient RH condition (Before 2017 January 9). There-

C2

fore, the depolarization ratio of the particles reported in this study was representative of that at relatively dry condition, not the ambient condition. Nevertheless, some water-soluble species could absorb water at very low relative humidity condition (Gibson et al., 2006). Take $\text{Ca}(\text{NO}_3)_2$ for instance, it would be deliquescent at relative humidity less than 10%, which means that they may still exist in liquid phase at relative humidity of $\sim 34\%$, though efflorescent process may reduce its sphericity. In Beijing, the major water soluble composition of the particles in coarse mode was nitrate, as reported by numbers of previous studies. Presuming the counterpart of nitrate was calcium, the water content could be calculated, and the influence of reduction in relative humidity in the measuring chamber could be also evaluated.

As shown in Fig.1, the mass concentration of H_2O as a result of deliquescent effect of nitrate in coarse mode was calculated. During the high cNO_3 and relative humidity condition, the water content of particle in coarse mode reduced by $\sim 40\%$ on average, accounting for less 10% of total mass of particle in coarse mode. The percentage reduction in diameter of particles was calculated to be less than 5%, implying that the evaporation of water in the measuring chamber did not change our main conclusion. As suggested, now we are planning an experiment using HTAAC (humidifying Tandem Aerodynamic Aerosol Classifier) in laboratory to check the relationship between depolarization ratio of dust particles and relative humidity, and the modification of morphology of dust particle due to reaction with nitric acid will be also considered.

Minor comments:

Line 24-25, page 1: When depolarization ratio decreases, does the particle become more spherical or more non-spherical? In the abstract a short introduction to this parameter should be included, though more information of this parameter can be found in page 3.

Reply: The information will be included in introduction section.

Line 53-54, page 2: A number of seminal papers (Krueger et al., 2003; Laskin et al.,

C3

2005; Tang et al., 2016) on HNO_3 - CaCO_3 reactions should be cited here.

Reply: The literatures will be cited in revised manuscript.

Line 55, page 2: For dust-cloud interactions, two studies (Sullivan et al., 2010; Tang et al., 2015) measured the CCN activity of CaCO_3 and its aging products and should be cited here.

Reply: The literatures will be cited in revised manuscript.

Line 361, page 12: please include proper literature.

Reply: Relevant literatures will be cited in revised manuscript.

Line 363, page 12: To support their claim, the author should cite previous studies (Sullivan et al., 2009; Ma et al., 2013; Gu et al., 2018) which showed the hygroscopicity of CaSO_4 is very low.

Reply: Relevant literatures will be cited in revised manuscript.

Line 347-349, page 11-12: A recent study (Wu et al., 2018) which explored the aerosol liquid water content and its impact on secondary particle formation should be cited and discussed here.â

Reply: Relevant literatures will be cited in revised manuscript.

Technical comments: Line 42, page 2: change "emit" to "emitted" or "have emitted".â
Line 99, page 4: change "was consecutively suffered from" to "consecutively suffered from".â
Line 152, page 5: change "note" to "noted".â
Line 175, page 6: please change "in consistent well" to "in good consistence".
Line 210, page 6: please change "Number of" to "A number of".â
Line 211, page 6: change "was" to "were".

Reply: All the Technical errors will be corrected in the revised manuscript, and we will check all grammar and expression in English throughout the content.

C5




Fig. 1. Temporal variations in mass concentration of nitrate in coarse mode (black line), relative humidity (gray line), calculated water content in coarse mode at ambient condition (red line with circ

C6