

1 **Replies to reviewer #2**

2 *At the outset, we thank the reviewer for positive and constructive comments that improved the*
3 *quality of the manuscript.*

4 **Comment:** Figure 5: Why CER of the ice show a decreasing trend and CER of water
5 showing an increasing trend over BOB beyond 30°C? Whereas over AS, both CER liquid and
6 Ice shows an increasing trend?

7 **Reply:** *The main reason for studying CER at different SST is to understand whether or not*
8 *the observed differences originated at the formation of cloud stage. For that, CER for water*
9 *is sufficient. Therefore, figure and text related to CER for ice are removed from the revised*
10 *manuscript.*

11 *Regarding reviewers' query, yes, there are some small differences in the variation of CER for*
12 *ice and water with SST above 30 °C, but they are not significant.*

13 **Comment:** Figure 5: Why CER of ice (water) shows a reverse trend beyond 30°C (28.5°C)
14 over AS and BoB.

15 **Reply:** *The CER depends on the ambient atmospheric aerosol concentration and availability*
16 *of water vapor. The variation of AOD with SST is substantial over the AS while it is marginal*
17 *over the BOB. As the SST increases AOD decreases and TCWV increases results in increase*
18 *in CER over the AS and is more prominent at higher SSTs (where the decrease of AOD with*
19 *SST is quite substantial). On the other hand, the decrease in AOD with SST is quite marginal*
20 *over BOB and in fact, AOD increases from 30 °C to 31 °C. Therefore, the CER for water*
21 *continuously increases with rapid increase beyond 28 °C over AS, while the increase is*
22 *marginal over BOB.*

23 **Comment:** Figures 2 and 5: Higher values of reflectivities beyond 8 km beyond 30°C over
24 AS is due to the higher values of CER liquid (Fig. 5)? That means higher convection over AS
25 than BOB?

26 Whether similar explanation holds good for LTS over AS?

27 **Reply:** *The differences in Z over AS and BOB at and above 8 km is very small (within 1 dBZ)*
28 *and not significant. Therefore, we are not attributing these to any physical or microphysical*
29 *processes.*

30
31
32
33
34
35
36
37
38

39

Referee #4

40 *At the outset, we thank the reviewer for positive and constructive comments that improved the*
41 *quality of the manuscript.*

42 **Comment:** This study investigated the variability in the vertical structure of precipitation as a
43 function of sea surface temperature using TRMM precipitation radar measurements. I think
44 the paper lacks focus, inadequate analysis, and insufficient literature review. The intent of the
45 paper digresses at some point by incorporating the aerosol/cloud radiation analysis without a
46 context jumbling both convective dynamics and radiative impacts of aerosols on clouds.
47 Given the scope, the section with aerosol and radiation properties is redundant. Most of the
48 analysis lacks context. Overall, the quality and the content of the present paper are poor.

49 **Reply:** *The aim of the present study is to understand differences in the variation of vertical*
50 *structure of precipitation with SST over the Arabian Sea and Bay of Bengal. SST being the*
51 *main driving force to trigger precipitating systems through air-sea interactions, the*
52 *occurrence of precipitation top height and intensity profiles (reflectivity) as a function of SST*
53 *are studied. Besides SST, the vertical structure can be modified by aerosols (or CCN, mostly*
54 *at the cloud formation stage) and thermodynamics of the ambient atmosphere. In the revised*
55 *manuscript, all these parameters are considered to explain the differences in the vertical*
56 *structure. Aerosols are considered only for understanding variation in cloud effective radius,*
57 *nevertheless their radiative effects (direct, indirect, etc.) are not considered in the present*
58 *study. Recent studies, indeed, have shown the impact of aerosols (PM_{10}) on the vertical*
59 *structure of precipitation (Gao et al., 2018 and references therein).*

60 *We have rewritten the introduction with more focus on the above aspects and highlighting the*
61 *known differences in various aspects/parameters over AS and BOB. The literature survey is*
62 *also improved considerably in the revised manuscript by adding appropriate references (Guo*
63 *et al. 2018; Nuijens et al. 2017; Weller et al. 2016; Sathiyamoorthy et al. 2013; Takayabu et*
64 *al. 2010; Bhat et al. 2001; Ramanathan et al. 2001; Gadgil 2000; Krishnamurti 1981;*
65 *Narayanan and Rao 1981;).*

66 **Comment:** Introduction lacks discussion on how Arabian Sea and Bay of Bengal regions are
67 distinctly different in its background state, which would help them explain the further
68 analysis on convective profiles. Though the authors have claimed to have studied the
69 “causative mechanisms” of SST with the vertical structure of precipitation in the
70 introduction, no suggestions based on the analysis performed have been discussed in the later
71 sections. Mere correlation doesn’t explain the causality, which needs carefully controlled
72 model experiments with a rigor to assess the confounding factors controlling the SST and
73 precipitation relationship.

74 **Reply:** *The introduction of the revised manuscript is modified by considering all the*
75 *suggestion of the reviewer. The role of the surrounding seas on the rainfall over the Indian*
76 *landmass is stated and the differences between the two seas are clearly mentioned with*
77 *proper references in the revised manuscript as follows:*

78 *Indian summer monsoon (ISM) is one of the most complex weather phenomena,*
79 *involving coupling between the atmosphere, land and ocean. At the boundary of the ocean*
80 *and atmosphere air-sea interactions play a key role for the coupled Earth system (Wu and*
81 *Kirtman 2005; Feng et al. 2018). SST – precipitation relations are the important measures*
82 *for the air-sea interactions on different temporal scales (Woolnough et al., 2000; Rajendran*
83 *et al. 2012). Recent studies (Wang et al. 2005; Rajeevan et al. 2012; Chaudhari et al. 2013;*
84 *2016; Weller et al. 2016; Feng et al. 2018) have shown that the simulation of ISM can be*
85 *improved with the exact representation of sea surface temperature (SST) - precipitation*
86 *relationship. SST modulates the meteorological factors that influence the formation and*
87 *evolution of different kinds of precipitating systems over tropical oceans (Gadgil et al. 1984;*
88 *Schumacher and Houze, 2003; Takayabu et al. 2010; Oueslati and Bellon 2015).*

89 *The studies dealing with SST and cloud/precipitation population considered whole*
90 *Indian Ocean as a single entity (Gadgil et al. 1984; Woolnough et al., 2000; Rajendran et al.*
91 *2012; Sabin et al. 2012; Meenu et al. 2012; Nair and Rajeev 2014; Roxy 2014; Nair et al.*
92 *2017). But in reality the BOB and the AS of Indian Ocean possesses distinctly different*
93 *features. The summer monsoon experiment (MONEX) showed the influence of the AS and the*
94 *BOB on the rainfall produced over the Indian sub-continent (Krishnamurti 1985; Houze and*
95 *Churchill 1987; Gadgil 2000; Bhat et al.2001) and also proved how these two seas are*
96 *different with respect to the other, in terms of SST, back ground atmosphere and the*
97 *occurrence of precipitating systems. The SST in the AS cools between 10°N and 20°N during*
98 *the monsoon season whereas warming is seen in other Oceans between the same latitudes*
99 *(Krishnamurthi 1981). SST variability is large over the AS than the BOB at seasonal and*
100 *intraseasonal scales (Sengupta et al. 2001; Roxy et al. 2013). The monsoonal winds are*
101 *stronger over the AS than BOB (Findlater 1969). Also, lower-tropospheric thermal*
102 *inversions are more frequent and stronger over the AS than BOB (Narayanan and Rao 1981;*
103 *Sathiyamoorthy et al. 2013). Thus, the atmospheric and sea surface conditions and in turn*
104 *the occurrence of different kinds of precipitating systems are quite different over the BOB*
105 *and the AS during the ISM period (June to September - JJAS). For instance, long-term*
106 *measurements of tropical rainfall measuring mission (TRMM) precipitation radar (PR) have*
107 *shown that shallow systems are more prevalent over the AS, while deeper systems occur*
108 *frequently over the BOB (Liu et al. 2007; Romatschke et al. 2010; Saikranthi et al. 2014;*
109 *Houze et al. 2015).*

110 *The aforementioned studies mainly focused on the morphology of vertical structure of*
111 *precipitation, but, none of them studied the variation of vertical structure of precipitation (in*
112 *terms of occurrence and intensity) with SST and the differences in the vertical structure over*
113 *AS and BOB. On the other hand, information on the vertical structure of precipitation is*
114 *essential for improving the accuracy of rainfall estimation (Fu and Liu 2001; Sunilkumar et*
115 *al. 2015), understanding the dynamical and microphysical processes of hydrometeor*
116 *growth/decay mechanisms (Houze 2004; Greets and Dejene 2005; Saikranthi et al. 2014;*
117 *Rao et al. 2016) and for improving the latent heating retrievals (Tao et al. 2006). SST being*
118 *the main driving force to trigger precipitating systems through air-sea interactions (Sabin et*
119 *al. 2012; Nuijens et al. 2017), can alter the vertical structure of precipitation (Oueslati and*
120 *Bellon 2015). Therefore, the present study aims to understand the variation of vertical*
121 *structure of precipitation (in terms of precipitation top height and intensity) with SST over the*
122 *AS and BOB. Besides the SST, vertical structure can be modified by aerosols (or CCN, mostly*
123 *at the cloud formation stage) and thermodynamics of the ambient atmosphere. For instance,*
124 *recent studies have shown the impact of surface aerosols (PM₁₀) in altering the vertical*
125 *structure of precipitation (Gao et al., 2018). All these parameters, therefore, are considered*
126 *in the present study to explain the differences in the vertical structure.*

127

128 **Comment:** Given the non-linear influence of sea surface temperature on the variability of
129 precipitation structure, it would be an oversimplification to look at the influence of SST on
130 the mean structure of radar echoes. It would have been interesting to classify the mean
131 structure further into different cloud types (e.g., shallow/congestus/deep/) and assess the
132 variability of these populations in terms of factors (e.g., winds, stability) that are co-
133 associated with SSTs. There are no insights been provided on why the differences in the
134 variabilities of vertical structure exist between AS and BOB. It is important to investigate if
135 more variability over the AS is due to fluctuations in the winds/SSTs or both. From figure 2,
136 it is evident that AS region has more seasonality in term of air-sea variables compared to
137 BOB. Given the influence of more variables, merely analyzing indirect relationships of
138 precipitation structure with SSTs would be futile. One way to analyze is to look at the
139 variability of large-scale parameters (e.g., stability, vertical velocity, wind speed) for a given
140 SST, and look at the cloud population in terms of these co-associated variables. By doing so,
141 one would prioritize the combination of factors that lead to different convection type. SST
142 influence on the clouds is of the first order; however, it is also important to show the temporal
143 variation, highlighting the seasonal evolution of cloud types collocated with SSTs and other
144 variables.

145 **Reply:** *We agree with the reviewer that all the forcing/controlling parameters (SST, winds,*
146 *vertical wind velocity, stability, etc.) need to be considered for understanding the vertical*
147 *structure of precipitation. We did the same in the revised version of the manuscript. Also, we*
148 *studied the vertical structure of two types of precipitation (deep and shallow) as suggested by*
149 *the reviewer. Since, stratiform rain is the trailing portions of convective complexes (Houze et*
150 *al. 2015) and is not directly driven by the SST, it's relation with SST is not dealt separately.*

151 **Comment:** The stability measure (LTS) used here is appropriate for stratiform clouds, which
152 may not be appropriate for convective clouds in these regions. One may use static stability
153 profiles instead.

154 **Reply:** *As suggested by the reviewer instead of LTS the static stability (profiles of θ_e) is used*
155 *in the revised manuscript to explain the convective strength as a function of SST.*

156

157

158

159

160

161

162

163

164

165

166

167

168

169

Replies to short comments

170 *At the outset thank Mr. B. Guha for reading our manuscript and suggesting comments.*

171 **Comment:** (a) The article title highlights aspect of the variability of vertical structure of
172 precipitation with sea surface temperature (SST). However, the authors explore the
173 relationships between the SST and other variables such as AOD, CER ice and CER liquid,
174 total column water vapour etc. that may not directly represent the vertical structure of
175 precipitation.

176 **Reply:** *The generation and growth of clouds and precipitating systems depend on the*
177 *triggering mechanisms (over Oceans, it is primarily SST) and ambient dynamical and*
178 *thermodynamical environment (Houze et al., 2015). Changes in SST have the potential of*
179 *altering the type of precipitating system and the vertical structure of precipitation (Oueslati*
180 *and Bellon 2015). Besides the SST, vertical structure can be modified by aerosols (or CCN,*
181 *mostly at the cloud formation stage) and thermodynamics of the ambient atmosphere. For*
182 *instance, recent studies have shown the impact of surface aerosols (PM₁₀) in altering the*
183 *vertical structure of precipitation (Gao et al., 2018 and references therein). We, therefore,*
184 *need to understand the observed variations exist at the cloud formation stage or manifested*
185 *during the descent of precipitation particles to the ground. The cloud effective radius (CER*
186 *for water) (depend on aerosols and TCWV) is a good proxy to understand the cloud*
187 *microphysical processes. While, vertical velocity, winds, stability parameters are considered*
188 *to depict the ambient atmosphere, which can alter the vertical structure of precipitation. All*
189 *these parameters are considered in the present study to understand the vertical structure of*
190 *precipitation over AS and BOB.*

191 **Comment:** (b) The figure 1 shows the regions considered in this study with background
192 colour representing the mean SST during SWM period over AS and BOB. It is clearly
193 evident that the regions of interest depict significant spatial heterogeneity in the SST (± 2
194 degrees C). In such a scenario, (in the figures 4, 5 and 6) I think the standard deviation should
195 be present in those figures.

196 **Reply:** *We wish to inform the reviewer that the segregation of SST data into different bins*
197 *(26° to 31 °C with 1 interval) is done not by averaging the spatial data, rather using 1° X 1°*
198 *gridded data. Therefore, there is no need to average the SST data. Instead, we provided*
199 *standard deviation/standard error of mean values for CER, AOD, TVWV and vertical profiles*
200 *of Z in the revised manuscript.*

201 **Comment:** (c) I would recommend to use MODIS level 2 data products for AOD, CER-ice
202 and CER-liquid for exploring the relationships between different variables. Further, the
203 authors have not mentioned from where the total column water vapour data was obtained.
204 Even the combined uncertainty from different sources of data (e.g., TRMM, MODIS and
205 ECMWF Interim Reanalysis) was not accounted for when establishing the relationships.

206 **Reply:** *The total column water vapor data are taken from the ERA-Interim reanalysis and this*
207 *information is included in the revised manuscript. The spatial resolutions of MODIS level-2*

208 *and ERA-Interim SST are different. Thus, to know the values of AOD and CER at different*
209 *SSTs, again the MODIS level-2 dataset needs to be regridded. Instead of regridding, we have*
210 *used equal spatial lengths MODIS level-3 and SST datasets.*

211 **Comment:** (d) It would be nice if the authors establish the mechanism on why the
212 contrasting relationships were observed over BOB and AS. The authors shall note that SST
213 depends on other factors such as turbidity of the sea water and sea surface albedo, which in
214 turn depend on other variables including wind speed and chlorophyll concentration. While
215 the authors have ignored these essential variables, the relationships with AOD, CER ice,
216 CER-liquid and total column water vapour alone cannot provide the variability in SST in the
217 regions of interest.

218 **Reply:** *We do agree that SST over open Oceans depends on many factors. But our interest is*
219 *not to show how precipitating systems alter the SST over the AS and BOB. Rather, we focused*
220 *on the variation of vertical structure of precipitation (in terms of precipitation top height and*
221 *intensity) with SST over the AS and the BOB and the factors responsible for the variations in*
222 *the vertical structure over both these oceans.*

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240 **Variability of vertical structure of precipitation with sea surface temperature over the**
241 **Arabian Sea and the Bay of Bengal as inferred by TRMM PR measurements**

242 **Kadiri Saikranthi¹, Basivi Radhakrishna², Thota Narayana Rao² and**
243 **Sreedharan Krishnakumari Satheesh³**

244 ¹ *Department of Earth and Climate Science, Indian Institute of Science Education and*
245 *Research (IISER), Tirupati, India.*

246 ² *National Atmospheric Research Laboratory, Department of Space, Govt. of India, Gadanki*
247 *- 517112, India.*

248 ³ *Divecha Centre for Climate Change, Centre for Atmospheric and Oceanic Sciences, Indian*
249 *Institute of Science, Bangalore - 560012, India.*

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269 **Address of the corresponding author**

270 Dr. K. Saikranthi,

271 Department of Earth and Climate Science,

272 Indian Institute of Science Education and Research (IISER),

273 Tirupati,

Andhra Pradesh, India.
Email: ksaikranthi@gmail.com

Abstract

274
275
276
277 Tropical ~~Rainfall Measuring Mission (TRMM) Precipitation Radar (PR) 2A25~~
278 ~~reflectivity profiles data during the period 1998–2013~~ rainfall measuring mission
279 precipitation radar measurements are used to ~~study~~ examine the ~~differences in the~~ variation of
280 vertical structure of precipitation ~~and its variation~~ with sea surface temperature (SST) over
281 the Arabian Sea (AS) and ~~the~~ Bay of Bengal (BOB). ~~Even though the AS and the BOB are~~
282 ~~parts of the Indian Ocean, they exhibit distinct features in vertical structure of precipitation~~
283 ~~and its variation with SST.~~ The variation of reflectivity and precipitation echo top ~~occurrence~~
284 with SST is remarkable over the AS but ~~trivial~~ small over the BOB. The ~~median~~ reflectivity
285 increases with SST ~~at all heights (from 26°C to 31°C) by ~1 dBZ and 4 dBZ above and~~
286 ~~below 106 km altitude, but the increase, respectively, over the AS while, its variation is~~
287 ~~prominent below the freezing level height < 0.5 dBZ over the BOB.~~ The transition from
288 shallow storms at lower SSTs (< 27°C) to deeper storms at higher SSTs is strongly associated
289 with the decrease in stability and mid-tropospheric wind shear over the AS. ~~On the other~~
290 ~~hand, irrespective of altitude, reflectivity profiles are same~~ Contrary, the storms are deeper at
291 all SSTs over the BOB. ~~To understand these differences, variation of aerosols, cloud due to~~
292 ~~weaker stability and water vapor with SST is studied over these seas, mid-tropospheric wind~~
293 ~~shear.~~ At lower SSTs ~~less than 27°C~~, the observed high aerosol optical depth (AOD) and low
294 total column water vapor (TCWV) over ~~the~~ AS results in small ~~Cloud-cloud~~ effective radius
295 (CER) ~~values~~ and ~~low~~ weaker reflectivity. As SST increases, AOD decreases and TCWV
296 increases, ~~which result in~~ leading to large CER and high reflectivity. ~~Over~~ The changes in
297 these parameters with SST are marginal over the BOB ~~the change in AOD, TCWV and hence~~
298 the CER with SST and reflectivity. The predominance of collision-coalescence process below
299 the bright band is ~~marginal.~~ Thus, responsible for the observed negative slopes in the

300 ~~reflectivity over both the seas. The observed~~ variations in reflectivity ~~profiles seem to be~~
301 ~~present from~~ are originated at the cloud formation stage ~~itself~~ over both the seas and these
302 variations are magnified during the descent of hydrometeors to ground.

Formatted: Indent: First line: 1.27 cm

303
304
305
306
307
308
309
310
311

312 1. Introduction

313 Indian summer monsoon (ISM - June through September) is one of the most complex
314 weather phenomena, involving coupling between the atmosphere, land and ocean. At the
315 boundary of the ocean and atmosphere air-sea interactions play a key role for the coupled
316 Earth system (Wu and Kirtman 2005; Feng et al. 2018). ~~SST~~The sea surface temperature
317 (SST) – precipitation relations are the important measures for the air-sea interactions on
318 different temporal scales (Woolnough et al., 2000; Rajendran et al. 2012). Recent studies
319 (Wang et al. 2005; Rajeevan et al. 2012; Chaudhari et al. 2013; ~~Chaudhari~~2016; Weller et al.
320 2016; Feng et al. 2018) have shown that the simulation of ISM can be improved with the
321 exact representation of ~~sea surface temperature (SST)~~ - precipitation relationship.

322 ~~The dynamics of Madden-Julian oscillation (MJO) campaign (DYNAMO) portrayed~~
323 ~~the importance of understanding the link between SST and convective initiation at MJO~~
324 ~~scales (Yoneyama et al. 2013). With known differences in SST between Western Pacific and~~

Formatted: Pattern: Clear (White)

325 ~~Indian Ocean, Barnes and Houze (2013) showed the occurrence of shallow systems~~
326 ~~maximized during the suppressed phases of MJO while the deep wide convective systems~~
327 ~~occurred during the active phases of MJO.~~ SST modulates the meteorological factors that
328 influence the formation and evolution of different kinds of precipitating systems over tropical
329 oceans (Gadgil et al. 1984; Schumacher and Houze, 2003; [Takayabu et al. 2010](#); Oueslati and
330 Bellon 2015).

Formatted: Pattern: Clear

331 The ~~relationships between the~~ studies dealing with SST and cloud/precipitation ~~have~~
332 ~~been studied in variety of contexts during the past three decades. The non-linear relationship~~
333 ~~of SST-precipitation/cloud occurrence~~ population considered whole Indian Ocean as a single
334 entity (Gadgil et al. 1984; Woolnough et al., 2000; Rajendran et al. 2012; Sabin et al. 2012;
335 Meenu et al. 2012; Nair and Rajeev 2014; Roxy 2014; [Nair et al. 2017](#)) is well documented
336 ~~over the Indian Ocean. The probability of organized convection increases with SST up to a~~
337 ~~critical value of $\sim 28^{\circ}\text{C}$ (Gadgil et al. 1984). Sabin et al. (2012) and Meenu et al. (2012)~~
338 ~~showed that the convection is no longer dependent on SST at SSTs greater than 30°C . Later,~~
339 ~~by considering the time lag between the SST and rainfall Roxy (2014) argued that this upper~~
340 ~~threshold can exceed till 31°C . Sengupta et al. (2001) showed that the intraseasonal~~
341 ~~variability of SST is not same over the entire Indian Ocean. Later, Roxy et al. (2013)~~
342 ~~estimated the time lag for SST and precipitation to be 2 and 5 weeks for). But in reality the~~
343 ~~Bay of Bengal (BOB) and the Arabian Sea (AS), respectively. Through this study they found~~
344 ~~that the response of precipitation to SST anomalies is faster over the AS than the BOB. Also,~~
345 ~~the summer-) of Indian Ocean possess distinctly different features. The~~ monsoon experiment
346 (MONEX) ~~showed the influence of the AS and the BOB on the rainfall produced over the~~
347 ~~Indian sub-continent (Krishnamurti 1985; Houze and Churchill 1987) and also proved and~~
348 Bay of Bengal monsoon experiment (BOBMEX) have shown how these two seas are

349 different with respect to ~~the each~~ other ~~oceans~~, in terms of SST, back ground atmosphere and
350 the occurrence of precipitating systems-

351 ~~Knowing the differences~~ (Krishnamurti 1985; Houze and Churchill 1987; Gadgil
352 2000; Bhat et al. 2001). The SST in the AS cools between 10 °N and 20 °N during the
353 monsoon season whereas warming is seen in other global Oceans between the same latitudes
354 (Krishnamurthi 1981). SST variability is large over the AS than the BOB at seasonal and
355 intraseasonal scales (Sengupta et al. 2001; Roxy et al. 2013). The monsoonal winds (in
356 particular the low-level jet) are stronger over the AS than BOB (Findlater 1969). Also, lower-
357 tropospheric thermal inversions are more frequent and stronger over the AS than BOB
358 (Narayanan and Rao 1981; Sathiyamoorthy et al. 2013). Thus, the atmospheric and sea
359 surface conditions over the AS and the BOB during June and September (JJAS) in turn the
360 occurrence of ~~various different~~ kinds of precipitating systems ~~over these two seas is studied in~~
361 Liu et al. (2007), Romatschke et al. (2010), Saikranthi et al. (2014), Houze et al. (are quite
362 different over the BOB and the AS during the ISM period. For instance, long-term
363 measurements of tropical rainfall measuring mission (TRMM) precipitation radar (PR) have
364 shown that 2015). These studies showed that the occurrence of shallow systems ~~is~~ are more
365 prevalent over the ~~Arabian Sea~~ AS, while deeper systems ~~are abundant in~~ occur frequently
366 over the BOB (Liu et al. 2007; Romatschke et al. 2010; Bay of Bengal. Recently, Saikranthi et
367 al. (2014, 2018; Houze et al. 2015.) showed that the observed differences in the occurrence
368 of various kinds of precipitating systems ~~is~~ exist even in El Niño and La Niña periods also,
369 but with variable magnitudes. Aforementioned

370 The aforementioned studies mainly ~~focussed~~ focused on the ~~variation of surface~~
371 rainfall, morphology of vertical structure of precipitation, ~~occurrence of cloudiness with SST~~
372 over the Indian Ocean. ~~But~~ but, none of them studied the variation of vertical structure of
373 precipitation (in terms of occurrence and intensity) with SST. ~~The strength of the convective~~

Formatted: Pattern: Clear (White)

Formatted: Pattern: Clear (White)

Formatted: Pattern: Clear (White)

Formatted: Pattern: Clear (White)

Formatted: Pattern: Clear (White)

Formatted: Pattern: Clear (White)

Formatted: Pattern: Clear (White)

374 ~~forcing strongly depends on SST (Sabin et al. 2012) and changes the differences in SST have~~
375 ~~the potential of altering the vertical structure over AS and BOB. On the other hand,~~
376 ~~information on the vertical structure of precipitation (Oueslati and Bellon 2015).~~

377 ~~The vertical structure of precipitation information~~ is essential for improving the
378 accuracy of rainfall estimation (Fu and Liu 2001; Sunilkumar et al. 2015), understanding the
379 dynamical and microphysical processes of hydrometeor growth/decay mechanisms (Houze
380 2004; Greets and Dejene 2005; Saikranthi et al. 2014; Rao et al. 2016) and ~~flash rates (Liu et~~
381 ~~al. 2012), and~~ for improving the latent heating retrievals (Tao et al. 2006). ~~Also, most of SST~~
382 ~~being the earlier studies dealing with SST and cloud/precipitation population considered~~
383 ~~whole Indian ocean as a single entity. But in reality the BOB and the AS of Indian ocean~~
384 ~~possesses distinctly different features, like SST and its variability over seasonal and~~
385 ~~intraseasonal scales (Sengupta main driving force to trigger precipitating systems through air-~~
386 ~~sea interactions (Sabin et al. 2012; Nuijens et al. 2017), can alter the 2001, Roxy et al. 2013),~~
387 ~~the monsoonal wind speeds (Findlater 1969) and also the type of rain (Liu et al. 2007;~~
388 ~~Romatschke et al. 2010; Saikranthi et al. 2014; Rao et al. 2016). Knowing the importance of~~
389 vertical structure of precipitation ~~and SST modulation of background atmospheric conditions,~~
390 ~~in (Oueslati and Bellon 2015). Therefore, the present study we have studied aims to~~
391 ~~understand~~ the variation of vertical structure of precipitation ~~with SST and their causative~~
392 ~~mechanisms over different regions of Indian ocean, in particular over (in terms of~~
393 ~~precipitation top height and intensity) with SST over the AS and BOB. Besides the SST,~~
394 ~~vertical structure can be modified by aerosols (or CCN, mostly at the cloud formation stage)~~
395 ~~and thermodynamics of the ambient atmosphere. For instance, recent studies have shown the~~
396 ~~BOB and impact of surface PM₁₀ aerosols in altering the AS. vertical structure of~~
397 ~~precipitation (Guo et al., 2018). All these parameters, therefore, are considered in the present~~
398 ~~study to explain the differences in the vertical structure.~~

Formatted: Font color: Custom
Color(RGB(34,34,34))

Formatted: Font color: Custom
Color(RGB(34,34,34))

Formatted: Font color: Custom
Color(RGB(34,34,34))

399 | ~~The present paper is organized as follows. Section 2 describes the data and method of~~
400 | ~~analysis. The variation of the vertical structure of precipitation with SST over BOB and AS is~~
401 | ~~studied in section 3. Section 4 discusses the factors influencing the variation of vertical~~
402 | ~~structure over BOB and AS. The results are summarized in Section 5.~~

403 | **2. Data**

404 | The present study utilizes 16 years (1998-2013) of ~~Tropical rainfall measuring~~
405 | ~~mission (TRMM) precipitation radar (PR) PR's~~ 2A25 (version 7) dataset ~~during the~~
406 | ~~southwest monsoon season (June to September). TRMM PR dataset,~~ comprising of vertical
407 | profiles of attenuation corrected reflectivity ~~with 17 dBZ as minimum detectable signal~~
408 | (Iguchi et al. 2009). ~~Comparing TRMM PR data with Kwajalein S-band radar data~~
409 | ~~Schumacher and Houze (2000) showed that TRMM PR misses 15% of the echo area~~
410 | ~~observed above 0°C levels due to the sensitivity threshold (17 dBZ). Through this study they~~
411 | ~~concluded that TRMM PR highly under samples weaker echoes from ice particles associated~~
412 | ~~with stratiform rain aloft but manages to capture most of the near-surface precipitation~~
413 | ~~accumulation), during the ISM.~~ The range resolution of TRMM-PR reflectivity profiles is
414 | 250 m with a horizontal footprint size of ~4.3 and 5 km before and after the boosting of its
415 | orbit ~~from 350 km to 403 km~~, respectively. It scans $\pm 17^\circ$ from nadir with a beam width of
416 | 0.71° covering a swath of 215 km (245 km after the boost). The uniqueness of TRMM-PR
417 | data ~~uniqueness~~ is its ability in pigeonholing the precipitating systems into convective,
418 | stratiform and shallow rain. This classification is based on two methods namely the
419 | horizontal method (H - method) and the vertical method (V - method) ~~using the bright band~~
420 | ~~identification and the reflectivity profile~~ (Awaka et al. 2009). The original TRMM-PR 2A25
421 | vertical profiles of attenuation corrected reflectivity are gridded to a three dimensional
422 | Cartesian coordinate system with a spatial resolution of $0.05^\circ \times 0.05^\circ$. The detailed
423 | methodology of interpolating the TRMM-PR reflectivity data into the 3D Cartesian grid is

424 discussed in Houze et al. (2007). This dataset is available at the University of Washington
425 website (<http://trmm.atmos.washington.edu/>).

426 To understand the observed variations in the vertical structure of precipitation in the
427 light of microphysics of clouds, Moderate Resolution Imaging Spectroradiometer (MODIS)
428 AQUA satellite level 3 data (MYD08) are considered. In particular, the daily atmospheric
429 products of aerosol optical depth (AOD) (Hubanks et al. 2008) ~~and~~ and cloud effective radius
430 (CER) ~~ice, and CER~~-liquid (Platnick et al. 2017) during the period 2003 and 2013 have been
431 used. MODIS AOD dataset is a collection of aerosol optical properties at 550 nm wavelength,
432 as well as particle size information. Level 2 MODIS AOD is derived from radiances using
433 either one of the three different algorithms, i.e., over ocean Remer et al. (2005) algorithm,
434 over land the Dark-Target (Levy et al. 2007) algorithm and for brighter land surfaces the
435 Deep-Blue (Hsu et al. 2004) algorithm. CER is nothing but the weighted mean of the size
436 distribution of cloud drops i.e., the ratio of third moment to second moment of the drop size
437 distribution. In the level 3 MODIS daily dataset, aerosol and cloud products of level 2 data
438 pixels with valid retrievals within a calendar day are first aggregated and gridded to a daily
439 average with a spatial resolution of $1^\circ \times 1^\circ$. For CER grid box values, CER values are
440 weighted by the respective ice/liquid water cloud pixel counts for the spatiotemporal
441 aggregation and averaging processes.

442 The background atmospheric structure ([winds and total column water vapor](#)) and SST
443 information are taken from the European Centre for Medium Range Weather Forecasting
444 (ECMWF) Interim Reanalysis (ERA) ([Dee et al. 2011](#)) ~~;~~ ERA-Interim runs 4DVAR
445 assimilation twice daily (00 and 12 UTC) to determine the most likely state of the atmosphere
446 at a given time (analysis). The consistency across variables in space and ~~in~~-time (during 12-
447 hour intervals) is thus ensured by the atmospheric model and its error characteristics as
448 specified in the assimilation. ERA-Interim is produced at T255 spectral resolution (about

449 0.75°, ~ 83 km) with a temporal resolution of 6h for upper air fields and 3h for surface fields.

450 ~~The performance of the data assimilation system and the strengths and limitations of ERA-~~
451 ~~Interim datasets are found in Dee et al. (2011).~~ The original 0.75° × 0.75° spatial resolution
452 gridded dataset is rescaled to a resolution of 0.125° × 0.125°. The temporal resolution of the
453 dataset used in the present study is 6h (00, 06, 12 and 18 UTC). The equivalent potential
454 temperature (θ_e) is estimated from the ERA-Interim datasets using the following formula
455 (Wallace and Hobbs 2006):

$$\theta_e = \theta \exp\left(\frac{L_v w_s}{C_p T}\right) \quad (1)$$

456 where θ is the potential temperature, L_v is the latent heat of vaporization, w_s is the saturation
457 mixing ratio, C_p is the specific heat at constant pressure and T is the absolute temperature.

Formatted: Indent: First line: 0 cm

458 The variation of vertical structure of precipitation with SST are studied by considering
459 the dataset between 63°E – 72°E and 8°N – 20°N over the AS and 83°E – 92°E and 8
460 °N – 21°N over the BOB. These regions of interest along with the SWMISM seasonal mean
461 SST over the two seas are depicted in Fig. 1. These regions are selected in such a way that the
462 costal influence on SST is eluded from the analysis. As ~~small amount of the~~ rainfall is
463 ~~observed scanty~~ over the western AS (west ~~to of~~ 63°E latitude) during SWMthe ISM
464 (Saikranthi et al. 2018), this region is also not considered in the present analysis. The
465 seasonal mean SST is higher over the BOB than in the AS by more than 1 °C during the
466 SWMISM season ~~corroborating the findings of, in agreement with~~ Shenoj et al. (2002). The
467 nearest space and time matched SST data from ERA-Interim are assigned to the TRMM-PR
468 and MODIS observations for further analysis.

469

470 **3. Variation of vertical structure of precipitation with SST**

471 The occurrence (in terms of %) of conditional precipitation echoes ($Z \geq 17$ dBZ) at
472 different altitudes as a function of SST over the AS and the BOB is shown in Fig. 2. The
473 variation of precipitation ~~echoes~~ echo occurrence frequency with SST is quite different over
474 both the seas. ~~It increases with increase in SST over The top of the AS, but remains nearly~~
475 ~~same over the BOB. Higher occurrence of~~ precipitation echoes extends to higher
476 heights/altitudes with increasing SST over the AS, while such variation is not quite evident
477 over the BOB. Precipitation echoes are confined to ≤ 8 km at lower SST ($< 28^\circ\text{C}$) over the
478 AS, but exhibits a gradual rise in height with increase in SST. ~~Confinement of Large~~
479 ~~population density of precipitation~~ echoes ~~to lower heights~~ at lower SST/altitudes is mainly
480 due to the abundant occurrence of shallow systems/storms over the AS (Saikranthi et al. 2014;
481 Rao et al. 2016). Interestingly, high the occurrence of precipitation echoes is seen at higher
482 heights/altitudes even at lower SSTs over the BOB, indicating the presence of deeper storms.
483 Such systems exist at all SST's over the BOB.

484 To examine the variation of reflectivity profiles with SST, median profiles of
485 reflectivity in each SST bin are computed over the AS and the BOB separately for deep and
486 shallow systems and are depicted in Figs. ~~3a & 3d~~ 3 & 4, respectively. The space- and time-
487 matched conditional reflectivity profiles are grouped into 1°C SST bins and then the median
488 is estimated at each height, only if the number of conditional reflectivity pixels (Figs. ~~3c &~~
489 ~~3f, 4c & 4f~~) is greater than 500. ~~It is clear from Figs. 3a & 3d that the~~ The median reflectivity
490 profiles corresponding to the deep systems are distinctly different over the AS and the BOB,
491 (Figs. 3a & 3d), even at the same SST. Over the AS, reflectivity ~~profiles show only of deep~~
492 systems at different SSTs shows small variations (≤ 1 dBZ) ~~with SST above (≥ 5 km) the~~
493 ~~melting region, (> 5 km), but~~ vary/ varies significantly (~ 4.5 dBZ) below the melting level ($<$
494 5 km). These variations in reflectivity profiles with SST are negligible (< 0.5 dBZ) over the
495 BOB. ~~Below both above and below~~ the melting ~~layer, the region.~~ The reflectivity increases

496 from 24~26.5 dBZ to ~~~28~~ 31 dBZ, with increase in SST from 26 °C to 30 °C over the AS,
497 but it is almost the same (~~~28dBZ~~ 30dBZ) at all SST's over the BOB. ~~The standard deviation~~
498 ~~of reflectivity also exhibits similar variation as that of median profiles with SST over the AS~~
499 ~~and the BOB. In general the~~ below the melting layer. The standard deviation of reflectivity,
500 representing the variability in reflectivity within the SST bin, is ~~larger over the BOB than the~~
501 ~~AS~~ similar at all SSTs over both the seas except for the 26 °C SST over AS. At this SST, the
502 standard deviation is lesser by ~ 1 dBZ than that of other SSTs.

503 ~~The median reflectivity profiles show a gradual increase with decreasing altitude from~~
504 ~~~10 km to 6 km and an abrupt enhancement is seen just below 6 km over both the seas. The~~
505 ~~sudden enhancement at the freezing level~~ The median reflectivity profiles of shallow storms
506 depicted in Figs. 4a & 4d also show a gradual increase in reflectivity from 20 dBZ to ~ 22
507 dBZ as SST changes from 26 °C to 31 °C at the precipitation top altitude over the AS and
508 don't show any variation with SST over the BOB. However at 1 km altitude, except at 26 °C
509 SST over the AS, the reflectivity variation with SST is not substantial over both the seas. The
510 standard deviation of reflectivity profiles show ~ 1 dBZ variation with SST (from 26 °C to
511 31°C) at all altitudes over the AS and don't show any variation over the BOB. The standard
512 deviation of reflectivity for shallow storms varies from 3 to 4 dBZ at the precipitation top
513 altitude and 4.5 to 5.3 dBZ at 1 km altitude over the AS while it shows ~ 4 dBZ at
514 precipitation top and ~ 5.5 dBZ at 1 km altitude over the BOB.

515 **4. Factors affecting the vertical structure of precipitation and their variability with SST**

516 The formation and evolution of precipitating systems over oceans depend on dynamical,
517 thermodynamical and microphysical factors, like SST, wind shear, vertical wind velocity,
518 stability, CER, etc., and need to be considered for understanding the vertical structure of
519 precipitation (Li and Min 2010; reamean et al. 2013; Chen et al. 2015; Shige and Kummerov
520 2016; Guo et al 2018).

521 **4.1. Dynamical and thermodynamical factors:**

522 To understand the role of stability/instability, θ_e values computed from (1) using the
523 ERA-Interim datasets during the ISM period over the AS and the BOB are averaged for a
524 season and are depicted in Figs. 5(a) & 5(b), respectively. The surface θ_e (at 1000 hPa) values
525 are larger over the BOB than those over AS for the same SST, indicating that the instability
526 and convective available potential energy (CAPE) could be higher over the BOB. Indeed,
527 higher CAPE is seen over the BOB (Fig. S1, calculated following Emanuel 1994) than AS at
528 all SSTs by a magnitude $> 300 \text{ J kg}^{-1}$. The θ_e increases with SST from 358 °K to 368 °K
529 from 27 °C to 31 °C and from 350 °K to 363 °K from 26 °C to 31 °C over the BOB and the
530 AS, respectively. The CAPE also increases with rise in SST over both the seas. To know the
531 stability of the atmosphere θ_e gradients are considered. Irrespective of SST, positive gradients
532 in θ_e are observed between 900 and 800 hPa levels over the AS indicating the presence of
533 strong stable layers. The strength of these stable layers decreases with increasing SST. These
534 stable layers are formed mainly due to the flow of continental dry warm air from Arabian
535 Desert and Africa above the maritime air causing temperature inversions below 750 hPa level
536 over the AS during the ISM period (Narayanan and Rao 1981). However over the BOB, such
537 temperature inversions are not seen in the lower troposphere.

538 To understand the effect of wind field on the vertical structure of precipitation,
539 profiles of ISM seasonal mean vertical wind velocity and vertical shear in horizontal wind at
540 various SSTs over the AS and the BOB are shown in Figs. 5(c), 5(d) & 5(e), 5(f)
541 respectively. The updrafts are prevalent at all SSTs throughout the troposphere over the BOB,
542 whereas downdrafts are seen in the mid-troposphere (between 200 and 600 hPa levels) up to
543 27 °C and updrafts in the entire troposphere at higher SSTs over the AS. Also, the magnitude
544 of the vertical wind velocity varies significantly with SST in the mid-troposphere over the
545 AS. Over the BOB, the magnitude of updrafts increases with altitude in the lower and middle

546 troposphere, but doesn't vary much with SST. In the mid-troposphere, updrafts are stronger
547 by $> 0.02 \text{ Pa S}^{-1}$ over the BOB than over the AS. The profiles shown in Fig. 5(e) & 5(f) are
548 the mean vertical shear in horizontal wind estimated following Chen et al. (2015) at different
549 levels with reference to 950 hPa level. The wind shear increases with increasing altitude at all
550 the SSTs up to 400 hPa, but the rate of increase is distinctly different between the AS and the
551 BOB at SSTs less than 28 °C and nearly the same at higher SSTs. The wind shear decreases
552 systematically with SST ($\sim 1.5 \text{ m s}^{-1}$ for 1° increase in SST) in the middle troposphere over
553 the AS while the change is minimal over the BOB ($\sim 2 \text{ m s}^{-1}$ for 27 °C and 31 °C).

554 Chen et al. (2015) highlighted the importance of mid-tropospheric wind shear in
555 generating mesoscale local circulations, like low-level cyclonic and upper-level anticyclonic
556 circulations. This feature is apparent over the AS, where down drafts are prevalent in mid-
557 upper troposphere and updrafts in the lower troposphere at lower SSTs. As SST increases, the
558 wind shear decreases and the updraft increases in the mid-troposphere. However, over the
559 BOB the wind shear is relatively weak when compared to the AS and hence the updrafts are
560 seen up to 200 hPa level at all SSTs. The weaker CAPE and stable mid-troposphere coupled
561 with upper- to mid- tropospheric downdrafts at lower SSTs over the AS inhibit the growth of
562 precipitating systems to higher altitudes and in turn precipitate in the form of shallow rain.
563 This result is in accordance with the findings of Shige and Kummerow (2016) that showed
564 the static stability at lower levels inhibits the growth of clouds and promotes the detrainment
565 of clouds over the Asian monsoon region and is considered as an important parameter in
566 determining the precipitation top height. As SST increases large CAPE and updrafts in the
567 middle troposphere collectively support the precipitating systems to grow to higher altitudes,
568 as evidenced in Fig. 2a. On the other hand, large CAPE and updrafts in the middle
569 troposphere prevalent over the BOB at all SSTs are conducive for the precipitating systems to
570 grow to higher altitudes as seen in Fig. 2b.

571 **4.2. Microphysical factors** ~~is primarily due to the aggregation of hydrometeors and~~
572 ~~change in dielectric factor from ice to water (Fabry and Zawadzki 1995; Rao et al. 2008; Cao~~
573 ~~et al. 2013). Below the bright band, the raindrops can grow by collision-coalescence process~~
574 ~~and reduce their size either by breakup or by evaporation processes. The collision-~~
575 ~~coalescence results in negative slope in the reflectivity profile, whereas breakup and~~
576 ~~evaporation results in positive slope (Liu and Zipser 2013; Cao et al. 2013; Saikranthi et al.~~
577 ~~2014). The observed negative slope in the median reflectivity profiles below the bright band~~
578 ~~region indicates the low-level hydrometeor growth over both the seas. This hydrometeor~~
579 ~~growth below melting region indicates the predominance of collision-coalescence process~~
580 ~~than the collision-breakup process over both the seas. The magnitude of the slope is nearly~~
581 ~~equal over both the seas, indicating that the rate of growth, on average, is nearly equal.~~

582 **4. Factors affecting the vertical variation of reflectivity with SST**

583 The formation and evolution of precipitating systems depends on the stability of the boundary
584 layer, dynamics and thermodynamics of the ambient atmosphere. To know the stability of the
585 marine boundary layer at various SSTs the lower tropospheric stability (LTS) is considered.
586 LTS is defined as the difference in potential temperature between 700 hPa (θ_{700}) and surface
587 (θ_0) i.e., $LTS = \theta_{700} - \theta_0$ that represents the strength of the inversion caps by the planetary
588 boundary layer (Wood and Bretherton 2006). The LTS values were computed from the ERA-
589 Interim temperature data during SWM season over the selected regions and are depicted in
590 Fig. 4(a). LTS decreases with SST up to 29°C and increases a little at further SSTs over both
591 the seas however when compared to the BOB the LTS values are larger over the AS at all
592 SSTs. The stability of the planetary boundary layer is very high at lower SSTs and as SST
593 increases the stability decreases drastically over the AS up to 29°C and increases a little at
594 further SSTs. On the other hand the variability in planetary boundary layer stability with SST
595 is trivial over the BOB. Also shown in Fig. 4(b) is the convective available potential energy

Formatted: Indent: First line: 0 cm

596 ~~(CAPE) at different SSTs over both the Seas. CAPE is calculated following Emanuel (1994).~~
597 ~~CAPE increases with rise in SST over both the seas while its magnitude is relatively large~~
598 ~~over the BOB than the AS at all SSTs. The large LTS and small CAPE values at lower SSTs~~
599 ~~over the AS don't allow the precipitating systems to grow to higher altitudes and in turn~~
600 ~~precipitate in the form of warm rain. As SST increases LTS decreases drastically and CAPE~~
601 ~~increases and hence the precipitating systems can grow to higher altitudes. Though LTS~~
602 ~~increases above 29°C the instability created by the large CAPE can penetrate the planetary~~
603 ~~boundary layer and favours the formation of deeper systems. On the other hand LTS values~~
604 ~~are lower and remain almost same at all SSTs and large CAPE values over the BOB are~~
605 ~~conducive for the precipitating systems to grow to higher altitudes as depicted in Fig. 2,~~

Formatted: Font: Bold

606 The observed differences in reflectivity profiles of precipitation with SST could be
607 originated at the cloud formation stage ~~or in the evolution stage or due to both. In order to~~
608 ~~understand this, the variation of mean CER for ice and liquid at different SST's over the AS~~
609 ~~and the BOB is depicted in Figs. 5a & 5b, respectively. The mean is calculated only when the~~
610 ~~number of data points is larger than 100 in each SST bin. It is evident from Figs. 5a & 5b that~~
611 ~~both CER ice and liquid increase with rise in SST substantially over the AS but the increase~~
612 ~~is marginal over the BOB. For example, as SST rises from 26°C to 31°C, the CER ice and~~
613 ~~liquid vary from 20 μm to 32 μm and 14.7 μm to 20.8 μm , respectively over the AS, whereas~~
614 ~~they vary, respectively, from 29 μm to 31 μm and 18.5 μm to 19.5 μm over the BOB. Also,~~
615 ~~the cloud droplets are small in size at lower SSTs and bigger at higher SSTs over the AS,~~
616 ~~whereas they are big over the BOB irrespective of SST. These smaller sized hydrometeors at~~
617 ~~low SSTs are responsible for the observed small reflectivities above the melting layer over~~
618 ~~the AS than the BOB as reflectivity is more sensitive to the particle size than the droplet~~
619 ~~concentration ($Z \propto D^6$). At higher SSTs, the CER values are approximately equal over both~~

620 ~~the seas and in turn the observed reflectivities (Fig. 5). This suggests that the variations seen~~
621 ~~in vertical profiles of reflectivity are originating in the cloud itself.~~

622 ~~Numerous studies have examined the aerosol effects on cloud formation through~~
623 ~~heterogeneous nucleation and precipitation itself or manifested during the evolution stage or~~
624 ~~due to both. Information on AOD and CER would be ideal to infer microphysical processes at~~
625 ~~the cloud formation stage. CER values are mainly controlled by the ambient aerosols~~
626 ~~concentration and the available moisture~~ (Twomey 1977; Albrecht 1989; Tao et al. 2012; and
627 Rosenfeld et al. 2014). For fixed liquid water content, as the concentration of aerosols
628 increases, the number of cloud drops increases and ~~droplet size reduces CER decreases~~
629 (Twomey 1977). ~~Utilizing the aircraft measurements over Indian sub-continent Ramanathan~~
630 ~~et al. (2001) showed that the cloud drop number density increase with increasing aerosol~~
631 ~~number density both over continental and maritime regions. Connolly et al. (2009), Li and~~
632 ~~Min (2010), Niemand et al. (2012), Creamean et al. (2013), and Fan et al. (2014) showed~~
633 ~~that dust also act as ice nuclei through heterogeneous nucleation and these ice nuclei directly~~
634 ~~change the ice nucleation processes that determine the initial number concentration and size~~
635 ~~distribution of ice crystals. Thus, to To understand the role variation of aerosols in AOD and~~
636 ~~TCWV and the observed variations in the resultant CER with SST, the seasonal mean AOD~~
637 ~~variation with and TCWV for different SST is bins are plotted in Fig Figs. 6a for the SWM. &~~
638 ~~6b. The mean and standard error are calculated only when the number of data points is more~~
639 ~~than 100 in each SST bin. AOD decreases from 0.62 to 0.31 with rise in SST from 26 °C to~~
640 ~~31 °C over the AS but only from 0.42 to 0.36 -as SST varies from 27 °C to 30 °C and then~~
641 ~~increases with rise in SST at higher SSTs over the BOB. Also shown in Fig. 6b is the The~~
642 ~~variation of total column water vapor (TCWV) with SST over both the seas. TCWV (Fig. 6b)~~
643 ~~shows a gradual increase with SST over the AS while it decreases initially from 27°C to~~
644 ~~28°C, and then increases with SST over the BOB. At a given SST the TCWV is more in the~~

645 ~~BOB than in the AS. More number of aerosols and relatively low TCWV over the AS results~~
646 ~~in large number of cloud drops with reduced size (Twomey 1977; Ramanathan 2001). These~~
647 ~~reduced size cloud drops are responsible for the observed small CER values at SSTs less than~~
648 ~~28°C. As SST rises the AOD decreases and TCWV increases such that the cloud particles~~
649 ~~grow in size which in turn increases CER. On the other hand, the change in AOD and TCWV~~
650 ~~(and as a result in CER) is not prominent with SST over the BOB, as seen in the Fig. 5. over~~
651 ~~the BOB. At a given SST the TCWV is more in the BOB (> 8 mm) than in the AS.~~

652 ~~To understand the transport of aerosols at low and mid levels the wind magnitudes~~
653 ~~and directions at 850 hPa and 500 hPa levels are shown in Fig. 7. The strong lower~~
654 ~~tropospheric winds produce sea salt particles as well as transport dust from the Horn of~~
655 ~~Africa and the mid tropospheric winds transport dust from the Arabian Desert over the AS~~
656 ~~(Li and Ramanathan 2002). On the other hand the continental aerosols from India landmass~~
657 ~~are transported to the BOB both at low and mid troposphere. The decrease in AOD and an~~
658 ~~increase in TCWV with SST result in an increase in CER (14.7 μm to 20.8 μm from 26°C to~~
659 ~~31°C) over the AS (Fig. 7). On the other hand, CER doesn't show much variation with SST~~
660 ~~(18.5 μm to 19.5 μm from 27°C to 31°C) over BOB due to smaller variations in AOD and~~
661 ~~TCWV. This also shows that the cloud droplets are smaller in size at lower SSTs over the~~
662 ~~AS than BOB, while they are bigger and nearly equal in size at higher SSTs. Since,~~
663 ~~reflectivity is more sensitive to the particle size ($Z \propto D^6$), the smaller-sized hydrometeors at~~
664 ~~lower SSTs over the AS yield weaker reflectivity than over the BOB (both for deep and~~
665 ~~shallow systems). As the SST increases, CER as well as the reflectivity increases over the~~
666 ~~AS. At higher SSTs, the CER values are approximately equal over both the seas and in turn~~
667 ~~the observed reflectivities (Figs. 3a & 4a). This suggests that the variations seen in the~~
668 ~~reflectivity are originated in the cloud formation stage itself.~~

669 The hydrometeors also evolve during their descent to the ground due to several
670 microphysical processes. These processes can be inferred from the vertical structure of
671 precipitation or vertical profiles of reflectivity. The median reflectivity profiles of deep
672 systems show a gradual increase from ~ 10 km to 6 km and an abrupt enhancement is seen
673 just below 6 km over both the seas (Figs. 3a & 3d). The sudden enhancement at the freezing
674 level (radar bright band) ~~Sathesh et al. (2006) showed an increase in AOD with increase in~~
675 latitude over the AS due to the dust advection from Arabia desert regions during SWM
676 season, whereas SST decreases with increase in the latitude. ~~is primarily due to the~~
677 aggregation of hydrometeors and change in dielectric factor from ice to water (Fabry and
678 Zawadzki 1995; Rao et al. 2008; Cao et al. 2013). Below the bright band, raindrops grow by
679 collision-coalescence process and reduce their size by either breakup and/or evaporation
680 processes. The collision-coalescence results in negative slope in the reflectivity profile,
681 whereas breakup and evaporation results in positive slope (Liu and Zipser 2013; Cao et al.
682 2013; Saikranthi et al. 2014; Rao et al. 2016). The observed negative slope ($\sim -0.3 \text{ dBZ km}^{-1}$)
683 in the median reflectivity profiles below the bright band indicates dominance of low-level
684 hydrometeor growth over both the seas. The magnitude of the slope decreases with SST over
685 the AS, while it is nearly equal at all SSTs over the BOB. It indicates the growth rate
686 decreases with SST over the AS and remains the same at all SSTs over the BOB. The median
687 reflectivity profiles of shallow systems also show negative slopes ($\sim -1 \text{ dBZ km}^{-1}$) at all SSTs
688 representing the predominance of low-level hydrometeor growth by collision-coalescence
689 processes over both the seas.

690 The present analysis shows that the observed reflectivity changes with SST over both
691 the seas originate at the cloud formation stage and magnify further during the descent of
692 hydrometeors to ground. ~~In other words the SST is low and AOD is high in northern AS~~
693 ~~whereas over the southern AS, SST is high and AOD is low. This contrasting spatial~~

694 ~~distribution of AOD and SST could cause a negative correlation between AOD and SST. To~~
695 ~~examine whether the observed decrease in AOD with increase in SST over the AS is due to~~
696 ~~the latitudinal variation of AOD or exists at all latitudes, we have segregated the data into 2°~~
697 ~~latitude bins and plotted the mean AOD with SST for all bins and is depicted in Fig. 8a. In~~
698 ~~spite of the magnitude, AOD variation with SST is nearly similar at all latitudes of the AS,~~
699 ~~i.e., the higher AOD is observed at lower SSTs and vice versa. On the other hand the~~
700 ~~latitudinal variation of AOD with SST over the BOB shown in Fig. 8b also show a decrease~~
701 ~~in AOD with SST up to 30°C but the magnitude of variation is trivial relative to the AS. As~~
702 ~~also depicted in Fig. 6a above 30°C AOD increases with SST over the BOB. This indicates~~
703 ~~that though there is a difference in magnitude of variation, AOD varies with SST over both~~
704 ~~the seas at all latitudes. This analysis is repeated using the multi-angle imaging~~
705 ~~spectroradiometer (MISR) dataset (which is not shown here) for small, medium large aerosol~~
706 ~~particles. Interestingly all three types also show a decrease in AOD with rise in SST over~~
707 ~~both the seas.~~

708 5. Conclusions

709 Sixteen years of TRMM-PR 2A25 reflectivity profiles and 11 years of MODIS AOD
710 and CER data are utilized to understand the differences in variation of vertical structure of
711 precipitation with SST over AS and BOB. ~~This analysis reveals that the variation of~~
712 ~~reflectivity with SST is~~Precipitation top height increases with SST over the AS indicating
713 that systems grow to higher altitudes with increase in SST while it is almost same at all SSTs
714 representing that the systems are deeper over the BOB. The decrease in stability and mid-
715 tropospheric wind shear with SST over the AS favour the formation of deeper system at
716 higher systems. However the low stability and small wind shear at all SSTs over the BOB
717 help the formation of deeper systems. The variation of reflectivity with SST is found to be
718 remarkable over the AS and marginal over the BOB. The reflectivity increases with rise in

Formatted: No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

719 SST over the AS and remains the same at all SSTs over the BOB. This change in reflectivity
720 over the AS is more prominent below the freezing level height (~ 4 dBZ) than the above (~ 1
721 dBZ). Over the AS, the abundance of aerosols and less moisture at SSTs < 27°C result in
722 high concentration of ~~small diameters~~smaller cloud droplets. As SST increases the aerosol
723 concentration decreases and moisture increases ~~such that~~leading to the formation of bigger
724 cloud droplets ~~are formed~~. Thus, the reflectivity increases with rise in SST over the AS. On
725 the other hand, AOD, TCWV and CER do not show substantial variation with SST over the
726 BOB and hence the change in reflectivity is small. Over the BOB, the mid troposphere is wet
727 and hydrometeor's size at the formation stage is nearly the same at all SSTs. ~~The evolution~~
728 of hydrometeors during their descent is also similar at all SST's, ~~as evidenced by nearly~~
729 similar. The collision-coalescence process is predominant below the bright band region over
730 both the seas and is responsible for the observed negative slope in the reflectivity profiles.

731 Acknowledgements

732 The authors would like to thank Prof. Robert Houze and his team for the interpolated 3D
733 gridded TRMM-PR dataset (<http://trmm.atmos.washington.edu>), ECMWF (<http://data-portal.ecmwf.int/>) team for providing the ERA-Interim dataset and MODIS
734 (<https://ladsweb.modaps.eosdis.nasa.gov/>) science team for providing the AOD and CER
735 dataset. The authors express their gratitude to Prof. J. Srinivasan for his fruitful discussions
736 and valuable suggestions in improving the quality of the manuscript. The corresponding
737 author would like to thank Department of Science & Technology (DST), India for providing
738 the financial support through the reference number DST/INSPIRE/04/2017/001185. We
739 thank the two referees for their critical comments in improving the quality of the manuscript.

741 References

742 Albrecht, B.A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227–
743 1230, 1989.

744 Awaka, J., Iguchi, T., and Okamoto, K.: TRMM PR standard algorithm 2A23 and its
745 performance on bright band detection, *J. Meteorol. Soc. Jpn.*, 87A, 31–52, 2009.

Formatted: Font: Times New Roman, 12 pt

746 ~~Bhat, G. S., Gadgil, S., Kumar, P. V. H., Kalsi, S. R., Madhusoodanan, P., Murty, V. S., Rao,~~

747 ~~C. V. P., Babu, V. R., Rao, L.V., Rao, R., Ravichandran, M., Reddy, K.G., Rao, P. S.,~~

Formatted: art_authors, Font: +Body (Calibri), 11 pt

748 ~~Sengupta, D., Sikka, D. R., Swain, J., and Vinayachandran, P. N.: BOBMEX: The Bay~~

Formatted: art_authors, Font: +Body (Calibri), 11 pt

749 ~~of Bengal Monsoon Experiment, *Bull. Amer. Meteor. Soc.*, Barnes, H. C., and Houze, Jr.~~

Formatted: Pattern: Clear

750 ~~R. A.: The precipitating cloud population of the Madden-Julian oscillation over the~~

751 ~~Indian and West Pacific Oceans. *J.* 82, 2217–2244, 2001.~~

752 ~~*Geophys. Res.*, 118, 6996–7023, doi:10.1002/jgrd.50375, 2013.~~

Formatted: Default Paragraph Font, Font: +Body (Calibri), 11 pt, Not Italic

753 Cao, Q., Hong, Y., Gourley, J. J., Qi, Y., Zhang, J., Wen, Y., and Kirstetter, P. E.: Statistical
754 and physical analysis of the vertical structure of precipitation in the mountainous west
755 region of the United States using 11+ years of space borne observations from TRMM
756 precipitation radar, *J. Appl. Meteorol. Climatol.*, 52, 408–424, 2013.

757 ~~Chaudhari, H. S., Pokhrel, S., Mohanty, S., and Saha, S. K.: Seasonal prediction of Indian~~

758 ~~summer monsoon in NCEP coupled and uncoupled model, *Theor. Appl. Climatol.*, 114,~~

759 ~~459–477, doi:10.1007/s00704-013-0854-8, 2013.~~

760 ~~Chaudhari, H. S., Pokhrel, S., Kulkarni, A., Hazra, A., and Saha, S. K.: Clouds-SST~~

761 ~~relationship and interannual variability modes of Indian summer monsoon in the~~

762 ~~context of clouds and SSTs: observational and modelling aspects, *Int. J. Climatol.*, doi:~~

763 ~~10.1002/joc.4664, 2016.~~

764 ~~Chaudhari, H. S., Pokhrel, S., Connolly, P. J., Möhler, O., Field, P. R., Saathoff, H., Burgess,~~

Formatted: art_authors, Font: +Body (Calibri), 11 pt

765 ~~R., Choulaton, T., and Gallagher, M.: Studies of heterogeneous freezing by three~~

766 ~~different desert dust samples. *Atmos. Chem. Phys.*, 9, 2805–2824, doi:10.5194/acp-9-~~

Formatted: Default Paragraph Font, Font: +Body (Calibri), 11 pt, Not Italic

767 ~~2805–2009, 2009.~~

768 Mohanty, S., and Saha, S. K.: Seasonal prediction of Indian summer monsoon in NCEP
769 coupled and uncoupled model, *Theor. Appl. Climatol.*, 114, 459–477,
770 [doi:10.1007/s00704-013-0854-8](https://doi.org/10.1007/s00704-013-0854-8), 2013.

771 Chen, Q., Fan, J., Hagos, S., Gustafson Jr., W. I., and Berg, L. K.: Roles of wind shear at
772 different vertical levels: Cloud system organization and properties, *J. Geophys. Res.*
773 *Atmos.*, 120, 6551–6574, 2015.

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt, Not Italic

774 Creamean, J. M., Suski, K. J., Rosenfeld, D., Cazorla, A., DeMott, P. J., Sullivan, R. C.,
775 White, A. B., Ralph, F. M., Minnis, P., Comstock, J. M., Tomlinson, J. M., Kimberly
776 A., and Prather, K. A.: Dust and biological aerosols from the Sahara and Asia influence
777 precipitation in the western U.S., *Science*, 339, 1572–1578,
778 [doi:10.1126/science.1227279](https://doi.org/10.1126/science.1227279), 2013.

779 Dee, D. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data
780 assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 553–597, 2011.

781 Emanuel, K. A.: Atmospheric convection. Oxford University Press, Oxford, 1994.

782 Fabry, F., and Zawadzki, I.: Long-term radar observations of the melting layer of
783 precipitation and their interpretation, *J. Atmos. Sci.*, 52, 838–851, 1995.

784 ~~Fan, J., Leung, L. R., DeMott, P. J., Comstock, J. M., Singh, B., Rosenfeld, D., Tomlinson, J.~~
785 ~~M., White, A., Prather, K. A., Minnis, P., Ayers, J. K., and Min, Q.: Aerosol impacts on~~
786 ~~California winter clouds and precipitation during CalWater 2011: local pollution versus~~
787 ~~long range transported dust, *Atmos. Chem. Phys.*, 14, 81–101,~~
788 ~~<https://doi.org/10.5194/acp-14-81-2014>, 2014.~~

Formatted: art_authors, Font: +Body
(Calibri), 11 pt

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt

789 Feng, X., Haines, K., Liu, C., de Boisséson, E., and Polo, I., Improved SST-precipitation
790 intraseasonal relationships in the ECMWF coupled climate reanalysis, *Geophys. Res.*
791 *Lett.*, 45, 3664–3672, 2018.

792 Findlater, J.: A major low-level air current near the Indian Ocean during the northern
793 summer, *Q. J. R. Meteorol. Soc.*, 95, 362–380, 1969.

794 Fu, Y., and Liu, G.: The variability of tropical precipitation profiles and its impact on
795 microwave brightness temperatures as inferred from TRMM data, *J. Appl. Meteorol.*,
796 40, 2130–2143, 2001.

797 Gadgil, S., Joseph, P. V., and Joshi, N. V.: Ocean atmosphere coupling over monsoonal
798 regions, *Nature*, 312, 141–143, 1984.

Formatted: Font: Not Bold

799 Gadgil, S.: Monsoon–ocean coupling. *Current Sci.*, 78, 309–323, 2000.

800 Geerts, B., and Dejene, T.: Regional and diurnal variability of the vertical structure of
801 precipitation systems in Africa based on space borne radar data, *J. Clim.*, 18, 893–916,
802 2005.

803 Guo, J., Liu, H., Li, Z., Rosenfeld, D., Jiang, M., Xu, W., Jiang, J. H., He, J., Chen, D., Min,
804 M., and Zhai, P.: Aerosol-induced changes in the vertical structure of precipitation: a
805 perspective of TRMM precipitation radar, *Atmos. Chem. Phys.*, **Houze, R. A18, 13329-**
806 **13343, <https://doi.org/10.5194/acp-18-13329-2018>, 2018.**

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt, Not Italic

807 ~~≡ Mesoscale convective systems, *Rev. Geophys.*, 42, RG4003, doi: 10.1029/2004RG000150,
808 2004.~~

809 Houze, R. A., and Churchill, D. D.: Mesoscale organization and cloud microphysics in a Bay
810 of Bengal depression, *J. Atmos. Sci.*, 44, 1845–1867, 1987.

811 ~~Houze, R. A., Wilton, D. C., and Smull, B. F.: Monsoon convection in the Himalayan region
812 as seen by the TRMM precipitation radar, *Q. J. R. Meteorol. Soc.*, 133, 1389–1411,
813 2007.~~

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt

814 ~~Houze, R. A.,~~ Rasmussen, K. L., Zuluaga, M. D., and Brodzik, S. R.: The variable nature of
815 convection in the tropics and subtropics: A legacy of 16 years of the Tropical rainfall
816 measuring mission satellite, *Rev. Geophys.*, 53, 994–1021, 2015.

Formatted: Font: Not Bold

817 [Houze, R. A., Wilton, D. C., and Smull, B. F.: Monsoon convection in the Himalayan region](#)
818 [as seen by the TRMM precipitation radar, *Q. J. R. Meteorol. Soc.*, **133**, 1389-1411,](#)
819 [2007.](#)
820 [Houze, R. A.: Mesoscale convective systems, *Rev. Geophys.*, **42**, RG4003, doi:](#)
821 [10.1029/2004RG000150, 2004.](#)

Formatted: Strong, Font: +Body (Calibri), 11 pt

Formatted: Default Paragraph Font, Font: +Body (Calibri), 11 pt

822 Hsu, N., Tsay, S., King, M., and Herman, J.: Aerosol properties over bright-reflecting source
823 regions, *Geosci. Remote Sens. IEEE Trans.*, **42**, 557–569, 2004.

824 Hubanks, P., King, M., Platnick, S., and Pincus, R.: MODIS atmosphere L3 gridded product
825 algorithm theoretical basis document collection 005 Version 1.1, Tech. Rep. ATBD-
826 MOD-30, NASA, 2008.

827 Iguchi, T., Kozu, T., Kwiatkowski, J., Meneghini, R., Awaka, J., and Okamoto, K.:
828 Uncertainties in the rain profiling algorithm for the TRMM precipitation radar, *J.*
829 *Meteor. Soc. Japan*, **87A**, 1–30, doi:10.2151/jmsj.87A.1, 2009.

830 Krishnamurti, T. N.: Summer monsoon experiment – A review. *Mon. Wea. Rev.*, **113**, 1590-
831 1626, 1985.

Formatted: Font: Not Bold

832 [Krishnamurti, T.: Cooling of the Arabian Sea and the onset-vortex during 1979. Recent](#)
833 [progress in equatorial oceanography: A report of the final meeting of SCOR](#)
834 [WORKING GROUP 47 in Venice, Italy, 1-12, 1981. \[Available from Nova Univ.,](#)
835 [Ocean Science Center, Dania, FL 33004\].](#)

836 Levy, R., Remer, L., Mattoo, S., Vermote, E., and Kaufman, Y.: Second-generation
837 operational algorithm: Retrieval of aerosol properties over land from inversion of
838 moderate resolution imaging spectroradiometer spectral reflectance, *J. Geophys. Res.*,
839 **112**, D13, doi:10.1029/2006JD007811, 2007.

840 Li, F., and Ramanathan, V.: Winter to summer monsoon variation of aerosol optical depth
841 over the tropical Indian Ocean, *J. Geophys. Res.*, 107(D16), doi:
842 10.1029/2001JD000949, 2002.

843 Li, R., and Min, Q.-L.: Impacts of mineral dust on the vertical structure of precipitation. *J.*
844 *Geophys. Res.*, 115, D09203, doi:10.1029/2009JD011925, 2010.

845 Liu, C., Zipser, E., and Nesbitt, S. W.: Global distribution of tropical deep convection:
846 Different perspectives using infrared and radar as the primary data source, *J. Climate,*
847 20, 489-503, 2007.

848 Liu, C., and Zipser, E. J.: Why does radar reflectivity tend to increase downward toward the
849 ocean surface, but decrease downward toward the land surface?, *J. Geophys. Res.*
850 *Atmos.*, 118, 135-148, doi: 10.1029/2012JD018134, 2013.

851 ~~Liu, C., Zipser, E., and Nesbitt, S. W.: Global distribution of tropical deep convection:~~
852 ~~Different perspectives using infrared and radar as the primary data source, *J. Climate,*~~
853 ~~20, 489-503, 2007.~~

854 Liu, C., Cecil, D., and Zipser, E. J.: Relationships between lightning flash rates and radar
855 reflectivity vertical structures in thunderstorms over the tropics and subtropics. *J.*
856 *Geophys. Res.*, doi:10.1029/2011JD017123, 2012.

Formatted: Emphasis, Font: +Body
(Calibri), 11 pt, Not Italic

857 Meenu, S., Parameswaran, K., and Rajeev, K.: Role of sea surface temperature and wind
858 convergence in regulating convection over the tropical Indian Ocean, *J. Geophys. Res.*
859 *Atmos.*, 117, D14102, 2012.

860 Nair, A. K. M., and Rajeev, K.: Multiyear CloudSat and CALIPSO observations of the
861 dependence of cloud vertical distribution on sea surface temperature and tropospheric
862 dynamics, *J. Clim.*, 27, 672-683, doi:10.1175/JCLI-D-13-00062.1, 2014.

863 ~~Nair, A.K.M., Rao, T. N., and Rajeev, K.: Role of cloud feedback in regulating the “pool of~~
864 ~~inhibited cloudiness” over the Bay of Bengal, *Meteorol. Atmos. Phys.*,~~
865 ~~<https://doi.org/10.1007/s00703-017-0560-7>, 2017.~~

866 ~~Niemand, M., Möhler, O., Vogel, B., Vogel, Narayanan, M. S., and Rao, B. M.: Detection of~~
867 ~~monsoon inversion by TIROS-N satellite, *Nature*, 294, 546-548, 1981.~~

868 ~~Nuijens, L., Emanuel, K., Masunaga, H., and L’Ecuyer, T.: Implications of warm rain in~~
869 ~~shallow cumulus and congestus clouds for large-scale circulations, *Surv. Geophys.*, 38,~~
870 ~~[1257-1282](https://doi.org/10.1007/s11069-017-0125-7), 2017.~~

871 ~~Hoose, C., Connolly, P., Klein, H., Bingemer, H., DeMott, P., Skrotzki, J., and Leisner,~~
872 ~~T.: A particle surface area-based parameterization of immersion freezing on desert dust~~
873 ~~particles, *J. Atmos. Sci.*, 69, 3077–3092, <https://doi.org/10.1175/JAS-D-11-0249.1>,~~
874 ~~2012.~~

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt

875 Oueslati, B., and Bellon, G.: The double ITCZ bias in CMIP5 models: interaction between
876 SST, large-scale circulation and precipitation. *Climate Clim. Dyn.*, 44, 585-607, 2015.

Formatted: Font: Not Bold

877 Platnick, S., et al.: The MODIS cloud optical and microphysical products: Collection 6
878 updates and examples from Terra and Aqua, *IEEE Trans. Geosci. Remote Sens.*, 55,
879 502–525, doi:10.1109/TGRS.2016.2610522, 2017.

880 Rajeevan, M., Unnikrishnan, C. K., and Preethi, B.: Evaluation of the ENSEMBLES multi-
881 model seasonal forecasts of Indian summer monsoon variability, *Clim. Dyn.*, 38, 2257–
882 2274, 2012.

883 Rajendran, K., Nanjundiah, R. S., Gadgil, S., and Srinivasan, J.: How good are the
884 simulations of tropical SST–rainfall relationship by IPCC AR4 atmospheric and
885 coupled models?, *J. Earth Sys. Sci.*, 121(3), 595–610, 2012.

886 ~~Ramanathan, V., Crutzen, P. J., Kiehl, J. T., Rosenfeld, D.: Aerosols, climate, and the~~
887 ~~hydrological cycle, *Science*, 294, 2119–2124, 2011.~~

888 Rao, T. N., Kirankumar, N. V. P., Radhakrishna, B., Rao, D. N., and Nakamura, K.:
889 Classification of tropical precipitating systems using wind profiler spectral moments.
890 Part I: Algorithm description and validation, *J. Atmos. Oceanic Technol.*, 25, 884–897,
891 2008.

892 Rao, T. N., Saikranthi, K., Radhakrisna, B., and Rao, S. V. B.: Differences in the
893 climatological characteristics of precipitation between active and break spells of the
894 Indian summer monsoon, *J. Clim.*, 29, 7797-7814, 2016.

895 Remer, L., Kaufman, Y., Tanr'e, D., Mattoo, S., Chu, D., Martins, J., Li, R., Ichoku, C.,
896 Levy, R., Kleidman, R., Eck, T., Vermote, E., and Holben, B.: The MODIS aerosol
897 algorithm, products, and validation, *J. Atmos. Sci.*, 62, 947–973, 2005.

898 Romatschke, U., Medina, S., and Houze, R. A.: Regional, seasonal, and diurnal variations of
899 extreme convection in the South Asian region, *J. Clim.*, 23, 419–439, 2010.

900 Rosenfeld, D., et al.: Global observations of aerosol-cloud-precipitation-climate interactions,
901 *Rev. Geophys.*, 52, 750-808, doi:10.1002/2013RG000441, 2014.

902 ~~Roxy, M.: Sensitivity of precipitation to sea surface temperature over the tropical summer~~
903 ~~monsoon region—and its quantification, *Clim. Dyn.*, 43, 1159–1169, 2014.~~

904 ~~Roxy, M.~~, Tanimoto, Y., Preethi, B., Terray, P., and Krishnan, R.: Intraseasonal SST-
905 precipitation relationship and its spatial variability over the tropical summer monsoon
906 region, *Clim. Dyn.*, 41, 45-61, 2013.

907 ~~Roxy, M.: Sensitivity of precipitation to sea surface temperature over the tropical summer~~
908 ~~monsoon region—and its quantification, *Clim. Dyn.*, 43, 1159–1169, 2014.~~

909 Sabin, T., Babu, C., and Joseph, P.: SST–convection relation over tropical oceans, *Int. J.*
910 *Climatol.* 33, 1424–1435, 2012.

911 Saikranthi, K., Radhakrishna, B., Satheesh, S. K., and Rao, T. N.: Spatial variation of
912 different rain systems during El Niño and La Niña periods over India and adjoining
913 ocean, *Clim. Dyn.*, 50, 3671-3685, doi: 10.1007/s00382-017-3833-4, 2018.

Formatted: Expanded by 0.2 pt,
Pattern: Clear (Custom
Color(RGB(252,252,252)))

914 Saikranthi, K., Rao, T. N., Radhakrishna, B., and Rao, S. V. B. : Morphology of the vertical
915 structure of precipitation over India and adjoining oceans based on long-term
916 measurements of TRMMPR, *J. Geophys. Res. Atmos.*, 119, 8433–8449, doi:
917 10.1002/2014JD021774, 2014.

918 Sathiyamoorthy, V., Mahesh, C., Gopalan, K., Prakash, S., Shukla, B. P., Mathur, A.:
919 Characteristics of low clouds over the Arabian Sea, *J. Geophys. Res.*
920 *Atmos.*, 118, 13489-13503, 2013.

Formatted: Emphasis, Font: +Body
(Calibri), 11 pt, Not Italic

921 ~~Radhakrishna, B., Satheesh, S. K., and Rao, T. N.: Spatial variation of different rain~~
922 ~~systems during El Niño and La Niña periods over India and adjoining ocean, *Clim.*~~
923 ~~*Dyn.*, 50, 3671-3685, doi: 10.1007/s00382-017-3833-4, 2018.~~

924 ~~Satheesh, S. K., Moorthy, K. K., Kaufman, Y. J., and Takemura, T.: Aerosol Optical depth,~~
925 ~~Physical properties and Radiative forcing over the Arabian Sea, *Meteorol. Atmos.*~~
926 ~~*Phys.*, 91, 45–62, doi:10.1007/s00703-004-0097-4, 2006.~~

927 ~~Schumacher, C. and Houze, R. A.: Comparison of radar data from the TRMM satellite and~~
928 ~~Kwajalein oceanic validation site. *J. Appl. Meteor.*, 39, 2151–2164, 2000.~~

Formatted: Strong, Font: +Body
(Calibri), 11 pt

Formatted: Default Paragraph Font,
Font: +Body (Calibri), 11 pt

929 Schumacher, C. and Houze, R. A.: Stratiform rain in the tropics as seen by the TRMM
930 precipitation radar, *J. Climate.*, 16, 1739–1756, 2003.

931 Sengupta, D., Goswami, B. N., and Senan, R.: Coherent intraseasonal oscillations of ocean
932 and atmosphere during the Asian summer monsoon, *Geophys. Res. Lett.*, 28, 4127–
933 4130, 2001.

934 Shenoi, S. S. C., Shankar, D., and Shetye, S. R.: Differences in heat budgets of the near-
935 surface Arabian Sea and Bay of Bengal: Implications for the summer monsoon, *J.*
936 *Geophys. Res.*, 107(C6), 3052, doi:10.1029/2000JC000679, 2002.

937 [Shige, S. and Kummerow, C.D.: Precipitation-Top Heights of Heavy Orographic Rainfall in](#)
938 [the Asian Monsoon Region, *J. Atmos. Sci.*, 73, 3009–3024, 2016.](#)

939 Sunilkumar, K., Rao, T. N., Saikranthi, K., and Rao, M. P.: comprehensive evaluation of
940 multisatellite precipitation estimates over India using gridded rainfall data, *J. Geophys.*
941 *Res. Atmos.*, 120, doi:10.1002/2015JD023437, 2015.

942 [Takayabu, Y. N., Shige, S., Tao, W., and Hirota, N.: Shallow and deep latent heating modes](#)
943 [over tropical Oceans observed with TRMM PR spectral latent heating Data, *J. Climate*,](#)
944 [23, 2030–2046, 2010.](#)

945 ~~Tao, W.-K., et al.: Retrieval of latent heating from TRMM measurements, *Bull. Am.*
946 *Meteorol. Soc.*, 87, 1555–1572, 2006.~~

947 Tao, W.-K., Chen, J.-P., Li, Z., Wang, C., and Zhang, C.: Impact of aerosols on convective
948 clouds and precipitation, *Rev. Geophys.*, 50, RG2001, doi:10.1029/2011RG000369,
949 2012.

950 [Tao, W.-K., et al.: Retrieval of latent heating from TRMM measurements, *Bull. Am.*
951 *Meteorol. Soc.*, 87, 1555–1572, 2006.](#)

952 Twomey, S.: The influence of pollution on the short wave albedo of clouds, *J. Atmos. Sci.*,
953 34, 1149–1152, 1977.

954 [Wallace, J. M., and Hobbs, P. V.: Atmospheric science: An introductory survey, Second](#)
955 [edition, Academic press, pp. 85, 2006.](#)

956 Wang, B., Ding, Q., Fu, X., Kang, I.-S., Jin, K., Shukla, J., and Doblus-Reyes, F.:
957 Fundamental challenge in simulation and prediction of summer monsoon rainfall,
958 *Geophys. Res. Lett.*, 32, L15711, doi:10.1029/2005GL022734, 2005.

959 ~~Wood, R., and Bretherton, C. S.: On the relationship between stratiform low cloud cover and~~
960 ~~lower tropospheric stability, *J. Clim.*, 19, 6425–6432, 2006.~~

961 ~~Weller, R. A., Farrar, J. T., Buckley, J., Mathew, S., Venkatesan, R., Lekha, J. S., Chaudhuri,~~
962 ~~D., Kumar, N. S., and Kumar, B. P.: Air-sea interaction in the Bay of Bengal,~~
963 ~~*Oceanography*, 29(2), 28–37, 2016.~~

964 Woolnough, S.J., Slingo, J.M., and Hoskins, B.J.: The relationship between convection and
965 sea surface temperature on intraseasonal timescales, *J. Climate*, 13, 2086–2104, 2000.

966 Wu, R., and Kirtman, B. P.: Roles of Indian and Pacific Ocean air–sea coupling in tropical
967 atmospheric variability, *Clim. Dyn.*, 25(2–3), 155–170, 2005.

Formatted: Widow/Orphan control,
Adjust space between Latin and Asian
text, Adjust space between Asian text
and numbers

968 ~~Yoneyama, K., Zhang, C., and Long, C. N.: Tracking pulses of the Madden–Julian~~
969 ~~oscillation, *Bull.*~~

970

971

972

973

974 ~~*Amer. Meteor. Soc.*, 94, 1871–1891, <https://doi.org/10.1175/BAMS-D-12-00157.1>, 2013.~~

Formatted: Pattern: Clear

975

Formatted: Indent: Left: 0 cm, First
line: 0 cm

976

977

978

979

980

981

982

983

Figure captions

984 **Figure 1:** Spatial distribution of SWMISM mean SST (in °C) obtained from ERA-Interim
985 reanalysis data over the AS (63°E-72°E & 8°N-20°N) and the BOB (83°E-92°E &
986 8°N-21°N). The regions considered in this analysis over these two seas are shown
987 with the boxes.

988 **Figure 2:** (a) and (b) represent the altitudinal distribution of occurrence of conditional
989 reflectivity (≥ 17 dBZ) as a function of SST with respect to precipitation occurrence at
990 that particular SST interval over the AS and the BOB, respectively.

991 **Figure 3:** (a), (d) and (b), (e) represent vertical profiles of median reflectivity correspond to
992 deep systems and their standard deviation (in dBZ) with SST over the AS (63°E-72°E
993 & 8°N-20°N) and the BOB (83°E-92°E & 8°N-21°N), respectively during the
994 SWMISM season. (c) and (f) show the number of conditional reflectivity pixels at
995 each altitude used for the estimation of the median and standard deviation.

996 ~~**Figure 4:** The variation of mean LTS with SST over the AS (63°E-72°E & 8°N-20°N) and~~
997 ~~the BOB (83°E-92°E & 8°N-21°N) during the SWM season.~~

998 ~~**Figure 4:** Same as Fig. 3 but for shallow precipitating systems.~~

999 **Figure 5:** (a) and (b), respectively, represent the ~~variation~~ vertical profiles of mean ~~CER_{ice}~~ θ_e
1000 (in ~~μmK~~) with SST over the AS and the BOB during the ISM season. (c) and ~~mean~~
1001 ~~CER_{liquid}~~ (in μm) with SST over the AS (63°E-72°E & 8°N-20°N) (d) and the BOB
1002 (83°E-92°E & 8°N-21°N) during the SWM season. (e) and (f) are same as (a) and (b)
1003 but for mean vertical velocity (in Pa s^{-1}) and wind gradient with reference to 950 hPa
1004 level (in m s^{-1}).

1005 **Figure 6:** (a) ~~The variation~~ Mean and standard error of ~~mean~~ AOD and (b) TCWV (in mm)
1006 with SST over the AS (63°E-72°E & 8°N-20°N) and the BOB during ISM.

Formatted: Font: Not Bold


1007 **Figure 7:** Variation of mean and standard error of CER liquid (in μm) with SST over the AS
1008 and the BOB (83°E 92°E & 8°N 21°N) during SWM. the ISM season.

1009 **Figure 7:** Winds during the SWM season at 850 hPa and 500 hPa levels. The shading colors
1010 represent the magnitude of the wind and arrow indicates the direction of the wind.


1011
1012
1013 **Figure 8:** Latitudinal variation (for every 2° latitude interval) of mean aerosol optical depth
1014 over Arabian Sea averaged over 63 72°E .

Formatted: Left, Indent: Left: 0 cm,
First line: 0 cm, Space After: 8 pt,
Line spacing: Multiple 1.08 li

1015
1016
1017 **Figures**
1018



1019 Formatted: Font: (Default) Times New Roman, 12 pt




Formatted: Font: (Default) Times New Roman, 12 pt

1020
 1021 **Figure 1:** Spatial distribution of SWMISM mean SST (in °C) obtained from ERA-Interim
 1022 reanalysis data over the AS (63°E-72°E & 8°N-20°N) and the BOB: (83°E-92°E &
 1023 8°N-21°N). The regions considered in this analysis over these two seas are shown
 1024 with the boxes.

1025
 1026
 1027
 1028
 1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038

1039

1040



1041

1042

1043 **Figure 2:** (a) and (b) represent the altitudinal distribution of occurrence of conditional
1044 reflectivity (≥ 17 dBZ) as a function of SST with respect to precipitation occurrence at
1045 that particular SST interval over the AS and the BOB, respectively.

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055


1056

1057

Formatted: Font: Not Bold, No underline

Formatted: Left, Line spacing: Multiple 1.08 li

1058
1059
1060
1061




1063

1064 **Figure 3:** (a), (d) and (b), (e) represent vertical profiles of median reflectivity correspond to
 1065 deep systems and their standard deviation (in dBZ) with SST over the AS (~~63°E-72°E~~
 1066 ~~& 8°N-20°N~~) and the BOB (~~83°E-92°E & 8°N-21°N~~), respectively during the
 1067 SWMISM season. (c) and (f) show the number of conditional reflectivity pixels at
 1068 each altitude used for the estimation of the median and standard deviation.

Formatted: Indent: Left: 0 cm,
Hanging: 1.25 cm


1069
1070
1071

Formatted: Left, Line spacing:
Multiple 1.08 li



1072

Formatted: Font: (Default) Times New Roman, 12 pt



1074

Formatted: Font: (Default) Times New Roman, 12 pt

1075

1076

1077

Figure 4: (a) The variation of mean LTS (in °K) with SST over the AS (63°E-72°E & 8°N-20°N) and the BOB (83°E-92°E & 8°N-21°N) during the SWM season. (b) Same as (a) Fig. 3 but for CAPE shallow precipitating systems.

1078


Formatted: Indent: Left: 0 cm, Hanging: 1.27 cm, Line spacing: Multiple 1.08 li

1079

Formatted: Left, Line spacing: Multiple 1.08 li

1080

1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107



Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Left, Line spacing: Multiple 1.08 li


Formatted: Font: Not Bold

Formatted: Left, Indent: Left: 0 cm, First line: 0 cm

Formatted: Indent: Left: 0 cm, Hanging: 1.25 cm

1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136

Figure 5: (a) and (b), respectively, represent the ~~variation~~vertical profiles of mean ~~CER~~ θ_e (in ~~μm~~ K) with SST over the AS and the BOB during the ISM season. (c) and (d) and (e) and (f) are same as (a) and (b) but for mean vertical velocity (in Pa s^{-1}) and wind gradient with reference to 950 hPa level (in m s^{-1}).



Formatted: Font: (Default) Times New Roman, 12 pt


Figure 6: Variation of mean and standard error of CER liquid (in μm) with SST over the AS (~~63°E 72°E & 8°N 20°N~~) and the BOB (~~83°E 92°E & 8°N 21°N~~) during the SWMISM season.

Formatted: Left, Indent: Left: 0 cm, First line: 0 cm, Space After: 8 pt, Line spacing: Multiple 1.08 li


Formatted: Left, Line spacing: Multiple 1.08 li

1137


1138



Formatted: Font: (Default) Times New Roman, 12 pt




Formatted: Font: (Default) Times New Roman, 12 pt



Formatted: Font: Not Bold

Formatted: Left, Indent: Left: 0 cm, First line: 0 cm



1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166 **Figure 67:** (a) ~~The variation~~ Mean and standard error of ~~mean~~ AOD and (b) TCWV (in mm)
1167 with SST over the AS (~~63°E 72°E & 8°N 20°N~~) and the BOB (~~83°E 92°E & 8°N~~
1168 ~~21°N~~) during SWMISM.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

1169

Formatted: Left, Indent: Left: 0 cm,
First line: 0 cm

1170

1171

1172

1173

1174

1175

1176

1177

Supplementary material

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190


Satheesh et al. (2006) showed an increase in AOD with increase in latitude over the AS due to the dust advection from Arabia desert regions during ISM season, whereas SST decreases with increase in the latitude. In other words the SST is low and AOD is high in northern AS whereas over the southern AS, SST is high and AOD is low. This contrasting spatial distribution of AOD and SST could cause a negative correlation between AOD and SST as depicted in Fig. 6a. To examine whether the observed decrease in AOD with increase in SST over the AS is due to the latitudinal variation of AOD or exists at all latitudes, we have segregated the data into 2° latitude bins and plotted the mean AOD with SST for all bins and is depicted in Fig. S2. In spite of the magnitude, AOD variation with SST is nearly similar at all latitudes of the AS, i.e., the higher AOD is observed at lower SSTs and vice versa (Fig. S2a). On the other hand the latitudinal variation of AOD with SST over the BOB shown in Fig. S2b also show a decrease in AOD with SST till 30 °C but the magnitude of variation is trivial relative to the AS. Also, as depicted in Fig. 6a AOD increases above 30 °C

1191 with SST over the BOB. This indicates that though there is a difference in magnitude of
1192 variation, AOD varies with SST over both the seas at all latitudes. This analysis is repeated
1193 using the multi-angle imaging spectroradiometer (MISR) dataset (which is not shown here)
1194 for small, medium large aerosol particles. Interestingly all three types also show a decrease in
1195 AOD with rise in SST over both the seas.

1196
1197 Satheesh, S. K., Moorthy, K. K., Kaufman, Y. J., and Takemura, T.: Aerosol Optical depth,
1198 physical properties and radiative forcing over the Arabian Sea, *Meteorol. Atmos.*
1199 *Phys.*, 91, 45–62, doi:10.1007/s00703-004-0097-4, 2006.

1200

1201



1202

1203 **Figure 7:** Winds during the SWM season at 850 hPa and 500 hPa levels. The shading colors
1204 represent the magnitude of the wind and arrow indicates the direction of the wind.

1205

1206


1207

1208

1209


Formatted: Font: (Default) Times New Roman, 12 pt

1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223



1224
1225
1226
1227
1228
1229

Formatted: Font: (Default) Times New Roman, 12 pt



Formatted: Font: (Default) Times New Roman, 12 pt


1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248

Figure S1: Variation of mean CAPE (in J kg⁻¹) with SST over the AS and the BOB during the ISM season.

1249

1250

Formatted: Font: (Default) Times New Roman, 12 pt



1251

1252 **Figure 8S2:** (a) and (b), respectively, represent latitudinal variation (for every 2° latitude
1253 interval) of mean AOD over the AS (between 63°E and 72°E) and the BOB (between
1254 83°E and 92°E).

1255