

Molecular characterization of organic aerosols in the Kathmandu Valley, Nepal: insights into primary and secondary sources

Xin Wan^{1,8}, Shichang Kang^{2,7}, Maheswar Rupakheti^{3,4}, Qianggong Zhang^{1,7}, Lekhendra Tripathi²,
5 Junming Guo², Pengfei Chen², Dipesh Rupakheti², Arnico K. Panday⁵, Mark G. Lawrence³, Kimitaka
Kawamura⁶, Zhiyuan Cong^{1,7*}

¹ Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan
Plateau Research, Chinese Academy of Sciences (CAS), Beijing 100101, China

10 ² State Key Laboratory of Cryospheric Science, Northwest Institute of Eco-Environment and Resources,
CAS, Lanzhou 730000, China

³ Institute for Advanced Sustainability Studies (IASS), Potsdam 14467, Germany

⁴ Himalayan Sustainability Institute (HIMSI), Kathmandu, Nepal

⁵ International Centre for Integrated Mountain Development (ICIMOD), Kathmandu, Nepal

15 ⁶ Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan

⁷ CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China

⁸ University of Chinese Academy of Sciences, Beijing 100039, China

*Corresponding author:

20 E-mail address: zhiyuancong@itpcas.ac.cn

Abstract:

Organic atmospheric aerosols in the Hindu Kush-Himalayan-Tibetan Plateau region are still poorly characterized. To better understand the sources and formation processes of the primary organic aerosols (POA) and secondary organic aerosol (SOA) in the foothills region of the central Himalaya, we studied atmospheric aerosol samples collected over a one-year period from April 2013 to April 2014 at the suburban site of Bode in the Kathmandu Valley. We measured major ions, organic carbon (OC), elemental carbon (EC), and various organic tracers emitted by specific sources. Tracer-based estimation methods were employed to characterize aerosol species, identify their likely sources, and apportion contributions from each source. The concentrations of OC and EC increased during winter with a maximum monthly average in January. Levoglucosan (an organic tracer for biomass burning), OC, and EC showed similar seasonal variations throughout the year. With an annual average concentration of $788 \pm 685 \text{ ng m}^{-3}$ (ranging from 58.8 to 3079 ng m^{-3}), levoglucosan was observed as the dominant species among all the analyzed organic tracers. Biomass burning contributed a significant fraction to OC, averaging $24.9 \pm 10.4\%$ during the whole year, and up to $36.3 \pm 10.4\%$ in the post-monsoon season. On an annual average basis, anthropogenic toluene-derived secondary OC accounted for 8.8% and biogenic secondary OC contributed 6.2% to total OC. The annual contribution of fugal-spores to OC was 3.2% with the maximum during the monsoon (5.9%). For plant debris, it accounted for 1.4% of OC during the monsoon. Therefore, OC is mainly associated with biomass burning and other anthropogenic activity in the Kathmandu Valley. Our findings are conducive to designing control measures to mitigate the heavy air pollution and its impacts in the Kathmandu Valley and surrounding area.

1. Introduction

The Indo-Gangetic Plain (IGP) region in South Asia, is a global air pollution hotspot. Atmospheric pollutants [e.g., organic carbon (OC), black carbon (BC) and SO₂] from South Asia have been increasing in recent decades (Ramanathan et al., 2005; Muzzini and Aparicio, 2013; Lawrence and Lelieveld, 2010). While these pollutants are of concern locally near the emission sources, they can also, in short span of time, be transported to rural and remote regions over a long distance. This results in an annually recurring regional scale haze, referred to as atmospheric brown clouds (ABC), that covers a large area from the Himalayan range to the Indian Ocean (Ramanathan et al., 2007). Until recently the emissions, types, levels, atmospheric transport and transformation, impacts and mitigation of various atmospheric pollutants were not well characterized in the vast mountain areas and the foothills region in South Asia. In this context, the international project of “A Sustainable Atmosphere for the Kathmandu Valley (SusKat)” was launched, aiming to comprehensively understand the causes of the severe air pollution in the region, and identifying appropriate solutions to reduce its impacts (Rupakheti et al., 2018). This paper presents analyses of samples collected as part of the SusKat field campaign.

The Kathmandu Valley, the capital region of Nepal, is considered as one of the most polluted cities in South Asia and the largest metropolitan region in the foothills of the Hindu Kush-Himalayas-Tibetan Plateau (HKHTP) region, facing rapid but unplanned urbanization, with the current population of approximately 4 million (Muzzini and Aparicio, 2013). Additionally, due to the bowl-shaped topography, the free flow of air is restricted, resulting in poor air quality (Panday and Prinn, 2009; Shakya et al., 2010; Mues et al., 2017). Giri et al. (2006) showed that the PM₁₀ concentrations in Kathmandu were about 2-4 times higher than the guidelines prescribed by the World Health Organization (WHO) (50 µg m⁻³ of 24-hour mean) (WHO, 2006). More recently, Shakya et al. (2017) reported that daily average mass concentrations of PM_{2.5} at seven locations in the Kathmandu Valley during 2014 were about 5 times higher than the WHO guidelines (25 µg m⁻³ of 24-hour mean) (WHO, 2006). Beside particulate matter, recent studies have pointed out that ground-level ozone (O₃) is also of concern (Mahata et al., 2017b; Bhardwaj et al., 2017). Putero et al. (2015) found O₃ levels at Pakanajol in the city center to exceed the WHO’s 8-hour maximum ozone guidelines of 100 µg m⁻³ on 125 days in a year, while Mahata et al. (2017b) reported such exceedance was for nearly 3 months at Bode (where sampling for this study was

70 conducted) and 6 months at Nagarkot, a hilltop site downwind of the Kathmandu Valley. The concentrations of acetonitrile and isoprene (precursor for both O₃ and secondary organic aerosol, SOA) investigated by Sarkar et al. (2016) in the Kathmandu Valley were comparable with the highest reported elsewhere in the world. Air pollution is a clear threat to human health (leading to respiratory disease, cardiovascular disease, cancer, etc.), agricultural productivity and revenues from tourism in the
75 Kathmandu Valley and surrounding regions (Putero et al., 2015;Shakya et al., 2016).

Carbonaceous aerosols [organic carbon (OC) and elemental carbon (EC)] are often a principal component of atmospheric aerosols and the atmospheric brown clouds (ABC) in South Asia (Ramanathan et al., 2005). Sources and chemical transformations of OC are complicated, including primary emissions (biomass and fossil fuel combustion, plant debris and soil dust) and secondary formation of the oxidative
80 products of precursor gases emitted from both biogenic and anthropogenic sources (Simoneit et al., 2004;Claeys et al., 2004;Ding et al., 2016a). While past studies of aerosol characteristics in the Kathmandu Valley and surrounding regions are limited, they demonstrated that OC was the main component of aerosols observed in the Kathmandu Valley (Shakya et al., 2017;Kim et al., 2015). These studies have focused on a few aerosol species or a limited number of organic compound classes (Kim et al., 2015;Chen et al., 2015;Sarkar et al., 2016). Analysis of organic aerosols at the molecular level has
85 been reported only at a rural Godavari site on the southern edge of the Kathmandu Valley (Stone et al., 2010;Stone et al., 2012). Overall, the composition and sources of OC are still poorly characterized.

Therefore, to overcome such research gaps, our study comprehensively investigates the organic molecular compositions of aerosols from the Kathmandu Valley, including anhydrosugars,
90 monosaccharides, sugar alcohols, phenolic compounds, resin acid, phthalate esters, and secondary organic aerosols based on the analysis of various organic tracers emitted from primary sources and secondary formation of organic aerosol. We also studied the seasonal variation and molecular distribution to decipher their abundances, understand their predominant sources (primary vs. secondary), and to quantify the contributions of the most abundant sources of the carbonaceous aerosols in the suburban environment
95 in the Himalayan foothills. Our work adds to the growing database on the chemical characteristics of organic aerosols in South Asia, particularly in the HKHTP region.

2. Materials and methods

2.1. Sampling site

The Kathmandu Valley is a round flat basin in the southern foothills of the Himalayas. The basin bottom has an elevation of approximately 1300 meters above sea level, and it is surrounded by a ring of green mountains, with elevation ranging from 1500 m to 2800 m above sea level (Panday and Prinn, 2009). Our sampling was performed for one year from April 2013 to April 2014 at Bode (27.67°N, 85.38°E, 1320 m above sea level), a suburban site to the east of Kathmandu city (Fig. 1). Bode is located in the Madhyapur-Thimi area in the eastern part of the Kathmandu Valley. There are two major wind flows in the Kathmandu Valley: (a) west to east, from Nagdhunga-Bhimdhunga mountain pass in the west to Nagarkot-Sanga mountain pass in the east, (b) south to north, from Bagmati River corridor to the northeast direction through the central-eastern part of the Valley. These two airflows meet around central-eastern part of the Valley and move eastward towards the Nagarkot-Sanga passes (Regmi et al., 2003; Panday and Prinn, 2009). The Bode area receives these two air flows, and hence it is downwind of Kathmandu city and Lalitpur or Patan city located in southwest, west and northwest direction during the daytime, and Bhaktapur city located in east and southeast during nighttime (Rupakheti et al., 2018; Mahata et al., 2017a; Bhardwaj et al., 2017; Sarkar et al., 2016). In addition, it is situated in a residential area with urban buildings and houses scattered across agricultural fields with paddy, wheat, corn and vegetable farms. Some small industries (plastics, electronics, wood, fabrics, etc.) and Bhaktapur Industrial Estate are located in the south-eastern direction of Bode, as well as several brick kilns that use low quality coal during January to April (Sarkar et al., 2016). The Tribhuvan international airport in the west of the Bode (~ 4 km) may have potential impacts when there is westerly wind. Approximately 1.5 and 7 km to the north, there are two reserve forests, consisting of a mix mainly broad-leaved deciduous trees and evergreen conifer trees (Department of Plant Resources, 2015). BC and O₃ measurements in the two major SusKat-ABC sites (Paknajol and Bode) in the Valley show similar levels (Putero et al., 2015; Mahata et al., 2017b). Therefore, Bode site can be taken as a representative site for the Kathmandu Valley (Rupakheti et al., 2018).

2.2 Sample collection

A medium-volume sampler (KC-120H, Laoshan Co., China) was placed on a building rooftop, about

125 20 m above the ground. We continuously collected total suspended particulates (TSP) for 23 h (day and
night time) every five days under the flow rate of 100 L min⁻¹. Overall, 82 aerosol samples were
successfully obtained using 90 mm diameter quartz filters (Whatman PLC, Maidstone, UK). The filters
were pre-baked at 550 °C for 6 h to remove all organic material and weighed by a microbalance
(sensitivity: ±0.01 mg) before and after sampling. Before each weighing, they were equilibrated for 24 h
130 at the constant temperature (25±3 °C) and humidity (30±5%) conditions. Finally, the filters were frozen
at -20 °C until laboratory analysis. Field blanks (one blank filter each month) were also collected, briefly
placing a filter into the sampler without drawing air to assess potential contamination. There may be
positive and negative artifacts during the sample handling/conditioning due to the adsorption/evaporation
processes of organic aerosols (Fu et al., 2010; Li et al., 2018; Boreddy et al., 2017; Oanh et al., 2016). In a
135 comparable study, Ding et al. (2013) reported the positive artifacts of 10–20% for OC and up to 16% for
organic tracers using a backup quartz filter placed behind the main quartz filter.

2.3 Chemical Analysis

The aerosol samples were analyzed for major ions, OC, EC, and organic molecular tracers in the
laboratory. Major ions (Ca²⁺, Na⁺, K⁺, Mg²⁺, NH₄⁺, Cl⁻, SO₄²⁻ and NO₃⁻) were measured using an ion
140 chromatography (Dionex, USA) with ICS-320 and ICS-1500 (Tripathy et al., 2017). The detection limit
for all the major ions was 0.01 µg m⁻³. They denoted less than 5% of the real sample concentrations in
the field blank filters (Tripathy et al., 2017). Non-sea-salt Ca²⁺ (nss-Ca²⁺) and K⁺ (nss-K⁺) was estimated
according to the method from George et al. (2008). OC and EC were determined by the DRI OC/EC
analyzer (Model 2001A, USA) according to the thermal/optical reflectance (TOR) method with the
145 IMPROVE-A protocol (Wan et al., 2015). The concentrations of OC and EC from field blank filters were
0.59±0.13 µg m⁻³ and 0.00 µg m⁻³, respectively. The OC data reported here were blank corrected.

Detailed analytical method of organic molecular tracers was described previously by Wan et al.
(2017). Briefly, small filter aliquots (1.13-3.39 cm²) were cut, spiked with appropriate amounts methyl-
β-D-xylanopyranoside (MXP, 99%, Sigma) and D₃-malic acid (DMA, CDN isotopes, 99%) as internal
150 recovery standards. The cut filters were then extracted three times with dichloromethane/methanol (2:1,
v/v) at room temperature for 30 minutes (20 ml each time). The solvent extracts in total of 60 ml were
combined and successively filtered with quartz wool, concentrated, blown to dryness using ultrapure

nitrogen gas and then reacted with 50 μl of 99% N,O-bis-(trimethylsilyl)trifluoroacetamide (BSTFA, with 1% trimethylsilyl chloride) and pyridine (v/v=2:1) at 70 $^{\circ}\text{C}$ for 3 h. After derivatization, 150 μl n-hexane was added to the derivatives. A trace gas chromatography coupled to a PolarisQ mass spectrometry detector (GC-MS, Thermo Scientific) was used for analysis. The GC instrument was equipped with a TG-5MS (30 m \times 0.25 mm I.D. \times 0.25 μm film thickness). The injection port was set for split/splitless injection. The oven temperature was initially held at 50 $^{\circ}\text{C}$ for 2 min, increased to 120 $^{\circ}\text{C}$ at 30 $^{\circ}\text{C min}^{-1}$, then to 300 $^{\circ}\text{C}$ at 6 $^{\circ}\text{C min}^{-1}$ and finally held for 16 min. The MS was operated in electron ionization (EI) mode at 70 eV with a scan range of 50-650 Da.

For quantitative analysis, calibration curves were established by using authentic standards that were processed as described above. For the quantification of target compounds that were no available standards, they were estimated by the following surrogate compounds: erythritol for 2-methylglyceric acid, 2-methyltetrols and C₅-alkene triols, cis-pinonic acid for 3-hydroxyglutaric and 3-methyl-1,2,3-butanetricarboxylic acids, pinic acid for β -caryophyllinic acid, azelaic acid for 2,3-dihydroxy-4-oxopentanoic acid (DHOPA). Recoveries for target compounds and MXP (Table S1) were more than 75%. The exception was for malic acid (50.3%-90.5%) and cis-pinonic acid (60.2%-81.8%). Duplicate analysis showed that the relative differences were less than 15%. The method detection limits were 0.04-0.13 ng m⁻³ (Table S1). The results reported in the current study were not corrected for the recoveries. Field blank filters were analyzed by the procedure used by the samples above, but no target compounds were detected.

Table 1 shows a list of chemical species (OC, EC and major ions) and molecular markers (from the sources of biomass burning, fungal spores, plant debris, plastic emission, biogenic and anthropogenic secondary formation) analyzed in this study. They include anhydrosugars (levoglucosan, mannosan and galactosan), primary saccharides (sucrose, glucose, trehalose and xylose) and sugar alcohols (mannitol, arabitol, sorbitol and erythritol), lignin and resin pyrolysis products (vanillic, syringic, p-hydroxybenzoic and dehydroabiatic acids), phthalate esters, isoprene tracers (2-methylglyceric acid, 2-methylthreitol, 2-methylerythritol, cis-2-methyl-1,3,4-trihydroxy-1-butene, 3-methyl-2,3,4-trihydroxy-1-butene and trans-2-methyl-1,3,4-trihydroxy-1-butene), monoterpene tracers (cis-pinonic, pinic, 3-hydroxyglutaric and 3-methyl-1,2,3-butanetricarboxylic acids), β -caryophyllinic and 2,3-dihydroxy-4-oxopentanoic acids.

2.4 Estimation of measurement uncertainty

Since there is no commercial standard available for most SOA tracers (except for cis-pinonic acid and pinic acid), the use of surrogate standards for quantification introduces additional error to the measurements. Error in analyte measurement (EA) is propagated from the standard deviation of the field blank (EFB), error in spike recovery (ER) and the error from surrogate quantification (EQ):

185

$$EA = \sqrt{EFB^2 + ER^2 + EQ^2}$$

190

EFB was 0 in this study due to SOA tracers that were not detected in the field blanks. The spike recoveries of surrogate standards were used to estimate the ER of tracers, ranging from 9.2% (erythritol) to 26.1% (cis-pinonic acid). According to Stone et al. (2012), there is an empirical approach to estimate EQ based on homologous series of atmospherically relevant compounds. The relative error introduced by each carbon atom (En) was estimated to be 15 %, each oxygenated functional group (Ef) to be 10% and alkenes (Ed) to be 60%. Therefore, the EQ are calculated as:

$$EQ = En\Delta n + Ef\Delta f + Ed\Delta d$$

where Δn , Δf and Δd are the difference of carbon atom number, oxygen-containing functional group and alkene functionality between a surrogate and an analyte, respectively.

195

The estimated uncertainties in tracer measurement is presented in Table S2. The EQ ranged from 15% (2-methyltetrols) to 120% (β -caryophyllenic acid) in this study. Propagated with the error in recovery, EA were estimated in the range of 17.6% to 122.4%.”

2.5 Meteorological parameters

200

The meteorological parameters such as air temperature (T), relative humidity (RH), atmospheric pressure (P), visibility (V), wind speed (WS) and direction (WD) were used in this study. They were derived from Tribhuvan International Airport (www.wunderground.com), which was located west of Bode (approximately 4 km). Mixing layer height (MLH) data was measured with a Vaisala ceilometer at Bode site (Mues et al., 2017). The meteorology of Kathmandu Valley and its surrounding regions is controlled by the South Asian monsoon circulations in the wet season (monsoon, June-September). Westerlies dominate the atmospheric circulation patterns during the dry seasons including pre-monsoon (March-May), post-monsoon (October-November) and winter (December-February) with limited precipitation (Pudasainee et al., 2006; Mues et al., 2017). Additionally, the valley is also influenced by

205

local mountain valley circulation (Mues et al., 2018).

3. Results and discussion

210 A statistical concentration summary of constituent major ions, OC, EC, and major organic compounds identified in TSP samples collected at the Bode site is presented in Table 1. Tracers for six classes of organic compounds were detected: anhydrosugars, monosaccharides, sugar alcohols, phenolic compounds and resin acid, phthalate esters, and secondary organic aerosol (SOA) tracers.

3.1 Aerosol loadings

215 The TSP samples at Bode site exhibited daily mass concentrations from 32.0 to 723 $\mu\text{g m}^{-3}$ with an average concentration of $255\pm 167 \mu\text{g m}^{-3}$ during April 2013 to April 2014 (Table 1). Putero et al. (2015) reported $195\pm 83 \mu\text{g m}^{-3}$ of online $\text{PM}_{2.5}$ concentration in Pakajol site (also one of SusKat-ABC sites), accounting for roughly 80% of TSP in our study. The TSP concentrations were comparable to those reported in other heavily polluted cities in South Asia, including Islamabad in Pakistan (Shah and Shaheen, 220 2008), Kolkata (Gupta et al., 2007) and Agra (Rajput and Lakhani, 2010) in India. Compared to the remote sites such as Manora Peak in the central Himalaya (Ram et al., 2010) and Lulang in the Tibetan Plateau (Wang et al., 2015), the TSP in Bode shows significantly higher mass concentrations. We found a clear seasonal variation in TSP mass concentrations (Fig. 2a) with a maximum during the pre-monsoon season while a minimum during the monsoon season, and an increasing trend from the post-monsoon to the winter season were observed. It generally corresponded to the build-up of the atmospheric brown clouds (ABCs), which engulfed most of South Asia and the Northern Indian Ocean, during the long dry season 225 extending from November to May (Ramanathan et al., 2005).

Meteorological parameters may also affect the TSP concentrations. The highest TSP concentration observed during the pre-monsoon can be caused by the fugitive dust which is been blown up by strong wind and the absence of wet-precipitation (Fig S1a and c). The lower TSP concentration in the monsoon 230 was likely related to increased precipitation (Fig S1c) after the onset of the South Asian monsoon. Nearly 80% of the annual precipitation falls during the monsoon season in the Kathmandu Valley, which flushes out pollutants from the atmosphere (Sharma et al., 2012; Tripathee et al., 2017). During winter, an inversion layer often occurs in the Kathmandu Valley owing to its bowl-shaped topography (Pudasainee

235 et al., 2006). The existence of an inversion layer with the lower temperature ($12.0\pm 2.41^\circ\text{C}$), wind speed ($2.86\pm 1.34\text{ km h}^{-1}$), and MLH ($0.34\pm 0.08\text{ km}$) (Mues et al., 2017) (Fig S1a, c and d) altogether reduced the pollution dispersion mechanism resulting in increased levels of pollutants close to the ground surface.

3.2 Major ions and OC/EC

240 Concentrations of eight major ions were measured in the aerosol samples from the Bode site. The total sum accounted for $17.1\% \pm 8.5\%$ of annual average TSP mass. Sulfate ranked the highest among them, with an annual mean of $10.8\pm 9.83\text{ }\mu\text{g m}^{-3}$, followed by Ca^{2+} ($7.96\pm 6.85\text{ }\mu\text{g m}^{-3}$), NH_4^+ ($5.92\pm 6.16\text{ }\mu\text{g m}^{-3}$), NO_3^- ($5.21\pm 4.35\text{ }\mu\text{g m}^{-3}$), Na^+ ($3.28\pm 1.58\text{ }\mu\text{g m}^{-3}$), K^+ ($2.43\pm 2.82\text{ }\mu\text{g m}^{-3}$), Cl^- ($2.15\pm 2.25\text{ }\mu\text{g m}^{-3}$) and Mg^{2+} ($0.61\pm 0.54\text{ }\mu\text{g m}^{-3}$). On average, the combination of SO_4^{2-} , NO_3^- and NH_4^+ , i.e. the secondary inorganic aerosols, constituted more than half (51.3%) of the total ionic concentrations. The Ca^{2+} alone
245 accounted for 22.1% of total ions.

Sulfate, ammonium and nitrate revealed a typical seasonality with the seasonally averaged concentrations ranked in the descending order of winter > pre-monsoon > post-monsoon > monsoon. This is consistent with the seasonal variation of the precursors NO_x , NO_2 and SO_2 , which are mainly caused by automobile exhaust, household cooking, and operation of the typical biomass co-fired brick kilns in
250 the Kathmandu Valley (Kondo et al., 2005; Kiros et al., 2016). Currently, nearly 50% of the total motor vehicles in Nepal (approximately 2.33 million) run on the Kathmandu Valley roads (DoTM, 2015; Mahata et al., 2017b). Diesel- or gasoline-powered generators (producing higher NO_x emissions) and garbage burning are other major sources of air pollution in Nepal during the sampling period, which can also emit many aerosol precursors (Stockwell et al., 2016).

255 Ions derived from crustal sources, such as Ca^{2+} and Mg^{2+} are related to the local fugitive dust sources such as road dusts and construction activities (Ram et al., 2010). Interestingly, good correlations between Ca^{2+} and SO_4^{2-} ($R^2 = 0.48$, $P < 0.001$), Ca^{2+} and NO_3^- ($R^2 = 0.58$, $P < 0.001$), Ca^{2+} and NH_4^+ ($R^2 = 0.62$, $P < 0.001$), Mg^{2+} and SO_4^{2-} ($R^2 = 0.61$, $P < 0.001$), Mg^{2+} and NO_3^- ($R^2 = 0.71$, $P < 0.001$), Mg^{2+} and NH_4^+ ($R^2 = 0.69$, $P < 0.001$) were observed (Table 2), which hinted that dust may co-exist with SO_4^{2-} , NO_3^- and
260 NH_4^+ in the Kathmandu Valley (Tripathee et al., 2017).

The annual mean concentrations of carbonaceous aerosols (OC: $38.7\pm 32.7\text{ }\mu\text{g m}^{-3}$ and EC: 9.92 ± 5.33

$\mu\text{g m}^{-3}$) accounted for $19.2\% \pm 5.48\%$ of TSP mass at the Bode site, which was higher than that of the major ions. OC alone accounted for $14.6\% \pm 4.81\%$ of the TSP mass. OC and EC showed much higher concentrations during winter and pre-monsoon seasons than that in monsoon season (Fig. 2b and c). In this study, we found that the daily OC/EC ratios ranged from 0.77 to 15.8, with an annual mean ratio of 3.78 ± 2.73 , and seasonal mean ratios of 4.44, 2.71, 3.31, and 5.86 during pre-monsoon, monsoon, post-monsoon, and winter, respectively (Table 1 and Fig. 2d). The OC/EC mass ratio of more than 2.0 indicates the presence of secondary organic matter or biomass burning aerosols (Cao et al., 2007). Their influence and contribution will be discussed in the following sections. The OC/EC ratios found in this study for the Kathmandu valley were similar to other sites in South Asia, like Lumbini (5.16 ± 2.09 , 2.41-10.03) (Wan et al., 2017), Delhi (5.86 ± 0.99 , 2.9-9.2) (Bisht et al., 2015) and Lahore (3.9 ± 1.6 , 1.5-7.2) (Alam et al., 2014).

3.3 Sugar compounds

3.3.1 Anhydrosugars

Levoglucosan (1,6-anhydro- β -D-glucopyranose) and its two isomers (mannosan and galactosan) have been used as an ideal molecular tracer for biomass burning emissions (Graham et al., 2002; Simoneit, 2002). They are exclusively produced from the pyrolysis of cellulose and hemicellulose. In the current study, levoglucosan was observed as the most abundant species among the individual compounds identified with an average concentration of $788 \pm 685 \text{ ng m}^{-3}$ (ranging from 58.8 to 3079 ng m^{-3}) (Table 1).

For the seasonality, levoglucosan showed significantly higher levels during winter, pre-monsoon and post-monsoon (Fig. 3a). Especially, the higher concentrations were recorded in winter ranging from 830 to 2395 ng m^{-3} (average: $1391 \pm 535 \text{ ng m}^{-3}$). It showed the comparable levels with other sites in the world, which were badly affected by the biomass burning emissions, e.g., New Delhi (1977 ng m^{-3}) and Raipur (2180 ng m^{-3}) in India (Li et al., 2014; Deshmukh et al., 2016), Tasmania ($4540 \pm 2480 \text{ ng m}^{-3}$) in Australia (Reisen et al., 2013) and Lumbini (1161 ± 1347) in Nepal (Wan et al., 2017). Our results were much higher than the aerosols ($20\text{-}372 \text{ ng m}^{-3}$) collected at a rural Godavari site, located on the southern edge of the Kathmandu Valley during 2006 (Stone et al., 2010). Good correlations exhibited between levoglucosan and OC ($R^2=0.79$, $P<0.001$), EC ($R^2=0.42$, $P<0.001$) and nss-K^+ ($R^2=0.35$, $P<0.01$) during

the campaign (Fig. 4). This indicates that OC and EC in Kathmandu Valley's aerosols are strongly related to biomass burning source (Kim et al., 2015).

The ratio of levoglucosan to mannosan (Lev/Man) has been applied to distinguish the possible biomass burning categories. Higher Lev/Man ratios were reported for emissions from combustion of hardwood (ranging from 12.9 to 35.4 with an average of 21.5 ± 8.3) and agricultural residues (range from 12.7 to 55.7 with an average of 32.6 ± 19.1) in previous studies (Engling et al., 2009; Sang et al., 2013). For the softwood burning, the average ratio was 4.0 ± 1.0 (ranging from 2.5 to 5.8). In this study, the average Lev/Man ratio was 16.3 ± 5.96 (ranging from 9.13 to 33.1 with only 9 samples <10). It can be inferred that the combustion of crop residues and hardwood is likely to be one of the major sources of atmospheric pollution in this region. A previous study also reported that the combustion of wood fuel for cooking and heating is common during wintertime in Nepal while there is much more crop residue combustion during pre-monsoon and post-monsoon seasons (Stockwell et al., 2016). This is not only a local but also a regional phenomenon; for example, Bhardwaj et al. (2017) and Wan et al. (2017) reported that the emissions from crop residue burning during the pre-monsoon and post-monsoon seasons from western India and eastern Pakistan impact the air quality in Nepal. Similarly, Rupakheti et al. (2017) also showed that the combustion of agricultural residues and forest fires over the northwestern IGP region are causes of the air pollution episodes over the foothills of the central Himalayas. In addition, brick kilns mainly operated during January-April, burned substantial quantities of low-grade coal, mixed crop wastes and firewood (Stone et al., 2010; Kim et al., 2015). Such emissions may also lead to the high levels of levoglucosan observed at Bode. We must point out that the incense burning in Kathmandu Valley may also influence the levoglucosan concentration.

3.3.2 Monosaccharides

Primary biological aerosol particle (PBAP) tracers, commonly known also as bioaerosols, were analyzed in the Bode aerosol samples, including five monosaccharides of glucose, fructose, trehalose, sucrose and xylose. PBAP is derived from fungal spores, vegetative debris, pollen, bacteria and viruses. Most of them can cause allergenic or immunotoxic effects on human health (Bauer et al., 2008; Myriokefalitakis et al., 2017).

In this study, total monosaccharides had an annual mean concentration of $298 \pm 127 \text{ ng m}^{-3}$. Glucose was the most abundant species ($124 \pm 60.0 \text{ ng m}^{-3}$), followed by fructose ($58.2 \pm 28.3 \text{ ng m}^{-3}$), sucrose ($48.3 \pm 27.4 \text{ ng m}^{-3}$), trehalose ($40.8 \pm 22.0 \text{ ng m}^{-3}$), and xylose ($26.5 \pm 18.1 \text{ ng m}^{-3}$) (Table 1). Except xylose, they all presented higher concentrations in pre-monsoon while being lower in winter (Fig 3h, i, j and k). There were significant linear correlations between glucose and fructose ($R^2 = 0.77$, $p < 0.001$), trehalose and glucose ($R^2 = 0.30$, $p < 0.001$), trehalose and fructose ($R^2 = 0.23$, $p < 0.001$), sucrose and glucose ($R^2 = 0.55$, $p < 0.001$), sucrose and fructose ($R^2 = 0.55$, $p < 0.001$), and sucrose and trehalose ($R^2 = 0.28$, $p < 0.001$) (Table 3). Therefore, the strong correlations indicated that they were derived from common sources, e.g. from local forests in the Kathmandu Valley during the period of high productivity of plants. In addition, the pollen produced from the flowering of local vegetation also largely contribute to glucose, fructose, trehalose, sucrose. The flowering of trees and crops peaks during the pre-monsoon season. The similar phenomenon was also reported in deciduous forests in northern Japan (Miyazaki et al., 2012).

Xylose has complex sources, including soils (Simoneit et al., 2004), microbiota (Wan and Yu, 2007), vegetation, bacteria (Cowie and Hedges, 1984) and biomass combustion (Zhu et al., 2015). It presents less abundant and only accounts for $6.90\% \pm 8.32\%$ of the total PBAP tracers identified in the Bode aerosols. For the seasonal pattern, it is characterized by waxing in winter ($47.4 \pm 24.3 \text{ ng m}^{-3}$) and waning in monsoon ($12.0 \pm 5.16 \text{ ng m}^{-3}$), which was different from the other primary monosaccharides (Table 1 and Fig. 2l). Close correlation between xylose and levoglucosan (the biomass burning tracer) was observed in our study (Fig. S2, $R^2 = 0.72$, $p < 0.001$), indicating that the emissions from the burning of biomass may largely contribute to xylose in Bode aerosols. A similar finding of xylose source (i.e. biomass burning) was also proposed by Zhu et al. (2015).

3.3.3 Sugar alcohols

Total concentration of sugar alcohols (arabitol, sorbitol, erythritol and mannitol), was $213 \pm 126 \text{ ng m}^{-3}$, and thus lower than that of total monosaccharides (Table 1). Mannitol ($86.9 \pm 55.3 \text{ ng m}^{-3}$) and arabitol ($68.4 \pm 39.8 \text{ ng m}^{-3}$) were the most abundant species of the sugar alcohols, followed by erythritol ($43.1 \pm 28.8 \text{ ng m}^{-3}$) and sorbitol ($14.2 \pm 8.02 \text{ ng m}^{-3}$). All of them exhibited monsoon maxima ($114 \pm 61.4 \text{ ng m}^{-3}$, $86.6 \pm 44.5 \text{ ng m}^{-3}$, $56.9 \pm 33.1 \text{ ng m}^{-3}$ and $17.9 \pm 9.31 \text{ ng m}^{-3}$, respectively) and winter minima (18.1 ± 6.02

ng m⁻³, 26.1±9.13 ng m⁻³, 5.82±2.72 ng m⁻³, 12.4±7.60 ng m⁻³, respectively) (Table 1 and Fig. 3m, n, o and p). They also showed significant correlations with each other, implying their common sources (Zhu et al., 2015;Fu et al., 2012). Mannitol and arabitol have been mostly associated with fungal spores, along with vegetation and mature leaves and algae (Liang et al., 2016;Myriokefalitakis et al., 2017;Yttri et al., 2007). Recent studies proposed that elevated concentrations of mannitol and arabitol were usually observed augmentation after rain events and also highly correlated with relative humidity (Zhang et al., 2010;Yue et al., 2016). Therefore, at Bode, sugar alcohols were likely emitted by plants in nearby forest and agriculture fields, especially during the monsoon with the higher relative humidity (Fig. S1b). In addition, the higher temperatures (Fig. S1a) were conducive for more active microbial activities. Notably, the levels of PBAP discussed above were much higher than other sites in the world (Zhu et al., 2015;Chen et al., 2013;Liang et al., 2016), indicating the strong fungal spore production in the Kathmandu Valley during the monsoon season.

3.4 Phenolic compounds and resin acid

Phenolic compounds (e.g., vanillic, syringic and p-hydroxybenzoic acids) derived from lignin pyrolysis and resin acid (e.g., dehydroabietic acid) from burning of conifer plants can be also used as biomarkers for biomass burning. Vanillic acid is dominant both in softwood and hardwood smoke (Myers-Pigg et al., 2016;Fu et al., 2010) while syringic acid is prevalent in hardwood smoke. Herbaceous plant smoke primarily contains p-anisaldehyde and p-anisic acid (e.g., p-hydroxybenzaldehyde and p-hydroxybenzoic acid). Dehydroabietic acid is a dominant compound in the total lipid material from pine wood smoke. Therefore, four pyrolysis products of lignin and resin acid (p-hydroxybenzoic, vanillic, syringic and dehydroabietic acids) were chosen as organic tracers in this study.

p-Hydroxybenzoic acid (19.8±12.3 ng m⁻³) was the predominant species, followed by dehydroabietic (13.8 ±6.19 ng m⁻³), vanillic (15.3±11.3 ng m⁻³) and syringic acids (17.1±13.7 ng m⁻³) (Table 1). They exhibited maximum concentrations during winter and pre-monsoon, decreased in monsoon and then increased from post-monsoon, which was consistent with the seasonal variation of levoglucosan (Fig 3d, e, f and g). There were also significant correlations of lignin and resin pyrolysis products with levoglucosan (cellulose pyrolysis products) (Fig. S3a, p-hydroxybenzoic acid and levoglucosan, R²=0.72, P<0.001; Fig. S3b, vanillic acid and levoglucosan, R²=0.86, P<0.001; Fig. S3c, syringic acid and

levoglucosan, $R^2=0.83$, $P<0.001$; Fig. S3d, levoglucosan and dehydroabietic acid, $R^2=0.63$, $P<0.001$). Such a result also shows that there are various biomass combustion sources in the valley.

375 The mass ratio of syringic to vanillic acids (syr/van) has recently been used to further discriminate the vegetation types burned (Fujii et al., 2015; Myers-Pigg et al., 2016; Wan et al., 2017). Myers-Pigg et al. (2016) reported that the syr/van ratios varied from 0.1 to 2.44 for combustion of hardwood and herbaceous angiosperm, while it was 0.01–0.24 for burning softwood. With regard to the aerosol samples from Kathmandu Valley, syr/van ratio was 0.94 ± 0.18 of an annual average ranging from 0.65 to 1.31, indicating that combustion of hardwood and herbaceous plant (including crop residue) are the most likely sources of biomass burning in the Valley. This finding is in agreement with the results obtained from the ratios of Lev/Man as discussed in Sect. 3.3.1.

380 Besides the information revealed by anhydrosugars discussed in section 3.3.1, lignin and resin biomarkers in the Kathmandu Valley's air further confirmed that biomass burning emissions is an important contributor to organic aerosols in this region, particularly during winter and pre-monsoon periods.

385 **3.5 Phthalate esters**

Phthalic acid esters or phthalates are extensively used as non-reactive plasticizers in the manufacture and processing of plastic products. They can be easily released into the environment from the matrix by evaporation due to their physically rather than chemically bonded to the polymer. Their potential carcinogenic and endocrine disrupting properties can affect human reproduction (Fu et al., 2010; Li et al., 2016). Diethyl, di-n-butyl and bis-(2-ethylhexyl) phthalates (DEP, DnBP and DEHP) were analyzed in current study. The annual average concentration of phthalates was 510 ± 230 ng m⁻³ (165-1520 ng m⁻³) (Table 1). They showed higher concentration in pre-monsoon (Fig. S4). Fu et al., (2010) reported similar concentrations of phthalates (the total of DEP, DnBP, dimethyl, diisobutyl and di-(2-ethylhexyl) phthalates) in an Indian urban site, with 553 ng m⁻³ (295-857 ng m⁻³) in summer and 303 ng m⁻³ (175-598 ng m⁻³) in winter. In South Asia, large quantities of municipal solid wastes containing plastic products are generally disposed of in open landfills. The open burning of plastics along with other municipal solid waste is common in Nepal, and thus can also release numerous phthalate compounds into the atmosphere.

3.6 SOA tracers

Emissions of volatile organic compounds from vegetation (VOCs) into the atmosphere, especially isoprene, monoterpenes and sesquiterpenes occurs in large amounts. These biogenic VOCs (B-VOCs) are crucial precursors of biogenic SOA. Globally, the emissions of B-VOCs (1150 TgC/yr) consisting of 44% isoprene and 11% monoterpenes are much higher than emissions of anthropogenic VOCs (only 110 TgC/yr) (Piccot et al., 1992;Guenther et al., 1995). It should be noted, besides biogenic emissions, combustion of biomass and fossil fuel also contribute to the isoprene, monoterpenes and sesquiterpenes (Jathar et al., 2014;Sarkar et al., 2016;Sarkar et al., 2017). The measurements of gaseous VOCs in winter (December 2012 to February 2013) air in the Kathmandu Valley during SusKat-ABC campaign also showed high levels of isoprene and it was attributed (at least during high isoprene periods) mostly to biogenic emissions (Sarkar et al., 2016;Sarkar et al., 2017). It is difficult to appropriately quantify the fractions of biogenic and anthropogenic emissions of these compounds, based on ambient measurement of these species alone without measurement of biomass burning tracers such as acetonitrile and furan. The budget of isoprene emissions (500 Tg y^{-1}) on a global scale is dominated by vegetation (Guenther et al., 2006). Therefore, in our study, we considered the oxidation products of isoprene, monoterpenes and sesquiterpenes as the tracers of biogenic emissions and attribute their main source as biogenic emissions. This may lead to some overestimation of their contributions to SOA formation.

3.6.1 Isoprene SOA tracers

Six isoprene-SOA (I-SOA) tracers were identified in the Bode aerosols: 2-methylglyceric acid (2-MGA), two diastereoisomeric 2-methyltetrols (2-methylthreitol and 2-methylerythritol, 2-MTLs) and three C5-alkene triols (cis-2-methyl-1,3,4-trihydroxy-1-butene, trans-2-methyl-1,3,4-trihydroxy-1-butene, and 3-methyl-2,3,4-trihydroxy-1-butene). Their total concentrations ranged from $38.8\text{-}444 \text{ ng m}^{-3}$ ($174 \pm 86.2 \text{ ng m}^{-3}$), with the maximum ($236 \pm 87.2 \text{ ng m}^{-3}$) in the monsoon (Table 1). During the post-monsoon and pre-monsoon, their concentrations were similar, and a little lower than those during the monsoon (Fig. 5d) while being the lowest during winter. Their seasonal variation was in agreement with the ambient temperature (Fig S1a), which can influence the isoprene emissions and the photochemical processes (Shen et al., 2015;Wang et al., 2008). The annual average concentration was higher than the

425 urban sites reported from Beijing (44.3 ng m^{-3}) and Kunming (108 ng m^{-3}) (Ding et al., 2016a), even one
to two orders of magnitude higher than that from global oceans and the Arctic (Hu et al., 2013). Among
I-SOA tracers, 2-MTLs were the major components ($51.0\% \pm 10.5\%$) (Fig. 6), with an annual average of
94.4 \pm 58.9 ng m^{-3} (ranging from 10.9 to 270 ng m^{-3}). Strong correlations were exhibited between the two
430 isomers during all the seasons (Fig. S5a), implying that they formed through the similar pathway (Shen
et al., 2015;Fu et al., 2010). The daily concentration of 2-MGA ranged from 7.10-79.0 ng m^{-3} with an
annual average of 34.2 \pm 14.8 ng m^{-3} . For C5-alkene triols, the average concentration was 45.0 \pm 29.4 ng m^{-3} .
They positively correlated with 2-MTLs (Fig. S5b), indicating they were also the oxidation products of
isoprene under low-NO_x conditions.

According to the reaction chamber results from Surratt et al. (2010), the formation mechanism of 2-
435 MGA remarkably differs from 2-MTLs. 2-MGA is produced under high-NO_x condition while 2-MTLs
are mainly formed under low-NO_x or NO_x-free conditions. The formation of 2-MGA can be enhanced
under lower RH condition, while it is opposite for 2-MTLs (Zhang et al., 2011). During the monsoon
season, due to the conducive conditions of high temperature, high RH (>70 %) (Fig. S1b), high solar
radiation and fully-grown plants, the isoprene emissions was large. In addition, NO_x during this season
440 was much lower than other seasons. Therefore, 2-MGA/2-MTLs ratios exhibited the lowest values
(0.20 \pm 0.08) in the aerosol samples during this wet season (Fig. 7). In contrast, 2-MGA/2-MTLs ratios
increased up to 0.95 in winter, owing to the lowest temperature and RH of the whole year (Fig. 7) and the
higher NO_x concentration in the Kathmandu Valley (Kondo et al., 2005;Kiros et al., 2016). NO_x from
anthropogenic sources (industry, transportation, biomass burning in the houses as well as in the field) and
445 meteorological conditions with reduced mixing layer heights in winter would also favor the formation of
2-MGA and subsequently increase the 2-MGA/2-MTLs ratio.

Positive correlations were observed between 2-MGA, SO₄²⁻ and NO₃⁻ (Fig. 8). Budisulistiorini et al.
(2017) investigated that the concentrations of B-SOA could significantly increase as the aerosol acidity
enhances based on the laboratory simulations and field observations. The remarkable influence of I-SOA
450 by SO₄²⁻ might be explained by the concerted nucleophilic addition to the key intermediates in the gas
phase (e.g., isoprene epoxydiols), which is the rate-determining step in SOA formation (Xu et al., 2015;Li
et al., 2018). Li et al. (2018) reported that SO₄²⁻ plays a significant part in promoting aqueous phase

oxidation of I-SOA tracers. There may be the similar effect of NO_3^- on the SOA formation that needs further research. Therefore, the increase of SO_4^{2-} and NO_3^- could effectively facilitate the ring-opening reaction of isoprene epoxydiols and the SOA formation. Thus, the higher 2-MGA in the Kathmandu Valley may be due to the abundant SO_4^{2-} and NO_3^- during pre-monsoon season when most of the brick kilns (more than 100) are operational. Our finding confirmed that the anthropogenic pollutants such as SO_2 and NO_x can be conducive to accelerating the oxidation of B-VOCs and enhancing the ambient concentrations of B-SOA.

3.6.2 Terpene SOA tracers

Besides isoprene tracers, we also measured four monoterpene oxidation products (M-SOA tracers), including cis-pinonic (PNA), pinic (PA), 3-hydroxyglutaric (3-HGA) and 3-methyl-1,2,3-butanetricarboxylic acids (MBTCA) (Claeys et al., 2007). They are produced through the photooxidation of monoterpenes with ozone and hydroxyl radical (Iinuma et al., 2004). The annual average concentration of the total M-SOA tracers was $59.3 \pm 24.6 \text{ ng m}^{-3}$ (Table 1). The concentration of M-SOA tracers was higher than those reported in the previous studies from the urban site in Kunming (annual average: $44.1 \pm 38.8 \text{ ng m}^{-3}$) (Ding et al., 2016b), three cities (Ohio, Michigan and California) in North America (summer: $30.4\text{--}60.6 \text{ ng m}^{-3}$) (Stone et al., 2009) and a forest site in Hyytiälä, Europe (summer: $15.1\text{--}33.3 \text{ ng m}^{-3}$) (Kourtchev et al., 2005).

For the seasonal variation, relatively high concentrations of M-SOA tracers occurred during pre-monsoon and post-monsoon seasons (Fig. 5e, f, g, h, and i). Interestingly, there is intensive biomass burning in Kathmandu Valley twice a year (forest fires and crop-residue fires during April to May, and crop-residue fires during October to November) discussed in section 3.3.1 and 3.4, which may have been associated with high concentrations of M-SOA tracers. During the fires, substantial amounts of aerosols and VOCs including isoprene and monoterpenes would generate, which can enhance the levels of B-SOA tracers (Ding et al., 2013; Yan et al., 2008; Jathar et al., 2014). Good correlations were observed between levoglucosan and the higher generation oxidation products (e.g., 3-HGA and MBTCA, $R^2=0.32$ and $R^2=0.53$ respectively) in the Bode aerosols (Fig. S6). The forests in the Kathmandu Valley consist of broad-leaved evergreen mixed forest of *Schima castanopsis* at the base, oak-laurel forest in the middle

480 (1800 to 2400 m a. s. l.) and oak forest at the top, while the conifer tree species *Pinus roxiburghii* (Khote Salla) and *Pinus wallichiana* (Gobre Salla) are also found (Department of Plant Resources, 2015; Sarkar et al., 2016). Monoterpenes were chiefly emitted from needle leaf trees (coniferous trees) (Kang et al., 2018). Therefore, it suggested that biomass-burning activities have had a significant influence on the atmospheric composition over Kathmandu Valley, especially for SOA tracers.

485 Sesquiterpenes are also among the biogenic SOA (B-SOA) precursors emitted from trees, which have been observed in the troposphere in a variety of field studies. Concentrations of β -caryophyllenic acid found in the Bode aerosols ranged from 1.53 to 18.5 ng m⁻³ with an average of 6.31 \pm 3.86 ng m⁻³. It shared the similar seasonal variation with M-SOA tracers and positively correlated with them, indicating the possible common emission pattern.

490 **3.6.3 Aromatic SOA tracer**

Anthropogenic SOA is also an important OC source. 2,3-dihydroxy-4-oxopentanoic acid, DHOPA is a tracer of anthropogenic SOA from aromatics. In this study, the level of DHOPA was higher during winter and pre-monsoon while lower during monsoon season (Fig. 5). Though the major emissions of aromatics come from solvent and fossil fuel use, biomass burning is also considered as an possible source in some sites of the world (Shen et al., 2015). There was a good correlation between DHOPA and levoglucosan (Fig. 9), especially during pre-monsoon with the value of R² at 0.73. This indicated that biomass-burning emission is an important source of DHOPA at the Bode site.

3.7 Estimation of the contributions of different sources to OC

As discussed above, both the primary and secondary sources have influence on OC in the atmospheric aerosols of the Kathmandu Valley. In this part, we will apply the tracer-based methods to estimate the contributions of different sources to OC. It should be noted here that tracer methods can provide a reasonable estimation, but uncertainties are introduced considering the site differences and the lack of representative source profiles for the given study location. The contribution evaluated from each source to OC in the current study is still inferable.

505 **3.7.1 Biomass burning-derived OC**

The ratio of levoglucosan to OC (Lev/OC) detected in source samples has been used in a wide range to quantitatively estimate the contribution from biomass burning to OC (Stone et al., 2012;Wan et al., 2017;Zhang et al., 2015), although the ratios in the BB source emissions vary among different types of biomass fuels and burning conditions (Mochida et al., 2010). Andreae and Merlet (2001) reported an average of 8.14% with a range from 8.0% to 8.2 % for Lev/OC from the burning sites of biofuel, crop residues, savanna, tropical forests, and so on. Zhang et al. (2007) reported that Lev/OC ratios ranged from 5.4%-11.8% (an average of 8.27%) in the combustion aerosols from cereal straw (wheat, corn and rice). Sheesley et al. (2003) reported an average of 7.94% of levoglucosan from the combustion of biomass (including rice straw, biomass briquettes, dried cow-dung patties, etc.) indigenous to South Asia. However, the ratio obtained from the hardwood burning in fireplaces and stoves in the US was 14%, which was applied at the background sites in Europe (Fine et al., 2004). Stone et al. (2012) used the Lev/OC ratio of $12\% \pm 0.2\%$ during the burning of acacia wood at Godavari in the Kathmandu Valley for the CBM profile source apportionment. The mean value of Lev/OC value of biomass burning from main biomass types was 10.1%. In this study, we choose the mostly used values of 8.14% for biomass burning estimation (Graham et al., 2002;Fu et al., 2014;Ho et al., 2014;Sang et al., 2011;Zhu et al., 2016;Mkoma et al., 2013). In addition, we also calculated the uncertainties of using different ratios (see Table S3). The diagnostic ratios among molecular tracers and OC (e.g., Lev/OC) from direct emissions are critical for more precise results. It's meaningful to understand the emission characteristics for individual OC emission categories, as well as in different locations, especially in South Asia.

Figures 10 and 11 present the monthly concentration variations of BB-OC and contribution of BB-OC to OC, respectively. Our estimation exhibited that BB-OC contributed $24.9 \pm 10.4\%$ (ranging from 6.32% to 61.5%) to OC throughout the year in Bode aerosols (Fig. 11a). This was higher than the study in Lumbini in Nepal ($19.8 \pm 19.4\%$) (Wan et al., 2017), and nearly twice of the BB-OC contribution to OC reported in Hong Kong (6.5%–11%) (13.1 %) and the Pearl River Delta in China (Sang et al., 2011;Ho et al., 2014). Moreover, the contribution of BB-OC to OC in current study maximized in the post-monsoon ($36.3 \pm 10.4\%$), higher than that in the pre-monsoon ($28.5 \pm 10.3\%$) and winter ($27.9 \pm 8.63\%$). These results indicate that biomass burning severely affect the air quality in the Kathmandu Valley, especially during the post-monsoon season. Similarly, Stone et al. (2010) reported $21 \pm 2\%$ of OC in $PM_{2.5}$ from Godavari

rural site in the outskirts of the Kathmandu Valley during 2006, was also attributed to the primary biomass
535 burning sources.

3.7.2 Plant-debris-OC and fungal spore-derived OC

Primary biological aerosol particle (PBAP) has been identified as an important source using tracers
(section 3.3.2). They are likely to have a big contribution to the aerosols in Bode. In order to reveal how
much they are contributing to organic aerosols, “total” plant debris was calculated based on glucose
540 following the equation (Puxbaum and Tenze-Kunit, 2003):

$$\text{Cellulose } (\mu\text{g}) = \text{D-Glucose } (\mu\text{g}) \times \text{GF} \times (1/\text{SY});$$

$$\text{Plant debris} = 2 \times \text{cellulose}.$$

where GF (0.90) is the glucose/cellulose weight conversion factor and SY is the saccharification
yield (0.717).

545 For OC fraction derived from fungal spores, it was estimated using mannitol levels according to the
studies from Bauer et al. (2008) and Holden et al. (2011), i.e., there was 1.7 pg mannitol and 13 pg OC
per spore.

As shown in Fig. 11a, fungal spore-derived OC and plant-debris-OC annually contribute to
3.15±2.86% and 1.42±1.03% of OC, respectively. The contributions were both higher in the monsoon
550 season, with 5.85±2.50% for fungal spore-derived OC and 2.29±0.79% for plant-debris-OC to OC,
respectively (Fig. 11c). During winter, the contributions were the lowest due to the inactive vegetation.
There are also some similar results from the literatures. For example, Zhu et al. (2016) reported the
contribution of plant debris to OC was 5.6% in nighttime and 4.6% in daytime respectively from aerosols
in a mid-latitude forest. Szidat et al. (2006) reported the plant debris contributed to 3.2% of OC during
555 summer in urban aerosols collected in Zurich, Switzerland. Fungal-spore-derived OC was the biggest
contributor to total OC of 3.1 % (0.03 %–19.8 %) in marine aerosols collected over the East China Sea
during 18 May to 12 June 2014 (Kang et al., 2018). The study in the aerosols of Brazil urban site showed
the mean contributions of fungal aerosol to OC was 8% (Emygdio et al., 2018). Liang et al. (2017)
reported the contributions of fungal spores to OC of 1.2 ± 0.7% and 3.5 ± 3.7% in aerosols from an urban
560 site and a rural site respectively during an entire year in Beijing, China. All above strengthened the

importance of plant-debris and fungal spores to the aerosol burden in the atmosphere.

3.7.3 Biogenic SOC and anthropogenic toluene SOC

Biogenic secondary organic carbon (B-SOC) and anthropogenic aromatic SOC (A-SOC) from the oxidation of isoprene, monoterpenes, sesquiterpene and toluene were assessed using the tracer-based method proposed by Kleindienst et al. (2007). This method has been applied successfully in many aerosol studies (Fu et al., 2010; Shen et al., 2015; Ding et al., 2016a). The mass fraction of tracer compounds in SOC (FSOC) for an individual precursor was calculate based on the smog chamber simulations. The calculation formula as following:

$$FSOC = \frac{\sum_i [tri]}{[SOC]}$$

where [tri] is the concentration of tracer i and [SOC] is the concentration of SOC. The conversion factors of FSOC were 0.155 ± 0.039 , 0.231 ± 0.111 , 0.0230 ± 0.0046 and $0.0079 \pm 0.0026 \mu\text{g } \mu\text{gC}^{-1}$ for isoprene, monoterpenes, sesquiterpene and toluene, respectively (Kleindienst et al., 2007).

The total calculated concentrations of B-SOC ranged from 0.41 to $2.77 \mu\text{g m}^{-3}$ with an annual average concentration of $1.36 \pm 0.49 \mu\text{g m}^{-3}$, a higher concentration of $1.43 \pm 0.48 \mu\text{g m}^{-3}$ in monsoon and lower concentration of $0.86 \pm 0.20 \mu\text{g m}^{-3}$ in winter (Fig 10g). The B-SOC/OC showed a higher average percentage of $10.1\% \pm 3.34\%$ in the monsoon season (Fig 11c), suggesting that B-SOC was an important sources to OC at the Bode during this period. During post-monsoon, B-SOC/OC declined to 5.36% (Fig 11d). The B-SOC/OC showed the lowest value of $1.52\% \pm 0.70\%$ in winter (Fig 11e), indicating that B-SOC had minor contributions to elevated OC in winter. The annual average concentration of A-SOC was $2.45 \pm 1.45 \mu\text{g m}^{-3}$, which is higher than the B-SOC. The highest A-SOC concentration was obtained in winter ($3.27 \pm 1.25 \mu\text{g m}^{-3}$) (Fig.10h). A-SOC was the second most important contributor to OC after BB-OC. It is not only derived from increased fossil fuel combustion and the subsequent oxidation, but also from biomass burning emissions.

In total, SOC (including B-SOC and A-SOC) reconstructed using the formula above in this section was $3.81 \pm 1.63 \mu\text{g m}^{-3}$, accounting for $15.0\% \pm 8.99\%$ of OC.

3.7.4 Possible sources of the unidentified OC

On the whole, biomass burning contributed one-fourth ($24.9\% \pm 10.4\%$) of the OC in Bode, followed by A-SOC ($8.82\% \pm 5.55\%$), B-SOC ($6.19\% \pm 4.49\%$), fungal-spores ($3.15\% \pm 2.86\%$) and plant-debris ($1.42\% \pm 1.03\%$) (Fig. 11a). Nevertheless, there is still part of OC (55.5%) that we were not able to be attributed to any specific sources based on the tracers analyzed in current study. There are partly uncertainties caused by the organic tracer analyses (estimation of measurement uncertainty was shown in Table S2). Furthermore, fossil fuel combustion and soil dust could be also notable fractions of OC in Bode aerosols. Additionally, low molecular weight (LMW) dicarboxylic acids from both primary and secondary sources also constitute a significant fraction of atmospheric organic aerosols (Kawamura and Bikkina, 2016). Humic-like substances and amines are another source of OC, but not well studied (Wu et al., 2018; Laskin et al., 2015). Therefore, the possible contributions of the unidentified OC (55.5%) from various sectors need further investigation, which is better to comprehensively understand the sources of South Asian aerosols and will be very useful for the targeted pollution control measures in this region.

4. Summary and conclusions

Field measurements of primary and secondary organic compounds in aerosols were conducted in Bode, a semi-urban site of the Kathmandu Valley, Nepal, from April 2013 to April 2014. A distinctive seasonality was observed for various aerosol species. Higher concentrations of OC, EC, anhydrosugars, phenolic compounds and resin acid were observed during the winter and pre-monsoon seasons, while their concentrations were lower during the monsoon season. Levoglucosan was the most abundant species among the individually identified tracers with an average concentration of 788 ng m^{-3} . We observed the highest abundances of monosaccharides during the pre-monsoon season and of sugar alcohols during the monsoon season, and lower levels in winter because of the reduced plant activities. I-SOA tracers represented a majority among B-SOA tracers with a maximum in the monsoon. The seasonal variation of M-SOA tracers was controlled by monoterpenes emission and biomass burning. DHOPA exhibited higher concentrations during the winter and pre-monsoon season.

The likely OC sources were further evaluated for their contributions to observed total OC using tracer-based methods. Biomass burning contributed a major fraction (24.9%) to OC in Bode, followed by

A-SOC (8.8%), B-SOC (6.2%), fungal spores (3.2%) and plant debris (1.4%). The highest contribution of BB-OC, 36.3%, occurred during post-monsoon season. A-SOC, B-SOC, fungal spores and plant debris all made larger contributions during the monsoon. The higher BB-OC and the A-SOC contributions imply that some biomass burning and anthropogenic components are widespread in the Kathmandu Valley and thus represent the main contributors affecting the regional air quality in the Kathmandu Valley region.

The present study clearly shows that the chemical constituents and sources of OC strongly vary with seasons, as a result of diverse air pollution sources in the valley across four seasons. The heavy biomass burning and the subsequent oxidant emissions are anticipated to cause larger contributions of B-SOC to OC. Understanding OC's climate impacts is a frontier area of research, because there is still a large uncertainty in the estimation of OC radiative forcing. Our study implies that since biomass burning is a major source of ambient OC, the fraction of OC that absorbs light (referred to as brown carbon) and also acts as cloud condensation nuclei, needs to be further studied in order to better understand radiative effects of OC on regional climate change. The current source contribution estimates from the tracer-based methods do not accurately evaluate the large temporal variations from all kinds of sources. Contributions from other sectors (ca 55.5%), including low molecular weight dicarboxylic acids (Kawamura and Sakaguchi, 1999; Kawamura and Bikkina, 2016), need further investigation to better understand the atmospheric aerosols from both urban and rural sources such as the Kathmandu Valley and other sites in the Himalayan foothills and the Indo-Gangetic Plain regions. These observations of the severe air pollution, particularly the particular matter pollution, provide valuable support for air pollution control measures, especially in determining which sources and sectors to first focus on the Kathmandu Valley and the surrounding region, in order to reduce the air pollution from being severe to become much cleaner in the near future. In addition, the current study based on the molecular level-source apportionment of OC in heavy polluted region from South Asia provides a much more specific quantification of source estimation for OC, which is different from previous studies based on the bulk carbonaceous aerosol using radiocarbon (^{14}C) measurements, PMF and CBM.

Data availability. Raw data are archived at the Institute of Tibetan Plateau Research, Chinese Academy

of Sciences, and are available on request by contacting the corresponding author.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments

645 The authors gratefully appreciate Shyam Kumar Newar and Bhogendra Kathayat for their assistance
in the sample collections, the staffs at Bode site in the Kathmandu Valley, and all individuals and groups
who participated in the SusKat-ABC field campaigns. We would like to thank senior scientist Karl Epton
Yttri from Norwegian Institute for Air Research for the helpful suggestions to reply the comments from
the referees. This study was supported by the Strategic Priority Research Program of Chinese Academy
650 of Sciences, Pan-Third Pole Environment Study for a Green Silk Road (Pan-TPE) (XDA20040501), the
National Natural Science Foundation of China (41522103, 41421061 and 41630754) and China
Postdoctoral Science Foundation (2018M630210). The coauthors from the IASS gratefully acknowledge
funding from the federal ministry of education and research (BMBF) and the Brandenburg state ministry
of science, research and culture (MWFK).

655 References

- Department of Transport Management (DoTM).: D. o. T. M.: Annual report of Ministry of Labor and
Transport Management, Government of Nepal, 2015.
- Alam, K., Mukhtar, A., Shahid, I., Blaschke, T., Majid, H., Rahman, S., Khan, R., and Rahman, N.: Source
apportionment and characterization of particulate matter (PM10) in urban environment of Lahore,
660 Aerosol and Air Quality Research, 14, 1851-1861, 2014.
- Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, Global
Biogeochemical Cycles, 15, 955-966, 10.1029/2000gb001382, 2001.
- Bauer, H., Claeys, M., Vermeylen, R., Schueller, E., Weinke, G., Berger, A., and Puxbaum, H.: Arabitol
and mannitol as tracers for the quantification of airborne fungal spores, Atmospheric Environment,
665 42, 588-593, 10.1016/j.atmosenv.2007.10.013, 2008.
- Bhardwaj, P., Naja, M., Rupakheti, M., Panday, A. K., Kumar, R., Mahata, K., Lal, S., Chandola, H. C.,
and Lawrence, M. G.: Variations in surface ozone and carbon monoxide in the Kathmandu Valley
and surrounding broader regions during SusKat-ABC field campaign: Role of local and regional
sources, Atmospheric Chemistry and Physics in review, 2017.
- 670 Bisht, D., Dumka, U., Kaskaoutis, D., Pipal, A., Srivastava, A., Soni, V., Attri, S., Sateesh, M., and Tiwari,
S.: Carbonaceous aerosols and pollutants over Delhi urban environment: temporal evolution,
source apportionment and radiative forcing, Science of the Total Environment, 521, 431-445, 2015.

- Boreddy, S. K. R., Kawamura, K., and Tachibana, E.: Long-term (2001-2013) observations of water-soluble dicarboxylic acids and related compounds over the western North Pacific: trends, seasonality and source apportionment, *Scientific Reports*, 7, 10.1038/s41598-017-08745-w, 2017.
- 675 Budisulistiorini, S. H., Nenes, A., Carlton, A. G., Surratt, J. D., McNeill, V. F., and Pye, H. O. T.: Simulating Aqueous-Phase Isoprene-Epoxydiol (IEPOX) Secondary Organic Aerosol Production During the 2013 Southern Oxidant and Aerosol Study (SOAS), *Environmental Science & Technology*, 51, 5026-5034, 10.1021/acs.est.6b05750, 2017.
- 680 Cao, J. J., Lee, S. C., Chow, J. C., Watson, J. G., Ho, K. F., Zhang, R. J., Jin, Z. D., Shen, Z. X., Chen, G. C., Kang, Y. M., Zou, S. C., Zhang, L. Z., Qi, S. H., Dai, M. H., Cheng, Y., and Hu, K.: Spatial and seasonal distributions of carbonaceous aerosols over China, *Journal of Geophysical Research-Atmospheres*, 112, 10.1029/2006jd008205, 2007.
- Chen, J., Kawamura, K., Liu, C.-Q., and Fu, P.: Long-term observations of saccharides in remote marine aerosols from the western North Pacific: A comparison between 1990-1993 and 2006-2009 periods, *Atmospheric Environment*, 67, 448-458, 10.1016/j.atmosenv.2012.11.014, 2013.
- 685 Chen, P., Kang, S., Li, C., Rupakheti, M., Yan, F., Li, Q., Ji, Z., Zhang, Q., Luo, W., and Sillanpaa, M.: Characteristics and sources of polycyclic aromatic hydrocarbons in atmospheric aerosols in the Kathmandu Valley, Nepal, *Science of the Total Environment*, 538, 86-92, 10.1016/j.scitotenv.2015.08.006, 2015.
- 690 Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, 303, 1173-1176, 10.1126/science.1092805, 2004.
- Claeys, M., Szmigielski, R., Kourtschev, I., Van der Veken, P., Vermeylen, R., Maenhaut, W., Jaoui, M., Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., and Edney, E. O.: Hydroxydicarboxylic Acids: Markers for Secondary Organic Aerosol from the Photooxidation of α -Pinene, *Environmental Science & Technology*, 41, 1628-1634, 10.1021/es0620181, 2007.
- 695 Cowie, G. L., and Hedges, J. I.: Carbohydrate sources in a coastal marine environment, *Geochimica Et Cosmochimica Acta*, 48, 2075-2087, 10.1016/0016-7037(84)90388-0, 1984.
- 700 Department of Plant Resources, N.: Bulletin of Department of Plant Resources, Nepal 37, Ministry of Forests and Soil Conservation, Kathmandu, 1-122, 2015.
- Deshmukh, D. K., Kawamura, K., and Deb, M. K.: Dicarboxylic acids, ω -oxocarboxylic acids, α -dicarbonyls, WSOC, OC, EC, and inorganic ions in wintertime size-segregated aerosols from central India: Sources and formation processes, *Chemosphere*, 161, 27-42, 2016.
- 705 Ding, X., Wang, X., Xie, Z., Zhang, Z., and Sun, L.: Impacts of Siberian Biomass Burning on Organic Aerosols over the North Pacific Ocean and the Arctic: Primary and Secondary Organic Tracers, *Environmental Science & Technology*, 47, 3149-3157, 10.1021/es3037093, 2013.
- Ding, X., He, Q.-F., Shen, R.-Q., Yu, Q.-Q., Zhang, Y.-Q., Xin, J.-Y., Wen, T.-X., and Wang, X.-M.: Spatial and seasonal variations of isoprene secondary organic aerosol in China: Significant impact of biomass burning during winter, *Scientific Reports*, 6, 10.1038/srep20411, 2016a.
- 710 Ding, X., Zhang, Y.-Q., He, Q.-F., Yu, Q.-Q., Shen, R.-Q., Zhang, Y., Zhang, Z., Lyu, S.-J., Hu, Q.-H., Wang, Y.-S., Li, L.-F., Song, W., and Wang, X.-M.: Spatial and seasonal variations of secondary organic aerosol from terpenoids over China, *Journal of Geophysical Research-Atmospheres*, 121, 14661-14678, 10.1002/2016jd025467, 2016b.

- 715 Emygdio, A. P. M., Andrade, M. d. F., Gonçalves, F. L. T., Engling, G., Zanetti, R. H. d. S., and Kumar, P.: Biomarkers as indicators of fungal biomass in the atmosphere of São Paulo, Brazil, *Science of The Total Environment*, 612, 809-821, 2018.
- Engling, G., Lee, J. J., Tsai, Y.-W., Lung, S.-C. C., Chou, C. C. K., and Chan, C.-Y.: Size-Resolved Anhydrosugar Composition in Smoke Aerosol from Controlled Field Burning of Rice Straw, *Aerosol Science and Technology*, 43, 662-672, 10.1080/02786820902825113, 2009.
- 720 Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the wood stove combustion of prevalent United States tree species, *Environmental Engineering Science*, 21, 705-721, 10.1089/ees.2004.21.705, 2004.
- Fu, P., Kawamura, K., Kobayashi, M., and Simoneit, B. R. T.: Seasonal variations of sugars in atmospheric particulate matter from Gosan, Jeju Island: Significant contributions of airborne pollen and Asian dust in spring, *Atmospheric Environment*, 55, 234-239, 10.1016/j.atmosenv.2012.02.061, 2012.
- Fu, P., Kawamura, K., Chen, J., and Miyazaki, Y.: Secondary production of organic aerosols from biogenic VOCs over Mt. Fuji, Japan, *Environmental Science & Technology*, 48, 8491-8497, 2014.
- Fu, P. Q., Kawamura, K., Pavuluri, C. M., Swaminathan, T., and Chen, J.: Molecular characterization of urban organic aerosol in tropical India: contributions of primary emissions and secondary photooxidation, *Atmospheric Chemistry and Physics*, 10, 2663-2689, 2010.
- 730 Fujii, Y., Kawamoto, H., Tohno, S., Oda, M., Iriana, W., and Lestari, P.: Characteristics of carbonaceous aerosols emitted from peatland fire in Riau, Sumatra, Indonesia (2): Identification of organic compounds, *Atmospheric Environment*, 110, 1-7, 2015.
- 735 George, S. K., Nair, P. R., Parameswaran, K., Jacob, S., and Abraham, A.: Seasonal trends in chemical composition of aerosols at a tropical coastal site of India, *Journal of Geophysical Research-Atmospheres*, 113, 10.1029/2007jd009507, 2008.
- Giri, D., Murthy, K. V., Adhikary, P. R., and Khanal, S. N.: Ambient air quality of Kathmandu valley as reflected by atmospheric particulate matter concentrations (PM10), *International Journal of Environmental Science & Technology*, 3, 403-410, 10.1007/bf03325949, 2006.
- 740 Graham, B., Mayol-Bracero, O. L., Guyon, P., Roberts, G. C., Decesari, S., Facchini, M. C., Artaxo, P., Maenhaut, W., Koll, P., and Andreae, M. O.: Water-soluble organic compounds in biomass burning aerosols over Amazonia - 1. Characterization by NMR and GC-MS, *Journal of Geophysical Research-Atmospheres*, 107, 10.1029/2001jd000336, 2002.
- 745 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *Journal of Geophysical Research-Atmospheres*, 100, 8873-8892, 10.1029/94jd02950, 1995.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics* 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.
- 750 Gupta, A. K., Karar, K., and Srivastava, A.: Chemical mass balance source apportionment of PM10 and TSP in residential and industrial sites of an urban region of Kolkata, India, *Journal of Hazardous Materials*, 142, 279-287, 10.1016/j.jhazmat.2006.08.013, 2007.
- 755 Ho, K. F., Engling, G., Ho, S. S. H., Huang, R., Lai, S., Cao, J., and Lee, S. C.: Seasonal variations of anhydrosugars in PM2.5 in the Pearl River Delta Region, China, *Tellus Series B-Chemical and*

Physical Meteorology, 66, 10.3402/tellusb.v66.22577, 2014.

- 760 Holden, A. S., Sullivan, A. P., Munchak, L. A., Kreidenweis, S. M., Schichtel, B. A., Malm, W. C., and Collett, J. L., Jr.: Determining contributions of biomass burning and other sources to fine particle contemporary carbon in the western United States, *Atmospheric Environment*, 45, 1986-1993, 10.1016/j.atmosenv.2011.01.021, 2011.
- Hu, Q.-H., Xie, Z.-Q., Wang, X.-M., Kang, H., He, Q.-F., and Zhang, P.: Secondary organic aerosols over oceans via oxidation of isoprene and monoterpenes from Arctic to Antarctic, *Scientific Reports*, 3, 10.1038/srep02280, 2013.
- 765 Iinuma, Y., Böge, O., Gnauk, T., and Herrmann, H.: Aerosol-chamber study of the α -pinene/O₃ reaction: influence of particle acidity on aerosol yields and products, *Atmospheric Environment*, 38, 761-773, 2004.
- Jathar, S. H., Gordon, T. D., Hennigan, C. J., Pye, H. O. T., Pouliot, G., Adams, P. J., Donahue, N. M., and Robinson, A. L.: Unspeciated organic emissions from combustion sources and their influence on the secondary organic aerosol budget in the United States, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 10473-10478, 10.1073/pnas.1323740111, 2014.
- 770 Kang, M., Fu, P., Kawamura, K., Yang, F., Zhang, H., Zang, Z., Ren, H., Ren, L., Zhao, Y., Sun, Y., and Wang, Z.: Characterization of biogenic primary and secondary organic aerosols in the marine atmosphere over the East China Sea, *Atmospheric Chemistry and Physics*, 18, 13947-13967, 10.5194/acp-18-13947-2018, 2018.
- 775 Kawamura, K., and Sakaguchi, F.: Molecular distributions of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics, *Journal of Geophysical Research-Atmospheres*, 104, 3501-3509, 10.1029/1998jd100041, 1999.
- 780 Kawamura, K., and Bikkina, S.: A review of dicarboxylic acids and related compounds in atmospheric aerosols: Molecular distributions, sources and transformation, *Atmospheric Research*, 170, 140-160, 2016.
- Kim, B. M., Park, J.-S., Kim, S.-W., Kim, H., Jeon, H., Cho, C., Kim, J.-H., Hong, S., Rupakheti, M., Panday, A. K., Park, R. J., Hong, J., and Yoon, S.-C.: Source apportionment of PM₁₀ mass and particulate carbon in the Kathmandu Valley, Nepal, *Atmospheric Environment*, 123, 190-199, 10.1016/j.atmosenv.2015.10.082, 2015.
- 785 Kiros, F., Shakya, K. M., Rupakheti, M., Regmi, R. P., Maharjan, R., Byanju, R. M., Naja, M., Mahata, K., Kathayat, B., and Peltier, R. E.: Variability of Anthropogenic Gases: Nitrogen Oxides, Sulfur Dioxide, Ozone and Ammonia in Kathmandu Valley, Nepal, *Aerosol and Air Quality Research*, 16, 3088-3101, 10.4209/aaqr.2015.07.0445, 2016.
- 790 Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenber, J. H., Lewis, C. W., Bhawe, P. V., and Edney, E. O.: Estimates of the contributions of biogenic and anthropogenic hydrocarbons to secondary organic aerosol at a southeastern US location, *Atmospheric Environment*, 41, 8288-8300, 10.1016/j.atmosenv.2007.06.045, 2007.
- 795 Kondo, A., Kaga, A., Imamura, K., Inoue, Y., Sugisawa, M., Shrestha, M. L., and Sapkota, B.: Investigation of air pollution concentration in Kathmandu valley during winter season, *Journal of Environmental Sciences*, 17, 1008-1013, 2005.
- Kourtchev, I., Ruuskanen, T., Maenhaut, W., Kulmala, M., and Claeys, M.: Observation of 2-methyltetrols

- and related photo-oxidation products of isoprene in boreal forest aerosols from Hyytiälä, Finland, *Atmospheric Chemistry and Physics*, 5, 2761-2770, 10.5194/acp-5-2761-2005, 2005.
- 800 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chemical Reviews*, 115, 4335, 2015.
- Lawrence, M. G., and Lelieveld, J.: Atmospheric pollutant outflow from southern Asia: a review, *Atmospheric Chemistry and Physics*, 10, 11017-11096, 10.5194/acp-10-11017-2010, 2010.
- 805 Li, J., Wang, G., Aggarwal, S. G., Huang, Y., Ren, Y., Zhou, B., Singh, K., Gupta, P. K., Cao, J., and Zhang, R.: Comparison of abundances, compositions and sources of elements, inorganic ions and organic compounds in atmospheric aerosols from Xi'an and New Delhi, two megacities in China and India, *Science of the Total Environment*, 476, 485-495, 10.1016/j.scitotenv.2014.01.011, 2014.
- 810 Li, J., Wang, G., Ren, Y., Wang, J., Wu, C., Han, Y., Zhang, L., Cheng, C., and Meng, J.: Identification of chemical compositions and sources of atmospheric aerosols in Xi'an, inland China during two types of haze events, *Science of the Total Environment*, 566, 230-237, 10.1016/j.scitotenv.2016.05.057, 2016.
- 815 Li, J., Wang, G., Wu, C., Cao, C., Ren, Y., Wang, J., Li, J., Cao, J., Zeng, L., and Zhu, T.: Characterization of isoprene-derived secondary organic aerosols at a rural site in North China Plain with implications for anthropogenic pollution effects, *Scientific Reports*, 8, 10.1038/s41598-017-18983-7, 2018.
- Liang, L., Engling, G., Du, Z., Cheng, Y., Duan, F., Liu, X., and He, K.: Seasonal variations and source estimation of saccharides in atmospheric particulate matter in Beijing, China, *Chemosphere*, 150, 365-377, 10.1016/j.chemosphere.2016.02.002, 2016.
- 820 Liang, L., Engling, G., Du, Z., Duan, F., Cheng, Y., Liu, X., and He, K.: Contribution of fungal spores to organic carbon in ambient aerosols in Beijing, China, *Atmospheric Pollution Research*, 8, 351-358, 10.1016/j.apr.2016.10.007, 2017.
- 825 Mahata, K. S., Panday, A. K., Rupakheti, M., Singh, A., Naja, M., and Lawrence, M. G.: Seasonal and diurnal variations in methane and carbon dioxide in the Kathmandu Valley in the foothills of the central Himalayas, *Atmospheric Chemistry and Physics* 17, 12573-12596, 10.5194/acp-17-12573-2017, 2017a.
- 830 Mahata, K. S., Rupakheti, M., Panday, A. K., Bhardwaj, P., Naja, M., Singh, A., Mues, A., Cristofanelli, P., Pudasainee, D., Bonasoni, P., and Lawrence, M. G.: Observation and analysis of spatio-temporal characteristics of surface ozone and carbon monoxide at multiple sites in the Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics Discussion*, 2017, 1-42, 10.5194/acp-2017-709, 2017b.
- Miyazaki, Y., Jung, J., Fu, P., Mizoguchi, Y., Yamanoi, K., and Kawamura, K.: Evidence of formation of submicrometer water-soluble organic aerosols at a deciduous forest site in northern Japan in summer, *Journal of Geophysical Research-Atmospheres*, 117, 10.1029/2012jd018250, 2012.
- 835 Mkoma, S. L., Kawamura, K., and Fu, P. Q.: Contributions of biomass/biofuel burning to organic aerosols and particulate matter in Tanzania, East Africa, based on analyses of ionic species, organic and elemental carbon, levoglucosan and mannosan, *Atmospheric Chemistry and Physics*, 13, 10325-10338, 10.5194/acp-13-10325-2013, 2013.
- 840 Mochida, M., Kawamura, K., Fu, P. Q., and Takemura, T.: Seasonal variation of levoglucosan in aerosols over the western North Pacific and its assessment as a biomass-burning tracer, *Atmospheric*

Environment, 44, 3511-3518, 10.1016/j.atmosenv.2010.06.017, 2010.

Mues, A., Rupakheti, M., Münkel, C., Lauer, A., Bozem, H., Hoor, P., Butler, T., and Lawrence, M. G.: Investigation of the mixing layer height derived from ceilometer measurements in the Kathmandu Valley and implications for local air quality, *Atmospheric Chemistry and Physics* 17, 8157-8176, 10.5194/acp-17-8157-2017, 2017.

845

Mues, A., Lauer, A., Lupascu, A., Rupakheti, M., Kuik, F., and Lawrence, M. G.: WRF and WRF-Chem v3.5.1 simulations of meteorology and black carbon concentrations in the Kathmandu Valley, *Geoscientific Model Development*, 11, 2067-2091, 10.5194/gmd-11-2067-2018, 2018.

Muzzini, E., and Aparicio, G.: *Urban Growth and Spatial Transition in Nepal: An Initial Assessment.*, World Bank Publications., 2013.

850

Myers-Pigg, A. N., Griffin, R. J., Louchouart, P., Norwood, M. J., Sterne, A., and Cevik, B. K.: Signatures of Biomass Burning Aerosols in the Plume of a Saltmarsh Wildfire in South Texas, *Environmental Science & Technology*, 50, 9308-9314, 10.1021/acs.est.6b02132, 2016.

Myriokefalitakis, S., Fanourgakis, G., and Kanakidou, M.: The Contribution of Bioaerosols to the Organic Carbon Budget of the Atmosphere, in: *Perspectives on Atmospheric Sciences*, Cham, 2017, 845-851.

855

Oanh, N. T. K., Hang, N. T., Aungsiri, T., Worrarat, T., and Danutawat, T.: Characterization of Particulate Matter Measured at Remote Forest Site in Relation to Local and Distant Contributing Sources, *Aerosol and Air Quality Research*, 16, 2671-2684, 10.4209/aaqr.2015.12.0677, 2016.

860

Panday, A. K., and Prinn, R. G.: Diurnal cycle of air pollution in the Kathmandu Valley, Nepal: Observations, *Journal of Geophysical Research-Atmospheres*, 114, 10.1029/2008jd009777, 2009.

Piccot, S. D., Watson, J. J., and Jones, J. W.: A global inventory of volatile organic compound emissions from anthropogenic sources, *Journal of Geophysical Research-Atmospheres*, 97, 9897-9912, 10.1029/92jd00682, 1992.

865

Pudasainee, D., Sapkota, B., Shrestha, M. L., Kaga, A., Kondo, A., and Inoue, Y.: Ground level ozone concentrations and its association with NO_x and meteorological parameters in Kathmandu valley, Nepal, *Atmospheric Environment*, 40, 8081-8087, 10.1016/j.atmosenv.2006.07.011, 2006.

Putero, D., Cristofanelli, P., Marinoni, A., Adhikary, B., Duchì, R., Shrestha, S. D., Verza, G. P., Landi, T. C., Calzolari, F., Busetto, M., Agrillo, G., Biancofiore, F., Di Carlo, P., Panday, A. K., Rupakheti, M., and Bonasoni, P.: Seasonal variation of ozone and black carbon observed at Paknajol, an urban site in the Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics*, 15, 13957-13971, 10.5194/acp-15-13957-2015, 2015.

870

Puxbaum, H., and Tenze-Kunit, M.: Size distribution and seasonal variation of atmospheric cellulose, *Atmospheric Environment*, 37, 3693-3699, 10.1016/s1352-2310(03)00451-5, 2003.

875

Rajput, N., and Lakhani, A.: Measurements of polycyclic aromatic hydrocarbons in an urban atmosphere of Agra, India, *Atmosfera*, 23, 165-183, 2010.

Ram, K., Sarin, M. M., and Hegde, P.: Long-term record of aerosol optical properties and chemical composition from a high-altitude site (Manora Peak) in Central Himalaya, *Atmospheric Chemistry and Physics*, 10, 11791-11803, 10.5194/acp-10-11791-2010, 2010.

880

Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J. T., Washington, W. M., Fu, Q., Sikka, D. R., and Wild, M.: Atmospheric brown clouds: Impacts on South Asian climate and hydrological cycle, *Proceedings of the National Academy of Sciences of the United States of America*, 102,

5326-5333, 10.1073/pnas.0500656102, 2005.

- 885 Ramanathan, V., Li, F., Ramana, M. V., Praveen, P. S., Kim, D., Corrigan, C. E., Nguyen, H., Stone, E.
A., Schauer, J. J., Carmichael, G. R., Adhikary, B., and Yoon, S. C.: Atmospheric brown clouds:
Hemispherical and regional variations in long-range transport, absorption, and radiative forcing,
Journal of Geophysical Research-Atmospheres, 112, 10.1029/2006jd008124, 2007.
- 890 Regmi, R. P., Kitada, T., and Kurata, G.: Numerical simulation of late wintertime local flows in
Kathmandu valley, Nepal: Implication for air pollution transport, Journal of Applied Meteorology,
42, 389-403, 2003.
- Reisen, F., Meyer, C. P., and Keywood, M. D.: Impact of biomass burning sources on seasonal aerosol air
quality, Atmospheric Environment, 67, 437-447, 2013.
- 895 Rupakheti, D., Adhikary, B., Praveen, P. S., Rupakheti, M., Kang, S., Mahata, K. S., Naja, M., Zhang, Q.,
Panday, A. K., and Lawrence, M. G.: Pre-monsoon air quality over Lumbini, a world heritage site
along the Himalayan foothills, Atmospheric Chemistry and Physics, 17, 11041-11063,
10.5194/acp-17-11041-2017, 2017.
- 900 Rupakheti, M., Panday, A. K., Lawrence, M. G., Kim, S. W., Sinha, V., Kang, S. C., Naja, M., Park, J. S.,
Hoor, P., Holben, B., Sharma, R. K., Mues, A., Mahata, K. S., Bhardwaj, P., Sarkar, C., Rupakheti,
D., Regmi, R. P., and Gustafsson, Ö.: Air pollution in the Himalayan foothills: overview of the
SusKat-ABC international air pollution measurement campaign in Nepal, Atmospheric Chemistry
and Physics, in preparation, 2018.
- 905 Sang, X.-F., Chan, C.-Y., Engling, G., Chan, L.-Y., Wang, X.-M., Zhang, Y.-N., Shi, S., Zhang, Z.-S.,
Zhang, T., and Hu, M.: Levoglucosan enhancement in ambient aerosol during springtime transport
events of biomass burning smoke to Southeast China, Tellus Series B-Chemical and Physical
Meteorology, 63, 129-139, 10.1111/j.1600-0889.2010.00515.x, 2011.
- Sang, X., Zhang, Z., Chan, C., and Engling, G.: Source categories and contribution of biomass smoke to
organic aerosol over the southeastern Tibetan Plateau, Atmospheric Environment, 78, 113-123,
2013.
- 910 Sarkar, C., Sinha, V., Kumar, V., Rupakheti, M., Panday, A., Mahata, K. S., Rupakheti, D., Kathayat, B.,
and Lawrence, M. G.: Overview of VOC emissions and chemistry from PTR-TOF-MS
measurements during the SusKat-ABC campaign: high acetaldehyde, isoprene and isocyanic acid
in wintertime air of the Kathmandu Valley, Atmospheric Chemistry and Physics, 16, 3979-4003,
2016.
- 915 Sarkar, C., Sinha, V., Sinha, B., Panday, A. K., Rupakheti, M., and Lawrence, M. G.: Source
apportionment of NMVOCs in the Kathmandu Valley during the SusKat-ABC international field
campaign using positive matrix factorization, Atmospheric Chemistry and Physics, 17, 8129-8156,
10.5194/acp-17-8129-2017, 2017.
- 920 Shah, M. H., and Shaheen, N.: Annual and seasonal variations of trace metals in atmospheric suspended
particulate matter in Islamabad, Pakistan, Water, air, and soil pollution, 190, 13-25, 2008.
- Shakya, K. M., Ziemba, L. D., and Griffin, R. J.: Characteristics and Sources of Carbonaceous, Ionic, and
Isotopic Species of Wintertime Atmospheric Aerosols in Kathmandu Valley, Nepal, Aerosol and
Air Quality Research, 10, 219-U213, 10.4209/aaqr.2009.10.0068, 2010.
- Shakya, K. M., Rupakheti, M., Aryal, K., and Peltier, R. E.: Respiratory Effects of High Levels of
Particulate Exposure in a Cohort of Traffic Police in Kathmandu, Nepal, Journal of Occupational

- 925 & Environmental Medicine, 58, 1, 2016.
- Shakya, K. M., Rupakheti, M., Shahi, A., Maskey, R., Pradhan, B., Panday, A., Puppala, S. P., Lawrence, M., and Peltier, R. E.: Near-road sampling of PM_{2.5}, BC, and fine-particle chemical components in Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics*, 17, 6503-6516, 10.5194/acp-17-6503-2017, 2017.
- 930 Sharma, R. K., Bhattarai, B. K., Sapkota, B. K., Gewali, M. B., and Kjeldstad, B.: Black carbon aerosols variation in Kathmandu valley, Nepal, *Atmospheric Environment*, 63, 282-288, 10.1016/j.atmosenv.2012.09.023, 2012.
- Sheesley, R. J., Schauer, J. J., Chowdhury, Z., Cass, G. R., and Simoneit, B. R. T.: Characterization of organic aerosols emitted from the combustion of biomass indigenous to South Asia, *Journal of Geophysical Research-Atmospheres*, 108, 10.1029/2002jd002981, 2003.
- 935 Shen, R. Q., Ding, X., He, Q. F., Cong, Z. Y., Yu, Q. Q., and Wang, X. M.: Seasonal variation of secondary organic aerosol tracers in Central Tibetan Plateau, *Atmospheric Chemistry and Physics*, 15, 8781-8793, 10.5194/acp-15-8781-2015, 2015.
- Simoneit, B. R. T.: Biomass burning - A review of organic tracers for smoke from incomplete combustion, *Applied Geochemistry*, 17, 129-162, 10.1016/s0883-2927(01)00061-0, 2002.
- 940 Simoneit, B. R. T., Elias, V. O., Kobayashi, M., Kawamura, K., Rushdi, A. I., Medeiros, P. M., Rogge, W. F., and Didyk, B. M.: Sugars - Dominant water-soluble organic compounds in soils and characterization as tracers in atmospheric particulate matter, *Environmental Science & Technology*, 38, 5939-5949, 10.1021/es0403099, 2004.
- 945 Stockwell, C. E., Christian, T. J., Goetz, J. D., Jayarathne, T., Bhawe, P. V., Praveen, P. S., Adhikari, S., Maharjan, R., DeCarlo, P. F., Stone, E. A., Saikawa, E., Blake, D. R., Simpson, I. J., Yokelson, R. J., and Panday, A. K.: Nepal Ambient Monitoring and Source Testing Experiment (NAMA_{STE}): emissions of trace gases and light-absorbing carbon from wood and dung cooking fires, garbage and crop residue burning, brick kilns, and other sources, *Atmospheric Chemistry and Physics*, 16, 11043-11081, 10.5194/acp-16-11043-2016, 2016.
- 950 Stone, E. A., Zhou, J., Snyder, D. C., Rutter, A. P., Mieritz, M., and Schauer, J. J.: A Comparison of Summertime Secondary Organic Aerosol Source Contributions at Contrasting Urban Locations, *Environmental Science & Technology*, 43, 3448-3454, 10.1021/es8025209, 2009.
- 955 Stone, E. A., Schauer, J. J., Pradhan, B. B., Dangol, P. M., Habib, G., Venkataraman, C., and Ramanathan, V.: Characterization of emissions from South Asian biofuels and application to source apportionment of carbonaceous aerosol in the Himalayas, *Journal of Geophysical Research-Atmospheres*, 115, 10.1029/2009jd011881, 2010.
- Stone, E. A., Nguyen, T. T., Pradhan, B. B., and Dangol, P. M.: Assessment of biogenic secondary organic aerosol in the Himalayas, *Environmental Chemistry*, 9, 263-272, 10.1071/en12002, 2012.
- 960 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6640-6645, 10.1073/pnas.0911114107, 2010.
- 965 Szidat, S., Jenk, T. M., Synal, H.-A., Kalberer, M., Wacker, L., Hajdas, I., Kasper-Giebl, A., and Baltensperger, U.: Contributions of fossil fuel, biomass-burning, and biogenic emissions to carbonaceous aerosols in Zurich as traced by ¹⁴C, *Journal of Geophysical Research: Atmospheres*,

111, doi:10.1029/2005JD006590, 2006.

970 Tripathee, L., Kang, S., Rupakheti, D., Cong, Z., Zhang, Q., and Huang, J.: Chemical characteristics of soluble aerosols over the central Himalayas: insights into spatiotemporal variations and sources, *Environmental Science and Pollution Research*, 24, 24454-24472, 10.1007/s11356-017-0077-0, 2017.

Wan, E. C. H., and Yu, J. Z.: Analysis of sugars and sugar polyols in atmospheric aerosols by chloride attachment in liquid chromatography/negative ion electrospray mass spectrometry, *Environmental Science & Technology*, 41, 2459-2466, 10.1021/es062390g, 2007.

975 Wan, X., Kang, S., Wang, Y., Xin, J., Liu, B., Guo, Y., Wen, T., Zhang, G., and Cong, Z.: Size distribution of carbonaceous aerosols at a high-altitude site on the central Tibetan Plateau (Nam Co Station, 4730 m a.s.l.), *Atmospheric Research*, 153, 155-164, 10.1016/j.atmosres.2014.08.008, 2015.

980 Wan, X., Kang, S., Li, Q., Rupakheti, D., Zhang, Q., Guo, J., Chen, P., Tripathee, L., Rupakheti, M., Panday, A. K., Wang, W., Kawamura, K., Gao, S., Wu, G., and Cong, Z.: Organic molecular tracers in the atmospheric aerosols from Lumbini, Nepal, in the northern Indo-Gangetic Plain: influence of biomass burning, *Atmospheric Chemistry and Physics*, 17, 8867-8885, 10.5194/acp-17-8867-2017, 2017.

985 Wang, W., Wu, M. H., Li, L., Zhang, T., Liu, X. D., Feng, J. L., Li, H. J., Wang, Y. J., Sheng, G. Y., Claeys, M., and Fu, J. M.: Polar organic tracers in PM_{2.5} aerosols from forests in eastern China, *Atmospheric Chemistry and Physics*, 8, 7507-7518, 2008.

Wang, X., Gong, P., Sheng, J., Joswiak, D. R., and Yao, T.: Long-range atmospheric transport of particulate Polycyclic Aromatic Hydrocarbons and the incursion of aerosols to the southeast Tibetan Plateau, *Atmospheric Environment*, 115, 124-131, 10.1016/j.atmosenv.2015.04.050, 2015.

WHO: Air quality guidelines: global update 2005, World Health Organization, 2006.

990 Wu, G., Wan, X., Gao, S., Fu, P., Yin, Y., Li, G., Zhang, G., Kang, S., Ram, K., and Cong, Z.: Humic-Like Substances (HULIS) in Aerosols of Central Tibetan Plateau (Nam Co, 4730 m asl): Abundance, Light Absorption Properties, and Sources, *Environmental Science & Technology*, 52, 7203-7211, 10.1021/acs.est.8b01251, 2018.

995 Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H., Hering, S. V., de Gouw, J., Baumann, K., Lee, S.-H., Nenes, A., Weber, R. J., and Ng, N. L.: Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States, *Proceedings of the National Academy of Sciences of the United States of America*, 112, 10.1073/pnas.1512279112, 2015.

1000 Yan, B., Zheng, M., Hu, Y. T., Lee, S., Kim, H. K., and Russell, A. G.: Organic composition of carbonaceous aerosols in an aged prescribed fire plume, *Atmospheric Chemistry and Physics*, 8, 6381-6394, 10.5194/acp-8-6381-2008, 2008.

Yttri, K. E., Dye, C., and Kiss, G.: Ambient aerosol concentrations of sugars and sugar-alcohols at four different sites in Norway, *Atmospheric Chemistry and Physics*, 7, 4267-4279, 2007.

1005 Yue, S., Ren, H., Fan, S., Sun, Y., Wang, Z., and Fu, P.: Springtime precipitation effects on the abundance of fluorescent biological aerosol particles and HULIS in Beijing, *Scientific Reports*, 6, 10.1038/srep29618, 2016.

Zhang, H., Surratt, J. D., Lin, Y. H., Bapat, J., and Kamens, R. M.: Effect of relative humidity on SOA

1010 formation from isoprene/NO photooxidation: enhancement of 2-methylglyceric acid and its corresponding oligoesters under dry conditions, *Atmospheric Chemistry and Physics*, 11, 6411-6424, 10.5194/acp-11-6411-2011, 2011.

Zhang, T., Engling, G., Chan, C.-Y., Zhang, Y.-N., Zhang, Z.-S., Lin, M., Sang, X.-F., Li, Y. D., and Li, Y.-S.: Contribution of fungal spores to particulate matter in a tropical rainforest, *Environmental Research Letters*, 5, 024010, 2010.

1015 Zhang, Y., Shao, M., Zhang, Y., Zeng, L., He, L., Zhu, B., Wei, Y., and Zhu, X.: Source profiles of particulate organic matters emitted from cereal straw burnings, *Journal of Environmental Sciences*, 19, 167-175, 10.1016/s1001-0742(07)60027-8, 2007.

Zhang, Y. L., Huang, R. J., El Haddad, I., Ho, K. F., Cao, J. J., Han, Y., Zotter, P., Bozzetti, C., Daellenbach, K. R., Canonaco, F., Slowik, J. G., Salazar, G., Schwikowski, M., Schnelle-Kreis, J., Abbaszade, G., Zimmermann, R., Baltensperger, U., Prevot, A. S. H., and Szidat, S.: Fossil vs. non-fossil sources of fine carbonaceous aerosols in four Chinese cities during the extreme winter haze episode of 2013, *Atmospheric Chemistry and Physics*, 15, 1299-1312, 10.5194/acp-15-1299-2015, 2015.

1025 Zhu, C., Kawamura, K., and Kunwar, B.: Organic tracers of primary biological aerosol particles at subtropical Okinawa Island in the western North Pacific Rim, *Journal of Geophysical Research-Atmospheres*, 120, 5504-5523, 10.1002/2015jd023611, 2015.

Zhu, C., Kawamura, K., Fukuda, Y., Mochida, M., and Iwamoto, Y.: Fungal spores overwhelm biogenic organic aerosols in a midlatitudinal forest, *Atmospheric Chemistry and Physics*, 16, 7497-7506, 10.5194/acp-16-7497-2016, 2016.

1030

1031 Table 1 Concentrations of TSP, major ions, OC, EC ($\mu\text{g m}^{-3}$) and molecular tracers in
 1032 the aerosols from Bode, Kathmandu Valley (ng m^{-3}).

Compounds	Annual			Pre-monsoon			Monsoon			Post-monsoon			Winter		
	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD
TSP	256	213	166	381	366	171	120	107	52.4	225	254	71.6	353	348	68.5
SO ₄ ²⁻	10.8	6.15	9.83	17.2	16	7.49	4.1	2.6	4.04	4.81	4.24	2.38	27.3	24.1	6.79
NO ₃ ⁻	5.21	3.8	4.35	8.82	8.41	4.41	2.34	1.85	1.65	3.52	3.65	0.92	9.47	8.11	4.43
NH ₄ ⁺	5.92	3.46	6.16	8.57	6.71	5.65	1.99	1.31	2.58	3.72	3.65	1.48	17.5	15.6	3.06
Ca ²⁺	7.96	5.82	6.85	11.6	8.98	8.97	4.47	3.59	3.66	6.21	6.01	1.51	13.8	15.6	5.02
K ⁺	2.43	1.5	2.82	3.87	2.42	4.27	1.15	0.94	0.96	1.97	2.03	0.46	4.03	3.39	1.45
Cl ⁻	2.15	1.18	2.25	2.72	2.27	1.83	0.73	0.67	0.32	1.68	1.78	0.4	6.94	7.87	2.23
Na ⁺	3.28	2.93	1.58	3.15	1.68	2.38	3.3	3.61	1.18	2.8	2.84	0.12	4.21	4.28	0.49
Mg ²⁺	0.61	0.39	0.54	0.95	0.67	0.7	0.32	0.24	0.25	0.41	0.4	0.07	1.07	1.09	0.39
OC	38.7	24.2	32.7	59.4	46.9	37.9	14.6	14.7	3.76	31.8	32.9	12.7	62.8	53.6	20.6
EC	9.92	9.34	5.33	14.4	13.85	5.24	5.61	4.76	1.8	9.37	9.95	2.19	11	9.89	3.48
OC/EC	3.78	3.09	2.37	4.44	3.29	3.23	2.71	2.54	0.69	3.31	3.01	0.93	5.86	5.3	1.75
Anhydrosugars															
Levoglucosan	788	631	685	1214	900	705	204	188	60.1	863	836	252	1391	1120	535
Galactosan	44.4	33.8	40.3	68.5	51.4	42.7	13.2	13.1	5.55	34.8	33.8	10.9	85.6	72.4	32.8
Mannosan	50.6	34.5	45.1	71.8	61.8	42.3	16	16.7	5.5	39.5	39.6	12.5	116	106	45
Subtotal	883	688	765	1354	974	781	233	218	68.2	937	922	273	1592	1268	611
Monosaccharides															
Glucose	124	114	60	137	118	65.1	129	122	47	143	149	48.7	39.9	38.7	12.5
Fructose	58.2	53.2	28.3	68.4	66.9	29.1	57.1	48.2	26.8	58.2	54.7	21	27	22.5	10.2
Trehalose	40.8	35.6	22	48	53.1	22.4	40.2	35.5	22.8	38.9	40.1	12.5	20.1	16.8	9.17
Sucrose	48.3	40.3	27.4	64.7	55.5	31.6	38.7	38.7	11.9	56.6	45	27.1	18.4	16.6	6.37
Xylose	26.5	20.5	18.1	37.8	30.9	20	13.2	13.4	5.68	24.9	26.5	9.33	38.6	37.5	14.1
Subtotal	298	285	127	356	333	141	278	249	93.5	322	318	104	144	135	35.8
Sugar alcohols															
Mannitol	86.9	77	55.3	84.6	78.8	38.7	114	102	61.4	63.4	53.6	34.6	18.1	19.3	6.02
Arabitol	68.4	60.6	39.8	68.9	65.3	30.3	86.6	68.9	44.5	42.6	33.5	22.5	26.1	24.7	9.13
Sorbitol	14.2	12.7	8.02	13.1	12.6	5.96	17.9	15.9	9.31	13.1	13.2	4.65	5.82	4.97	2.72
Erythritol	43.1	36.5	28.8	35.8	34	15.4	56.9	47.5	33.1	48.6	39.1	31.8	12.4	10.2	7.6
Subtotal	213	192	126	202	198	84.8	275	245	143	168	151	80.1	62.5	61.8	19.7
Total sugars	1394	1206	813	1913	1570	919	787	727	249	1427	1367	327	1798	1462	651
Phenolic compounds and resin acid															
Vanillic acid	15.3	11.3	11.3	20.8	15.8	12.9	7.1	6.84	1.94	14.3	12.3	4.95	26.9	30.5	9.37
Syringic acid	17.1	11.6	13.7	23.7	17.1	15.6	7.82	7.52	2.67	13.7	12.5	4.54	32	37.8	12.3
p-Hydroxybenzoic acid	19.8	15.5	12.3	26.1	19.3	16.7	14.5	13	4.57	14.2	14.6	3.76	23.6	23.1	9.15
Dehydroabietic acid	13.8	12.6	6.19	16.3	15.4	6.52	10.3	9.7	2.32	10.9	9.3	3.1	21.4	20.1	6.76
Subtotal	66.1	49.7	41.4	86.9	67	49.2	39.7	39.6	9.22	53	47.5	15.6	104	112.8	36.2

1033

Compounds	Annual			Pre-monsoon			Monsoon			Post-monsoon			Winter		
	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD
Phthalate esters															
Diethyl (DEP)	16.6	15.6	8.41	19.4	17.5	11.6	15.4	15.5	4.42	14.9	13	6.25	12.9	9.3	6.49
Di-n-butyl (DnBP)	56.2	48.5	25.6	63.6	59.8	30.7	52.2	48.4	20.9	55.4	45.3	22.8	46.4	41.3	21.8
Bis-(2-ethylhexyl) (DEHP)	438	378	200	495	466	239	407	377	162	431	353	177	361	322	170
Subtotal	510	444	230	578	545	276	474	445	184	501	408	204	420	372	196
Isoprene tracers															
2-Methylglyceric acid	34.2	30.2	14.8	45.9	48.8	15.2	25.1	24.5	6.91	36.2	34.9	9.86	25	24.7	8.56
2-Methylthreitol	30.4	27.3	19.5	22	22.8	10	45	44.9	20.6	27.4	26.3	7.39	8.4	7.9	3.83
2-Methylerythritol	64.1	58.3	39.6	45	44.5	20.4	97.3	98.9	38	53	52	9.78	18.5	18.7	4.63
2-Methylterols ^a	94.4	84.3	58.9	67	68.1	30.4	142	142	58.3	80.5	78.3	17	27	26.6	8.39
C5-Alkene triols ^b	45	39.2	29.4	30.4	28.8	13.4	68.8	69.1	30	35.1	30.6	18	17.5	15.3	4.88
Subtotal	174	160	86.2	144	150	53.4	236	220	87.2	152	145	39.3	69.5	72.4	19.3
Monoterpene tracers															
cis-Pinonic acid	26	24.5	11.6	32.3	30	13.8	21	19.1	8.04	28	29.1	5.92	20.6	20	8.01
Pinic acid	11.9	10.8	4.48	11.4	10.4	3.78	12.7	12.3	5.07	13.7	15.6	5.25	8.83	8.67	0.92
3-Hydroxyglutaric acid	10.6	9.16	6.85	13.9	12.1	7.75	5.75	4.88	3.01	16	15.4	4.18	11.5	10	4.72
3-MBTCA ^c	10.8	9.53	7.36	16.9	15.1	7.32	5.14	4.3	3.12	11.9	9.61	3.63	9.3	8.89	3.75
Subtotal	59.3	55.6	24.6	74.6	72.1	28.3	44.6	42.6	14.4	69.5	65.7	12.6	50.2	49.6	10.8
Sesquiterpene tracer															
β -Caryophyllenic acid	6.43	5.35	3.93	8.61	7.15	4.29	3.66	2.85	2	8.35	7.2	3.02	7.16	6.6	2.76
Total B-SOA tracers ^d	136	234	50.4	133	229	43.9	156	275	53.9	131	217	26.2	75.6	129	14.2
Toluene tracer															
DHOPA ^e	19.4	16.5	11.5	22.8	19	15.5	15	13.4	5.69	17.3	18.2	5.94	25.8	23	9.84
Total SOA tracers ^f	259	249	94.7	250	242	84.9	299	286	101	247	243	48.2	153	162	29.4

1034 ^a Sum of 2-methylthreitol and 2-methylerythritol.

1035 ^b C5-Alkene triols: 3-Methyl-2,3,4-trihydroxy-1-butene, cis-2-Methyl-1,3,4-trihydroxy-1-butene and trans-2-Methyl-1,3,4-trihydroxy-1-butene.

1036 ^c 3-MBTCA: 3-methyl-1,2,3-butanetricarboxylic acid.

1037 ^d Sum of 2-methylglyceric acid, 2-methylterols, C5-Alkene triols, cis-pinonic acid, pinic acid, 3-hydroxyglutaric acid and 3-MBTCA.

1038 ^e DHOPA: 2,3-dihydroxy-4-oxopentanoic acid.

1039 ^f Sum of 2-methylglyceric acid, 2-methylterols, C5-Alkene triols, cis-pinonic acid, pinic acid, 3-hydroxyglutaric acid and 3-MBTC and 2,3-dihydroxy-4-oxopentanoic acid.

1040

1041

1042

1043

1044 Table 2 Linear correlation coefficients (R^2) among major ions and OC, EC in aerosols in Bode, Kathmandu Valley

	SO ₄ ²⁻	NO ₃ ⁻	NH ₄ ⁺	Ca ²⁺	nss-Ca ²⁺	Mg ²⁺	nss-Mg ²⁺	K ⁺	nss-K ⁺	Cl ⁻	Na ⁺	OC	EC
SO ₄ ²⁻	1.00												
NO ₃ ⁻	0.78**	1.00											
NH ₄ ⁺	0.87**	0.69**	1.00										
Ca ²⁺	0.48**	0.58**	0.62**	1.00									
nss-Ca ²⁺	0.48**	0.58**	0.62**	1.00**	1.00								
Mg ²⁺	0.61**	0.71**	0.69**	0.91**	0.91**	1.00							
nss-Mg ²⁺	0.65**	0.76**	0.70**	0.88**	0.88**	1.00**	1.00						
K ⁺	0.40**	0.59**	0.55**	0.78**	0.78**	0.85**	0.83**	1.00					
nss-K ⁺	0.40**	0.60**	0.55**	0.77**	0.77**	0.85**	0.83**	1.00**	1.00				
Cl ⁻	0.67**	0.58**	0.76**	0.41**	0.40**	0.48**	0.50**	0.37**	0.37**	1.00			
Na ⁺	0.05*	0.08*	0.23**	0.48**	0.48**	0.37**	0.28**	0.40**	0.39**	0.09*	1.00		
OC	0.59**	0.61**	0.57**	0.32**	0.32**	0.48**	0.53**	0.36**	0.36**	0.48**	0.01	1.00	
EC	0.35**	0.43**	0.24**	0.12**	0.12**	0.23**	0.27**	0.16**	0.16**	0.24**	0.02	0.36**	1.00

1045

1046 *: P<0.1.

1047 **: P<0.001.

1048 Table 3 Linear correlation coefficients (R^2) among monosaccharides and sugar alcohols in aerosols from Bode, Kathmandu Valley

	Glucose	Fructose	Sucrose	Trehalose	Xylose	Mannitol	Arabitol	Sorbitol	Erythritol
Glucose	1								
Fructose	0.77***	1							
Sucrose	0.55***	0.55***	1						
Trehalose	0.30***	0.23***	0.28***	1					
Xylose	0.001	0.06*	0.19***	0.05*	1				
Mannitol	0.51***	0.51***	0.23***	0.14***	0.002 ^a	1			
Arabitol	0.44***	0.50***	0.22***	0.16***	0.0005	0.77***	1		
Sorbitol	0.51***	0.53***	0.20***	0.13**	0.0002	0.83***	0.68***	1	
Erythritol	0.46***	0.40***	0.16***	0.13**	0.007	0.77***	0.62***	0.77***	1


1049

1050 a Negative values indicate negative correlations.

1051 * $p < 0.1$

1052 ** $p < 0.01$.

1053 *** $p < 0.001$.



1054

1055 **Fig. 1.** Location of measurement site: (a) Kathmandu Valley, (b) urban measurement site at Bode in
1056 Kathmandu Valley, (c) air pollution observed from the Bode site in the afternoon.

1057




Fig. 2. Monthly variations of TSP, OC, EC, OC/EC ratios at Bode site, Kathmandu Valley during April 2013-April 2014.





Fig. 3. Monthly variations of biomass burning tracers, monosaccharides and sugar alcohols at Bode site, Kathmandu Valley during April 2013-April 2014.




1065

1066


1067

Fig. 4. Correlations between (a) levoglucosan and OC, (b) levoglucosan and EC, (c) levoglucosan and nss-K⁺ in Bode aerosols during the sampling period (April 2013 to April 2014).




1068

1069 **Fig. 5.** Monthly variations of B-SOA tracers, total isoprene tracers, total monoterpene tracers, β -
 1070 caryophyllenic acid, total B-SOA tracers and DHPOA at Bode site, Kathmandu Valley during April
 1071 2013-April 2014.



1072


1073 **Fig. 6.** The percentage contributions of the isoprene SOA tracers to the total during different seasons in
 1074 the atmospheric aerosols from Kathmandu.



1075


1076

Fig. 7. Ratios of 2-MGA/2-MTLs during different seasons in Bode, Kathmandu.




1077

1078 **Fig. 8.** Concentration correlation between (a) 2-methylglyceric acid (2-MGA) and SO₄²⁻, (b) 2-
 1079 methylglyceric acid and NO₃⁻ in the aerosols from Bode, Kathmandu.




1080

1081 **Fig. 9.** Correlation between 2,3-dihydroxy-4-oxopentanoic acid (DHOPA) and levoglucosan in Bode
 1082 aerosols during the sampling period (April 2013 to April 2014).



1083

1084 **Fig. 10.** Monthly concentration variations of (a) BB-OC, (b) plant-debris-OC, (c) fungal-spore-OC, (d)
 1085 SOC-isoprene (I-SOC), (e) SOC-monoterpenes (M-SOC), (f) SOC-sequiterpene (SOC-C), (g) SOC-
 1086 isoprene+monoterpenes+sequiterpene (B-SOC), (h) SOC-toluene (A-SOC), and (i) total SOC that were
 1087 estimated using a tracer-based method at Bode site, Kathmandu Valley during April 2013-April 2014.



1088

1089

1090

1091

Fig. 11. Pie-charts showing contributions from different sources to organic carbon based on the estimation of tracer-method in Bode, Kathmandu Valley: (a) annual, (b) pre-monsoon, (c) monsoon, (d) post-monsoon and (e) winter.