

# Molecular characterization of organic aerosols in the Kathmandu Valley, Nepal: insights into primary and secondary sources

Xin Wan<sup>1,8</sup>, Shichang Kang<sup>2,7</sup>, Maheswar Rupakheti<sup>3,4</sup>, Qiangong Zhang<sup>1,7</sup>, Lekhendra Tripathi<sup>2</sup>,  
5 Junming Guo<sup>2</sup>, Pengfei Chen<sup>2</sup>, Dipesh Rupakheti<sup>2</sup>, Arnico K. Panday<sup>5</sup>, Mark G. Lawrence<sup>3</sup>, Kimitaka  
Kawamura<sup>6</sup>, Zhiyuan Cong<sup>1,7\*</sup>

<sup>1</sup> Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan  
Plateau Research, Chinese Academy of Sciences (CAS), Beijing 100101, China

10 <sup>2</sup> State Key Laboratory of Cryospheric Science, Northwest Institute of Eco-Environment and Resources,  
CAS, Lanzhou 730000, China

<sup>3</sup> Institute for Advanced Sustainability Studies (IASS), Potsdam 14467, Germany

<sup>4</sup> Himalayan Sustainability Institute (HIMSI), Kathmandu, Nepal

<sup>5</sup> International Centre for Integrated Mountain Development (ICIMOD), Kathmandu, Nepal

15 <sup>6</sup> Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan

<sup>7</sup> CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China

<sup>8</sup> University of Chinese Academy of Sciences, Beijing 100039, China

\*Corresponding author:

20 E-mail address: zhiyuancong@itpcas.ac.cn

## Abstract:

Organic atmospheric aerosols in the Hindu Kush-Himalayan-Tibetan Plateau region are still poorly characterized. To better understand the chemical characteristics and sources of organic aerosols in the foothill region of the central Himalaya, the atmospheric aerosol samples were collected in Bode, a suburban site of the Kathmandu Valley (KV) over a one-year period from April 2013 to April 2014. Various molecular tracers from specific sources of primary organic aerosols (POA) and secondary organic aerosols (SOA) were determined. Tracer-based estimation methods were employed to apportion contributions from each source. The concentrations of organic carbon (OC) and element carbon (EC) increased during winter with a maximum monthly average in January. Levoglucosan (a molecular tracer for biomass burning, BB) was observed as the dominant species among all the analyzed organic tracers and its annual average concentration was  $788 \pm 685 \text{ ng m}^{-3}$  (ranging from 58.8 to 3079  $\text{ng m}^{-3}$ ). Isoprene-SOA (I-SOA) represented a high concentration among biogenic-SOA tracers. For the seasonality, anhydrosugars, phenolic compounds, resin acid and aromatic SOA tracer showed similar seasonal variations with OC, and EC while monosaccharides, sugar alcohols and I-SOA tracers showed lower levels during winter. BB contributed a significant fraction to OC, averaging  $24.9 \pm 10.4\%$  during the whole year, and up to  $36.3 \pm 10.4\%$  in the post-monsoon season. On an annual average basis, anthropogenic toluene-derived secondary OC accounted for 8.8% and biogenic secondary OC contributed 6.2% to total OC. The annual contribution of fungal-spores to OC was 3.2% with the maximum during the monsoon season (5.9%). For plant debris, it accounted for 1.4% of OC during the monsoon. Therefore, OC is mainly associated with BB and other anthropogenic activity in the KV. Our findings are conducive to designing effective measures to mitigate the heavy air pollution and its impacts in the KV and surrounding area.

## 1. Introduction

South Asia, especially the Indo-Gangetic Plain (IGP) region, is a global air pollution hotspot. Atmospheric pollutants [e.g., organic carbon (OC), black carbon (BC), gaseous pollutants, etc.] from South Asia have been increasing during recent decades (Ramanathan et al., 2005; Muzzini and Aparicio, 2013; Lawrence and Lelieveld, 2010). While these pollutants are of concern locally near the emission sources, they can also, in short span of time, be transported to rural and remote regions over a long distance. This results in an annually recurring regional scale haze, referred to as atmospheric brown clouds (ABC), covers a large area from the Himalayan range to the Indian Ocean (Ramanathan et al., 2007). Until recently the emissions, types, levels, atmospheric transport and transformation, impacts and mitigation of various atmospheric pollutants were not well characterized in the vast mountain areas and the foothill region in South Asia. In this context, the international project of “A Sustainable Atmosphere for the Kathmandu Valley (SusKat)” was launched, aiming to comprehensively understand the causes of the severe air pollution in the region, and identifying appropriate solutions to reduce its impacts (Rupakheti et al., 2018). This paper presents analyses of samples collected as part of the SusKat field campaign.

The Kathmandu Valley (KV), the capital region of Nepal, is considered one of the most polluted regions over South Asia and the largest metropolitan region in the foothills of the Hindu Kush-Himalayas-Tibetan Plateau (HKHTP) region, facing rapid but unplanned urbanization, with the current population of approximately 4 million (Muzzini and Aparicio, 2013). Additionally, the bowl-shaped topography restrict the free flow of air, resulting in poor air quality (Pudasainee et al., 2006; Panday and Prinn, 2009). Giri et al. (2007) showed  $PM_{10}$  concentrations in Kathmandu were about 2-4 times higher than the guidelines prescribed by the World Health Organization (WHO) ( $PM_{10}$  24-hour mean:  $50 \mu g m^{-3}$ ) (WHO, 2006). More recently, Shakya et al. (2017) reported that daily mean  $PM_{2.5}$  concentrations at seven locations in the KV during 2014 were about 5 times higher than the WHO guidelines ( $PM_{2.5}$  24-hour mean:  $25 \mu g m^{-3}$ ) (WHO, 2006). Beside particulate matter, recent studies have pointed out that ground-level ozone ( $O_3$ ) is also of concern (Mahata et al., 2017b; Bhardwaj et al., 2017). Ozone levels at Pakanajol in the city center exceeded the WHO’s 8-hour maximum  $O_3$  guidelines of  $100 \mu g m^{-3}$  on 125 days in a year (Putero et al., 2015), while Mahata et al. (2017b) reported such exceedance was for nearly 3 months at Bode (where sampling for this study was conducted) and 6 months at Nagarkot, a hilltop site downwind

of the KV. The concentrations of acetonitrile and isoprene (precursor for both O<sub>3</sub> and secondary organic aerosol, SOA) investigated by Sarkar et al. (2016) in the KV were comparable with the highest reported elsewhere in the world. Air pollution is a clear threat to human health (leading to respiratory disease, cardiovascular disease, cancer, etc.), agricultural productivity and revenues from tourism in the KV and surrounding regions (Putero et al., 2015;Shakya et al., 2016).

Carbonaceous aerosols (OC and BC/EC, element carbon) are often a principal component of atmospheric aerosols and the atmospheric brown clouds (ABC) over South Asia (Wester et al., 2019). Sources and chemical transformations of OC are complicated, including primary emission sources (e.g. biomass/biofuel burning and fossil fuel combustion, plant debris, soil dust, etc.) and secondary formation of the oxidative products from precursor gases produced from both biogenic and anthropogenic compounds (Simoneit, 2002;Claeys et al., 2004;Fu et al., 2010). Although previous studies on organic aerosol characteristics in the KV and surrounding regions are limited, they demonstrated that OC was the main component of aerosols (Shakya et al., 2017;Kim et al., 2015). However, they have focused on a few aerosol species or a handful of organic compound classes (Chen et al., 2015;Sarkar et al., 2016). Only at a rural site, Godavari on the southern edge of the KV, analysis of organic aerosols at the molecular level has been reported (Stone et al., 2010;Stone et al., 2012). Overall, the composition and sources of OC are still poorly characterized.

Therefore, to overcome such research gaps, our study comprehensively investigates the organic molecular compositions of aerosols from the KV, including anhydrosugars, monosaccharides, sugar alcohols, phenolic compounds, resin acid, phthalic acid esters, and secondary organic aerosols produced from primary emission sources and secondary formation. We also studied the seasonal variation and molecular distribution to decipher their abundances, understand their predominant sources (primary vs. secondary), and to evaluate contributions of different sources to the carbonaceous aerosols in the suburban environment in the Himalayan foothills. Our current work enriches the database of the chemical characteristics of organic aerosols in South Asia, particularly in the HKHTP region.

## **2. Materials and methods**

### **2.1. Sampling site**

The KV is a round flat basin with the bottom of an elevation of approximately 1300 meters above sea level (a.s.l.) in the southern foothills of the Himalayas. It is encircled by green mountains (elevation: 1500 m to 2800 m a.s.l.) (Panday and Prinn, 2009). Our sampling was performed during April 2013 to April 2014 in Bode (27.67°N, 85.38°E, 1320 m a.s.l.), a suburban site to the east of Kathmandu city in the valley (Fig. 1). There are two major wind flows in the KV: (a) west to east, from Nagdhunga-Bhimdhunga mountain pass in the west to Nagarkot-Sanga mountain pass in the east, (b) south to north, from Bagmati River corridor to the northeast direction through the central-eastern part of the Valley. These two airflows meet around central-eastern part of the Valley and move eastward towards the Nagarkot-Sanga passes (Panday and Prinn, 2009). The Bode area receives these two air flows, and hence it is downwind of Kathmandu city and Lalitpur or Patan city located in southwest, west and northwest direction during the daytime, and Bhaktapur city located in east and southeast during nighttime (Bhardwaj et al., 2017; Mahata et al., 2017a; Rupakheti et al., 2018). In addition, it is situated in a residential area with urban buildings and houses scattered across agricultural fields with paddy, wheat, corn and vegetable farms. Some small industries (plastics, electronics, wood, fabrics, etc.) and Bhaktapur Industrial Estate are located in the south-eastern direction of Bode, as well as several brick kilns that use low quality coal during January to April (Sarkar et al., 2016). The Tribhuvan international airport in the west of Bode (~4 km) may have potential impacts when there is westerly wind. Approximately 1.5 and 7 km to the north, there are two reserve forests, consisting of a mix mainly broad-leaved deciduous trees and evergreen conifer trees (Department of Plant Resources, 2015). BC and O<sub>3</sub> measurements in the two major SusKat-ABC sites (Paknajol and Bode) in the Valley show similar levels (Putero et al., 2015; Mahata et al., 2017b). Therefore, Bode site can be taken as a representative site for the KV (Rupakheti et al., 2018).

## 2.2 Sample collection

The total suspended particulates (TSP) were continuously collected for 23 h (day and night time) every five days by a medium-volume sampler (model: KC-120H, Laoshan Co., China), which was installed on the rooftop of a building, approximately 20 m above ground. The flow rate was 100 L min<sup>-1</sup>. Overall, 82 aerosol samples were successfully obtained using 90 mm diameter quartz filters (Whatman PLC, UK). The filters were pre-baked (550 °C, 6 h) to remove all organic material and weighed by a microbalance (sensitivity: ±0.01 mg) before and after sampling. Before each weighing, they were

equilibrated at constant temperature ( $25\pm 3$  °C) and humidity ( $30\pm 5\%$ ) conditions for 24 h. Finally, the filters were preserved at 20 °C below zero until laboratory analysis. Field blanks (one blank filter each month) were also collected, briefly putting a filter onto the instrument without drawing air to assess potential contamination. There may be positive and negative artifacts during the sample handling/conditioning due to the adsorption/evaporation processes of organic aerosols (Li et al., 2018; Boreddy et al., 2017; Oanh et al., 2016). In a comparable study, Ding et al. (2013) reported the positive artifacts for OC and organic tracers were 10–20% and up to 16%, respectively.

### 2.3 Chemical Analysis

The aerosol samples were analyzed for major ions, OC, EC, and organic molecular tracers in the laboratory. Major ions ( $\text{Ca}^{2+}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$ ,  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$ ) were measured using an ion chromatography (Dionex, USA) with ICS-320 and ICS-1500 (Tripathy et al., 2017). The detection limit (LOD) of all the major ions was  $0.01 \mu\text{g m}^{-3}$ . They denoted less than 5% of the real sample concentrations in the field blank filters (Tripathy et al., 2017). Non-sea-salt  $\text{Ca}^{2+}$  (nss- $\text{Ca}^{2+}$ ) and  $\text{K}^+$  (nss- $\text{K}^+$ ) was estimated according to the method from George et al. (2008). OC and EC were determined by a thermal/optical reflectance analyzer (Model 2001A, USA) (Wan et al., 2015). The field blank OC ( $0.59\pm 0.13 \mu\text{g m}^{-3}$ ) were subtracted from the filter samples. EC in the field blank sample was  $0.00 \mu\text{g m}^{-3}$ .

Detailed analytical method of organic molecular tracers was described previously by Wan et al. (2017). A trace gas chromatography coupled to a PolarisQ mass spectrometry detector (GC-MS, Thermo Scientific) was used for analysis. Briefly, small filter portions ( $1.13\text{--}3.39 \text{ cm}^2$ ) were cut, spiked with appropriate amounts of methyl- $\beta$ -D-xylanopyranoside (MXP, 99%, Sigma) and  $D_3$ -malic acid (DMA, CDN isotopes, 99%) as internal recovery standards. Each filter portion was then extracted three times with a mixture of 20-ml dichloromethane/methanol (2:1, v/v) at room temperature for 30 minutes. The solvent extracts in total of 60 ml were combined and successively filtered through quartz wool, concentrated, dried over ultrapure nitrogen gas and then reacted with 50  $\mu\text{l}$  of 99% *N,O*-bis-(trimethylsilyl)trifluoroacetamide (BSTFA, with 1% trimethylsilyl chloride) and pyridine (v/v=2:1) at 70 °C for 3 h. n-Hexane of 150  $\mu\text{l}$  was added after derivatization. A TG-5MS ( $30 \text{ m} \times 0.25 \text{ mm} \times 0.25 \mu\text{m}$ ) was used for separation according to the GC temperature program. The oven temperature was initially

held at 50 °C for 2 min, increased to 120 °C at 30 °C min<sup>-1</sup>, then to 300 °C at 6 °C min<sup>-1</sup> and finally held  
155 for 16 min. The MS was operated in electron ionization (EI) mode at 70 eV with a scan range of 50-650  
Da.

For quantitative analysis, authentic standards processed as the samples above were used to establish  
the calibration curves. To quantitate the target compounds that were no available standards, surrogate  
compounds were used as following: erythritol for 2-methylglyceric acid (2-MGA), 2-methyltetrols (2-  
160 MTLs) and C<sub>5</sub>-alkene triols, cis-pinonic acid (PNA) for 3-hydroxyglutaric acid (3-HGA) and 3-methyl-  
1,2,3-butanetricarboxylic acid (MBTCA), pinic acid (PA) for β-caryophyllinic acid (β-CA), azelaic acid  
for 2,3-dihydroxy-4-oxopentanoic acid (DHOPA). Recoveries for target tracers and MXP (Table S1) were  
more than 75%. The exception was for malic acid (50.3%-90.5%) and cis-pinonic acid (60.2%-81.8%).  
The relative differences based on duplicate analysis were less than 15%. The method detection limits  
165 (MDL) were 0.04-0.13 ng m<sup>-3</sup> (Table S1). Our results were not corrected for the recoveries. No target  
compounds were detected in the field blank filters using the same procedure with the samples.

Table 1 shows a list of chemical species (OC, EC and major ions) and molecular markers (from the  
sources of biomass burning (BB), fungal spores, plant debris, plastic emission, biogenic and  
anthropogenic secondary formation) analyzed in this study. They include anhydrosugars (levoglucosan,  
170 mannosan and galactosan), primary saccharides (sucrose, trehalose, glucose and xylose) and sugar  
alcohols (mannitol, arabitol, sorbitol and erythritol), lignin and resin pyrolysis products (vanillic, syringic,  
*p*-hydroxybenzoic and dehydroabietic acids), phthalic acid esters, isoprene tracers (2-MGA, 2-MTLs of  
methylthreitol and 2-methylerythritol, C<sub>5</sub>-alkene triols of cis-2-methyl-1,3,4-trihydroxy-1-butene, 3-  
methyl-2,3,4-trihydroxy-1-butene and trans-2-methyl-1,3,4-trihydroxy-1-butene), monoterpene tracers  
175 (PNA, PA, 3-HGA and MBTCA), β-CA and DHOPA.

## 2.4 Estimation of measurement uncertainty

The application of surrogate standards for the quantification of most SOA tracers excluding PNA  
and PA) could cause additional errors to the measurements. Error in analyte measurement (EA) is  
propagated from the standard deviation of the field blank (EFB), error in spike recovery (ER) and error  
180 from surrogate quantification (EQ):

$$EA = \sqrt{EFB^2 + ER^2 + EQ^2}$$

EFB was 0 in this study due to the undetectable SOA tracers in the field blanks. To estimate the ER of tracers, the spike recoveries of surrogate standards within the range of 9.2% (erythritol) to 26.1% (PNA) were used. EQ was estimated by an empirical approach according to Stone et al. (2012). The relative error introduced by each carbon atom (En), oxygenated functional group (Ef) and alkenes (Ed) was estimated to be 15 %, 10% and 60%, respectively. Therefore, the EQ are calculated as:

$$EQ = E_n \Delta_n + E_f \Delta_f + E_d \Delta_d$$

where  $\Delta_n$ ,  $\Delta_f$  and  $\Delta_d$  are the difference between a surrogate and an analyte of carbon atom number, oxygen-containing functional group and alkene functionality, respectively.

Currently, EQ was calculated in the range of 15% (2-MTLs) to 120% ( $\beta$ -CA) and the estimation of EA ranged from 17.6% to 122.4%. The estimated uncertainties for the measurement of the SOA tracers is presented in Table S2.

## 2.5 Meteorological parameters

The meteorological parameters (e.g., temperature (T), relative humidity (RH), etc.) used in this study were derived from Tribhuvan International Airport ([www.wunderground.com](http://www.wunderground.com)), which was located west of Bode (approximately 4 km). Mixing layer height (MLH) data was measured with a Vaisala ceilometer at Bode site (Mues et al., 2017). The meteorology of KV and its surrounding regions is controlled by the South Asian monsoon circulations in the wet season (monsoon, June-September). Westerlies dominate the atmospheric circulation patterns during the dry seasons including pre-monsoon (March-May), post-monsoon (October-November) and winter (December-February) with limited precipitation (Pudasainee et al., 2006; Mues et al., 2017). Additionally, it is also influenced by local mountain valley circulation (Mues et al., 2018).

## 3. Results and discussion

A statistical concentration summary of major ions, OC, EC, and organic tracers identified in TSP samples collected at the Bode site is presented in Table 1. Tracers for six classes of organic compounds were detected: anhydrosugars, monosaccharides, sugar alcohols, phenolic compounds and resin acid, phthalic acid esters, and secondary organic aerosol (SOA) tracers.



### 3.1 Aerosol loadings

The TSP samples at Bode site exhibited daily mass concentrations from 32.0 to 723  $\mu\text{g m}^{-3}$  ( $255 \pm 167$   $\mu\text{g m}^{-3}$ ) during April 2013 to April 2014 (Table 1). Putero et al. (2015) reported  $195 \pm 83$   $\mu\text{g m}^{-3}$  of online  $\text{PM}_{2.5}$  concentration in Pakajol site (also one of SusKat-ABC sites), accounting for roughly 80% of TSP in our study. The TSP concentrations were comparable to those reported over other heavily polluted regions in South Asia, including Islamabad in Pakistan (Shah and Shaheen, 2008), Kolkata (Gupta et al., 2007) and Agra (Rajput and Lakhani, 2010) in India. Compared to the remote sites such as Lulang in the Tibetan Plateau (Wang et al., 2015) and Manora Peak in the central Himalaya (Ram et al., 2010), the TSP in Bode shows significantly higher mass concentrations. We found a clear seasonal variation in TSP mass concentrations (Fig. 2a), higher in pre-monsoon season ( $381 \pm 366$   $\mu\text{g m}^{-3}$ ) and winter ( $353 \pm 348$   $\mu\text{g m}^{-3}$ ) while lower in monsoon period ( $120 \pm 107$   $\mu\text{g m}^{-3}$ ) which was nearly half of post-monsoon season ( $256 \pm 213$   $\mu\text{g m}^{-3}$ ). It generally corresponded to the build-up of the atmospheric brown clouds (ABCs), which engulfed most of South Asia and the Northern Indian Ocean extending from November to May (Ramanathan et al., 2005).

Meteorological parameters may also affect the TSP concentrations. The highest TSP concentration observed during the pre-monsoon can be caused by the fugitive dust which is been blown up by strong wind and the absence of wet-precipitation (Fig S1a and c). The lower TSP concentration in the monsoon was likely related to increased precipitation (Fig S1c) after the onset of the South Asian monsoon. During this season, nearly 80% of the annual precipitation falls in the KV, which flushes out pollutants from the atmosphere (Tripathee et al., 2017; Wester et al., 2019). During winter, an inversion layer often occurs in the KV owing to its bowl-shaped topography (Pudasainee et al., 2006). The existence of an inversion layer with the lower temperature ( $12.0 \pm 2.41^\circ\text{C}$ ), wind speed ( $2.86 \pm 1.34$   $\text{km h}^{-1}$ ), and MLH ( $0.34 \pm 0.08$  km) (Mues et al., 2017) (Fig S1a, c and d) altogether reduced the pollution dispersion mechanism resulting in increased levels of pollutants close to the ground surface.

### 3.2 Major ions and OC/EC

Concentrations of eight major ions were measured in the aerosol samples from the Bode site. The total sum accounted for  $17.1\% \pm 8.5\%$  of annual average TSP mass. Sulfate ranked the highest among

235 them (annual mean:  $10.8 \pm 9.83 \mu\text{g m}^{-3}$ ), followed by  $\text{Ca}^{2+}$  ( $7.96 \pm 6.85 \mu\text{g m}^{-3}$ ),  $\text{NH}_4^+$  ( $5.92 \pm 6.16 \mu\text{g m}^{-3}$ ),  $\text{NO}_3^-$  ( $5.21 \pm 4.35 \mu\text{g m}^{-3}$ ),  $\text{Na}^+$  ( $3.28 \pm 1.58 \mu\text{g m}^{-3}$ ),  $\text{K}^+$  ( $2.43 \pm 2.82 \mu\text{g m}^{-3}$ ),  $\text{Cl}^-$  ( $2.15 \pm 2.25 \mu\text{g m}^{-3}$ ) and  $\text{Mg}^{2+}$  ( $0.61 \pm 0.54 \mu\text{g m}^{-3}$ ). On average, the combination of  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , i.e. the secondary inorganic aerosols, constituted more than half (51.3%) of the total ionic concentrations. The  $\text{Ca}^{2+}$  alone accounted for 22.1% of total ions.

240 Sulfate, ammonium and nitrate revealed a typical seasonality with the seasonally averaged concentrations ranked in the descending order of winter > pre-monsoon > post-monsoon > monsoon. This is consistent with the seasonal variation of the precursors  $\text{NO}_x$ ,  $\text{NO}_2$  and  $\text{SO}_2$ , which are mainly caused by automobile exhaust, household cooking, and emissions from brick kilns co-fired with biomass in the KV (Kiros et al., 2016; Wester et al., 2019). Currently, nearly 50% of the total motor vehicles in Nepal  
245 (approximately 2.33 million) run on the KV roads (DoTM, 2015; Mahata et al., 2017b). Diesel- or gasoline-powered generators (producing higher  $\text{NO}_x$  emissions) and garbage burning are other major pollution sources in Nepal during the sampling period, which can also emit much aerosol precursors (Stockwell et al., 2016).

Ions derived from crustal sources, such as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  are related to the local fugitive dust sources  
250 such as unpaved roads and construction activities (Ram et al., 2010). Interestingly, good correlations were found for  $\text{Ca}^{2+}$  and  $\text{SO}_4^{2-}$  ( $R^2 = 0.48$ ,  $P < 0.001$ ),  $\text{NO}_3^-$  ( $R^2 = 0.58$ ,  $P < 0.001$ ) and  $\text{NH}_4^+$  ( $R^2 = 0.62$ ,  $P < 0.001$ ), and for  $\text{Mg}^{2+}$  and  $\text{SO}_4^{2-}$  ( $R^2 = 0.61$ ,  $P < 0.001$ ),  $\text{NO}_3^-$  ( $R^2 = 0.71$ ,  $P < 0.001$ ) and  $\text{NH}_4^+$  ( $R^2 = 0.69$ ,  $P < 0.001$ ), respectively (Table 2), which hinted that dust may co-exist with  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$  and  $\text{NH}_4^+$  in the KV (Tripathee et al., 2017).

255 Carbonaceous aerosols (OC:  $38.7 \pm 32.7 \mu\text{g m}^{-3}$  and EC:  $9.92 \pm 5.33 \mu\text{g m}^{-3}$ ) accounted for  $19.2\% \pm 5.48\%$  of TSP mass through the sampling period at the Bode site, which was higher than that of the major ions. OC alone accounted for  $14.6\% \pm 4.81\%$  of the TSP mass. During winter and pre-monsoon seasons, OC and EC showed much higher concentrations than that during wet season (Fig. 2b and c). In this study, we found that the daily OC to EC mass ratios (OC/EC) varied from 0.77 to 15.8 (annual mean:  $3.78 \pm 2.73$ )  
260 and seasonal mean ratios of 4.44, 2.71, 3.31, and 5.86 during pre-monsoon, monsoon, post-monsoon, and winter seasons, respectively (Table 1 and Fig. 2d). The OC/EC ratios of more than 2.0 indicates the BB aerosols or the formation of secondary organic matter (Cao et al., 2007). Their influence and contribution

will be discussed in the following sections. The OC/EC ratios found in this study for the KV were similar to other sites in South Asia, like Lumbini ( $5.16 \pm 2.09$ , 2.41-10.03) (Wan et al., 2017), Delhi ( $5.86 \pm 0.99$ , 2.9-9.2) (Bisht et al., 2015) and Lahore ( $3.9 \pm 1.6$ , 1.5-7.2) (Alam et al., 2014).

### 3.3 Sugar compounds

#### 3.3.1 Anhydrosugars

Anhydrosugars of levoglucosan (1,6-anhydro- $\beta$ -D-glucopyranose) and its two isomers (mannosan and galactosan) have been used as ideal molecular tracers for BB emissions (Simoneit, 2002; Bhattarai et al., 2019). They are exclusively emitted from the combustion and pyrolysis of cellulose and hemicelluloses. In the current study, the annual average concentration of levoglucosan was  $788 \pm 685$  ng  $m^{-3}$ , ranging from 58.8 to 3079 ng  $m^{-3}$ , which was the dominant species of the total identified tracer compounds (Table 1).

For the seasonality, levoglucosan showed significantly higher levels during winter, pre-monsoon and post-monsoon seasons (Fig. 3a). Especially, higher concentrations were recorded in winter varying from 830 to 2395 ng  $m^{-3}$  (annual mean:  $1391 \pm 535$  ng  $m^{-3}$ ). It showed comparable levels with other sites in the world, which were badly affected by the BB emissions, e.g., New Delhi (1977 ng  $m^{-3}$ ) (Li et al., 2014) and Raipur (2180 ng  $m^{-3}$ ) in India (Deshmukh et al., 2016), Tasmania ( $4540 \pm 2480$  ng  $m^{-3}$ ) in Australia (Reisen et al., 2013) and Lumbini ( $1161 \pm 1347$ ) in Nepal (Wan et al., 2017). Our results were much higher than the aerosols (20-372 ng  $m^{-3}$ ) collected at a rural Godavari site (Stone et al., 2010), located on the southern edge of the KV during 2006. Good correlations exhibited between levoglucosan and OC ( $R^2=0.79$ ,  $P<0.001$ ), EC ( $R^2=0.42$ ,  $P<0.001$ ) and nss- $K^+$  ( $R^2=0.35$ ,  $P<0.01$ ) during the sampling campaign (Fig. 4). This indicates that OC and EC in KV's aerosols are strongly related to BB source (Kim et al., 2015).

The ratio of levoglucosan to mannosan (Lev/Man) has been applied to distinguish the possible categories of biomass burnt. Previously, higher Lev/Man ratios were reported for emissions from combustion of hardwood (ranging from 12.9 to 35.4 with an average of  $21.5 \pm 8.3$ ) and agricultural residues (range from 12.7 to 55.7 with an average of  $32.6 \pm 19.1$ ) (Sang et al., 2013; Bhattarai et al., 2019). For the softwood burning, the average ratio was  $4.0 \pm 1.0$  (ranging from 2.5 to 5.8). In current study, the

290 annual mean ratio of Lev/Man was  $16.3 \pm 5.96$  ranging from 9.13 to 33.1 with only 9 samples  $< 10$ . It can  
be inferred that the combustion of crop residues and hardwood is likely to be one of the major sources of  
atmospheric pollution in this region. A previous study also reported that the combustion of wood fuel for  
cooking and heating is common during wintertime in Nepal while there is much more crop residue  
combustion during both pre- and post-monsoon seasons (Stockwell et al., 2016). This is not only a local  
295 but also a regional phenomenon; for example, Bhardwaj et al. (2017) and Wan et al. (2017) pointed out  
emissions from crop residue burning during the pre- and post-monsoon periods from western India and  
eastern Pakistan impact the air quality in Nepal. Similarly, Rupakheti et al. (2017) also showed that the  
combustion of agricultural residues and forest fires over the northwestern IGP region are causes of the air  
pollution episodes over the foothills of the central Himalayas. In addition, brick kilns mainly operated  
300 during January-April, burned substantial quantities of low-grade coal, mixed crop wastes and firewood  
(Kim et al., 2015; Wester et al., 2019). Such emissions may also lead to the high levels of levoglucosan  
observed at Bode. We must point out that the incense burning in KV may also influence the levoglucosan  
concentration.

### 3.3.2 Monosaccharides

305 Primary biological aerosol particle (PBAP) tracers, commonly known also as bioaerosols, were  
analyzed in the Bode aerosol samples, including five monosaccharides of glucose, fructose, trehalose,  
sucrose and xylose. PBAP is derived from fungal spores, vegetative debris, pollen, bacteria and viruses.  
Most of them can cause adverse effects (e.g., allergenic reaction) on human health (Bauer et al.,  
2008; Myriokefalitakis et al., 2017).

310 In current work, total monosaccharides had an annual mean concentration of  $298 \pm 127$  ng m<sup>-3</sup>.  
Glucose was the predominant species among monosaccharides ( $124 \pm 60.0$  ng m<sup>-3</sup>), followed by fructose  
( $58.2 \pm 28.3$  ng m<sup>-3</sup>), sucrose ( $48.3 \pm 27.4$  ng m<sup>-3</sup>), trehalose ( $40.8 \pm 22.0$  ng m<sup>-3</sup>), and xylose ( $26.5 \pm 18.1$  ng  
m<sup>-3</sup>) (Table 1). Except xylose, they all presented higher concentrations in pre-monsoon period while being  
lower in winter (Fig 3h, i, j and k). There were significant linear correlations between glucose and fructose  
315 ( $R^2 = 0.77$ ,  $p < 0.001$ ), trehalose and glucose ( $R^2 = 0.30$ ,  $p < 0.001$ ), trehalose and fructose ( $R^2 = 0.23$ ,  
 $p < 0.001$ ), sucrose and glucose ( $R^2 = 0.55$ ,  $p < 0.001$ ), sucrose and fructose ( $R^2 = 0.55$ ,  $p < 0.001$ ), and  
sucrose and trehalose ( $R^2 = 0.28$ ,  $p < 0.001$ ) (Table 3). Therefore, the strong correlations indicated that they

were derived from common sources, e.g. from local forests in the KV during the period of high productivity of plants. In addition, the pollen produced from the flowering of local vegetation also largely contribute to glucose, fructose, trehalose, sucrose. The flowering of trees and crops peaks during the pre-monsoon season. The similar phenomenon was also reported in deciduous forests in northern Japan (Miyazaki et al., 2012).

Xylose has complex sources, including soils (Simoneit et al., 2004), microbiota (Wan and Yu, 2007), vegetation, bacteria (Cowie and Hedges, 1984) and biomass combustion (Zhu et al., 2015). It presents less abundant and only accounts for  $6.90\% \pm 8.32\%$  of the total PBAP tracers identified in the Bode aerosols. For the seasonal pattern, it is characterized by waxing in winter ( $47.4 \pm 24.3 \text{ ng m}^{-3}$ ) and waning in monsoon season ( $12.0 \pm 5.16 \text{ ng m}^{-3}$ ), which was different from the other primary monosaccharides (Table 1 and Fig. 2l). Close correlation between xylose and levoglucosan (the BB tracer) was observed in our study (Fig. S2,  $R^2=0.72$ ,  $p<0.001$ ), indicating that the emissions from the burning of biomass may largely contribute to xylose in Bode aerosols. A similar finding of xylose source (i.e. BB) was also proposed by Zhu et al. (2015).

### 3.3.3 Sugar alcohols

Total concentration of sugar alcohols (arabitol, sorbitol, erythritol and mannitol) was  $213 \pm 126 \text{ ng m}^{-3}$ , and thus lower than that of total monosaccharides (Table 1). Mannitol ( $86.9 \pm 55.3 \text{ ng m}^{-3}$ ) and arabitol ( $68.4 \pm 39.8 \text{ ng m}^{-3}$ ) showed higher concentrations, followed by erythritol ( $43.1 \pm 28.8 \text{ ng m}^{-3}$ ) and sorbitol ( $14.2 \pm 8.02 \text{ ng m}^{-3}$ ). All of them exhibited monsoon maxima ( $114 \pm 61.4 \text{ ng m}^{-3}$ ,  $86.6 \pm 44.5 \text{ ng m}^{-3}$ ,  $56.9 \pm 33.1 \text{ ng m}^{-3}$  and  $17.9 \pm 9.31 \text{ ng m}^{-3}$ , respectively) and winter minima ( $18.1 \pm 6.02 \text{ ng m}^{-3}$ ,  $26.1 \pm 9.13 \text{ ng m}^{-3}$ ,  $5.82 \pm 2.72 \text{ ng m}^{-3}$ ,  $12.4 \pm 7.60 \text{ ng m}^{-3}$ , respectively) (Table 1 and Fig. 3m, n, o and p). They also showed significant correlations with each other, implying their common sources (Zhu et al., 2015). Mannitol and arabitol have been mostly associated with fungal spores, along with vegetation and mature leaves and algae (Yttri et al., 2007; Myriokefalitakis et al., 2017). Recent studies proposed elevated concentrations of mannitol and arabitol were usually observed augmentation after rain events and also highly correlated with relative humidity (Yue et al., 2016; Zhu et al., 2016). Therefore, at Bode, sugar alcohols were likely emitted by plants in nearby forest and agriculture fields, especially during the

345 monsoon with the higher relative humidity (Fig. S1b). In addition, the higher temperatures (Fig. S1a) were conducive for more active microbial activities. Notably, the levels of PBAP discussed above were much higher than other sites in the world (Zhu et al., 2015; Liang et al., 2016; Yttri et al., 2007), indicating the strong fungal spore production in the KV during the wet season.

### 3.4 Phenolic compounds and resin acid

350 Phenolic compounds (e.g., vanillic, syringic and *p*-hydroxybenzoic acids) derived from lignin pyrolysis and resin acid (e.g., dehydroabietic acid) from burning of conifer plants can be also used as biomarkers for BB. Syringic acid is prevalent in hardwood smoke while vanillic acid is dominant both in softwood and hardwood smoke (Myers-Pigg et al., 2016; Wan et al., 2019). Herbaceous plant smoke primarily contains *p*-anisic acid and *p*-anisaldehyde (e.g., *p*-hydroxybenzoic acid and *p*-  
355 hydroxybenzaldehyde). Dehydroabietic acid is a dominant compound in the total lipid material from pine wood smoke. Therefore, three pyrolysis products of lignin (vanillic, syringic and *p*-hydroxybenzoic acids) and one pyrolysis of resin acid (dehydroabietic acid) were chosen as organic markers in this study.

*p*-Hydroxybenzoic acid ( $19.8 \pm 12.3 \text{ ng m}^{-3}$ ) was the predominant species, the second was dehydroabietic acid ( $13.8 \pm 6.19 \text{ ng m}^{-3}$ ), and then vanillic ( $15.3 \pm 11.3 \text{ ng m}^{-3}$ ) and syringic acids  
360 ( $17.1 \pm 13.7 \text{ ng m}^{-3}$ ) (Table 1). They exhibited maximum concentrations during winter and pre-monsoon periods, while decreased during wet season and then increased from post-monsoon period, which was consistent with the seasonal variation of levoglucosan (Fig 3d, e, f and g). There were also significant correlations of lignin and resin pyrolysis products with levoglucosan (cellulose pyrolysis products) (Fig. S3a, *p*-hydroxybenzoic acid and levoglucosan,  $R^2=0.72$ ,  $P<0.001$ ; Fig. S3b, vanillic acid and levoglucosan,  $R^2=0.86$ ,  $P<0.001$ ; Fig. S3c, syringic acid and levoglucosan,  $R^2=0.83$ ,  $P<0.001$ ; Fig. S3d, levoglucosan and dehydroabietic acid,  $R^2=0.63$ ,  $P<0.001$ ). Such a result also shows that there are various biomass combustion sources in the valley.

The concentration ratio of syringic to vanillic acids (Syr/Van) has recently been used to further discriminate the vegetation types burned (Fujii et al., 2015; Myers-Pigg et al., 2016; Wan et al., 2019).  
370 Previous study documented the Syr/Van ratios ranged from 0.1 to 2.44 for combustion of hardwood and herbaceous angiosperm, while it varied from 0.01 to 0.24 for burning softwood (Myers-Pigg et al., 2016). Regarding the aerosol samples from KV, the Syr/Van ratio was  $0.94 \pm 0.18$  of an annual average ranging

from 0.65 to 1.31, indicating that combustion of hardwood and herbaceous plant (including crop residues) are the most likely sources of BB in the Valley. This finding agrees with the results obtained from the Lev/Man ratios discussed in Sect. 3.3.1.

Besides the information revealed by anhydrosugars discussed in section 3.3.1, lignin and resin biomarkers further confirmed that BB emissions play a significant role in contributing to organic aerosols in the KV, particularly during winter and pre-monsoon periods.

### 3.5 Phthalic acid esters

Phthalates or phthalic acid esters are extensively utilized as key additives in the manufacture and processing of plastic products. As they are physically rather than chemically bonded to the polymer, they can be easily released into the environment. There are potential adverse effects on ecological system and human health due to their toxicity, e.g., carcinogenicity and endocrine disruption (Fu et al., 2010; Li et al., 2016). Diethyl (DEP), di-n-butyl (DnBP) and bis-(2-ethylhexyl) (DEHP) phthalates were analyzed in current study. The annual average concentration of phthalates was  $510 \pm 230 \text{ ng m}^{-3}$  ( $165\text{-}1520 \text{ ng m}^{-3}$ ) (Table 1). They showed higher concentration during pre-monsoon (Fig. S4). Similar concentrations of phthalates (the total of DEP, DnBP, dimethyl, diisobutyl and di-(2-ethylhexyl) phthalates) was also investigated in an Indian urban site, with  $553 \text{ ng m}^{-3}$  ( $295\text{-}857 \text{ ng m}^{-3}$ ) in May and  $303 \text{ ng m}^{-3}$  ( $175\text{-}598 \text{ ng m}^{-3}$ ) during winter (Fu et al., 2010). In South Asia, large quantities of municipal solid wastes containing plastic products are generally disposed of in open landfills. The open burning of plastics along with other municipal solid waste is common in Nepal, and thus can also release numerous phthalate compounds into the atmosphere.

### 3.6 SOA tracers

Emissions of volatile organic compounds from vegetation (VOCs) into the atmosphere, especially isoprene, monoterpenes and sesquiterpenes occurs in large amounts. These biogenic VOCs (B-VOCs) are crucial precursors of biogenic SOA. Globally, the emissions of B-VOCs ( $1150 \text{ TgC/yr}$ ) consisting of 44% isoprene and 11% monoterpenes are much higher than emissions of anthropogenic VOCs (only  $110 \text{ TgC/yr}$ ) (Guenther et al., 1995). It should be noted, besides biogenic emissions, combustion of biomass and fossil fuel also contribute to the isoprene, monoterpenes and sesquiterpenes (Jathar et al., 2014; Sarkar

400 et al., 2016;Sarkar et al., 2017). The investigation of gaseous VOCs during winter (December 2012 to  
February 2013) air in the KV during SusKat-ABC campaign also showed high levels of isoprene and it  
was attributed (at least during high isoprene periods) mostly to biogenic emissions (Sarkar et al.,  
2016;Sarkar et al., 2017). It is difficult to appropriately quantify the fractions of biogenic and  
405 anthropogenic emissions of these compounds, based on ambient measurement of these species alone  
without measurement of BB tracers such as acetonitrile and furan. The budget of isoprene emissions (500  
Tg y<sup>-1</sup>) on a global scale is dominated by vegetation (Guenther et al., 2006). Therefore, in our study, we  
considered the oxidation products of isoprene, monoterpenes and sesquiterpenes as the tracers of biogenic  
emissions and attribute their main source as biogenic emissions. This may lead to some overestimation  
of their contributions to SOA formation.

### 410 3.6.1 Isoprene SOA tracers

Six isoprene-SOA (I-SOA) tracers were identified in the Bode aerosols: 2-MGA, two  
diastereoisomeric 2-methyltetrols (2-MTLs) and three C5-alkene triols. Their total concentrations ranged  
from 38.8-444 ng m<sup>-3</sup> (174±86.2 ng m<sup>-3</sup>) with the maximum (236±87.2 ng m<sup>-3</sup>) in the monsoon season  
(Table 1). During the post-monsoon and pre-monsoon periods, their concentrations were similar, and a  
415 little lower than those during the monsoon (Fig. 5d) while being the lowest during winter. Their seasonal  
variation was in agreement with the ambient temperature (Fig S1a), which can influence the isoprene  
emissions and the photochemical processes (Shen et al., 2015;Wang et al., 2008). The annual average  
concentration was higher than the urban sites reported from Beijing (44.3 ng m<sup>-3</sup>) and Kunming (108 ng  
m<sup>-3</sup>) (Ding et al., 2016a), even one to two orders of magnitude higher than that from global oceans and  
420 the Arctic (Hu et al., 2013). Among I-SOA tracers, 2-MTLs were the major components (51.0% ± 10.5%)  
(Fig. 6), with an annual average of 94.4±58.9 ng m<sup>-3</sup> (ranging from 10.9 to 270 ng m<sup>-3</sup>). Strong correlations  
were exhibited between the two isomers during all the seasons (Fig. S5a), implying that they formed  
through the similar pathway (Shen et al., 2015;Fu et al., 2010). The daily concentration of 2-MGA ranged  
from 7.10-79.0 ng m<sup>-3</sup> with an annual average of 34.2±14.8 ng m<sup>-3</sup>. For C5-alkene triols, the average  
425 concentration was 45.0±29.4 ng m<sup>-3</sup>. They positively correlated with 2-MTLs (Fig. S5b), indicating they  
were also the oxidation products of isoprene under low-NOx conditions.



According to the reaction chamber results from Surratt et al. (2010), the formation mechanism of 2-MGA remarkably differs from 2-MTLs. 2-MGA is formed under high-NO<sub>x</sub> condition while 2-MTLs are mainly produced under low-NO<sub>x</sub> or NO<sub>x</sub>-free conditions. The formation of 2-MGA can be enhanced under lower RH condition, while it is opposite for 2-MTLs (Zhang et al., 2011). During the monsoon season, due to the conducive conditions of high temperature, high RH (>70 %) (Fig. S1b), high solar radiation and fully-grown plants, the isoprene emissions was large. In addition, NO<sub>x</sub> during this season was much lower than other seasons. Therefore, 2-MGA/2-MTLs ratios exhibited the lowest values (0.20±0.08) in the aerosol samples during this wet season (Fig. 7). In contrast, 2-MGA/2-MTLs ratios increased up to 0.95 in winter, owing to the lowest temperature and RH of the whole year (Fig. 7) and the higher NO<sub>x</sub> concentration in the KV (Kondo et al., 2005;Kiros et al., 2016). NO<sub>x</sub> from anthropogenic sources (industry, transportation, BB in the houses as well as in the field) and meteorological conditions with reduced mixing layer heights in winter would also favor the formation of 2-MGA and subsequently increase the 2-MGA/2-MTLs ratio.

Positive correlations were observed between 2-MGA, SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> (Fig. 8). Budisulistiorini et al. (2017) investigated that the concentrations of B-SOA could significantly increase as the aerosol acidity enhances based on the laboratory simulations and field observations. The significant influence of I-SOA by SO<sub>4</sub><sup>2-</sup> might be explained by the concerted nucleophilic addition to the key intermediates in the gas phase (e.g., isoprene epoxydiols), which is the rate-determining step in SOA formation (Xu et al., 2015;Li et al., 2018). Li et al. (2018) reported that SO<sub>4</sub><sup>2-</sup> plays an important part in promoting aqueous phase oxidation of I-SOA tracers. There may be the similar effect of NO<sub>3</sub><sup>-</sup> on the SOA formation that needs further research. Therefore, the increase of SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> could effectively facilitate the ring-opening reaction of isoprene epoxydiols and the SOA formation. Thus, the higher 2-MGA in the KV may be due to the abundant SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> during pre-monsoon season when most of the brick kilns (more than 100) are operational. Our finding demonstrated the anthropogenic pollutants (e.g., SO<sub>2</sub>, NO<sub>x</sub>, etc.) can be conducive to accelerating the oxidation of B-VOCs and enhancing the ambient concentrations of B-SOA.

### 3.6.2 Terpene SOA tracers

Besides isoprene tracers, we also measured four monoterpene oxidation products (M-SOA tracers), including PNA, PA, 3-HGA and MBTCA (Claeys et al., 2007). They are produced through the

455 photooxidation of monoterpenes with ozone and hydroxyl radical (Iinuma et al., 2004). The annual  
average concentration of the total M-SOA tracers was  $59.3 \pm 24.6 \text{ ng m}^{-3}$  (Table 1). The concentration of  
M-SOA tracers was higher than those investigated in previous studies from the urban site in Kunming  
(annual average:  $44.1 \pm 38.8 \text{ ng m}^{-3}$ ) (Ding et al., 2016b), three cities (Ohio, Michigan and California) in  
North America (summer:  $30.4\text{--}60.6 \text{ ng m}^{-3}$ ) (Stone et al., 2009) and a forest site in Hyytiälä, Europe  
460 (summer:  $15.1\text{--}33.3 \text{ ng m}^{-3}$ ) (Kourtchev et al., 2005).

For the seasonal variation, relatively high concentrations of M-SOA tracers occurred during pre-  
monsoon and post-monsoon seasons (Fig. 5e, f, g, h, and i). Interestingly, there is intensive BB in KV  
twice a year (forest fires and crop-residue fires during April to May, and crop-residue fires during October  
to November) discussed in section 3.3.1 and 3.4, which may have been associated with high  
465 concentrations of M-SOA tracers. During the fires, substantial amounts of aerosols and VOCs including  
isoprene and monoterpenes would generate, which can enhance the levels of B-SOA tracers (Jathar et al.,  
2014;Ding et al., 2013). Good correlations were obtained between the BB tracer i.e. levoglucosan and the  
higher generation oxidation products (e.g., 3-HGA and MBTCA,  $R^2=0.32$  and  $R^2=0.53$  respectively) in  
the Bode aerosols (Fig. S6). The forests in the KV mainly consist of broad-leaved evergreen mixed forest,  
470 oak-laurel forest and oak forest as well as the conifer tree species (Department of Plant Resources,  
2015;Sarkar et al., 2016). Monoterpenes were mainly released from coniferous trees (Kang et al., 2018).  
Therefore, it suggested that the atmospheric aerosol compositions especially of SOA tracers over KV  
maybe significantly affected by BB activities.

Sesquiterpenes (e.g.,  $\beta$ -caryophyllene) are also among the biogenic SOA (B-SOA) precursors  
475 emitted from trees, which have been observed in the troposphere in a lot of studies.  $\beta$ -CA is the tracer of  
 $\beta$ -caryophyllene SOA and its concentration in the Bode aerosols was  $6.31 \pm 3.86 \text{ ng m}^{-3}$  with the range of  
 $1.53$  to  $18.5 \text{ ng m}^{-3}$ . It shared the similar seasonal variation with M-SOA tracers and positively correlated  
with them, indicating the possible common emission pattern.

### 3.6.3 Aromatic SOA tracer

480 Anthropogenic SOA is also an important OC source. DHOPA is a tracer of anthropogenic SOA from  
aromatics. In this study, DHOPA showed higher levels in winter and pre-monsoon periods while lower

in monsoon season (Fig. 5). Though the major emissions of aromatics come from fossil sources, BB is also considered as an possible source in some sites of the world (Shen et al., 2015). There was a good correlation between DHOPA and levoglucosan (Fig. 9), especially during pre-monsoon with the value of  $R^2$  at 0.73. This indicated BB emissions are also significant source of DHOPA in Bode.

### 3.7 Estimation of the contributions of different sources to OC

As discussed above, both the primary and secondary sources have influence on OC in the atmospheric aerosols of the KV. In this part, we will apply the tracer-based methods to evaluate the different sources' contributions to OC. It should be noted here that tracer methods can provide a reasonable estimation, but uncertainties are introduced considering the site differences and the lack of representative source profiles for the given study location. The contribution evaluated from each source to OC in the current study is still inferable.

#### 3.7.1 BB-derived OC

Levoglucosan to OC ratios (Lev/OC) detected in source samples has been used in a wide range for quantitative estimation of the BB contribution to OC (Stone et al., 2012;Zhang et al., 2015;Wan et al., 2017), although the ratios vary among different types of biomass burnt and combustion conditions (Bhattacharai et al., 2019). An average of 8.14% (8.0% to 8.2 %) for Lev/OC from the burning sites of biofuel, savanna, crop residues, tropical forests, and so on was documented in (Andreae and Merlet, 2001). Zhang et al. (2007) obtained the Lev/OC ratios ranging from 5.4%-11.8% (an average of 8.27%) from the aerosols of cereal straw (wheat, corn and rice) combustion. Sheesley et al. (2003) reported an average of 7.94% of levoglucosan from the combustion of biomass (including rice straw, biomass briquettes, dried cow-dung patties, etc.) indigenous to South Asia. However, the ratio acquired from the hardwood burning in fireplaces and stoves in the US was 14%, which was applied at the background sites in Europe (Fine et al., 2004). Stone et al. (2012) used the Lev/OC ratio of  $12\% \pm 0.2\%$  during the burning of acacia wood at Godavari in the KV for the CBM profile source apportionment. The mean value of Lev/OC value of BB from main biomass types was 10.1%. In this study, we choose the mostly used values of 8.14% for BB estimation (Graham et al., 2002;Fu et al., 2014;Ho et al., 2014;Sang et al., 2011;Zhu et al., 2016;Mkoma et al., 2013). In addition, the uncertainties of using different ratios were also calculated (see

Table S3). The diagnostic ratios among molecular tracers and OC (e.g., Lev/OC) from direct emissions  
510 are critical for more precise results. It's meaningful to understand the emission characteristics for  
individual OC emission categories, as well as in different locations, especially in South Asia.

Figures 10 and 11 present the monthly concentration variations of BB-OC and contribution of BB-  
OC to OC, respectively. Current estimation exhibited BB-OC contributed  $24.9\pm 10.4\%$  (varying from 6.32%  
515 to 61.5%) to OC throughout the year in Bode aerosols (Fig. 11a). This was higher than the study in  
Lumbini in Nepal ( $19.8\pm 19.4\%$ ) (Wan et al., 2017), and nearly twice of the BB-OC contribution to OC  
reported in Hong Kong (6.5%–11%) (13.1 %) (Sang et al., 2011) and the Pearl River Delta in China (Ho  
et al., 2014). Moreover, the contribution of BB-OC to OC in current study maximized in the post-monsoon  
season ( $36.3\pm 10.4\%$ ), higher than that in the pre-monsoon ( $28.5\pm 10.3\%$ ) and winter ( $27.9\pm 8.63\%$ ) periods.  
These results indicate that BB severely affect the air quality in the KV, especially during the post-monsoon  
520 period. Similarly, Stone et al. (2010) reported  $21\pm 2\%$  of OC in  $PM_{2.5}$  from Godavari rural site in the  
outskirts of the KV during 2006, was also attributed to the primary BB sources.

### 3.7.2 Plant-debris-OC and fungal-spore-derived OC

Primary biological aerosol particle (PBAP) has been identified as an important source using tracers  
(section 3.3.2). They are likely to have a big contribution to the aerosols in Bode. In order to reveal how  
525 much they are contributing to organic aerosols, “total” plant debris was calculated based on glucose  
following the equation (Zhu et al., 2016; Puxbaum and Tenze-Kunit, 2003):

$$\text{Cellulose } (\mu\text{g}) = \text{D-glucose } (\mu\text{g}) \times \text{GF} \times (1/\text{SY});$$

$$\text{Plant debris} = 2 \times \text{cellulose}.$$

where GF (0.90) is the glucose/cellulose weight conversion factor and SY (0.717) is the  
530 saccharification yield.

For OC fraction derived from fungal spores, it was estimated using mannitol levels according to the  
studies from Bauer et al. (2008) and Holden et al. (2011), i.e., there was 1.7 pg mannitol and 13 pg OC  
per spore.

As shown in Fig. 11a, fungal spore-derived OC and plant-debris-OC annually contribute to  
535  $3.15\pm 2.86\%$  and  $1.42\pm 1.03\%$  of OC, respectively. The contributions were both higher in the monsoon

season, with  $5.85\pm 2.50\%$  for fungal spore-derived OC and  $2.29\pm 0.79\%$  for plant-debris-OC to OC, respectively (Fig. 11c). During winter, the contributions were the lowest due to the inactive vegetation. There are also some similar results from the literatures. For example, Zhu et al. (2016) reported the plant debris contribution to OC was 5.6% and 4.6% during nighttime and daytime respectively from aerosols in a mid-latitude forest. Szidat et al. (2006) reported the plant debris contributed to 3.2% of OC during summer in urban aerosols collected in Zurich, Switzerland. The contributions of fungal aerosol to OC was 8% in the aerosols from a Brazil urban site (Eymygdio et al., 2018). Liang et al. (2017) reported the fungal aerosol contributions of  $3.5 \pm 3.7\%$  in aerosols from a rural site in Beijing, China. In marine aerosols, the fungal spores were documented the major contributor to total OC with 3.1% (0.03%–19.8%) over the East China Sea (Kang et al., 2018). All above strengthened the importance of plant-debris and fungal spores to the aerosol burden in the atmosphere.

### 3.7.3 Biogenic SOC and anthropogenic toluene SOC

Biogenic secondary organic carbon (B-SOC) and anthropogenic aromatic SOC (A-SOC) from the oxidation of isoprene, monoterpenes, sesquiterpene and toluene were assessed using the tracer-based method proposed by Kleindienst et al. (2007). This method has been applied successfully in numerous aerosol studies (Shen et al., 2015; Ding et al., 2016a; Kang et al., 2018). The mass fraction of tracer compounds in SOC (FSOC) for an individual precursor was calculated based on the smog chamber simulations. The calculation formula as following:

$$FSOC = \frac{\sum_i [tri]}{[SOC]}$$

where [tri] is the concentration of tracer *i* and [SOC] is the concentration of SOC. The conversion factors of FSOC were  $0.155\pm 0.039$ ,  $0.231\pm 0.111$ ,  $0.0230\pm 0.0046$  and  $0.0079\pm 0.0026 \mu\text{g } \mu\text{gC}^{-1}$  for isoprene, monoterpenes, sesquiterpene and toluene, respectively (Kleindienst et al., 2007).

The total calculated concentrations of B-SOC varied from 0.41 to  $2.77 \mu\text{g m}^{-3}$  with an annual mean concentration of  $1.36\pm 0.49 \mu\text{g m}^{-3}$ , a higher concentration of  $1.43\pm 0.48 \mu\text{g m}^{-3}$  in monsoon and lower concentration of  $0.86\pm 0.20 \mu\text{g m}^{-3}$  in winter (Fig 10g). The B-SOC/OC showed a higher average percentage of  $10.1\% \pm 3.34\%$  in the monsoon season (Fig 11c), indicating B-SOC was an important OC

sources in Bode during this period. During post-monsoon, B-SOC/OC declined to 5.36% (Fig 11d). The B-SOC/OC showed the lowest value of  $1.52\% \pm 0.70\%$  in winter (Fig 11e), indicating that B-SOC had minor contributions to elevated OC in winter. The annual average concentration of A-SOC was  $2.45 \pm 1.45 \mu\text{g m}^{-3}$ , which is higher than the B-SOC. The highest A-SOC concentration was obtained in winter ( $3.27 \pm 1.25 \mu\text{g m}^{-3}$ ) (Fig.10h). A-SOC was the second most important contributor to OC after BB-OC. It is not only derived from increased fossil fuel combustion and the subsequent oxidation, but also from BB emissions.

In total, SOC (including B-SOC and A-SOC) reconstructed using the formula above in this section was  $3.81 \pm 1.63 \mu\text{g m}^{-3}$ , accounting for  $15.0\% \pm 8.99\%$  of OC.

### 3.7.4 Possible sources of the unidentified OC

On the whole, BB contributed one-fourth ( $24.9\% \pm 10.4\%$ ) of the OC in Bode, followed by A-SOC ( $8.82\% \pm 5.55\%$ ), B-SOC ( $6.19\% \pm 4.49\%$ ), fungal-spores ( $3.15\% \pm 2.86\%$ ) and plant-debris ( $1.42\% \pm 1.03\%$ ) (Fig. 11a). Nevertheless, there is still part of OC (55.5%) that we were not able to be attributed to any specific sources based on the tracers analyzed in current study. There are partly uncertainties caused by the organic tracer analyses (estimation of measurement uncertainty was shown in Table S2). Furthermore, fossil fuel combustion and soil dust could be also notable fractions of OC in Bode aerosols. Additionally, low molecular weight (LMW) dicarboxylic acids from both primary and secondary sources is also a remarkable contributor to atmospheric organic aerosols (Kawamura and Bikkina, 2016). Humic-like substances and amines can constitute another fraction of OC, but not well studied (Wu et al., 2018;Laskin et al., 2015). Therefore, the possible contributions of the unidentified OC (55.5%) from various sectors need further investigation, which is better to comprehensively understand the sources of South Asian aerosols and will be very useful for the targeted pollution control measures in this region.

## 4. Summary and conclusions

Field measurements of atmospheric aerosols were conducted in a semi-urban site (Bode) of the KV, Nepal, from April 2013 to April 2014. The organic tracers from primary and secondary organic aerosols were determined. A distinctive seasonality was observed for various aerosol species. Higher

concentrations of OC, EC, anhydrosugars, phenolic compounds and resin acid were observed in winter and pre-monsoon seasons, while their concentrations were lower during the wet (monsoon) period. Levoglucosan was the dominant species of the total identified tracers with an average concentration of 788 ng m<sup>-3</sup>. We observed high abundances of monosaccharides in the pre-monsoon season and of sugar alcohols in the wet period, and lower levels in winter because of the reduced plant activities. I-SOA tracers represented a majority among B-SOA tracers with a maximum in the monsoon. The seasonal variation of M-SOA tracers was controlled by monoterpenes emission and BB. DHOPA exhibited higher concentrations during the winter and pre-monsoon seasons.

The likely OC sources were further evaluated for their contributions to observed total OC using tracer-based methods. BB-OC contributed a major fraction (24.9%) to OC in Bode, followed by A-SOC (8.8%), B-SOC (6.2%), fungal spores (3.2%) and plant debris (1.4%). The highest contribution of BB-OC, 36.3%, occurred during post-monsoon season. A-SOC, B-SOC, fungal spores and plant debris all made larger contributions during the monsoon. The higher BB-OC and the A-SOC contributions imply that some BB and anthropogenic components are widespread in the KV and thus represent the main contributors affecting the regional air quality in the KV region.

The present study clearly shows that the chemical constituents and sources of OC strongly vary with seasons, as a result of diverse air pollution sources in the valley across four seasons. The heavy BB and the subsequent oxidant emissions are anticipated to cause larger contributions of B-SOC to OC. Understanding OC's climate impacts is a frontier area of research, because a large uncertainty still exists in the estimation of OC radiative forcing. Our study implies that since BB is a major source of ambient OC, the fraction of OC that absorbs light (referred to as brown carbon) and also acts as cloud condensation nuclei, needs to be further studied in order to better understand radiative effects of OC on regional climate change. The current source contribution estimates from the tracer-based methods do not accurately evaluate the large temporal variations from all kinds of sources. Contributions from other sectors (ca 55.5%), including low molecular weight dicarboxylic acids (Kawamura and Sakaguchi, 1999; Kawamura and Bikkina, 2016), need further investigation to better understand the atmospheric aerosols from both urban and rural sources such as the KV and other sites in the Himalayan foothills and the Indo-Gangetic Plain regions. These observations of the severe air pollution, particularly the particular matter pollution,

provide valuable support for air pollution control measures, especially in determining which sources and sectors to first focus on the KV and the surrounding region, in order to reduce the air pollution from being severe to become much cleaner in the near future. In addition, the current study based on the molecular level-source apportionment of OC in heavy polluted region from South Asia provides a much more specific quantification of source estimation for OC, which is different from previous studies based on the bulk carbonaceous aerosol using radiocarbon ( $^{14}\text{C}$ ) measurements, PMF and CBM.

There are additional improvements for future studies to be addressed in the supporting information file. The key recommendations are as follows: (i) much more tracers need to be identified to explain the other sources of organic aerosols in the KV; (ii) the conversion factors of tracers to organic carbon from local emissions are critical for more precise source apportionments and therefore, a number of studies on the emission characteristics will be valuable; (iii) comprehensive methods (e.g., carbon isotope and modeling) need to be integrated together for the source apportionment of organic aerosols in the KV; (iv) the influences of BB on the formation of secondary organic aerosols could be further studied, especially during the heavily polluted dry season, with additional simultaneous measurements of precursors (e.g.,  $\text{NO}_x$  and  $\text{O}_3$ ),  $\text{PM}_{2.5}$  and so on at the same time; (v) to better understand the atmospheric processes of various chemical species, investigations of size-segregated aerosols are especially needed in the heavy polluted KV.

*Data availability.* Raw data are archived at the Institute of Tibetan Plateau Research, Chinese Academy of Sciences, and are available on request by contacting the corresponding author.

*Competing interests.* The authors declare that they have no conflict of interest.

*Author contributions.* All authors contributed to the final version of this article. XW analyzed the organic molecular tracers and wrote the paper under the supervision of ZC. SK, QZ and ZC organized the campaign. JG and PC worked with the instruments and collected the aerosol samples. PC and LT analyzed the OC, EC and major ions data. MR, KK, ML, AP and DR provided advice and feedback throughout the drafting and submission process.



## Acknowledgments

The authors gratefully appreciate Shyam Kumar Newar and Bhogendra Kathayat for their assistance in the sample collections, the staffs at Bode site in the Kathmandu Valley, and all individuals and groups who participated in the SusKat-ABC field campaigns, and the supports from Kathmandu Center for Research and Education, Chinese Academy of Sciences-Tribhuvan University. We would like to thank senior scientist Karl Epton Yttri from Norwegian Institute for Air Research for the helpful suggestions to reply the comments from the referees. This study was supported by the Strategic Priority Research Program of Chinese Academy of Sciences, Pan-Third Pole Environment Study for a Green Silk Road (Pan-TPE) (XDA20040501), the National Natural Science Foundation of China (41522103, 41421061 and 41630754) and China Postdoctoral Science Foundation (2018M630210). The coauthors from the IASS gratefully acknowledge funding from the federal ministry of education and research (BMBF) and the Brandenburg state ministry of science, research and culture (MWFK).

## References

- Department of Transport Management (DoTM).: Annual report of Ministry of Labor and Transport Management, Government of Nepal, 2015.
- Alam, K., Mukhtar, A., Shahid, I., Blaschke, T., Majid, H., Rahman, S., Khan, R., and Rahman, N.: Source apportionment and characterization of particulate matter (PM<sub>10</sub>) in urban environment of Lahore, *Aerosol and Air Quality Research*, 14, 1851-1861, 2014.
- Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochemical Cycles*, 15, 955-966, 10.1029/2000gb001382, 2001.
- Bauer, H., Claeys, M., Vermeylen, R., Schueller, E., Weinke, G., Berger, A., and Puxbaum, H.: Arabitol and mannitol as tracers for the quantification of airborne fungal spores, *Atmospheric Environment*, 42, 588-593, 10.1016/j.atmosenv.2007.10.013, 2008.
- Bhardwaj, P., Naja, M., Rupakheti, M., Panday, A. K., Kumar, R., Mahata, K., Lal, S., Chandola, H. C., and Lawrence, M. G.: Variations in surface ozone and carbon monoxide in the Kathmandu Valley and surrounding broader regions during SusKat-ABC field campaign: Role of local and regional sources, *Atmospheric Chemistry and Physics in review*, 2017.
- Bhattacharai, H., Saikawa, E., Wan, X., Zhu, H., Ram, K., Gao, S., Kang, S., Zhang, Q., Zhang, Y., Wu, G., Wang, X., Kawamura, K., Fu, P., and Cong, Z.: Levoglucosan as a tracer of biomass burning: Recent progress and perspectives, *Atmospheric Research*, 220, 20-33, 2019.
- Bisht, D., Dumka, U., Kaskaoutis, D., Pipal, A., Srivastava, A., Soni, V., Attri, S., Sateesh, M., and Tiwari, S.: Carbonaceous aerosols and pollutants over Delhi urban environment: temporal evolution, source apportionment and radiative forcing, *Science of the Total Environment*, 521, 431-445, 2015.
- Boreddy, S. K. R., Kawamura, K., and Tachibana, E.: Long-term (2001-2013) observations of water-soluble dicarboxylic acids and related compounds over the western North Pacific: trends, seasonality and source apportionment, *Scientific Reports*, 7, 10.1038/s41598-017-08745-w, 2017.
- Budisulistiorini, S. H., Nenes, A., Carlton, A. G., Surratt, J. D., McNeill, V. F., and Pye, H. O. T.: Simulating Aqueous-Phase Isoprene-Epoxydiol (IEPOX) Secondary Organic Aerosol Production During the 2013 Southern Oxidant and Aerosol Study (SOAS), *Environmental Science & Technology*, 51, 5026-5034, 10.1021/acs.est.6b05750, 2017.

- 685 Cao, J. J., Lee, S. C., Chow, J. C., Watson, J. G., Ho, K. F., Zhang, R. J., Jin, Z. D., Shen, Z. X., Chen, G. C., Kang, Y. M., Zou, S. C., Zhang, L. Z., Qi, S. H., Dai, M. H., Cheng, Y., and Hu, K.: Spatial and seasonal distributions of carbonaceous aerosols over China, *Journal of Geophysical Research-Atmospheres*, 112, 10.1029/2006jd008205, 2007.
- Chen, P., Kang, S., Li, C., Rupakheti, M., Yan, F., Li, Q., Ji, Z., Zhang, Q., Luo, W., and Sillanpaa, M.: Characteristics and sources of polycyclic aromatic hydrocarbons in atmospheric aerosols in the Kathmandu Valley, Nepal, *Science of the Total Environment*, 538, 86-92, 10.1016/j.scitotenv.2015.08.006, 2015.
- 690 Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, 303, 1173-1176, 10.1126/science.1092805, 2004.
- Claeys, M., Szmigielski, R., Kourtchev, I., Van der Veken, P., Vermeylen, R., Maenhaut, W., Jaoui, M., Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., and Edney, E. O.: Hydroxydicarboxylic Acids: Markers for Secondary Organic Aerosol from the Photooxidation of  $\alpha$ -Pinene, *Environmental Science & Technology*, 41, 1628-1634, 10.1021/es0620181, 2007.
- 695 Cowie, G. L., and Hedges, J. I.: Carbohydrate sources in a coastal marine environment, *Geochimica Et Cosmochimica Acta*, 48, 2075-2087, 10.1016/0016-7037(84)90388-0, 1984.
- Department of Plant Resources, N.: Bulletin of Department of Plant Resources, Nepal 37, Ministry of Forests and Soil Conservation, Kathmandu, 1-122, 2015.
- Deshmukh, D. K., Kawamura, K., and Deb, M. K.: Dicarboxylic acids,  $\omega$ -oxocarboxylic acids,  $\alpha$ -dicarbonyls, WSOC, OC, EC, and inorganic ions in wintertime size-segregated aerosols from central India: Sources and formation processes, *Chemosphere*, 161, 27-42, 2016.
- 700 Ding, X., Wang, X., Xie, Z., Zhang, Z., and Sun, L.: Impacts of Siberian Biomass Burning on Organic Aerosols over the North Pacific Ocean and the Arctic: Primary and Secondary Organic Tracers, *Environmental Science & Technology*, 47, 3149-3157, 10.1021/es3037093, 2013.
- 705 Ding, X., He, Q.-F., Shen, R.-Q., Yu, Q.-Q., Zhang, Y.-Q., Xin, J.-Y., Wen, T.-X., and Wang, X.-M.: Spatial and seasonal variations of isoprene secondary organic aerosol in China: Significant impact of biomass burning during winter, *Scientific Reports*, 6, 10.1038/srep20411, 2016a.
- Ding, X., Zhang, Y.-Q., He, Q.-F., Yu, Q.-Q., Shen, R.-Q., Zhang, Y., Zhang, Z., Lyu, S.-J., Hu, Q.-H., Wang, Y.-S., Li, L.-F., Song, W., and Wang, X.-M.: Spatial and seasonal variations of secondary organic aerosol from terpenoids over China, *Journal of Geophysical Research-Atmospheres*, 121, 14661-14678, 10.1002/2016jd025467, 2016b.
- 710 Emygdio, A. P. M., Andrade, M. d. F., Gonçalves, F. L. T., Engling, G., Zanetti, R. H. d. S., and Kumar, P.: Biomarkers as indicators of fungal biomass in the atmosphere of São Paulo, Brazil, *Science of The Total Environment*, 612, 809-821, 2018.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the wood stove combustion of prevalent United States tree species, *Environmental Engineering Science*, 21, 705-721, 10.1089/ees.2004.21.705, 2004.
- 715 Fu, P., Kawamura, K., Chen, J., and Miyazaki, Y.: Secondary production of organic aerosols from biogenic VOCs over Mt. Fuji, Japan, *Environmental Science & Technology*, 48, 8491-8497, 2014.
- Fu, P. Q., Kawamura, K., Pavuluri, C. M., Swaminathan, T., and Chen, J.: Molecular characterization of urban organic aerosol in tropical India: contributions of primary emissions and secondary photooxidation, *Atmospheric Chemistry and Physics*, 10, 2663-2689, 2010.
- 720 Fujii, Y., Kawamoto, H., Tohno, S., Oda, M., Iriana, W., and Lestari, P.: Characteristics of carbonaceous aerosols emitted from peatland fire in Riau, Sumatra, Indonesia (2): Identification of organic compounds, *Atmospheric Environment*, 110, 1-7, 2015.
- 725 George, S. K., Nair, P. R., Parameswaran, K., Jacob, S., and Abraham, A.: Seasonal trends in chemical composition of aerosols at a tropical coastal site of India, *Journal of Geophysical Research-Atmospheres*, 113, 10.1029/2007jd009507, 2008.
- Giri, D., Murthy, V. K., Adhikary, P. R., and Khanal, S. N.: Cluster analysis applied PM10 concentration data for determination of sources and spatial patterns in ambient air-quality of Khatmandu Valley, *Current Science*, 93, 684-688, 2007.
- Graham, B., Mayol-Bracero, O. L., Guyon, P., Roberts, G. C., Decesari, S., Facchini, M. C., Artaxo, P., Maenhaut, W., Koll, P., and Andreae, M. O.: Water-soluble organic compounds in biomass burning aerosols over Amazonia - 1. Characterization by NMR and GC-MS, *Journal of Geophysical Research-Atmospheres*, 107, 10.1029/2001jd000336,
- 730

2002.

- 735 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A.,  
Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural  
volatile organic compound emissions, *Journal of Geophysical Research-Atmospheres*, 100, 8873-8892,  
10.1029/94jd02950, 1995.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene  
emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and  
Physics* 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.
- 740 Gupta, A. K., Karar, K., and Srivastava, A.: Chemical mass balance source apportionment of PM10 and TSP in residential and  
industrial sites of an urban region of Kolkata, India, *Journal of Hazardous Materials*, 142, 279-287,  
10.1016/j.jhazmat.2006.08.013, 2007.
- Ho, K. F., Engling, G., Ho, S. S. H., Huang, R., Lai, S., Cao, J., and Lee, S. C.: Seasonal variations of anhydrosugars in PM2.5  
in the Pearl River Delta Region, China, *Tellus Series B-Chemical and Physical Meteorology*, 66,  
10.3402/tellusb.v66.22577, 2014.
- 745 Holden, A. S., Sullivan, A. P., Munchak, L. A., Kreidenweis, S. M., Schichtel, B. A., Malm, W. C., and Collett, J. L., Jr.:  
Determining contributions of biomass burning and other sources to fine particle contemporary carbon in the western  
United States, *Atmospheric Environment*, 45, 1986-1993, 10.1016/j.atmosenv.2011.01.021, 2011.
- Hu, Q.-H., Xie, Z.-Q., Wang, X.-M., Kang, H., He, Q.-F., and Zhang, P.: Secondary organic aerosols over oceans via oxidation  
of isoprene and monoterpenes from Arctic to Antarctic, *Scientific Reports*, 3, 10.1038/srep02280, 2013.
- 750 Iinuma, Y., Böge, O., Gnauk, T., and Herrmann, H.: Aerosol-chamber study of the  $\alpha$ -pinene/O<sub>3</sub> reaction: influence of particle  
acidity on aerosol yields and products, *Atmospheric Environment*, 38, 761-773, 2004.
- Jathar, S. H., Gordon, T. D., Hennigan, C. J., Pye, H. O. T., Pouliot, G., Adams, P. J., Donahue, N. M., and Robinson, A. L.:  
Unspeciated organic emissions from combustion sources and their influence on the secondary organic aerosol budget  
in the United States, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 10473-  
755 10478, 10.1073/pnas.1323740111, 2014.
- Kang, M., Fu, P., Kawamura, K., Yang, F., Zhang, H., Zang, Z., Ren, H., Ren, L., Zhao, Y., Sun, Y., and Wang, Z.:  
Characterization of biogenic primary and secondary organic aerosols in the marine atmosphere over the East China  
Sea, *Atmospheric Chemistry and Physics*, 18, 13947-13967, 10.5194/acp-18-13947-2018, 2018.
- 760 Kawamura, K., and Sakaguchi, F.: Molecular distributions of water soluble dicarboxylic acids in marine aerosols over the  
Pacific Ocean including tropics, *Journal of Geophysical Research-Atmospheres*, 104, 3501-3509,  
10.1029/1998jd100041, 1999.
- Kawamura, K., and Bikkina, S.: A review of dicarboxylic acids and related compounds in atmospheric aerosols: Molecular  
distributions, sources and transformation, *Atmospheric Research*, 170, 140-160, 2016.
- 765 Kim, B. M., Park, J.-S., Kim, S.-W., Kim, H., Jeon, H., Cho, C., Kim, J.-H., Hong, S., Rupakheti, M., Panday, A. K., Park, R.  
J., Hong, J., and Yoon, S.-C.: Source apportionment of PM10 mass and particulate carbon in the Kathmandu Valley,  
Nepal, *Atmospheric Environment*, 123, 190-199, 10.1016/j.atmosenv.2015.10.082, 2015.
- Kiros, F., Shakya, K. M., Rupakheti, M., Regmi, R. P., Maharjan, R., Byanju, R. M., Naja, M., Mahata, K., Kathayat, B., and  
Peltier, R. E.: Variability of Anthropogenic Gases: Nitrogen Oxides, Sulfur Dioxide, Ozone and Ammonia in  
Kathmandu Valley, Nepal, *Aerosol and Air Quality Research*, 16, 3088-3101, 10.4209/aaqr.2015.07.0445, 2016.
- 770 Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberg, J. H., Lewis, C. W., Bhave, P. V., and Edney, E. O.: Estimates of  
the contributions of biogenic and anthropogenic hydrocarbons to secondary organic aerosol at a southeastern US  
location, *Atmospheric Environment*, 41, 8288-8300, 10.1016/j.atmosenv.2007.06.045, 2007.
- Kondo, A., Kaga, A., Imamura, K., Inoue, Y., Sugisawa, M., Shrestha, M. L., and Sapkota, B.: Investigation of air pollution  
concentration in Kathmandu valley during winter season, *Journal of Environmental Sciences*, 17, 1008-1013, 2005.
- 775 Kourtchev, I., Ruuskanen, T., Maenhaut, W., Kulmala, M., and Claeys, M.: Observation of 2-methyltetrols and related photo-  
oxidation products of isoprene in boreal forest aerosols from Hyytiälä, Finland, *Atmospheric Chemistry and Physics*,  
5, 2761-2770, 10.5194/acp-5-2761-2005, 2005.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chemical Reviews*, 115, 4335, 2015.
- 780 Lawrence, M. G., and Lelieveld, J.: Atmospheric pollutant outflow from southern Asia: a review, *Atmospheric Chemistry and  
Physics*, 10, 11017-11096, 10.5194/acp-10-11017-2010, 2010.

- Li, J., Wang, G., Aggarwal, S. G., Huang, Y., Ren, Y., Zhou, B., Singh, K., Gupta, P. K., Cao, J., and Zhang, R.: Comparison of abundances, compositions and sources of elements, inorganic ions and organic compounds in atmospheric aerosols from Xi'an and New Delhi, two megacities in China and India, *Science of the Total Environment*, 476, 485-495, 10.1016/j.scitotenv.2014.01.011, 2014.
- 785 Li, J., Wang, G., Ren, Y., Wang, J., Wu, C., Han, Y., Zhang, L., Cheng, C., and Meng, J.: Identification of chemical compositions and sources of atmospheric aerosols in Xi'an, inland China during two types of haze events, *Science of the Total Environment*, 566, 230-237, 10.1016/j.scitotenv.2016.05.057, 2016.
- Li, J., Wang, G., Wu, C., Cao, C., Ren, Y., Wang, J., Li, J., Cao, J., Zeng, L., and Zhu, T.: Characterization of isoprene-derived secondary organic aerosols at a rural site in North China Plain with implications for anthropogenic pollution effects, *Scientific Reports*, 8, 10.1038/s41598-017-18983-7, 2018.
- 790 Liang, L., Engling, G., Du, Z., Cheng, Y., Duan, F., Liu, X., and He, K.: Seasonal variations and source estimation of saccharides in atmospheric particulate matter in Beijing, China, *Chemosphere*, 150, 365-377, 10.1016/j.chemosphere.2016.02.002, 2016.
- Liang, L., Engling, G., Du, Z., Duan, F., Cheng, Y., Liu, X., and He, K.: Contribution of fungal spores to organic carbon in ambient aerosols in Beijing, China, *Atmospheric Pollution Research*, 8, 351-358, 10.1016/j.apr.2016.10.007, 2017.
- 795 Mahata, K. S., Panday, A. K., Rupakheti, M., Singh, A., Naja, M., and Lawrence, M. G.: Seasonal and diurnal variations in methane and carbon dioxide in the Kathmandu Valley in the foothills of the central Himalayas, *Atmospheric Chemistry and Physics* 17, 12573-12596, 10.5194/acp-17-12573-2017, 2017a.
- Mahata, K. S., Rupakheti, M., Panday, A. K., Bhardwaj, P., Naja, M., Singh, A., Mues, A., Cristofanelli, P., Pudasainee, D., 800 Bonasoni, P., and Lawrence, M. G.: Observation and analysis of spatio-temporal characteristics of surface ozone and carbon monoxide at multiple sites in the Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics Discussion*, 2017, 1-42, 10.5194/acp-2017-709, 2017b.
- Miyazaki, Y., Jung, J., Fu, P., Mizoguchi, Y., Yamanoi, K., and Kawamura, K.: Evidence of formation of submicrometer water-soluble organic aerosols at a deciduous forest site in northern Japan in summer, *Journal of Geophysical Research-Atmospheres*, 117, 10.1029/2012jd018250, 2012.
- 805 Mkoma, S. L., Kawamura, K., and Fu, P. Q.: Contributions of biomass/biofuel burning to organic aerosols and particulate matter in Tanzania, East Africa, based on analyses of ionic species, organic and elemental carbon, levoglucosan and mannosan, *Atmospheric Chemistry and Physics*, 13, 10325-10338, 10.5194/acp-13-10325-2013, 2013.
- Mues, A., Rupakheti, M., Munkel, C., Lauer, A., Bozem, H., Hoor, P., Butler, T., and Lawrence, M. G.: Investigation of the mixing layer height derived from ceilometer measurements in the Kathmandu Valley and implications for local air quality, *Atmospheric Chemistry and Physics* 17, 8157-8176, 10.5194/acp-17-8157-2017, 2017.
- 810 Mues, A., Lauer, A., Lupascu, A., Rupakheti, M., Kuik, F., and Lawrence, M. G.: WRF and WRF-Chem v3.5.1 simulations of meteorology and black carbon concentrations in the Kathmandu Valley, *Geoscientific Model Development*, 11, 2067-2091, 10.5194/gmd-11-2067-2018, 2018.
- 815 Muzzini, E., and Aparicio, G.: *Urban Growth and Spatial Transition in Nepal: An Initial Assessment.*, World Bank Publications., 2013.
- Myers-Pigg, A. N., Griffin, R. J., Louchouart, P., Norwood, M. J., Sterne, A., and Cevik, B. K.: Signatures of Biomass Burning Aerosols in the Plume of a Saltmarsh Wildfire in South Texas, *Environmental Science & Technology*, 50, 9308-9314, 10.1021/acs.est.6b02132, 2016.
- 820 Myriokefalitakis, S., Fanourgakis, G., and Kanakidou, M.: The Contribution of Bioaerosols to the Organic Carbon Budget of the Atmosphere, in: *Perspectives on Atmospheric Sciences*, Cham, 2017, 845-851.
- Oanh, N. T. K., Hang, N. T., Aungsiri, T., Worrarat, T., and Danutawat, T.: Characterization of Particulate Matter Measured at Remote Forest Site in Relation to Local and Distant Contributing Sources, *Aerosol and Air Quality Research*, 16, 2671-2684, 10.4209/aaqr.2015.12.0677, 2016.
- 825 Panday, A. K., and Prinn, R. G.: Diurnal cycle of air pollution in the Kathmandu Valley, Nepal: Observations, *Journal of Geophysical Research-Atmospheres*, 114, 10.1029/2008jd009777, 2009.
- Pudasainee, D., Sapkota, B., Shrestha, M. L., Kaga, A., Kondo, A., and Inoue, Y.: Ground level ozone concentrations and its association with NO<sub>x</sub> and meteorological parameters in Kathmandu valley, Nepal, *Atmospheric Environment*, 40, 8081-8087, 10.1016/j.atmosenv.2006.07.011, 2006.
- 830 Putero, D., Cristofanelli, P., Marinoni, A., Adhikary, B., Duchi, R., Shrestha, S. D., Verza, G. P., Landi, T. C., Calzolari, F.,

Busetto, M., Agrillo, G., Biancofiore, F., Di Carlo, P., Panday, A. K., Rupakheti, M., and Bonasoni, P.: Seasonal variation of ozone and black carbon observed at Paknajol, an urban site in the Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics*, 15, 13957-13971, 10.5194/acp-15-13957-2015, 2015.

835 Puxbaum, H., and Tenze-Kunit, M.: Size distribution and seasonal variation of atmospheric cellulose, *Atmospheric Environment*, 37, 3693-3699, 10.1016/s1352-2310(03)00451-5, 2003.

Rajput, N., and Lakhani, A.: Measurements of polycyclic aromatic hydrocarbons in an urban atmosphere of Agra, India, *Atmosfera*, 23, 165-183, 2010.

840 Ram, K., Sarin, M. M., and Hegde, P.: Long-term record of aerosol optical properties and chemical composition from a high-altitude site (Manora Peak) in Central Himalaya, *Atmospheric Chemistry and Physics*, 10, 11791-11803, 10.5194/acp-10-11791-2010, 2010.

Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J. T., Washington, W. M., Fu, Q., Sikka, D. R., and Wild, M.: Atmospheric brown clouds: Impacts on South Asian climate and hydrological cycle, *Proceedings of the National Academy of Sciences of the United States of America*, 102, 5326-5333, 10.1073/pnas.0500656102, 2005.

845 Ramanathan, V., Li, F., Ramana, M. V., Praveen, P. S., Kim, D., Corrigan, C. E., Nguyen, H., Stone, E. A., Schauer, J. J., Carmichael, G. R., Adhikary, B., and Yoon, S. C.: Atmospheric brown clouds: Hemispherical and regional variations in long-range transport, absorption, and radiative forcing, *Journal of Geophysical Research-Atmospheres*, 112, 10.1029/2006jd008124, 2007.

Reisen, F., Meyer, C. P., and Keywood, M. D.: Impact of biomass burning sources on seasonal aerosol air quality, *Atmospheric Environment*, 67, 437-447, 2013.

850 Rupakheti, D., Adhikary, B., Praveen, P. S., Rupakheti, M., Kang, S., Mahata, K. S., Naja, M., Zhang, Q., Panday, A. K., and Lawrence, M. G.: Pre-monsoon air quality over Lumbini, a world heritage site along the Himalayan foothills, *Atmospheric Chemistry and Physics*, 17, 11041-11063, 10.5194/acp-17-11041-2017, 2017.

855 Rupakheti, M., Panday, A. K., Lawrence, M. G., Kim, S. W., Sinha, V., Kang, S. C., Naja, M., Park, J. S., Hoor, P., Holben, B., Sharma, R. K., Mues, A., Mahata, K. S., Bhardwaj, P., Sarkar, C., Rupakheti, D., Regmi, R. P., and Gustafsson, Ö.: Air pollution in the Himalayan foothills: overview of the SusKat-ABC international air pollution measurement campaign in Nepal, *Atmospheric Chemistry and Physics*, in preparation, 2018.

860 Sang, X.-F., Chan, C.-Y., Engling, G., Chan, L.-Y., Wang, X.-M., Zhang, Y.-N., Shi, S., Zhang, Z.-S., Zhang, T., and Hu, M.: Levoglucosan enhancement in ambient aerosol during springtime transport events of biomass burning smoke to Southeast China, *Tellus Series B-Chemical and Physical Meteorology*, 63, 129-139, 10.1111/j.1600-0889.2010.00515.x, 2011.

Sang, X., Zhang, Z., Chan, C., and Engling, G.: Source categories and contribution of biomass smoke to organic aerosol over the southeastern Tibetan Plateau, *Atmospheric Environment*, 78, 113-123, 2013.

865 Sarkar, C., Sinha, V., Kumar, V., Rupakheti, M., Panday, A., Mahata, K. S., Rupakheti, D., Kathayat, B., and Lawrence, M. G.: Overview of VOC emissions and chemistry from PTR-TOF-MS measurements during the SusKat-ABC campaign: high acetaldehyde, isoprene and isocyanic acid in wintertime air of the Kathmandu Valley, *Atmospheric Chemistry and Physics*, 16, 3979-4003, 2016.

Sarkar, C., Sinha, V., Sinha, B., Panday, A. K., Rupakheti, M., and Lawrence, M. G.: Source apportionment of NMVOCs in the Kathmandu Valley during the SusKat-ABC international field campaign using positive matrix factorization, *Atmospheric Chemistry and Physics*, 17, 8129-8156, 10.5194/acp-17-8129-2017, 2017.

870 Shah, M. H., and Shaheen, N.: Annual and seasonal variations of trace metals in atmospheric suspended particulate matter in Islamabad, Pakistan, *Water, air, and soil pollution*, 190, 13-25, 2008.

Shakya, K. M., Rupakheti, M., Aryal, K., and Peltier, R. E.: Respiratory Effects of High Levels of Particulate Exposure in a Cohort of Traffic Police in Kathmandu, Nepal, *Journal of Occupational & Environmental Medicine*, 58, 1, 2016.

875 Shakya, K. M., Rupakheti, M., Shahi, A., Maskey, R., Pradhan, B., Panday, A., Puppala, S. P., Lawrence, M., and Peltier, R. E.: Near-road sampling of PM<sub>2.5</sub>, BC, and fine-particle chemical components in Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics*, 17, 6503-6516, 10.5194/acp-17-6503-2017, 2017.

Sheesley, R. J., Schauer, J. J., Chowdhury, Z., Cass, G. R., and Simoneit, B. R. T.: Characterization of organic aerosols emitted from the combustion of biomass indigenous to South Asia, *Journal of Geophysical Research-Atmospheres*, 108, 10.1029/2002jd002981, 2003.

880 Shen, R. Q., Ding, X., He, Q. F., Cong, Z. Y., Yu, Q. Q., and Wang, X. M.: Seasonal variation of secondary organic aerosol

tracers in Central Tibetan Plateau, *Atmospheric Chemistry and Physics*, 15, 8781-8793, 10.5194/acp-15-8781-2015, 2015.

Simoneit, B. R. T.: Biomass burning - A review of organic tracers for smoke from incomplete combustion, *Applied Geochemistry*, 17, 129-162, 10.1016/s0883-2927(01)00061-0, 2002.

885 Simoneit, B. R. T., Elias, V. O., Kobayashi, M., Kawamura, K., Rushdi, A. I., Medeiros, P. M., Rogge, W. F., and Didyk, B. M.: Sugars - Dominant water-soluble organic compounds in soils and characterization as tracers in atmospheric particulate matter, *Environmental Science & Technology*, 38, 5939-5949, 10.1021/es0403099, 2004.

890 Stockwell, C. E., Christian, T. J., Goetz, J. D., Jayarathne, T., Bhawe, P. V., Praveen, P. S., Adhikari, S., Maharjan, R., DeCarlo, P. F., Stone, E. A., Saikawa, E., Blake, D. R., Simpson, I. J., Yokelson, R. J., and Panday, A. K.: Nepal Ambient Monitoring and Source Testing Experiment (NAMASTE): emissions of trace gases and light-absorbing carbon from wood and dung cooking fires, garbage and crop residue burning, brick kilns, and other sources, *Atmospheric Chemistry and Physics*, 16, 11043-11081, 10.5194/acp-16-11043-2016, 2016.

895 Stone, E. A., Zhou, J., Snyder, D. C., Rutter, A. P., Mieritz, M., and Schauer, J. J.: A Comparison of Summertime Secondary Organic Aerosol Source Contributions at Contrasting Urban Locations, *Environmental Science & Technology*, 43, 3448-3454, 10.1021/es8025209, 2009.

Stone, E. A., Schauer, J. J., Pradhan, B. B., Dangol, P. M., Habib, G., Venkataraman, C., and Ramanathan, V.: Characterization of emissions from South Asian biofuels and application to source apportionment of carbonaceous aerosol in the Himalayas, *Journal of Geophysical Research-Atmospheres*, 115, 10.1029/2009jd011881, 2010.

900 Stone, E. A., Nguyen, T. T., Pradhan, B. B., and Dangol, P. M.: Assessment of biogenic secondary organic aerosol in the Himalayas, *Environmental Chemistry*, 9, 263-272, 10.1071/en12002, 2012.

Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6640-6645, 10.1073/pnas.0911114107, 2010.

905 Szidat, S., Jenk, T. M., Synal, H.-A., Kalberer, M., Wacker, L., Hajdas, I., Kasper-Giebl, A., and Baltensperger, U.: Contributions of fossil fuel, biomass-burning, and biogenic emissions to carbonaceous aerosols in Zurich as traced by 14C, *Journal of Geophysical Research: Atmospheres*, 111, doi:10.1029/2005JD006590, 2006.

910 Tripathee, L., Kang, S., Rupakheti, D., Cong, Z., Zhang, Q., and Huang, J.: Chemical characteristics of soluble aerosols over the central Himalayas: insights into spatiotemporal variations and sources, *Environmental Science and Pollution Research*, 24, 24454-24472, 10.1007/s11356-017-0077-0, 2017.

Wan, E. C. H., and Yu, J. Z.: Analysis of sugars and sugar polyols in atmospheric aerosols by chloride attachment in liquid chromatography/negative ion electrospray mass spectrometry, *Environmental Science & Technology*, 41, 2459-2466, 10.1021/es062390g, 2007.

915 Wan, X., Kang, S., Wang, Y., Xin, J., Liu, B., Guo, Y., Wen, T., Zhang, G., and Cong, Z.: Size distribution of carbonaceous aerosols at a high-altitude site on the central Tibetan Plateau (Nam Co Station, 4730 m a.s.l.), *Atmospheric Research*, 153, 155-164, 10.1016/j.atmosres.2014.08.008, 2015.

920 Wan, X., Kang, S., Li, Q., Rupakheti, D., Zhang, Q., Guo, J., Chen, P., Tripathee, L., Rupakheti, M., Panday, A. K., Wang, W., Kawamura, K., Gao, S., Wu, G., and Cong, Z.: Organic molecular tracers in the atmospheric aerosols from Lumbini, Nepal, in the northern Indo-Gangetic Plain: influence of biomass burning, *Atmospheric Chemistry and Physics*, 17, 8867-8885, 10.5194/acp-17-8867-2017, 2017.

Wan, X., Kawamura, K., Ram, K., Kang, S., Loewen, M., Gao, S., Wu, G., Fu, P., Zhang, Y., Bhattarai, H., and Cong, Z.: Aromatic acids as biomass-burning tracers in atmospheric aerosols and ice cores: A review, *Environmental Pollution*, 247, 216-228, 2019.

925 Wang, W., Wu, M. H., Li, L., Zhang, T., Liu, X. D., Feng, J. L., Li, H. J., Wang, Y. J., Sheng, G. Y., Claeys, M., and Fu, J. M.: Polar organic tracers in PM<sub>2.5</sub> aerosols from forests in eastern China, *Atmospheric Chemistry and Physics*, 8, 7507-7518, 2008.

Wang, X., Gong, P., Sheng, J., Joswiak, D. R., and Yao, T.: Long-range atmospheric transport of particulate Polycyclic Aromatic Hydrocarbons and the incursion of aerosols to the southeast Tibetan Plateau, *Atmospheric Environment*, 115, 124-131, 10.1016/j.atmosenv.2015.04.050, 2015.

930 Wester, P., Mishra, A., Mukherji, A., and Shrestha, A. B. e.: The Hindu Kush Himalaya Assessment: Mountains, Climate

Change, Sustainability and People, Springer Nature Switzerland AG, Cham, 2019.

WHO: Air quality guidelines: global update 2005, World Health Organization, 2006.

935 Wu, G., Wan, X., Gao, S., Fu, P., Yin, Y., Li, G., Zhang, G., Kang, S., Ram, K., and Cong, Z.: Humic-Like Substances (HULIS) in Aerosols of Central Tibetan Plateau (Nam Co, 4730 m asl): Abundance, Light Absorption Properties, and Sources, *Environmental Science & Technology*, 52, 7203-7211, 10.1021/acs.est.8b01251, 2018.

940 Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H., Hering, S. V., de Gouw, J., Baumann, K., Lee, S.-H., Nenes, A., Weber, R. J., and Ng, N. L.: Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States, *Proceedings of the National Academy of Sciences of the United States of America*, 112, 10.1073/pnas.1512279112, 2015.

Yttri, K. E., Dye, C., and Kiss, G.: Ambient aerosol concentrations of sugars and sugar-alcohols at four different sites in Norway, *Atmospheric Chemistry and Physics*, 7, 4267-4279, 2007.

Yue, S., Ren, H., Fan, S., Sun, Y., Wang, Z., and Fu, P.: Springtime precipitation effects on the abundance of fluorescent biological aerosol particles and HULIS in Beijing, *Scientific Reports*, 6, 10.1038/srep29618, 2016.

945 Zhang, H., Surratt, J. D., Lin, Y. H., Bapat, J., and Kamens, R. M.: Effect of relative humidity on SOA formation from isoprene/NO photooxidation: enhancement of 2-methylglyceric acid and its corresponding oligoesters under dry conditions, *Atmospheric Chemistry and Physics*, 11, 6411-6424, 10.5194/acp-11-6411-2011, 2011.

Zhang, Y., Shao, M., Zhang, Y., Zeng, L., He, L., Zhu, B., Wei, Y., and Zhu, X.: Source profiles of particulate organic matters emitted from cereal straw burnings, *Journal of Environmental Sciences*, 19, 167-175, 10.1016/s1001-0742(07)60027-8, 2007.

950 Zhang, Y. L., Huang, R. J., El Haddad, I., Ho, K. F., Cao, J. J., Han, Y., Zotter, P., Bozzetti, C., Daellenbach, K. R., Canonaco, F., Slowik, J. G., Salazar, G., Schwikowski, M., Schnelle-Kreis, J., Abbaszade, G., Zimmermann, R., Baltensperger, U., Prevot, A. S. H., and Szidat, S.: Fossil vs. non-fossil sources of fine carbonaceous aerosols in four Chinese cities during the extreme winter haze episode of 2013, *Atmospheric Chemistry and Physics*, 15, 1299-1312, 10.5194/acp-15-1299-2015, 2015.

955 Zhu, C., Kawamura, K., and Kunwar, B.: Organic tracers of primary biological aerosol particles at subtropical Okinawa Island in the western North Pacific Rim, *Journal of Geophysical Research-Atmospheres*, 120, 5504-5523, 10.1002/2015jd023611, 2015.

960 Zhu, C., Kawamura, K., Fukuda, Y., Mochida, M., and Iwamoto, Y.: Fungal spores overwhelm biogenic organic aerosols in a midlatitudinal forest, *Atmospheric Chemistry and Physics*, 16, 7497-7506, 10.5194/acp-16-7497-2016, 2016.

962 Table 1 Concentrations of TSP, major ions, OC, EC ( $\mu\text{g m}^{-3}$ ) and molecular tracers in  
 963 the aerosols from Bode, Kathmandu Valley ( $\text{ng m}^{-3}$ ).

Compounds	Annual			Pre-monsoon			Monsoon			Post-monsoon			Winter		
	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD
TSP	256	213	166	381	366	171	120	107	52.4	225	254	71.6	353	348	68.5
SO <sub>4</sub> <sup>2-</sup>	10.8	6.15	9.83	17.2	16	7.49	4.1	2.6	4.04	4.81	4.24	2.38	27.3	24.1	6.79
NO <sub>3</sub> <sup>-</sup>	5.21	3.8	4.35	8.82	8.41	4.41	2.34	1.85	1.65	3.52	3.65	0.92	9.47	8.11	4.43
NH <sub>4</sub> <sup>+</sup>	5.92	3.46	6.16	8.57	6.71	5.65	1.99	1.31	2.58	3.72	3.65	1.48	17.5	15.6	3.06
Ca <sup>2+</sup>	7.96	5.82	6.85	11.6	8.98	8.97	4.47	3.59	3.66	6.21	6.01	1.51	13.8	15.6	5.02
K <sup>+</sup>	2.43	1.5	2.82	3.87	2.42	4.27	1.15	0.94	0.96	1.97	2.03	0.46	4.03	3.39	1.45
Cl <sup>-</sup>	2.15	1.18	2.25	2.72	2.27	1.83	0.73	0.67	0.32	1.68	1.78	0.4	6.94	7.87	2.23
Na <sup>+</sup>	3.28	2.93	1.58	3.15	1.68	2.38	3.3	3.61	1.18	2.8	2.84	0.12	4.21	4.28	0.49
Mg <sup>2+</sup>	0.61	0.39	0.54	0.95	0.67	0.7	0.32	0.24	0.25	0.41	0.4	0.07	1.07	1.09	0.39
OC	38.7	24.2	32.7	59.4	46.9	37.9	14.6	14.7	3.76	31.8	32.9	12.7	62.8	53.6	20.6
EC	9.92	9.34	5.33	14.4	13.85	5.24	5.61	4.76	1.8	9.37	9.95	2.19	11	9.89	3.48
OC/EC	3.78	3.09	2.37	4.44	3.29	3.23	2.71	2.54	0.69	3.31	3.01	0.93	5.86	5.3	1.75
<b>Anhydrosugars</b>															
Levoglucosan	788	631	685	1214	900	705	204	188	60.1	863	836	252	1391	1120	535
Galactosan	44.4	33.8	40.3	68.5	51.4	42.7	13.2	13.1	5.55	34.8	33.8	10.9	85.6	72.4	32.8
Mannosan	50.6	34.5	45.1	71.8	61.8	42.3	16	16.7	5.5	39.5	39.6	12.5	116	106	45
Subtotal	883	688	765	1354	974	781	233	218	68.2	937	922	273	1592	1268	611
<b>Monosaccharides</b>															
Glucose	124	114	60	137	118	65.1	129	122	47	143	149	48.7	39.9	38.7	12.5
Fructose	58.2	53.2	28.3	68.4	66.9	29.1	57.1	48.2	26.8	58.2	54.7	21	27	22.5	10.2
Trehalose	40.8	35.6	22	48	53.1	22.4	40.2	35.5	22.8	38.9	40.1	12.5	20.1	16.8	9.17
Sucrose	48.3	40.3	27.4	64.7	55.5	31.6	38.7	38.7	11.9	56.6	45	27.1	18.4	16.6	6.37
Xylose	26.5	20.5	18.1	37.8	30.9	20	13.2	13.4	5.68	24.9	26.5	9.33	38.6	37.5	14.1
Subtotal	298	285	127	356	333	141	278	249	93.5	322	318	104	144	135	35.8
<b>Sugar alcohols</b>															
Mannitol	86.9	77	55.3	84.6	78.8	38.7	114	102	61.4	63.4	53.6	34.6	18.1	19.3	6.02
Arabitol	68.4	60.6	39.8	68.9	65.3	30.3	86.6	68.9	44.5	42.6	33.5	22.5	26.1	24.7	9.13
Sorbitol	14.2	12.7	8.02	13.1	12.6	5.96	17.9	15.9	9.31	13.1	13.2	4.65	5.82	4.97	2.72
Erythritol	43.1	36.5	28.8	35.8	34	15.4	56.9	47.5	33.1	48.6	39.1	31.8	12.4	10.2	7.6
Subtotal	213	192	126	202	198	84.8	275	245	143	168	151	80.1	62.5	61.8	19.7
Total sugars	1394	1206	813	1913	1570	919	787	727	249	1427	1367	327	1798	1462	651
<b>Phenolic compounds and resin acid</b>															
Vanillic acid	15.3	11.3	11.3	20.8	15.8	12.9	7.1	6.84	1.94	14.3	12.3	4.95	26.9	30.5	9.37
Syringic acid	17.1	11.6	13.7	23.7	17.1	15.6	7.82	7.52	2.67	13.7	12.5	4.54	32	37.8	12.3
<i>p</i> -Hydroxybenzoic acid	19.8	15.5	12.3	26.1	19.3	16.7	14.5	13	4.57	14.2	14.6	3.76	23.6	23.1	9.15
Dehydroabietic acid	13.8	12.6	6.19	16.3	15.4	6.52	10.3	9.7	2.32	10.9	9.3	3.1	21.4	20.1	6.76
Subtotal	66.1	49.7	41.4	86.9	67	49.2	39.7	39.6	9.22	53	47.5	15.6	104	112.8	36.2



Compounds	Annual			Pre-monsoon			Monsoon			Post-monsoon			Winter		
	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD	Mean	Median	SD
<b>Phthalic acid esters</b>															
Diethyl (DEP)	16.6	15.6	8.41	19.4	17.5	11.6	15.4	15.5	4.42	14.9	13	6.25	12.9	9.3	6.49
Di-n-butyl (DnBP)	56.2	48.5	25.6	63.6	59.8	30.7	52.2	48.4	20.9	55.4	45.3	22.8	46.4	41.3	21.8
Bis-(2-ethylhexyl) (DEHP)	438	378	200	495	466	239	407	377	162	431	353	177	361	322	170
Subtotal	510	444	230	578	545	276	474	445	184	501	408	204	420	372	196
<b>Isoprene tracers</b>															
2-Methylglyceric acid	34.2	30.2	14.8	45.9	48.8	15.2	25.1	24.5	6.91	36.2	34.9	9.86	25	24.7	8.56
2-Methylthreitol	30.4	27.3	19.5	22	22.8	10	45	44.9	20.6	27.4	26.3	7.39	8.4	7.9	3.83
2-Methylerythritol	64.1	58.3	39.6	45	44.5	20.4	97.3	98.9	38	53	52	9.78	18.5	18.7	4.63
2-Methylterols <sup>a</sup>	94.4	84.3	58.9	67	68.1	30.4	142	142	58.3	80.5	78.3	17	27	26.6	8.39
C5-Alkene triols <sup>b</sup>	45	39.2	29.4	30.4	28.8	13.4	68.8	69.1	30	35.1	30.6	18	17.5	15.3	4.88
Subtotal	174	160	86.2	144	150	53.4	236	220	87.2	152	145	39.3	69.5	72.4	19.3
<b>Monoterpene tracers</b>															
cis-Pinonic acid	26	24.5	11.6	32.3	30	13.8	21	19.1	8.04	28	29.1	5.92	20.6	20	8.01
Pinic acid	11.9	10.8	4.48	11.4	10.4	3.78	12.7	12.3	5.07	13.7	15.6	5.25	8.83	8.67	0.92
3-Hydroxyglutaric acid	10.6	9.16	6.85	13.9	12.1	7.75	5.75	4.88	3.01	16	15.4	4.18	11.5	10	4.72
3-MBTCA <sup>c</sup>	10.8	9.53	7.36	16.9	15.1	7.32	5.14	4.3	3.12	11.9	9.61	3.63	9.3	8.89	3.75
Subtotal	59.3	55.6	24.6	74.6	72.1	28.3	44.6	42.6	14.4	69.5	65.7	12.6	50.2	49.6	10.8
<b>Sesquiterpene tracer</b>															
$\beta$ -Caryophyllenic acid	6.43	5.35	3.93	8.61	7.15	4.29	3.66	2.85	2	8.35	7.2	3.02	7.16	6.6	2.76
Total B-SOA tracers <sup>d</sup>	136	234	50.4	133	229	43.9	156	275	53.9	131	217	26.2	75.6	129	14.2
<b>Toluene tracer</b>															
DHOPA <sup>e</sup>	19.4	16.5	11.5	22.8	19	15.5	15	13.4	5.69	17.3	18.2	5.94	25.8	23	9.84
Total SOA tracers <sup>f</sup>	259	249	94.7	250	242	84.9	299	286	101	247	243	48.2	153	162	29.4

965 <sup>a</sup> Sum of 2-methylthreitol and 2-methylerythritol.

966 <sup>b</sup> C5-Alkene triols: 3-Methyl-2,3,4-trihydroxy-1-butene, cis-2-Methyl-1,3,4-trihydroxy-1-butene and trans-2-Methyl-1,3,4-trihydroxy-1-butene.

967 <sup>c</sup> 3-MBTCA: 3-methyl-1,2,3-butanetricarboxylic acid.

968 <sup>d</sup> Sum of 2-methylglyceric acid, 2-methylterols, C5-Alkene triols, cis-pinonic acid, pinic acid, 3-hydroxyglutaric acid and 3-MBTCA.

969 <sup>e</sup> DHOPA: 2,3-dihydroxy-4-oxopentanoic acid.

970 <sup>f</sup> Sum of 2-methylglyceric acid, 2-methylterols, C5-Alkene triols, cis-pinonic acid, pinic acid, 3-hydroxyglutaric acid and 3-MBTC and 2,3-dihydroxy-4-oxopentanoic acid.

971

972

973

974

975 Table 2 Linear correlation coefficients ( $R^2$ ) among major ions and OC, EC in aerosols in Bode, Kathmandu Valley

	SO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	NH <sub>4</sub> <sup>+</sup>	Ca <sup>2+</sup>	nss-Ca <sup>2+</sup>	Mg <sup>2+</sup>	nss-Mg <sup>2+</sup>	K <sup>+</sup>	nss-K <sup>+</sup>	Cl <sup>-</sup>	Na <sup>+</sup>	OC	EC
SO <sub>4</sub> <sup>2-</sup>	1.00												
NO <sub>3</sub> <sup>-</sup>	0.78**	1.00											
NH <sub>4</sub> <sup>+</sup>	0.87**	0.69**	1.00										
Ca <sup>2+</sup>	0.48**	0.58**	0.62**	1.00									
nss-Ca <sup>2+</sup>	0.48**	0.58**	0.62**	1.00**	1.00								
Mg <sup>2+</sup>	0.61**	0.71**	0.69**	0.91**	0.91**	1.00							
nss-Mg <sup>2+</sup>	0.65**	0.76**	0.70**	0.88**	0.88**	1.00**	1.00						
K <sup>+</sup>	0.40**	0.59**	0.55**	0.78**	0.78**	0.85**	0.83**	1.00					
nss-K <sup>+</sup>	0.40**	0.60**	0.55**	0.77**	0.77**	0.85**	0.83**	1.00**	1.00				
Cl <sup>-</sup>	0.67**	0.58**	0.76**	0.41**	0.40**	0.48**	0.50**	0.37**	0.37**	1.00			
Na <sup>+</sup>	0.05*	0.08*	0.23**	0.48**	0.48**	0.37**	0.28**	0.40**	0.39**	0.09*	1.00		
OC	0.59**	0.61**	0.57**	0.32**	0.32**	0.48**	0.53**	0.36**	0.36**	0.48**	0.01	1.00	
EC	0.35**	0.43**	0.24**	0.12**	0.12**	0.23**	0.27**	0.16**	0.16**	0.24**	0.02	0.36**	1.00

976

977 \*: P<0.1.

978 \*\*: P<0.001.

979 Table 3 Linear correlation coefficients ( $R^2$ ) among monosaccharides and sugar alcohols in aerosols from Bode, Kathmandu Valley

	Glucose	Fructose	Sucrose	Trehalose	Xylose	Mannitol	Arabitol	Sorbitol	Erythritol
Glucose	1								
Fructose	0.77***	1							
Sucrose	0.55***	0.55***	1						
Trehalose	0.30***	0.23***	0.28***	1					
Xylose	0.001	0.06*	0.19***	0.05*	1				
Mannitol	0.51***	0.51***	0.23***	0.14***	0.002 <sup>a</sup>	1			
Arabitol	0.44***	0.50***	0.22***	0.16***	0.0005	0.77***	1		
Sorbitol	0.51***	0.53***	0.20***	0.13**	0.0002	0.83***	0.68***	1	
Erythritol	0.46***	0.40***	0.16***	0.13**	0.007	0.77***	0.62***	0.77***	1


980

981 a Negative value indicate negative correlation.

982 \* $p < 0.1$

983 \*\* $p < 0.01$ .


984 \*\*\* $p < 0.001$ .




985

986 **Fig. 1.** Location of measurement site: (a) Kathmandu Valley, (b) urban measurement site at Bode in  
987 Kathmandu Valley, (c) air pollution observed from the Bode site in the afternoon.

988




989  
 990 **Fig. 2.** Monthly variations of TSP, OC, EC, OC/EC ratios at Bode site, Kathmandu Valley during  
 991 April 2013-April 2014.




**Fig. 3.** Monthly variations of biomass burning tracers, monosaccharides and sugar alcohols at Bode site, Kathmandu Valley during April 2013-April 2014.

992  
993  
994



995


996 **Fig. 4.** Correlations between (a) levoglucosan and OC, (b) levoglucosan and EC, (c) levoglucosan and  
 997 nss-K<sup>+</sup> in Bode aerosols during the sampling period (April 2013 to April 2014).



998


999 **Fig. 5.** Monthly variations of B-SOA tracers, total isoprene tracers, total monoterpene tracers,  $\beta$ -  
 1000 caryophyllenic acid, total B-SOA tracers and DHPOA at Bode site, Kathmandu Valley during April  
 1001 2013-April 2014.





1002


1003 **Fig. 6.** The percentage contributions of the isoprene SOA tracers to the total during different seasons in  
 1004 the atmospheric aerosols from Kathmandu.



1005


1006

**Fig. 7.** Ratios of 2-MGA/2-MTLs during different seasons in Bode, Kathmandu.




1007

1008 **Fig. 8.** Concentration correlation between (a) 2-methylglyceric acid (2-MGA) and  $\text{SO}_4^{2-}$ , (b) 2-  
1009 methylglyceric acid and  $\text{NO}_3^-$  in the aerosols from Bode, Kathmandu.




1010

1011 **Fig. 9.** Correlation between 2,3-dihydroxy-4-oxopentanoic acid (DHOPA) and levoglucosan in Bode  
 1012 aerosols during the sampling period (April 2013 to April 2014).



1013

1014 **Fig. 10.** Monthly concentration variations of (a) BB-OC, (b) plant-debris-OC, (c) fungal-spore-OC, (d)  
 1015 SOC-isoprene (I-SOC), (e) SOC-monoterpenes (M-SOC), (f) SOC-sequiterpene (SOC-C), (g) SOC-  
 1016 isoprene+monoterpenes+sequiterpene (B-SOC), (h) SOC-toluene (A-SOC), and (i) total SOC that were  
 1017 estimated using a tracer-based method at Bode site, Kathmandu Valley during April 2013-April 2014.



1018

1019

1020

1021

**Fig. 11.** Pie-charts showing contributions from different sources to organic carbon based on the estimation of tracer-method in Bode, Kathmandu Valley: (a) annual, (b) pre-monsoon, (c) monsoon, (d) post-monsoon and (e) winter.