Impact of upstream moisture structure on a back-building convective precipitation system in south-eastern France during HyMeX IOP13

By K. O. Lee et al.

Reply to the referees' comments

In the following, the comments made by the referees appear in black, while our replies are in red, and the proposed modified text in the typescript is in blue.

Referee #1 comments

General Comments

I thank the authors for considering my comments and for making an effort to modify their manuscript accordingly. My major concern about the first draft was related to the unclear explanation of the impact of free-atmospheric moisture on cold-pool intensity. From the current draft, I understand that the weakening of the cold pool (leading to the observed reduction of precipitation in the FAxM experiments) was caused mainly by the reduced moisture supply upstream of the convective system in the 1-2 km layer, which led to lower rainfall rates. This seems a reasonable explanation, and it is well supported by the new Figure 16. Although the authors don't mention it explicitly, it can be concluded that (in the FAxM experiments) the expected enhancement of rainfall evaporation in the drier 1-2 km layer was relatively unimportant.

I think it now makes sense to recommend the manuscript for publication. However, a few minor aspects should still be fixed before acceptance.

The authors are grateful for reviewer's interest in this study and the many helpful suggestions for improving this manuscript. Replies to each minor comments are listed below.

Minor comments

1) P4, L29: Reference to "Collela and Woodward, 1984". I believe the family name of the first author should be changed to "Colella".

Yes, indeed. This is now corrected (P4, L29) also in the list of reference (P19, L12).

2) P5, L23: "Bubble initiation". The authors are describing the control run here, so there should be no sensitivity bubble. If "bubble" is used improperly as a synonym of "convective ascent" (or of similar expressions), please fix. Corrected by removing the "bubble initiation" in the sentence as:

"The intense precipitation (\geq 15 mm per 15 min, red circle in Fig. 2b) is first simulated at 1400 UTC north of Marseille (5.6°E, 43.5°N) [...]"

3) P14, L4-6: "While the dried air mass in the 1–2 km layer is advected toward upstream the convection, it mixes with the moist air below, reducing the total moisture below 2 km". This sentence is somehow confusing. Mixing between the 0-1 km layer and the 1-2 km layer cannot reduce the total moisture below 2 km (mixing redistributes a quantity, but it can't remove it from the system). Advection of relatively dry air, instead, does reduce moisture. I think the authors mean the following: Considering the region immediately upstream of convective ascents, the total moisture below 2 km in the FAxM simulations is lower than in CNTL (or decreases over time), due to the advection of the sensitivity bubble containing dry air at heights between 1 and 2 km. Please check if this interpretation is correct and, if so, clarify the text.

Thanks, this is exactly what we meant. For the sake of clarity, we have corrected the sentence as you have suggested (P14, L3-5).

"[...]Considering the region immediately upstream of convective ascents, the total moisture below 2 km in the FA×M simulations is lower than in CNTL, due to the advection of the sensitivity bubble containing dry air at heights between 1 and 2 km."

4) P39, L3: Caption of F16. This figure shows the temporal evolution of area-averaged rainfall rate and virtual potential temperature. The boundaries of the area are reported to be: 5.5–7.5°E, 43–44.5°N. Is the northern boundary actually 43.5 °N? 44.5 seems to be too far to the north.

The northern boundary of 44.5°N is correct. Most of the precipitation produced by CNTL, FA×P, and FA×M stay in the south of 44.5°N, and it is evidenced by Figures 2 and 13.

1	Impact of upstream moisture structure on a back-building convective precipitation
2	system in southeastern France during HyMeX IOP13
3	
4	Keun-Ok Lee ¹ , Cyrille Flamant ² , Fanny Duffourg ³ , Véronique Ducrocq ³ , and Jean-Pierre
5	Chaboureau ¹
6	
7	¹ Laboratoire d'Aérologie, Université de Toulouse, CNRS, UPS, Toulouse, France
8	² LATMOS/IPSL, CNRS, Sorbonne Université and Université Paris-Saclay, Paris, France
9	³ CNRM, UMR 3589, Université de Toulouse, Météo-France & CNRS, Toulouse, France

- 10
- 11

ABSTRACT

12 The present study examines the impact of the environmental moisture structure in the lower troposphere (below 2 km above sea level, ASL) on the precipitation development, observed in southern France during intensive 13 observation period (IOP) 13 of the first Special Observation Period of the Hydrological cycle in the 14 Mediterranean Experiment (HyMeX SOP-1), through a series of sensitivity experiments using the non-15 hydrostatic numerical model Meso-NH. The control simulation (CNTL) and all the other 12 sensitivity 16 experiments examined in this study succeed in reproducing a heavy precipitation event (HPE) in the coastal 17 18 mountainous region of Var in southeastern France as observed. The sensitivity experiments are designed to 19 investigate the response of the HPE to the variability of the water vapour content upstream in the moist marine 20 atmospheric boundary layer (MABL) and the drier air above. The comparisons between CNTL and the 12 sensitivity experiments show how the life cycle of precipitation associated with the HPE, but also the upstream 21 flow (over the sea), is modified, even for moisture contents changes of only 1 g kg⁻¹ below 2 km ASL. Within 22 23 the low-level wind convergence between southerlies and southwesterlies, a small increase of moisture content 24 in the MABL prolongs moderate precipitation (\geq 5 mm in 15 min) and enlarges the area of weak precipitation $(\geq 1 \text{ mm in } 15 \text{ min})$. The moistening in the 1–2 km ASL layer, just above the MABL, prolongs the duration of 25 moderate precipitation, for a similar total precipitation amount as in CNTL. The drier MABL and 1-2 km ASL 26 layer shorten the life-time of precipitation and reduced the total precipitation amount with respect to CNTL. 27 28 We also found that the moisture in the MABL has a stronger impact on producing enhanced precipitation (both

in terms of amount and intensity) than the moisture just above (1-2 km ASL). Also it is worth noting that adding moisture in the MABL does not necessarily lead to enhanced precipitation amount. In moistening MABL, the duration of moderate precipitation increases with increasing moisture as does the area covered by weak precipitations, while the area covered by the intense precipitation ($\geq 30 \text{ mm}$) decreases. Despite a simplified moisture-profile modification approach, this study suggests that moisture structure in the lower troposphere is a key for accurate prediction at short-term range of the timing and location of precipitation in the coastal mountainous region in southern France.

- 8
- 9

Running title: Impact of upstream moisture structure on HPE

10

11 **1. Introduction**

Most of the Mediterranean countries face heavy precipitation events (HPEs), especially during autumn. More than 100 mm of precipitation in less than 6 hours is not uncommon in these regions, and such rainfall accumulations often cause flash-floods with large material damages and important human losses. These highimpact events are most often due to quasi-stationary Mesoscale Convective System (MCS) (*e.g.* Nuissier et al. 2008; Romero et al. 1999; Trapero et al. 2013a, 2013b; Barthlott and Davolio, 2015).

During the autumn, the warm Mediterranean is a significant source of moistening and heating for the 17 18 marine atmospheric boundary layer (MABL) through latent and sensible heat fluxes (Duffourg and Ducrocq, 2011). The southerly low-level marine winds that prevail during HPEs over the northwestern Mediterranean 19 transport this moist and conditionally unstable air toward the coastal mountainous region (Ricard et al. 2012). 20 Such low-level conditions are favoured by typical synoptic patterns. Nuissier et al. (2011) found that the 21 22 synoptic-scale pattern propitious to HPEs over the French Mediterranean region consist of an upper-level trough located west of France and an upper-level ridge over central Europe. MCSs preferably develop eastward 23 of a slow-evolving disturbance associated with an upper-level trough and leading a southwesterly diffluent 24 flow. Such a synoptic pattern favours a persistent low-level moist and conditionally unstable marine flow 25 26 directed towards the costal mountainous regions.

When this conditionally unstable low-level flow impinges on some of the mountain range foothills that border the Western Mediterranean, a back-building MCS can be triggered and renewed repeatedly at the same location, as long as the same low-level and upper-level conditions persist. Orographic lifting has been largely

proposed in past studies as a mechanism for triggering HPEs over the western Mediterranean (e.g. Buzzi et al. 1 1998; Houze, 1993; Jansa et al. 2001; Lin, 1993; Romero et al. 2015; Rotunno and Ferretti, 2001; Smith, 1979; 2 3 Trapero et al. 2013a, 2013b; Miglietta and Rotunno, 2014). Furthermore, a low-level cold pool forming under 4 a MCS can also lift the impinging ambient low-level flow at its leading edge (Durocq et al., 2008) or modify the low-level circulation locally and enhance convergence areas (Duffourg et al., 2016). Furthermore, other 5 convection triggering mechanisms, stemming from the low-level marine flows interaction with the complex 6 terrain of the western Mediterranean, have been highlighted (e.g. Ducrocq et al., 2016). The dynamical and 7 8 thermodynamical characteristics of the low-level flows are decisive with respect to the triggering mechanisms 9 involved (Bresson et al. 2012).

10 The water vapour transported by the marine flow is a crucial ingredient of Mediterranean HPEs. Therefore, the realistic representation of its spatio-temporal variability in numerical weather prediction models 11 is critical for HPE forecast. Predicting the initiation of convection in cloud resolving models can also be highly 12 dependent on very accurate estimates of water vapor within and just above the boundary layer (e.g. Crook, 13 1996; Ducrocq et al., 2002; Weckwerth et al., 2004; Bielli et al., 2012). This requires relevant moisture 14 observations in the inflow region or within the convective storms, particularly over the sea. These are difficult 15 to obtain on a regular basis in order to properly constrain numerical weather prediction models in terms of 16 17 moisture.

The first Special Observation Period (SOP-1, Ducrocq et al. 2014) of the Hydrological cycle in the 18 Mediterranean Experiment (HyMeX) took place in autumn 2012, aiming to improve our knowledge of the 19 origin and transport pattern of moist air masses in pre-convective conditions and determine the link between 20 these air masses and HPEs. During Intensive Observation Period 13 (IOP13, 14 October 2012) of HyMeX 21 SOP-1, back-building MCSs developed in southeastern France shortly after 1300 UTC (Duffourg et al., 2018). 22 The synoptic situation of IOP13 is characterised by an upper-level trough associated with a surface disturbance 23 situated over north-western France. Ahead of the associated cold front, a southwesterly low-level flow over 24 25 the Mediterranean French coast brings a warm marine air mass inland. Thanks to aircraft water vapour measurement acquired over the Gulf of Lion (northwestern Mediterranean) during IOP13, Duffourg et al. 26 (2018) detailed the moisture structure upstream the MCSs which was characterised by a moist conditionally 27 unstable MABL (below 1 km above sea level, ASL) topped by dry air masses just above (1-2 km ASL). They 28 emphasized the importance of the low level moisture for feeding the convective systems observed during this 29

HPE and the significance of the dry air above for strengthening the associated cold pools. The goal of the
 present study is to further investigate the impact of moist air in the boundary layer and surrounding dry air
 masses on the development and evolution of MCSs.

The impact of the environmental moisture structure on the development of a convective system (which produced 70 mm of precipitation in 6 hours), observed a few kilometres southwest offshore of the Marseille and Var coastlines in southern France, is investigated here through a series of sensitivity experiments using the French non-hydrostatic numerical research model Meso-NH (Lac et al., 2018) at the horizontal grid spacing of 2.5-km. The article is organized as follows: Section 2 describes the experimental design, the results of the control and sensitivity simulations are presented in sections 3 and 4, and then summarised in section 5 with concluding remarks.

11

12 **2. Numerical simulation**

13 2.1. Meso-NH model configuration

Meso-NH (Lac et al. 2018), mesoscale non-hydrostatic model, has already shown its capability to simulate 14 HPEs in both real and idealized frameworks (Argence et al., 2008; Nuissier et al., 2008; Ducrocq et al., 2008; 15 Clark and Chaboureau, 2010; Bresson et al., 2012). For this study, the same set-up as in Duffourg et al. (2016) 16 and 2018), that proved to be able to reproduce realistic convective systems, was used. It includes the same 17 simulation domain over southeastern France and the north-western Mediterranean (Fig. 1, 200-km × 200-km 18 19 domain encompassing the precipitating systems and their marine low-level moisture-supplying flow), the same horizontal grid spacing of 2.5 km, the same vertical grid with 55 stretched vertical levels (Gal-Chen and 20 21 Somerville, 1975) up to 20 km and the same parametrisation schemes: the bulk 1-moment mixed microphysical scheme (Caniaux et al., 1994; Pinty and Jabouille, 1998) combining a three-class ice parametrisation with a 22 Kessler's scheme for the warm processes, a 1D-turbulence parametrisation based on a 1.5-order closure (Cuxart 23 et al., 2000) of the turbulent kinetic energy equation with the Bougeault and Lacarrere (1989) mixing length, 24 25 the Pergaud et al. (2009) Eddy Diffusivity Mass Flux scheme for shallow convection, the Rapid Radiation Transfer Model (Mlawer et al., 1997) for radiation and the surface model SURFEX (Masson et al. 2013). 26 Transport scheme for momentum variables is a WENO (weighted essentially non-oscillatory) scheme (Shu 27 and Osher, 1988) while other variables are transported with the PPM (Piecewise Parabolic Method) scheme 28 29 (Collella and Woodward, 1984). An open wave radiation condition (Carpenter, 1982) combined with a five1 grid-point relaxation flow scheme (Davies, 1976) are applied at the boundaries.

The control simulation (referred to as CNTL in the following) was initialized at 0900 UTC on 14 October 2012. As in Duffourg et al. (2018), the initial and lateral boundary conditions are provided by the 3-hourly Application of Research to Operational at Mesoscale-West Mediterranean (AROME-WMED) analyses (Fourrié et al., 2015). AROME-WMED is a dedicated version of the French operational convection-permitting numerical prediction system AROME (Seity et al., 2011), set up specifically for the HyMeX field campaigns and covering the western Mediterranean.

8

9 2.2. Control simulation

During IOP13, convection initiation is observed around 1200 UTC, over the sea, a few kilometres offshore the 10 11 coast of southeastern France. It develops into organised convective systems (not shown) in the surroundings of Marseille and along the coast of the Var region (geographical positions are given in Figure 1). A maximum 12 6-hour rainfall accumulation of 67 mm is recorded over land under the convective line anchored along the Var 13 14 coast between 1400 UTC and 1900 UTC on 14 October 2012. The CNTL simulation succeeds in reproducing heavy precipitation over southeastern France. The CNTL simulation starts showing evidence of weak 15 precipitation (< 5 mm per 15 min) and low reflectivity (values than 30 dBZ) around 1200 UTC over the sea 16 (not shown). Then the convection (highlighted by reflectivity values exceeding 45 dBZ, Lee et al., 2012) 17 18 initiates upstream (yellow star, Fig. 2b), about 25 km further south than observed at 1200 UTC and maximum 19 15-minute precipitation over 5 mm is simulated approximately 93 km from the initiation at 1230 UTC (Fig. 20 2b). It develops preferably toward Marseille, then around the Argens valley region and the east coastal Var region (Figs. 2b and 3a–b), similarly as observed. The intense precipitation (≥ 15 mm per 15 min, red circle 21 22 in Fig. 2b) is first simulated at 1400 UTC north of Marseille (5.6°E, 43.5°N) in a location distant of about 95 km from the bubble initiation. The convective regions are simulated in the Argens valley region and the east 23 24 coastal Var region from 1400 to 1600 UTC (Figs. 3b-d). The simulated amount of rainfall accumulation is realistic, with 74.2 mm in 6 hours associated with the convective line over the eastern Var coast (longitude 25 6.5°E, Figure 2a). 26

Figure 4 displays the hourly evolution simulated by CNTL of the equivalent potential temperature, θ_e , at 925 hPa and virtual potential temperature, θ_v , at the first model level (about 10 m), together with the horizontal wind and the vertical velocity at 925 hPa level during the period between 1300 and 1600 UTC.

1 During this period, the marine warm and moist low-level air ($\theta_e > 322$ K) progresses north-eastwards. At 1300 UTC (Fig. 4a) it reaches the Var coast and convective ascents are localized along the coast on the mountain 2 sides. From 1400 UTC on (Figs. 4b–d), the cold air ($\theta_v < 291$ K) formed by evaporative cooling under the 3 intensifying precipitation spreads out over the plains (Fig. 3b-d and 2b). At 1400 UTC very high CAPE 4 (Convective Available Potential Energy) values (i.e. in excess of of 1590 J kg⁻¹) are simulated around the Var 5 coast in CNTL, in good agreement with the CAPE of 1600 J kg⁻¹ derived from radiosounding measurements 6 in Marseille at 1500 UTC (not shown). The convection develops rapidly on the leading edge of the cold air 7 8 channelled in the Argens valley (geophysical positions are given in Figure 1) (Fig. 3c), then it propagates to the eastern Var coast after 1600 UTC (Fig. 3d). Convection is quasi-stationary along the eastern Var coastal 9 region (Fig. 4d). The cold pool plays a major role in determining the location of the precipitation (Duffourg et 10 al., 2018). 11

To characterize the air mass supplying the precipitating system, backward trajectories of air parcels 12 reaching the top of the simulated convective ascents are performed. They show that the convective ascents are 13 fed by the southwesterly marine flow all similarly along the red dashed line drawn in Figure 1. The vertical 14 projection of the backward trajectories (not shown) provides evidence that the flow is confined to the lowest 15 1000 m above the sea surface before being lifted up to the top of the troposphere within 1 hour in the convective 16 ascents of the precipitating system. The low-level flow brings a moist (about 10 g kg⁻¹) and warm (θ_e about 17 322 K) air mass from the north-east of the Balearic Islands at 0900 UTC to the southeastern French coast, 18 19 where it feeds the convective ascent. Another set of backward trajectories is computed to identify the air mass involved in the formation of the cold pool under the intense precipitation in the eastern Var coastal region. All 20 trajectories follow a similar pathway represented by a blue dashed line in Figure 1. Moreover, the air mass 21 involved in cold-air pool formation comes mainly from a dry layer between 1 and 2 km ASL, just above the 22 moist boundary layer, with water vapour mixing ratios (WVMRs) ranging between 3 and 8 g kg⁻¹ (not shown). 23 The water vapour contents upstream of the HPE of IOP13 (over the sea; 3.5–5.5°E, 41–42.5°N) at 0600 24 UTC on 14 October 2012 of AROME-WMED analyses data are compared with that retrieved by the Special 25 Sensor Microwave/Imager (SSM/I) at 0625 UTC of the same day. The horizontal distribution of integrated 26 water vapour (IWV) values between the AROME-WMED analyses and SSM/I over the north-western 27 Mediterranean are in broad agreement, i.e. moister air to the northeast of Balearic Islands and drier air offshore 28 of the southwest French coast (not shown) showing a mean IWV difference of 0.7 kg m^{-2} , with averaged values 29

of 18.2 kg m⁻² in AROME-WMED analyses Vs 18.9 kg m⁻² in SSM/I. In AROME-WMED analyses, near 90 %
 of the IWV concentrates below 2 km ASL, i.e. 69.5% below 1 km ASL, 21 % between 1 and 2 km ASL.

In the following, we will analyse the precipitating pattern and characteristics simulated over the Var coastal region, and their sensitivity to the related upstream moisture environment by comparisons of results between CNTL and 12 sensitivity experiments with varying water vapour content in lower troposphere (below 2 km ASL).

7

8 2.3. Initial conditions of sensitivity experiments

9 To study the sensitivity of IOP13 precipitation to the moisture structure in the low troposphere, two sets of experiments, in which the initial WVMR profiles are modified in the moist MABL (below 1 km ASL) and in 10 11 the dry layer above (1-2 km ASL), are carried out, namely the MBL (marine boundary layer) and FA (free atmosphere) experiments, respectively (Table 1 and Figures 1 and 5). The WVMR profiles are modified in the 12 areas involved in the moisture supply of the precipitating system and in the cold pool formation. Then the 13 14 impact on the precipitation amount and location is analysed. The location and extent of the "sensitivity 15 bubbles" are defined based on the results of backward trajectory analyses (red and blue dashed lines in Fig. 1) done using CNTL. For the six MBL experiments, the "sensitivity bubble" is located over the sea, centred about 16 230 km from the Marseille coastline (red-coloured cross in Fig. 1), with an ellipsoidal shape with a short axis 17 of 80 km and a long axis of 160 km length (red ellipse in Fig. 1). Three MBL experiments were designed, with 18 WVMR values between 100 and 1000 m ASL in the "sensitivity bubble" being increased by 1, 2, and 5 g kg⁻¹ 19 up to the saturation limit (Figure 5a and Table 1a). In other words, the water vapour value at saturation with 20 respect to liquid water was calculated at each altitude and used as an upper threshold of the modified WVMR. 21 22 The sensitivity bubbles are referred to as MBL1P, MBL2P and MBL5P experiments, respectively, in the following. Likewise, three MBL experiments were designed with WVMR values between 100 and 1000 m 23 ASL in the "sensitivity bubble" being decreased by 1, 2, and 5 g kg⁻¹ (MBL1M, MBL2M and MBL5M, 24 respectively, see Figure 5a). 25

For the six FA experiments (Table 1b), the "sensitivity bubble" is centred at about 150 km from the Marseille coastline (blue-coloured cross in Fig. 1), and has an ellipsoidal shape with a short axis of 60 km and a long axis of 120 km length (blue ellipse in Fig. 1). In these experiments, the WVMR values between 1000 and 2000 m ASL in the "sensitivity bubble" are increased (decreased) by 1, 2, and 5 g kg⁻¹ (Fig. 5b) keeping

the minimum WVMR value to 0 g kg⁻¹, and they are referred to as FA1P (FA1M), FA2P (FA2M) and FA5P (FA5M) experiments, respectively. For FA2M and FA5M, the bottom threshold of 0 g kg⁻¹ is used for reformulating the WVMR profiles. Other environmental conditions in MBL and FA experiments are the same as in CNTL. Using backward trajectory analysis, we can assess that the origins of air parcels feeding the convective system and of air parcels feeding the cold pools are within the sensitivity bubbles of the FA and MBL experiments, respectively, (ellipsoids in Fig. 1) and that the trajectories do not deviate significantly from one experiment to the next, even though some differences exist.

8 The magnitude of the WVMR anomalies introduced in the "sensitivity bubbles" are based on recent studies which confronted the quality of the AROME-WMED analyses in the lower troposphere with WVMR 9 profiling instruments operating over the Mediterranean, in particular using ground-based and airborne water 10 vapour lidars (e.g. Chazette et al., 2016; Lee et al., 2016; 2017). Chazette et al. (2016) evidenced that the root 11 mean square error between AROME and lidar WVMRs in the vicinity of Balearic Islands below 2 km ASL 12 range between 1 and 1.6 g kg⁻¹, while Lee et al. (2016) and Lee et al. (2017) highlighted differences as large 13 as 3-4 g kg⁻¹ in the MABL upstream of MCSs over the Tyrrhenian Sea and the Balearic Sea, respectively. In 14 this study, experiments with ± 1 and 2 g kg⁻¹ were conducted to understand the impact of water vapour 15 uncertainty in the vicinity of Balearic Island on precipitation downstream, while the experiments with $\pm 5 \text{ g kg}^-$ 16 ¹ are done to see such an impact in a more extreme environment. 17

To investigate the WVMR impact on the location, intensity and duration of precipitation, we consider 18 19 several indicators, such as: 1) the maximum of the 6-hour accumulated precipitation amount (RR_{max}), the 20 domain-averaged total sum of the 6-hour rainfall accumulation (RR_{sum}), and the RR_{sum} produced over land (RR_{land}) and over the sea (RR_{sea}) in the fixed area (3.5-9°E, 40.5-45°N) from 1200 UTC on 14 October 2012 21 to understand the impact of WVMR variability on precipitation amount, 2) the horizontal area (km²) of 6-hour 22 precipitation $\geq 1 \text{ mm}(AR_{01})$ and $\geq 30 \text{ mm}(AR_{30})$ in CNTL, and its deviation (%) from those in the MBL and 23 FA experiments, and 3) the duration of precipitation (\geq 5 mm) over the land (D_{land}), duration of precipitation \geq 24 5 mm per 15 min (D_{RR05}), and duration of precipitation \geq 15 mm per 15 min (D_{RR15}). Those indicators are 25 presented in Figures 6 and 12 for the different sensitivity experiments (MBL and FA, respectively). Figure 7 26 displays the 6-hour rainfall accumulation simulated in the MBL sensitivity experiments, and Figure 8 show 27 the evolution of the maximum 15-min accumulated precipitation amount (RR_{max}) every 15 min during 6 hours, 28 highlighting the differences in time, intensity and location of the precipitation. Figure 9 shows the temporal 29

evolution of the maximum of CAPE in the "sensitivity bubble" every 15 min from 0930 to 1445 UTC in CNTL,
MBL×M (i.e. MBL1M, MBL2M, and MBL5M), and MBL×P (i.e. MBL1P, MBL2P, and MBL5P). Figures 13
and 14 show the same parameters as Figs. 7 and 8, respectively but for FA experiments. Note that the weak
precipitation amounts (i.e. less than 5 mm per 15 min) are not displayed in Figs. 8 and 14 to focus on active
convective system. Also note that the analysis of the precipitation characteristics is conducted for 6 hours (from
1200 to 1800 UTC) when the convective systems affect the inland and coastal regions. Precipitation continues
over the sea after 1800 UTC in all experiments, as observed.

8

9 **3.** Sensitivity to moisture in the marine boundary layer (MBL experiments)

10 *3.1. Dried marine boundary layer*

11 Decreasing the moisture contents in the MABL in MBL2M and MBL5M leads to less precipitation with both decreasing RR_{max} and RR_{sum} from CNTL to MBL×M (grey and white bars in Fig. 6a), while the maximum 12 precipitation accumulation remains similar in MBL1M with 75.7 mm. The reduction of total precipitation seen 13 14 over land (RR_{land} , between -0.5 and -3.4 mm) is more significant than that seen over the sea (RR_{sea} , between 15 -0.1 to -0.6 mm) in MBL×M (Table 2a). The duration of precipitation and their horizontal extent are also reduced in MBL×M experiments. The duration of intense precipitation (≥ 15 mm in 15 min, red dots in Fig. 16 17 8b and c; D_{RR15} , black bar in Fig. 6c) are shortened by a factor of 0.75 and 0.38 in MBL2M and MBL5M, respectively, compared to CNTL (90 min and 45 min, instead of 120 min). Similarly, the time for precipitation 18 \geq 5 mm (D_{RR05}, grey bar in Fig. 6c) is shortened in MBL1M and MBL2M by one hour and half compared to 19 20 CNTL (435 min, Fig. 6c).

The horizontal distribution of the 6-hour accumulated precipitation simulated in MBL×M experiments 21 is displayed in Figures 7a-c. It confirms that the precipitation accumulation is reduced, especially for the 22 largest amounts along the eastern coastal Var region. Approximately 25 mm less precipitation is simulated 23 along the eastern coast of Var in MBL1M than in CNTL (area enclosed in solid line, Fig. 7a). In MBL2M and 24 MBL5M, 6-hour accumulation precipitation along the eastern coastal region remains mainly lower than 35 25 mm, whereas larger accumulations are simulated in mountainous region. Figure 6b shows that the area with 26 27 RR_{max} in excess of 30 mm (AR₃₀, grey bars) is reduced by 4.4%, 35.0%, and 65.7% in MBL1M, MBL2M, and MBL5M, respectively, when compared to CNTL, whereas the deviation of area affected by the weak 28

precipitation, RR_{max} in excess of 1 mm (AR₀₁, white bars) is reduced only by 2.8%, 3.7%, and 1.7 % in
 MBL1M, MBL2M, and MBL5M, respectively.

3 The sensitivity bubble of MBL experiments travels north-eastward over the sea as in CNTL, and it keeps its ellipse shape but with a slightly reduced horizontal size due to lateral mixing with the ambient air mass 4 during its advection. When it arrives near the Var coast, the convection initiated at 1200 UTC (yellow star in 5 Fig. 8a-c). The temporal evolution of the location and amounts of RR_{max} (displayed in Figure 8a-c) shows 6 that, in the MBL×M experiments, precipitation (RR_{max} in excess of 5 mm) starts at the same time and location 7 8 as in CNTL (at 1230 UTC) to the southwestern coast of Var, about 95 km distant from its initiation. It then shifts towards Marseille and later to the eastern coastal region of Var. In MBL2M and MBL5M (Fig. 8b-c), 9 the largest 15-min rainfall accumulations stay longer west of 6°E than in CNTL (Fig. 2b). The intense 10 precipitation (\geq 15 mm per 15 min) starts 3 hour later in MBL5M than in CNTL, MBL1M, and MBL2M. The 11 convection at 1800 UTC is located at the eastern coast of Var, as in CNTL (square in Fig. 2b) and MBL×M 12 (square in Fig. 8a-c), and then stays 2-3 additional hours in this region. 13

To understand the reduction of the precipitation amount (RR_{max} and RR_{sum}) and of the precipitating 14 15 duration along the eastern Var coast, Figure 9 shows the domain-averaged CAPE values (using a rising air parcel having its initial height about 20 m) within the sensitivity bubble upstream the precipitation in CNTL 16 (solid line) and MBL×M (dashed lines) from 0930 UTC to 1445 UTC. At 1000 UTC, the CAPE is reduced by 17 about 35, 67, and 95 % in MBL1M, MBL2M, and MBL5M, respectively, compared to CNTL (1083 J kg⁻¹). 18 19 In MBL1M, MBL2M, and MBL5M, CAPE values increase gradually until 1445 UTC but remain lower than CNTL. At 1400 UTC, the spatial distribution of CAPE values less than 1000 J kg⁻¹) is highlighted offshore of 20 21 the Var coast where the sensitivity bubble is located (i.e. $5.2-6^{\circ}E$, $42.4-43^{\circ}N$, dashed ellipsoid in Figure 10b) in MBL2M. It is worth noting that higher CAPE values (> 1400 J kg⁻¹) are displayed in the same region in 22 CNTL (Figure 10a). 23

Figure 11a–c shows that the warm air mass over the Mediterranean and the coastal region ($\theta_e > 322$ K at 925 hPa) is reduced in MBL×M at 1500 UTC in comparison to CNTL (Fig. 4c). The extent of cold air mass ($\theta_v < 291$ K at the first model level) formed along the foothill of mountain by evaporation of the intense precipitation is also reduced in MBL×M (Figure 11a–c). Moreover in MBL2M and MBL5M (Figure 11b–c), the cold pool weakened further with the weakened vertical motion at the southern edge of the cold pool. This combination of decreased CAPE and weakened cold pool which are induced by the weakened precipitation 1 (Figs. 7a–c) is found around the Argens valley region in particular. After 1600 UTC, the cold air mass ($\theta_v <$ 2 291 K) intensifies along the eastern Var coastal region showing the intensified vertical motion at its southern 3 edge (not shown) and the location of RR_{max} (red dots, ≥ 15 mm) shifts to the north-eastern Var coast (Fig. 4 8a–c).

In summary, a decrease in moisture in the MABL results in a reduction of the precipitation amount intensity, and in a shortening of the precipitation duration. The reduction of precipitation is highlighted both around the Argens valley region and over the eastern coast of the Var region because the lifted air is too dry to reach condensation along the coast on the mountainsides with reduced CAPE values, while larger precipitation remains in the mountainous region by the continuous supplies of low-level moist air as CNTL.

10

11 *3.2. Moistened marine boundary layer*

Figure 6d does not show significant differences between MBL1P and CNTL except for an increase of RR_{max} 12 (from 74.2 mm in CNTL to 83mm in MBL1P) and a decrease of D_{land} (from 315 min in CNTL to 240 min in 13 14 MBL1P). In MBL2P, and MBL5P experiments, the increase of the moisture content in the MABL (0.1–1 km ASL) induces more precipitation over the sea than in CNTL. For instance, excesses of 1.3 mm for RR_{land} and 15 of 1.9 mm for RR_{land} are produced in MBL2P with respect to CNTL (RR_{land} of 14.2 mm, RR_{sea} of 1.5 mm) 16 17 (Table 2a). The D_{RR05} (grey bar in Figure 6f) is increased significantly with 480 min, and 660 min in MBL2P 18 and MBL5P instead of 435 min in CNTL, while D_{land} (white bar) is reduced in MBL2P (270 min), and MBL5P 19 (180 min) instead of 315 min in CNTL. Besides, the horizontal area covered by precipitation in excess of 1 mm (AR₀₁, white bar in Figure 6e) which is enlarged by 17.4% and 26.0% in MBL2P, and MBL5P, 20 respectively, while AR_{30} (grey bar) is reduced by 30.5% and 51.5%, respectively. This indicates more 21 22 widespread but weaker precipitation in MBL2P and MBL5P simulations. For MBL2P, RR_{sum} is not significantly modified while RRmax increases to 81.8 mm, more than 74.2 mm in CNTL, and for MBL5P, RRsum 23 decreases from 7.8 mm in CNTL to 6.6 mm, while RR_{max} decreases. 24

The 6-hour accumulated precipitation in MBL1P experiment displayed in Figure 7d confirms that in MBL1P, the largest accumulation on the eastern Var coast is increased (consistently with the increase of RR_{max} , Fig. 6d) and slightly shifted offshore (consistently with the reduced D_{land} , Fig. 6f) (area enclosed by solid line, Fig. 7d). This is consistently seen in the temporal evolution of the location and amounts of the maximum of 15-min accumulated rainfall in Figure 8d, which shows that this accumulation is due to a stationary system blocked over the Var coast, similarly as in CNTL. The precipitating system is blocked by a cold air pool (θ_v < 291 K) over the Argens valley as shown by Figure 11d. In MBL×P experiments (Fig. 8d–f), precipitation is initiated over the sea at 1200 UTC, and then the bubbles travel along similar pathways over the sea as in CNTL. However the inland precipitation starts later (e.g. 15, 45, and 90 min later in MBL1P, MBL2P, and MBL5P, respectively) than in CNTL. This late onset of inland precipitation is also consistent with more widespread precipitation over the sea in MBL2P and MBL5P (Fig. 7e, f).

For MBL2P, Figure 7e still shows a large accumulation on the eastern Var coast due to a stationary system blocked by a cold air pool (Figure 11e) but also an increase of weak precipitation over the sea consistently with Figure 6 results. The precipitation pattern in MBL5P shown by Figures 7f and 8f is quite different from that in CNTL, with more widespread precipitation over the sea (around 6°E, 42°N) and with much less precipitation (64 mm) along the eastern coast of Var (east of 6°E).

The large accumulation of precipitation on the eastern Var coast and the increase of weak precipitation 12 over the sea in MBL1P and MBL2P can be explained by the consistently high CAPE values upstream the 13 precipitation area from 0930 UTC to 1445 UTC compared to CNTL (Figure 9). With the high CAPE values, 14 Figures 8d-f also show that precipitation in MBL×P simulations starts earlier over the sea than in CNTL 15 (Figure 2b) (consistently with the increased D_{RR05}). At 1000 UTC, 1 and 2 g kg⁻¹ increase in moisture in the 16 MABL increases upstream CAPE values by 29 and 55.5 % in MBL1P and MBL2P, respectively, compared to 17 CNTL (1083 J kg⁻¹). At 1400 UTC (Figure 10c), the high CAPE (\geq 1300 J kg⁻¹) in MBL2P is seen offshore of 18 the Var coast (6°E, 42.5°N, dashed ellipsoid in Figure 10c). Figure 11d-e shows that the marine air mass 19 advected towards the coast is warmer and moister ($\theta_e > 324$ K) in MBL1P and MBL2P than in CNTL (Fig. 20 4a-b). The widespread and weaker precipitation over the sea seen more particularly in MBL5P is associated 21 22 to a less organized precipitating system when the moisture content in the MABL is increased. The lesser degree of organization of the convective system in MBL5P is related to the absence of a cold pool (Figure 11f). CAPE 23 values higher than 1500 J kg⁻¹ lead to an increase of the degree of instability in the upstream environment 24 (blue solid line in Figure 9). 25

In summary, increasing the WVMR in the MABL enables an earlier initiation of convection over the sea. Within the warm and moist air mass advected towards the Var region, a small increase of moisture content in MABL favours convection triggering. Precipitation is weaker, more scattered and widespread, especially over the sea, for a WVMR increase exceeding 2 g kg⁻¹ in MABL. This precipitation pattern was similarly seen 1 in an environment of very moist boundary layer of Bresson et al. (2012).

2

3 4. Sensitivity to moisture between 1 and 2 km ASL (FA experiments)

4 4.1. Dried lower troposphere between 1 and 2 km ASL

Decreasing moisture contents in the dry layer at 1-2 km ASL in FA×M (FA1M, FA2M, and FA5M) reduces 5 6 the total accumulated precipitation (RR_{sum} equal to about 7.6 mm, instead of 7.8 mm in CNTL, Figure 12a). However the total amount of precipitation simulated in the FA×M experiments remains larger than for MBL×M 7 8 experiments with a drier MABL (e.g. RR_{sum} equal to 7.5 mm, 7.2 mm and 5.8 mm for MBL1M, MBL2M, and 9 MBL5M, respectively). This indicates that the precipitation amount is more sensitive to the moist air in the 10 MABL (below 1 km ASL) than the air just above (1-2 km ASL). The duration of precipitation (D_{R05} and D_{land}) 11 is also reduced (Fig. 12c). Correspondingly RR_{land} is reduced to between -0.5 and -0.7 mm with respect to the 12 value of 14.2 mm in CNTL (Table 2b). The area affected by precipitation (AR_{01}) is reduced as well, especially for accumulation ≥ 30 mm (AR₃₀) and for a change of WVMR larger than 2 g kg⁻¹ (Fig. 12b). 13

The horizontal distributions of the 6-hour accumulated precipitation in FA×M experiments are displayed 14 15 in Figure 13a-c. The maximum of precipitation located along the eastern Var coast (closed circle by solid line) is about 15–20 mm smaller than in CNTL. The temporal evolution of the location and amounts of the maximum 16 of 15-min accumulated rainfall every 15 min in FA×M experiments is displayed in Fig. 14a-c. The sensitivity 17 bubbles of FA experiments travel north-eastward over the sea as similar as seen in CNTL and MBL, but the 18 horizontal extent is slightly reduced by mixing with the ambient air mass during the travel. The convection 19 initiates about 20 km offshore of the Var coast (yellow star, Fig. 14a–c). In FA×M experiments, $RR_{max} \ge 5 \text{ mm}$ 20 21 starts about at the same time (or slightly later) than in CNTL. However, precipitation is initiated at a more realistic location, closer to Marseille (about 5.5°E), when the bubble arrives, especially in FA2M and FA5M, 22 instead of the southern tip of the Var region (about 6°E) of CNTL (Figure 2b). 23

Figure 15a–c displays the hourly evolution simulated by FA×M simulations of θ_e at 925 hPa and θ_v at the first model level (about 10 m high), together with the horizontal wind and the vertical velocity at 500 m ASL at 1500 UTC. It shows a weaker cold pool with higher θ_v values (~1.5 K) along the east coastal region of Var (6.2–7.2°E) in FA×M simulations (Fig. 15a–c), compared to CNTL (Fig. 4c). The ascents (green areas) at the southern edge of the cold pool ($\theta_v \le 291$ K) are consistently slightly reduced in FA×M, compared to CNTL. Figure 16 shows that the rainfall intensity in FA×M simulation (dashed lines) is reduced by about 0.1–0.5 mm

 hr^{-1} compared to CNTL (black line) between 1200 and 1430 UTC, and the rainfall intensity $\ge 0.1 \text{ mm }hr^{-1}$ 1 starts about 45 min later (1300 UTC, dashed line) than in CNTL (1215 UTC, black line). Under the weakened 2 3 precipitation, evaporation rate is reduced. The corresponding values of θ_{v} in the FA×M simulations are increased by about 0.2–0.5 K with respect to CNTL (dashed line, Fig. 16b). While the dried air mass in the 4 1-2 km layer is advected toward upstream the convection, it mixes with the moist air below, reducing the total 5 moisture below 2 km. Considering the region immediately upstream of convective ascents, the total moisture 6 below 2 km in the FA×M simulations is lower than in CNTL, due to the advection of the sensitivity bubble 7 8 containing dry air at heights between 1 and 2 km. With the reduced moisture in the lower troposphere, the 9 triggered convection near the Var coast produces less intensive rainfall and the weakened cold pool. This is consistent with the reduction of the rainfall accumulation over the Var coast in FA×M simulations. Modifying 10 the moisture content in the dry layer at 1-2 km ASL does not change much the CAPE values upstream the 11 precipitation (not shown). The CAPE values in both CNTL and FA×M slightly increased over time from 1083 12 to 1650 J kg⁻¹ in the location of the sensitivity bubble as it mixed with the adjacent moister air (not shown). 13

In summary, decreasing the moisture content in the dry layer between 1 and 2 km ASL reduces the total amount of precipitation as well as the area affected by the precipitation and the duration of the precipitating episode. As the MABL is nearly saturated, the convection triggers easily along the coast but develops less intensively. The maximum of precipitation located along the eastern Var coast is reduced, corresponding to a weakened cold pool and weakened ascents at its southern boundary. This shows that around the coast, the dryness in the 1–2 km ASL layer is not a major ingredient for the convection development and the cold pool generation when the lowermost layer is nearly saturated.

21

4.2. Moistened lower troposphere between 1 and 2 km ASL

The experiments FA×P (FA1P, FA2P and FA5P) with increased moisture content in the free troposphere between 1–2 km ASL produced a similar total precipitation accumulation (RR_{sum} about 7.7 mm in Fig. 12d) to CNTL (7.8 mm). The duration of precipitation D_{RR05} is prolonged from 435 min in CNTL to 465 min, 510 min, and 510 min, in FA1P, FA2P, and FA5P respectively (Fig. 12f). However, the duration of precipitation over land (D_{land}) and the duration of more intense precipitation (D_{RR15}, \geq 15 mm in 15 min) are shortened in FA×P experiments than in CNTL. Table 2b shows the reduced RR_{land} values between -0.7 and -1.5 mm in FA×P, with respect to CNTL (1.5 mm), as well as the increased RR_{sea} values between +0.6 and +1.4 mm. The areas of precipitation AR_{01} and AR_{30} are not much modified in FA1P and FA2P simulations and both are slightly

2 reduced in FA5P (Fig. 12e).

The horizontal distribution of the 6-hour accumulated precipitation in FA×P experiments is displayed in Figure 13d–f. The precipitation pattern is not much modified, but the maximum precipitation amount over the eastern coastal region of Var is slightly shifted offshore and reduced mainly in FA2P and FA5P (with 65.8 mm and 65.9 mm instead of 74.2 mm in CNTL, east of 6°E, Figure 13e and f).

The temporal evolution of the location and amount of the maximum of 15-min accumulated rainfall 7 8 every 15 min in FA×P experiments (Fig. 14d-f) shows that increasing the moisture contents in the dry layer between 1 km and 2 km ASL induces an initiation of the precipitation about 1 hour earlier than in CNTL 9 (consistently with the increase of D_{RR05}). The convection initiates near the Var coast, as in CNTL, however the 10 horizontal extent of the convective region (reflectivity \geq 45 dBZ) is relatively large compared to CNTL. The 11 precipitation starts at 1200 UTC about 50 km offshore the southern Var coast, when the bubble reaches this 12 region ~140 km distant from the location of initiation, with larger 15-min accumulation values (10-20 mm, 13 black and red dots in Fig. 14e-f, compared to grey dots < 10 mm in CNTL in Fig. 2b). 14

Regarding the large precipitation around the southern Var coast and its enlarged horizontal extent of the 15 convective region in FA2P and FA5P, Figure 15e–f shows the large affected area by the cold pool ($\theta_{\nu} < 291$ 16 K) compared to CNTL (Fig. 4a) at 1500 UTC. Figure 16a shows the enhanced rainfall intensities (about 17 $0.2-2.5 \text{ mm hr}^{-1}$) in FA×P simulations (colored solid lines) compared to CNTL (black line) between 1200 and 18 19 1430 UTC, especially FA5P produces the precipitation about 45 min earlier in time. Under the intensifying precipitation, the values of θ_v in FA×P simulations are about 0.2–1.7 K reduced, indicating the intensified cold 20 21 pool formed by evaporative cooling than in CNTL (Fig. 16b). As the moistened air mass in the 1-2 km layer 22 mixed with the moist air below during its advection towards the region of upstream convection, the total moisture below 2 km ASL increased. Thanks to the moistened lower troposphere, the convection triggered 23 near the Var coast produces more intensive rainfall and correspondingly more intense cold pool. Also the 24 25 moistened air masses in the lower troposphere initiate precipitation earlier while the precipitating area, particularly over the sea, is enlarged. In FA×P, the duration of the entire episode is increased, but the intensity 26 of the precipitation is reduced (red dots in Fig. 14d-f; D_{RR15} decreases in Fig. 12f). It also appears that the 27 precipitation is less stationary. 28

29

In summary, increasing the moisture content in the dry lower troposphere between 1 and 2 km ASL

enlarges the precipitating area, particularly over the sea. With the moistened air mass in the 1–2 km ASL layer, similar CAPE values ($\geq 1050 \text{ J kg}^{-1}$) were calculated offshore of the Var coast in the FA×P experiments and in CNTL. In the environment with similar instability but with further moistened conditions in the lower troposphere, the triggered convection further intensifies with enlarged horizontal extent of cold pool and strengthened ascents at its southern boundary.

6

7 **5. Summary**

8 The present study examines the impact of the environmental moisture structure of the lower troposphere (below 9 2 km ASL) on the precipitation development and organisation, observed in southern France during IOP13 of the HyMeX SOP-1, through a series of sensitivity experiments using the non-hydrostatic numerical model 10 Meso-NH. The moisture structure upstream of the IOP13 HPE is characterised by a moist conditionally 11 unstable MABL (below 1 km ASL) topped by a dry air mass just above (1-2 km ASL). A CNTL simulation 12 13 shows that the moisture supply to the precipitating system is provided by the moist air mass of the MABL, while the dry air mass between 1 and 2 km ASL is involved in the formation of a cold pool. Focusing on these 14 two layers, namely (1) a moist layer from 0.1 to 1 km ASL, and (2) a dry layer from 1 to 2 km ASL, 12 15 sensitivity experiments were carried out to study the influence of upstream moisture structure in the lower 16 troposphere on convection development occurred in southern France. 17

The control simulation (CNTL), as well as all the other sensitivity experiments examined in this study, 18 19 succeed in reproducing heavy precipitation in the coastal mountainous region of Var in southeastern France. Through comparisons between CNTL and the 12 sensitivity experiments, we show how the life cycle of 20 precipitation is modified even for moisture contents changes as small as 1 g kg⁻¹ below 2 km ASL. The results 21 are summarized schematically in Figure 167. Increasing the moisture content by 1, 2 or 5 g kg⁻¹ in the MABL 22 (0.1-1 km ASL) (see Fig. 17a) induces an earlier initiation of precipitation offshore of the Var region (light 23 24 blue area) with an increase of CAPE values, as well as longer lasting precipitations. A small increase of 25 moisture content in the warm and moist MABL increases the degree of instability. In the environment with an increase of the MABL moisture content lower than 2 g kg⁻¹, precipitation developing offshore arrives to the 26 coastal region of Var with the low-level southwesterlies warm and moist air and anchors over the coasts as it 27 is blocked by a cold pool (blue line). For an increase of the moisture content exceeding 2 g kg⁻¹, the area of 28 29 intense precipitation around the coastal region is reduced and the precipitating system is less stationary because

1 of a weakened cold pool.

Similarly, moistening the layer at 1–2 km ASL (see Fig. 17c), just above the MABL, increases the humidity in the lowermost 2 km ASL by mixing in the environment with a similar instability as in CNTL. With more moisture in lower troposphere, the triggered convection further intensifies which produces an enlargement of the horizontal extent of cold pools (dark blue area). Also it produces precipitation earlier offshore (light blue area), but does not increase the total amount of precipitation much. Precipitation lasts longer and affects a larger area, particularly over the sea.

8 A drier MABL (Fig. 17b) shortens the life-time of precipitation and reduces the total precipitation amount. For instance, a 2 g kg⁻¹ decreased moisture content in the MABL results in a reduction of about 10 % 9 in terms of total precipitation in the coastal regions of Var (closed area with blue broken line), and in a reduction 10 of about 67 % in term of CAPE. A drier lower troposphere in the 1-2 km ASL layer (see Fig. 17d) also 11 contributes to reduce the precipitating area (light blue area) with rainfall accumulation located more inland 12 rather than over the coastal region because of a weakened cold pool. Also it is true that dried air mass in the 13 1-2 km ASL initiates precipitation at a more accurate location, about 25 km closer to Marseille. Decreasing 14 the moisture content in MABL has a stronger impact on precipitation both in terms of amount and intensity 15 16 than decreasing the moisture content in the layer just above (1-2 km ASL). The dryness in the 1-2 km ASL layer is not a major ingredient for the convection development and the cold pool generation when the 17 lowermost layer is nearly saturated. 18

19 Despite a simplified moisture profile modification approach, this study suggests that moisture structure in lower troposphere (below 2 km ASL) is a key for an accurate prediction of the timing and location of 20 21 precipitation in the coastal mountainous region (e.g. the Var region) in southern France. At the same time, this 22 study shows the importance of accurate moisture content (amount and profile) in the initial field to reproduce realistic convective systems, emphasizing the importance of high-resolution and three-dimensional moisture 23 observation upstream the HPE, especially over the sea. This study focused on the impact of moisture contents 24 on precipitation development in southern France, and the HPE which occurred during IOP13 of the HyMeX 25 SOP-1. It would be interesting to consider other regions in the Mediterranean basin by analysing additional 26 HPE cases of SOP-1 when interesting moisture structure including the presence of dry air masses above 2 km 27 ASL upstream of the HPE were identified, e.g., IOP13 in south Italy (Lee et al., 2016) and IOP8 in north-east 28 of Iberian plateau (Bouin et al., 2017). By accumulating the event scale analysis in other regions of the 29

Mediterranean basin, we can expand our knowledge what is the general impact of upstream water vapour on precipitation (e.g. categories of synoptic conditions). Another approach is to use stable water isotopologue data to disentangle the various moisture sources, i.e. evaporation over the sea, advected moisture upstream of the HPEs in the Mediterranean (Sodemann et al., 2017).

5

6 Acknowledgements

This work was supported by the French Agence Nationale de la Recherche (ANR) via the IODA-MED Grant
ANR-11-BS56-0005, the MUSIC grant ANR-14-CE01-014 and the MISTRALS/HyMeX programme. This
work was partly supported by DRIHM.

10

11 **References**

- Argence, S., Lambert, D., Richard, E., Chaboureau, J.P., and Söhne, N.: Impact of initial condition
 uncertainties on the predictability of heavy rainfall in the Mediterranean: a case study. Q. J. R. Meteorol.
 Soc, 134, 1775–1788, doi: 10.1002/qj.314, 2008.
- Barthlott, C. and Davolio, S.: Mechanisms initiating heavy precipitation over Italy during the HyMeX Special
 Observation Period 1: A numerical case study using two mesoscale models. Q. J. R. Meteorol. Soc, doi:
 10.1002/qj.2630, 2015.
- Bielli, S., Grzeschik, M., Richard, E., Flamant, C., Champollion, C., Kiemle, C., Dorninger, M., and
 Brousseau, P.: Assimilation of water-vapour airborne lidar observations: impact study on the COPS
 precipitation forecasts. Q. J. R. Meteorol. Soc., 138, 1652–1667, 2012.
- Bougeault, P. and Lacarrère, P.: Parameterization of orography-induced turbulence in a meso-beta-scale model.
 Mon. Weather Rev. 117(8): 1872–1890, 1989.
- Bouin, M. N., Redelsperger, J. L., and Leveaupin, B. C.: Processes leading to deep convection and sensitivity
 to sea-state representation during HyMeX IOP8 heavy precipitation event. Q. J. R. Meteorol. Soc, doi:
 10.1002/qj.3111, 2017.
- Bresson, E., Ducrocq, V., Nuissier, O., Ricard, D., and de Saint-Aubin C.: Idealized numerical simulations of
 quasi-stationary convective systems over the Northwestern Mediterranean complex terrain. Q. J. R.
 Meteorol. Soc. 138:1751–1763, doi: 10.1002/qj.1911, 2012.

1	Buzzi, A., Tartaglione, N., and Malguzzi, P.: Numerical simulations of the 1994 Piemont flood: Role of the
2	orography and moist processes. Mon. Weather Rev. 126: 2369-2383, 1998.
3	Carpenter K.: Note on the paper: Radiational condition for the lateral boundaries of limited-area numerical
4	models by MJ Miller and AJ Thorpe. Q. J. R. Meteorol. Soc. 108:717-719, 1982.
5	Caniaux, G., Redelsperger, J. L., and Lafore, J. P.: A numerical study of the stratiform region of a fast-moving
6	squall line. J. Atmos. Sci. 51(14): 2046-2074, 1994.
7	Clark, H. and Chaboureau, J.P.: Uncertainties in short-term forecasts of Mediterranean heavy precipitation
8	event: Assessment with satellite observations. J. Geophys. Res., 115, D22213, 2010.
9	Chazette, P., Flamant, C., Raut, J. C., Totems, J., and Shang, X.: Tropical moisture enriched storm tracks
10	over the Mediterranean and their link with intense rainfall in the Cevennes-Vivarais area during
11	HyMeX. Q. J. R. Meteorol. Soc. 142: 320-334, doi:10.1002/qj.2674, 2016.
12	Collella, P. and Woodward, P. R.: The piecewise parabolic method (PPM) for gas dynamical simulations. J.
13	Comput. Phys. 54: 174-201, doi:10.1016/0021-9991(84)90143-8, 1984.
14	Crook, N. A.: Sensitivity of moist convection forced by boundary layer processes to low-level thermodynamic
15	fields. Mon. Wea. Rev., 124, 1767–1785, 1996.
16	Cuxart, J., Bougeault, P., and Redelsperger, J. L.: A turbulence scheme allowing for mesoscale and large-eddy
17	simulations. Q. J. R. Meteorol. Soc. 126(562): 1-30, doi: 10.1002/qj.49712656202, 2000.
18	Davies, H.: A lateral boundary formulation for multi-level prediction models. Q. J. R. Meteorol. Soc.
19	102:405–418, 1976.
20	Ducrocq, V., Braud, I., Davolio, S., Ferretti, R., Flamant, C., Jansa, A., Kalthoff, N., Richard, E., Taupier-
21	Letage, I., Ayral, P.A., Belamari, S., Berne, A., Borga, M., Boudevillain, B., Bock, O., Boichard, J. L.,
22	Bouin, M. N., Bousquet, O., Bouvier, C., Chiggiato, J., Ciimini, D., Corsmeier, U., Coppola, L.,
23	Cocquerez, P., Defer, E., Delanoë, J., Di Girolamo, P., Doerenbecher, A., Drobinski, P., Dufournet, Y.,
24	Fourrié, N., Gourley, J.J., Labatut, L., Lambert, D., Le Coz, J., Marzano, F.S., Molinié, G., Montani, A.,
25	Nord, G., Nuret, M., Ramage, K., Rison, W., Roussot, O., Said, F., Schwarzenboeck, A., Testor, P., Van
26	Baelen, J., Vincendon, B., Aran, M., and Tamayo, J.: HyMeX-SOP1: The Field Campaign Dedicated to
27	Heavy Precipitation and Flash Flooding in the Northwestern Mediterranean. Bull. Am. Meteorol. Soc.
28	95: 1083-1100, doi:10.1175/BAMS-D-12-00244.1, 2014.
29	Ducrocq, V., Davolio, S., Ferretti, R., Flamant, C., Santaner, V. H., Kalthoff, N., Richard, E., and Wernli, H.:

- Introduction to the HyMeX Special Issue on 'Advances in understanding and forecasting of heavy
 precipitation in the Mediterranean through the HyMeX SOP1 field campaign'. Q. J. R. Meteorol. Soc.
 142, Issue S1, 1–6, 2016.
- Ducrocq, V., Nuissier, O., Ricard, D., Lebeaupin, C., and Thouvenin, R.: A numerical study of three
 catastrophic precipitating events over southern France. II: Mesoscale triggering and stationarity factors.
 Q. J. R. Meteorol. Soc. 134: 131–145, 2008.
- Ducrocq, V., Ricard, D., Lafore, J. P., and Orain, F.: Storm-scale numerical rainfall prediction for five
 precipitating events over France: On the importance of the initial humidity field. Weather and
 Forecasting 17: 1236–1256, 2002.
- Duffourg, F. and Ducrocq, V.: Origin of the moisture feeding the heavy precipitating systems over southeastern
 France. Nat. Hazards Earth Syst. Sci. 11: 1163–1178, 2011.
- Duffourg, F., Lee, K. O., Ducrocq, V., Flamant, C., Chazette, P., and Di Girolamo, P.: Role of moisture patterns
 in the backbuilding formation of HyMeX IOP13 Heavy Precipitating Systems. Q. J. R. Meteorol. Soc.
 14 144: 291–303, doi:10.1002/qj.3201, 2018.
- Duffourg, F., Nuissier, O., Ducrocq, V., Flamant, C., Chazette, P., Delanoë, J., Doerenbecher, A., Fourrié, N.,
 Di Girolamo, P., Lac, C., Legain, D., Martinet, M., Saïd, F., and Bock, O.: Offshore deep convection
 initiation and maintenance during the HyMeX IOP 16a Heavy Precipitation Event. Q. J. R. Meteorol.
 Soc. 142 (S1): 259–274, doi:10.1002/qj.2725, 2016.
- 19 Fourrié, N., Bresson, E., Nuret, M., Jany, C., Brousseau, P., Doerenbecher, A., Kreitz, M., Nuissier, O.,
- Sevault, E., Bénichou, H., Amodei, M., and Pouponneau, F.: Arome-wmed, a real-time mesoscale model
 designed for the HyMeX special observation periods. Geosci. Model Dev. 8(2): 1801–1856,
 doi:10.5194:/gmdd-8-1801-2015, 2015.
- Gal-Chen, T. and Somerville, R. C. J.: On the use of a coordinate transformation for the solution of the NavierStokes equations. J. Comput. Phys. 17: 209–228, doi:10.1016/0021-9991(75)90037-6, 1975.
- Houze, R. J.: Cloud Dynamics, International Geophysics Vol. 53. Academic Press: New York, NY, 1993.
- 26 Jansa, A., Genoves, A., Picornell, M. A., Campins, J., Riosalido, R., and Carretero, O.: Western Mediterranean
- 27 cyclones and heavy rain. Part 2: Statistical approach. Meteorol. Appl. 8: 43–56, doi:10.1017/
 28 S1350482701001049, 2001.
- 29 Lac, C., Chaboureau, J. P., Masson, V., Pinty, J. P., Tulet, P., Escobar, J., Leriche, M., Barthe, C., Aouizerats,

1	B., Augros, C., Aumond, P., Auguste, F., Bechtold, P., Berthet, S., Bielli, S., Bosseur, F., Caumont, O.,
2	Cohard, J. M., Colin, J., Couvreux, F., Cuxart, J., Delautier, G., Dauhut, T., Ducrocq, V., Filippi, J. B.,
3	Gazen, D., Geoffroy, O., Gheusi, F., Honnert, R., Lafore, J. P., Lebeaupin, Brossier C., Libois, Q., Lunet,
4	T., Mari, C., Maric, T., Mascart, P., Mogé, M., Molinié, G., Nuissier, O., Pantillon, F., Peyrillé, P.,
5	Pergaud, J., Perraud, E., Pianezze, J., Redelsperger, J. L., Ricard, D., Richard, E., Riette, S., Rodier, Q.,
6	Schoetter, R., Seyfried, L., Stein, J., Suhre, K., Taufour, M., Thouron, O., Turner, S., Verrelle, A., Vié,
7	B., Visentin, F., Vionnet, V., and Wautelet, P.: Overview of the Meso-NH model version 5.4 and its
8	applications. Geosci. Model Dev. 11, 1929–1969, https://doi.org/10.5194/gmd-11-1929-2018, 2018.
9	Lee, K. O., Flamant, C., Ducrocq, V., Duffourg, F., Fourrié, N., and Davolio, S.: Convective initiation and
10	maintenance processes of two back-building mesoscale convective systems leading to heavy
11	precipitation events in Southern Italy during HyMeX IOP 13. Q. J. R. Meteorol. Soc, 142: 2623-2635,
12	doi: 10.1002/qj.2978, 2016.
13	Lee, K. O., Flamant, C., Ducrocq, V., Duffourg, F., Fourrié, N., Delanoë, J., and Bech, J.: Initiation and
14	development of a mesoscale convective system in the Ebro River Valley and related heavy precipitation
15	over northeastern Spain during HyMeX IOP15a. Q. J. R. Meteorol. Soc, 143: 942-956, doi:
16	10.1002/qj.2851, 2017.
17	Lee, K. O., Uyeda, H., Shimizu, S., and Lee, D. I.: Dual-Doppler radar analysis of the enhancement of a
18	precipitation system on the northern side of Mt. Halla, Jeju Island, Korea on 6 July 2007. Atmos. Res.,
19	118, 133–152, 2012.
20	Lin, Y.: Orographic effects on airflow and mesoscale weather systems over Taiwan. Terr. Atmos. Ocean 4:
21	381–420, 1993.
22	Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Bardu, A., Boone, A.,
23	Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.C., Carrer, D., Decharme, B., Delire, C., Donier, S.,
24	Essaouini, K., Gibelin, A. L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E.,
25	Lafaysse, M., Lafont, S., Lebeaupin, BC., Lemonsu, A., Mahfouf, J.F., Marguinaud, P., Mokhtari, M.,
26	Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet,
27	V., and Voldoire, A.: The surfex v7.2 land and ocean surface platform for coupled or offline simulation
28	of earth surface variables and fluxes. Geosci. Model Dev. 6(4): 929-960, doi:10.5194/gmd-6-929-2013,
29	2013.

- Meyers, M. P., Demott, P. J., and Cotton, W. R.: New primary ice-nucleation parameterizations in an explicit
 cloud model. J. Appl. Meteor., 31, 708–721, 1992.
- Miglietta, M. and Rotunno, R.: Numerical simulations of sheared conditionally unstable flows over a mountain
 ridge, J. Atmos. Sci., 71, 1747–1762, 2014.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for
 inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave. J. Geophys.
 Res., 102D, 16663–16682, 1997.
- Nuissier, O., Ducrocq, V., Ricard, D., Lebeaupin, Brossier C., and Anquetin, S.: A numerical study of three
 catastrophic precipitating events over southern France. I: Numerical framework and synoptic
 ingredients. Q. J. R. Meteorol. Soc. 134: 111–130, 2008.
- Nuissier, P., Joly, B., Joly, A., Ducrocq, V., and Arbogast, P.: A statistical downscaling to identify the large scale circulation patterns associated with heavy precipitation events over southern France. Q. J. R.
 Meteorol. Soc. 137: 1812–1827, doi:10.1002/qj.866, 2011.
- Pergaud, J., Masson, V., Malardel, S., and Couvreax, F.: A parameterization of dry thermals and shallow cumuli
 for mesoscale Numerical Weather Prediction. Bound.-Lay. Meteorol. 132(1): 83–106,
 doi:10.1007/s10546-009-9388-0, 2009.
- Pinty, J. P. and Jabouille, P.: A mixed-phased cloud parametrization for use in a mesoscale non-hydrostatic
 model: Simulations of a squall line and of orographic precipitation. In: Proc. Of the Conference on Cloud
 Physics. Amer. Meteorol. Soc, Boston: Everett, WA, USA, 17–21 Aug. 1998. Pp. 217–220, 1998.
- Ricard, D., Ducrocq, V., and Auger, L.: A climatology of the mesoscale environment associated with heavily
 precipitating events over a northwestern mediterranean area. J. Appl. Meteorol. Climatol. 51: 468–488,
 doi:10.1175/JAMC-D-11-017.1, 2012.
- Rotunno, R. and Ferretti, R.: Mechanisms of intense Alpine rainfall. J. Atmos. Sci. 58: 1732–1749, 2001.
- Romero, R., Ramis, C., and Homar, V.: On the severe convective storm of 29th October 2013 in the Balearic
 Islands: Observational and numerical study. Q. J. R. Meteorol. Soc, doi:10.1002/qj.2429, 2015.
- Romero, R., Sumner, G., Ramis, C., and Genoves, A.: A classification of the atmospheric circulation patterns
 producing significant daily rainfall in the Spanish Mediterranean area. Int. J. Climatol. 19: 765–785,
 1999.
- 29 Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bernard, P., Bouttier, F., Lac, C., Masson, V.: The AROME-

- France convective-scale operational model. Mon. Weather Rev. 139, 976–991, 2011.
- Shu, C. W. and Osher, S.: Efficient implementation of essentially non-oscillatory shock-capturing schemes.
 Journal of Computational Physics 77(2): 439–471, 1988.
- 4 Smith, R.: The influence of mountains on the atmosphere. Adv. Geophys. 21: 87–230, 1979.
- Sodemann, H., Aemisegger, F., Pfahl, S., Bitter, M., Corsmeier, U., Feuerle, T., Graf, P., Hankers, R., Jsiao, G.,
 Schulz, H., Wieser, A., and Wernli, H.: The stable isotopic composition of water vapour above Corsica
 during the HyMeX SOP1 campaign: insight into vertical mixing processes from lower-tropospheric
 survey flights. Atmos. Chem. Phys., 17, 6125–6151, 2017.
- 9 Trapero, L., Bech, J., and Lorente, J.: Numerical modelling of heavy precipitation events over Eastern
 10 Pyrenees: Analysis of orographic effects. Atmos. Res. 123: 368–383, 2013a.
- Trapero, L., Bech, J., Duffourg, F., Esteban, P., and Lorente, J.: Mesoscale numerical analysis of the historical
 November 1982 heavy precipitation event over Andorra (Eastern Pyrenees). Nat. Hazards Earth Syst.
 Sci. 13: 2969–2990, 2013b.
- Weckwerth, T., Parsons, D. B., Koch, S. E., Moore, J. A., LeMone, M. A., Demoz, B. B., Flamant, C., Geerts,
 B., Wang, J., and Feltz, W. F.: An overview of the International H2O Project (IHOP-2002) and some
 preliminary highlight. Bull. Am. Meteorol. Soc. 85: 253–277, 2004.

Table 1. List of sensitivity experiment and WVMR modifications with respect to the control simulation (CNTL).

(a) Exp.	MBL1M	MBL2M	MBL5M	MBL1P	MBL2P	MBL5P
Max WVMR difference	$-1 \mathrm{g kg^{-1}}$	$-2 \mathrm{g kg^{-1}}$	$-5 \mathrm{g kg^{-1}}$	$+1 \mathrm{g kg^{-1}}$	$+2 \mathrm{g} \mathrm{kg}^{-1}$	$+5 \mathrm{g kg^{-1}}$
Altitudes modified			0.1 to 1 km above sea level			
(b) Exp.	FA1M	FA2M	FA5M	FA1P	FA2P	FA5P
(b) Exp. Max WVMR difference	FA1M -1 g kg ⁻¹	FA2M -2 g kg ⁻¹	FA5M 5 g kg ⁻¹	FA1P +1 g kg ⁻¹	FA2P +2 g kg ⁻¹	FA5P +5 g kg ⁻¹

Table 2. Deviation of domain-averaged 6-hour total accumulated precipitation amount over land (RR_{land}) and over sea

 (RR_{sea}) for the MBL and FA experiments with respect to the CNTL (RR_{land} of 14.2 mm, RR_{sea} of 1.5 mm) within the fixed

8 area (3.5–9°E, 40.5–45°N) from 1200 UTC on 14 October 2012.

(a) Exp.	MBL1M	MBL2M	MBL5M	MBL1P	MBL2P	MBL5P
RR _{land}	-0.5	-0.8	-3.4	-0.7	-1.9	-3.2
RR _{sea}	-0.1	-0.4	-0.6	+0.6	+1.3	+0.8
(b) Exp.	FA1M	FA2M	FA5M	FA1P	FA2P	FA5P
RRland	-0.6	-0.5	-0.7	-0.7	-1.2	-1.5
RR _{sea}	+0.2	+0.2	+0.3	+0.6	+1.1	+1.4





Figure 1. Topography and domain (black box) considered in numerical simulations using Meso-NH with a resolution of 2.5 km. The centre location of the initial "sensitivity bubble" set in MBL and FA experiments is depicted by red- and blue- coloured cross mark, respectively, while the horizontal range of the initial bubbles are marked by an ellipse-shaped solid line of the same colour. The red dashed line indicates the horizontal projection of backward trajectories of some air parcels taken in the upper part of the convective systems of the eastern Var coast, and the blue dashed line depicts the horizontal projection of backward trajectories of some air parcels taken in the cold pool generated by the convective systems over the eastern Var coast.





Figure 2. (a) 6-hour accumulated precipitation at 18 UTC, and (b) evolution every 15 min between 12 UTC and 18 UTC of the locations of maximum 15-min accumulated rainfall amount (RR_{max}) simulated by the control run (CNTL) on 14 October 2012. Also shown is the topography (see Figure 1 for the scale). The location of first convection initiation (reflectivity over 45 dBZ) is marked by a yellow star, and the last precipitating point at 18 UTC is marked by a square pointed by an arrow in (b).



2

Figure 3. Radar reflectivity (dBZ) and horizontal winds simulated in CNTL at 2000 m ASL at (a) 1300 UTC, (b) 1400

4 UTC, (c) 1500 UTC and (d) 1600 UTC on 14 October 2012.



Figure 4. Equivalent potential temperature θ_e at 925 hPa (K, red areas), virtual potential temperature θ_v at the first model level (K, blue areas below 291 K), vertical motion at 500 m ASL (green areas above 0.5 m s⁻¹) and horizontal wind at 925 hPa (arrows) simulated by CNTL at (a) 1400 UTC, (b) 1500 UTC, (c) 1600 UTC and (d) 1700 UTC on 14 October 2012.

- 7
- 8



Figure 5. Vertical profiles of WVMR in the centre of the initial "sensitivity bubble" used in (a) MBL experiments, and (b) FA experiments. Black solid lines are vertical profiles of the CNTL run at the same grid point as in MBL and FA. Solid grey lines represent experiments in which WVMR was augmented, while dashed grey lines represent experiments for which WVMR was reduced. The centre location of the initial "sensitivity bubble" used in MBL and FA are shown in Figure 1.



3 Figure 6. Results of CNTL, MBL×M (upper panel, a–c) and MBL×P (bottom panel, d–f) experiments: (a) and (d) maximum 6-4 hour accumulated precipitation amount (RR_{max}), and the domain-averaged total sum of the 6-h accumulated precipitation amount 5 (RR_{sum}, mm) from 1200 UTC on 14 October 2012, (b) and (e) the deviation of areas (km²) of RR_{max} \ge 1 mm (AR01) and RR_{max} 6 \ge 30 mm (AR30) in MBL to ones in CNTL, and (c) and (f) duration of precipitation (\ge 5 mm) over the land (D_{land}), duration of 7 precipitation \ge than 5 mm per 15 min (D_{RR05}), duration of intense precipitation \ge 15 mm per 15 min (D_{RR15}). The RR_{sum} and 8 duration of precipitation were calculated at a fixed area of latitude of 40.5–45°N, longitude of 3.5–9°E where the sensitivity 9 bubble passed though are used, respectively.



Figure 7. Distribution of the 6-hour accumulated precipitation simulated by (a) MBL1M, (b) MBL2M, (c) MBL5M, (d)
MBL1P, (e) MBL2P, and (f) MBL5P at 18 UTC on 14 October 2012. Black contour line shows the coast of southern
France. Ellipsoid in (a)–(c) indicates the area with less precipitation than CNTL, while the ellipsoid in (d)–(e) shows the
shifted precipitation area to the offshore region.





Figure 8. Evolution of the location of the maximum 15-min accumulated rainfall for 6 hours (every 15 min from 1200
UTC on 14 October 2012) simulated by (a) MBL1M, (b) MBL2M, (c) MBL5M, (d) MBL1P, (e) MBL2P, and (f) MBL5P.
Also shown is the topography (see Figure 1 for the scale). The location of first convection initiation is marked by a yellow
star, and the last precipitating point at 18 UTC is marked by a square pointed by an arrow.





2

3 Figure 11. Equivalent potential temperature θ_e at 925 hPa (K, red areas), virtual potential temperature θ_v at the first model level (K, blue areas below 292 K), vertical motion at 500 m ASL (green areas above 0.5 m s⁻¹, arrow) and horizontal wind 4 5 at 925 hPa simulated by (a) MBL1M, (b) MBL2M, (c) MBL5M, (d) MBL1P, (e) MBL2P, and (f) MBL5P at 1500 UTC 6 on 14 October 2012.



Figure 12. As Figure 6 but for CNTL, FA×M (upper panel, a–c) and FA×P (bottom panel, d–f) experiments.



Figure 13. Same as Figure 7 for (a) FA1M, (b) FA2M, (c) FA5M, (d) FA1P, (e) FA2P, and (f) FA5P. In (a)–(c), the reduced
precipitation around the coast is indicated by closed ellipsoid with solid line.



Figure 14. Same as Figure 8 for (a) FA1M, (b) FA2M, (c) FA5M, (d) FA1P, (e) FA2P, and (f) FA5P.





Figure 15. Same as Figure 11 for (a) FA1M, (b) FA2M, (c) FA5M, (d) FA1P, (e) FA2P, and (f) FA5P at 1500 UTC on 14

October 2012.



Figure 16. Temporal evolution of domain-averaged rainfall intensity (mm hr⁻¹) and virtual potential temperature (θ_v) at the first model level in a fixed domain of 5.5–7.5°E, 43–44.5°N where most of the precipitation produced by CNTL (black line), FA×P (solid lines), and FA×M (dashed lines) from 1115 to 1800 UTC on 14 October 2012.





Figure 17. Schematics summarizing the main features of the impact of upstream moisture in MBL in (a) and (b), and in the free atmosphere (FA) layer between 1 and 2 km ASL in (c) and (d) on back-building heavy precipitation in southeastern France during HyMeX IOP 13. Dark-blue arrows in (a) and (c) and light-blue arrows in (b) and (d) indicate the moistened and dried upstream air mass, respectively. In (a)–(d), black arrows and a broken line indicate the low-level wind and the synoptic-scale wind convergence line, respectively. The light-blue region depicts the weak precipitation area while the dark-blue region shows the intense precipitation area. The blue solid line illustrates the southern edge of cold pool while the closed area with blue dashed line in (b) and (d) indicates the absence of intense offshore rainfall.