Ambient measurement of shipping emissions in Shanghai port areas

Xinning Wang¹, Yin Shen¹, Yanfen Lin¹, Jun Pan¹, Yan Zhang², Peter K.K. Louie³, Mei Li⁴, Qingyan Fu¹

5 ¹ Shanghai Environmental Monitoring Center, Shanghai 200030, P.R. China.

² Department of Environmental Science and Engineering, Fudan University, Shanghai 200433, P.R. China.

⁴ Institute of Mass Spectrometer and Atmospheric Environment, Jinan University, Guangzhou, 510632, China.

Correspondence to: Qingyan Fu (qingyanf@semc.gov.cn)

10

Abgreact. Growing shipping activities in port areas have generated negative impacts on climate, air quality and human health. To better evaluate the environmental impact of shipping emissions, in the summer of 2016 ambient air quality measurement was carried out at Shanghai port, one of the busiest port in the world. Gaseous (NO, NO₂, SO₂, O₃) and

- (15) particulate concentrations (PM_{2.5}), particle sizes and chemical composition of individual shipping emission particles were continuously monitored for 3 months. High temporal resolution data show that shipping emissions is a major culprit of local air pollution problem. Shipping emission plumes were clearly observed using online measurement in port area. The SO₂ and Vanadium particles numbers were found to correlate best with shipping emissions in Shanghai port. Single particle mass spectra of fresh shipping emission were identified based on the dominant peaks of Sulfate, EC and indicative metals of V, Ni,
- Fe and Ca, and nitrate peaks in aged particles. Fresh particles from ship emission mainly concentrated in ultra-fine size range where their number contributions are more apparent than their mass. Advanced measurement conducted in the present study show that in port region shipping emissions contributed 36.4 % SO₂, 0.7 % NO, 5.1 % NO₂, -0.9 % O₃, 5.9 % PM_{2.5}, 49.5 % Vanadium particles if land-based emissions were included, and 57.2 % SO₂, 71.9 % NO, 30.4 % NO₂, -16.6 % O₃, 27.6 % PM_{2.5}, 77.0 % Vanadium particles if land-based emissions were excluded.
- 25 Keywords

Shipping emission; Shanghai port; emission source contribution; SPAMS

1 Introduction

S Ship emission constitutes an important source of gaseous and particulate pollution world wide. Growing shipping activities in recent years are attracting much attention to assessing its impact on environment and health (Fuglestvedt et al., 2009). For

30 emissions from sea-going vessels in pristine marine environment, it is found that ship emission affect clouds properties along



³ Hong Kong Environmental Protection Department, Hong Kong, P.R. China.

cruising route (Petzold et al., 2008;Coggon et al., 2012), which is directly relevant to earth radiation budget and climate issues. In portside or coastal regions, ship emissions could have a negative impact on the air quality at variant degrees in these areas or eities (Donateo et al., 2014;Liu et al., 2017). growing contributions of air pollution from ship emissions, its negative effects on human health of coastal residents has been studied (Corbett et al., 2007).

- 5 The typical fuel that ships burn is Residual Fuel Oil (RFO) usually has high sulfur content. Combustion of RFO in ship engines produces high concentration of gaseous and particulate pollutants including NOx, SO2, Elemental Carbon (EC), Organic Carbon (OC), sulfate and trace metals. Emission Factors of these pollutants from various ship types have been determined to develop emission inventories (Moldanova et al., 2013;Buffaloe et al., 2014;Cappa et al., 2014). In ambient measurement, however, the chemical and physical attributes of ship emissions are critical for identifying ship emission and
- 10 assess their impacts (Murphy et al., 2009). Owing to more stringent regulations toward ship emissions by restricting sulfur content in fuel, the detection of ship emissions relying only on individual tracers is unreliable because of the changing composition RFO in different areas. To better identify ship emission in this context, multi-components characterizations including both gaseous and particulate measurements are found necessary in ambient studies (Viana et al., 2009;Xiao et al., 2018).
- 15 In Yangtze River Delta (YRD) region in China the shipping activities increase significantly due to intensified international trades in recent years. The accompanying potential environmental and health problems from shipping in YRD are well recognized (Fu et al., 2017;Zhang et al., 2017;Chen et al., 2018). Global distributions of ship trades in indicate that South and Eastern China Sea region have the highest emission densities (Johansson et al., 2017). As shown in an emission inventory in China, shipping traffics emitted about 1.3 Tg SO2, 1.9 Tg NOx and 0.16 Tg PM in 2013, with NOx and PM
- 20 being equivalent to ~ 34 % and 29 % of total mobile vehicle emissions in China (Fu et al., 2017). To cope with severe air pollution caused by shipping emissions, Shanghai government has initiated Domestic Emission Control Areas (DECA) in YRD to control air pollution from shipping activities. At present stage, by way of YRD DECA regulations, the sulfur content of any fuel used on board while berthing at Shanghai port shall not exceed 0.5 % (m/m), except for the first hour after arrival and the last hour before departure, which has taken effect on April 1, 2016. More stringent regulations of sulfur content 1 limitations of 0.5 % and eventually 0.1 % in DECA are in preparation.
- Despite the high emissions level from ship traffics in YRD, ambient measurements on ship emissions in this area are very limited except studies on ship emission inventories or modeling (Liu et al., 2017;Zhang et al., 2017;Chen et al., 2018). In order to eonstraint the uncertainties inherent to inventories or simulations studies it is essential that the validation is compared with the actual measurement data (Zhao et al., 2013). In the summer of 2016, an in-site experimental
- 30 eharacterization focusing on ship emissions were performed at Shanghai Port. Gaseous and particulate matters concentrations were achieved by online monitoring for 3 months to identify and characterize the ship emissions in port areas. Based on the measurement data, quantitative assessment of the contribution from ship emissions on portside air quality was performed. ping emission aerosol particles were characterized by a single particle aerosol mass spectrometer (SPAMS) which was deployed at the same site in parallel to the gaseous measurement. The SPAMS were utilized to identify ship
 - 2

emission aerosol composition and size with high temporal resolution, which is useful in detecting fast transient ship plumes, as demonstrated previously (Healy et al., 2009;Ault et al., 2010). In addition, the ship emission particle signatures obtained here is valuable in SPAMS source apportionment in future studies. The present study represents an advanced comprehensive characterization of gaseous and particulate ship emissions in YRD and serves to provide essential scientific supporting information for future evidence-based ship emission control policies formulation.

5

2 Experimental

2.1 Sampling site

The Waigaoqiao Port (31.337° N, 121.665° E) locates in the northeast of Shanghai city (Fig. 1) and is the largest port in China. The port has about 7 km of docks (3 km north section and 4 km south section). In 2016 the port has yearly traffic of

- 10 367 M-tons of goods and container volume of 37.13 million TEU. Ship categories in port consist of container vessel (62.4 %), tug (18.6 %), oil tanker (9.0 %), bulk (1.8 %), Ro-Ro (1.7 %) and other ships (6.5 %) (private data from authority). A power plant and a shipbuilding factory reside between the north and south section of port, which have their own docks. The in-port air monitoring station locates on the south bank of Yangtze River, 400 m away from the nearest dock. The outlets of main sampling tube, fixed on the roof of the station, are 3.5m above the ground. To the south and west of site there
- 15 are intense road traffics of container trucks in surrounding regions and the Shanghai outer ring. Traffic emissions in this direction have important influences to air pollution in monitoring site when inland wind prevails. The impact of land traffic emission will be discussed in following sections.

2.2 Gaseous, PM2.5 and peripheral measurement

In port gaseous concentrations of NO-NO₂-NO_x, SO₂, and O₃ were measured continuously from Jun-21 to Sep-21, 2016. The gaseous pollutant monitors are a suit of Thermal Scientific analyzers (NO-NO₂-NO_x, model 42i; SO₂, model 43i; O₃, 49i) installed in the monitoring station. Calibration and maintenance of instruments were regularly performed according to the requirement of relevant national standards in China. The PM_{2.5} concentrations are monitored with the oscillating microbalance method (Thermo TEOM 1405-F). Ambient conditions (temperature, humidity, pressure, wind speed and direction) are monitored by a meteorological monitor on the rooftop of the station, which is about 3.5m above the ground.

25 Data from all instruments are set to 5 min resolution and is managed in a customized database. The pollution level in Shanghai city area, including gaseous and $PM_{2.5}$ concentrations, are monitored continuously by 9 national air quality monitoring stations. Pollutants averaged concentrations at these 9 stations during the same monitoring period were included as a reference.

2.3 Single particle aerosol mass spectrometer (SPAMS)

During the period from Jun-21 to Sep-21, 2016, a SPAMS (HeXin Analytical Instrument Co., Ltd., China) concurrently characterized single particle chemical composition and size of ambient aerosol in real time (Li et al., 2011). Operation principle of SPAMS is briefly described here. Ambient aerosol is drawn into SPAMS vacuum region through a critical

- 5 orifice with limited aerosol flow. Aerosol particles then enter an aerodynamic focusing lens (AFL) where they are focused into a thin beam with transiting velocities as a function of particle aerodynamic size. In the SPAMS sizing region particles consecutively encounter two continuous laser beams (532 nm wavelength), reflect light and generate signals in two PMTs. The time lag between two PMT signals is used to calculate particle velocity and to trigger the third laser (266 nm wavelength) at appropriate time to ionize the same particle. Chemical composition of ionized particle is measured by a dual polar time-of-
- 10 flight mass spectrometer. The time lags between two PMTs of PSL particles of known size are used to calibrate the aerodynamic size of ambient particles. Particle size, dual polar mass spectra, particle reflect signals from two PMTs are saved for each particle. A PM_{2.5} cyclone was placed at the outlet of sampling tube to obtain the PM_{2.5} cut-point before detection by SPAMS.

2.4 Data analysis

- 15 The high temporal resolution of SPAMS makes it very suitable to couple with online gaseous data to identify ship emissions. The quick fluctuations of gaseous concentrations, shifting of wind and the arrival of particle plumes, were well registered in the high resolution measurement by SPAMS. In addition, present study takes advantage of the unique power of SPAMS to identify individual shipping emission particles based on particle fingerprint. Mass spectral pattern of shipping emission particles were first identified and then were utilized to extract them out. The temporal trends, size distribution, chemical composition, and wind rose of the searched particles could be examined in further detail.
- During sampling of 3 months a large amount of particles were chemically analyzed by SPAMS (>2.3 million mass spectra). The identification of shipping emission particles from 2.3 million total particles were based on a combined method of peak searching and algorithm clustering to achieve better performance. Specifically, the SPAMS data are pre-analyzed by visually inspecting individual particle mass spectra to identify MS patterns of shipping emission particles during ship plumes. The
- concurrent SO2 concentrations were utilized to locate ship emission plumes when sharp SO₂ peaks occurred, which is typical for RFO combustions (Murphy et al., 2009;Merico et al., 2016). Compared with non-plumes period, the most important indicating peaks occurred at V⁺(51), VO⁺(67), Fe⁺(56), Ni⁺(58) and serial peaks of elemental carbon at C_n⁺(n=1,2,3...,12) in the positive mass spectrum. The obtained mass spectra of shipping emission particles are similar as detected previously in other port regions (Ault et al., 2009;Healy et al., 2009;Ault et al., 2010). In this study the Vanadium mass peaks (peak V⁺(51))
- and VO⁺(67))) were determined to be a prerequisite to indicate ship particles during plumes. Further notes on this particle identification method from shipping emission are seen in the supplementary material. Peak searching method of criteria of m/z = 51 and 67 (i.e., only the existence of mass peak at 51 and 67, no peak area limitation) was firstly applied to search all

possible candidates from all analyzed particles. This search criterion is not too stringent because particles producing organic peaks at the same nominal mass (e.g. $C_4H_3^+(51)$, $C_4H_3O^+(67)$) could interfere and may enter into searched clusters. Then the ART-2a algorithm (Vigilance=0.85; Learning=0.05; Iteration=20) was applied to the searched clusters to generate subclusters of particles. By inspecting composition, size and wind rose patterns of sub-clusters, a small fraction of outlier particles from non-shipping emission sources were thus picked out and discarded.

3 Results and discussions

3.1 Identification and statistics of ship emission plumes

In the vicinity of port, the measured ship emission pollutants concentrations often produce sharp peaks in relatively short period (Fig. 2). The sharp peaks are produced by ship emission plumes corresponding to shipping activities such as arrival, hoteling and departure, which typically persist for a few hours. The measured SO₂, NO, NO₂, O₃ and PM_{2.5} concentrations during a typical period (Aug 27 -29) are shown in Figure 2 to illustrate several ship plumes. For comparison purpose the averaged SO₂ concentration in Shanghai city and Vanadium particle number concentration during the same period are included. It is clear that during plumes tracking period, the SO₂ concentration peaks were well correlated with Vanadium particles number concentration as detected by SPAMS. This result is in good agreement with previous characterizations

15 studies in portside (Healy et al., 2009;Ault et al., 2010;Merico et al., 2016). During the whole sampling period the Shanghai city SO₂ concentration matched very well with ambient SO₂ concentration in port area during non-plumes periods. This is also consistent with the observation data that pollution plumes arising from shipping emission superimpose on background and regional SO₂.

As typical combustion products, the NO and NO₂ concentrations also show corresponding elevated concentrations during

- 20 plumes under favorable wind fields (Fig. 2). However, during the whole study NO and NO₂ are more importantly influenced by land traffics emissions (mostly from transportation diesel trucks) when the inland wind prevails. The shipping emission NO_x plumes reached sampling site have been slightly aged. To understand the aging effect, when wind direction is in port sector ($300^{\circ}-0^{\circ}-120^{\circ}$), the averaged NO/NO₂ ratio is 0.6 (mostly fall in 0.1~4.5 range), lower than typical ratio of 4 at ship exhaust (Alföldy et al., 2013), suggesting the oxidation of primary NO into NO₂ have occurred for some time (1~50 min,
- based on wind speed measurement and transportation distance). This result is evidenced by the apparent consumption of O_3 in plumes as shown in Figure 2, commonly termed as titration effect between NO_x and O₃. The PM_{2.5} mass concentration did not show as apparent response as that of SO₂ during ship emission plumes, despite that the typical Vanadium particles had reached the site, as shown. The obscure response of PM_{2.5} to ship plumes is explained by the fine particle sizes in relatively fresh shipping emissions, as discussed in following section.
- 30 Taking into consideration of all the relevant factors, the present study defines ship plume periods by ways of SO₂ concentrations and Vanadium particle number concentrations. For SO₂, a minimum threshold of Δ_{SO2} = SO₂(Port)-SO₂(Shanghai) > 5 ug/m³ is applied to indicate ship plumes. For shipping emission particles, the number concentration of

Vanadium particle is considered because in some cases the SO₂ peaks are absent or obscure as typical fresh Vanadium particles are indeed mount up quickly. This is probably caused by anchored ships using oil of low Sulfur contend (<0.5 % m/m) to comply with new regulations in Shanghai port, which came into force on April 1, 2016. In this study the threshold of Vanadium particle detection speed in ship plumes are set to $C_V > 25$ particles/hour. That is, ship plumes are defined as either $\Delta_{SO2} > 5\mu g/m^3$ or $C_V > 25$ particles/hour.

- In total about 210 shipping emission plumes were captured during the sampling campaign. Table 1 summarizes the statistics on pollutants concentration of SO₂, NO, NO₂, O₃, PM_{2.5} in port area and Shanghai city during the sampling period. Vanadium particles number concentrations are represented by particle detection speeds by SPAMS. It is stated that the SPAMS detection speed are positively correlated with particle number concentrations in ambient air, but should not be
- 10 explained as absolute number concentrations without correction for SPAMS efficiency (Wenzel et al., 2003). Statistics are performed on pollution level in plume periods and in non-plumes. To separate influences from land sources (principally traffics), non-plume periods during wind from port direction are calculated in Table 1. In general, the port site has concentrations of combustion products SO₂ and NO_x much higher than that in Shanghai city regions (Table 1). For SO₂ its concentrations in non-plumes are comparable with that in Shanghai city, regardless of wind
- 15 direction, representing a background SO₂ level. However, NO_x concentrations in non-plumes from port sector wind are significantly lower than from land directions. This result suggests that shipping activities are the main source of SO₂ plumes in port. For NO_x, due to the specific geographic proximity of Waigaoqiao port, its concentration is significantly influenced by land-based traffics. In general the ozone concentrations in port are lower than Shanghai urban region by 13-33 %. The lower O₃ level around portside is consistent with higher SO₂ and NO_x concentrations in this area to consume O₃. For PM_{2.5},
- 20 its concentrations in port area are slightly lower than Shanghai city either in plume or non-plume periods, although Vanadium particles concentrations in plumes are about 4 times higher than in non-plumes (Table 1). Longer period of $PM_{2.5}$ data at the same station supports the lower $PM_{2.5}$ concentration compared to Shanghai city, suggesting that it is a general trend at this port site. This is probably caused by the clean air advection from the sea and the wiping out effect of adjacent water surface through particle sedimentation. The Vanadium particle number fraction in total particles, as measured by 25 CRAMO is a local back of 20%
- 25 SPAMS, is clearly larger (6.7 % on average) in portside than in the urban area in Shanghai (1-2 %) (Liu et al., 2017).

3.2 Characterization of single particle properties from ship emissions

3.2.1 Background and fresh shipping emission particles in port

5

Particle size and chemical characterization from ship emission was performed by a SPAMS (Li et al., 2011) in parallel with gaseous measurement. The close proximity of the monitoring site to docks makes it possible to detect very fresh ship emission particles in addition to aged ones (Healy et al., 2009). By using single particle characterization techniques, fresh or 'pure' ship emission particles can be separated from background or aged aerosol based on single particle signatures. It was observed that the fresh ship emission particles in plumes are characterized by the absence of or very low nitrate (-62NO₃ in

negative spectra) signal in mass spectra, a pattern commonly found in combustion source characterizations (Spencer et al., 2006;Toner et al., 2006). In another respect, the temporal pattern, wind rose and size of nitrate containing Vanadium particles were disproportionately distributed compared with fresh ones, which distinguished themselves as background ship emission particles (Healy et al., 2009;Ault et al., 2010) (Fig. 3). The mass spectra of fresh ship emission particles and aged

- 5 ones are shown in Figure 3. The dominant peaks in mass spectra of fresh ship emission particles are Sulfate (-97HSO₄⁻), EC (C_n^+/C_n^- , n are integers), and Vanadium (51V⁺, 67VO⁺) peaks, reflecting the major components found in ship emission particles (Moldanova et al., 2013;Becagli et al., 2012;Murphy et al., 2009). Except for the nitrate peak (-62NO₃⁻), other mass spectral patterns of background and fresh ship emission particles are similar (Fig. 3). Although OC is an important composition of ship emission particles (Lack et al., 2009), the organic mass peaks in SPAMS spectra are insignificant
- 10 compared with metal and EC peaks. It maybe owing to the fact that SPAMS is not so sensitive to nic species due to the low ionization efficiencies in laser ionization (Ulbrich et al., 2009). However, there is an organic cluster of particles were indeed identified in ship plumes, although of minor number fractions, as discussed in subsequent section.
 The wind rose distribution of fresh and background Vanadium particles are shown in figure 3. It is clear that fresh and
- background Vanadium particles have different size and wind rose patterns (Fig. 3, 4). The fresh Vanadium particles have wind rose pattern which runs almost parallel direction with riverbanks (300°-0°-120°). This is strong evidence that ships are the most predominant source of fresh vanadium particles in Shanghai port. Background Vanadium particles, however, did
- not show any prominent source directions and displayed nearly uniform concentrations in all directions. It is reasonable to assume that the aged Vanadium particles are background particles which have undergone atmospheric processing in local or regional scale.
- 20 The size distributions of Vanadium particles as shown in Figure 4 indicates fresh Vanadium particles with dominate particle numbers in small size range (<0.5 um), compared with background ones. Although SPAMS detection efficiency declines in smaller size range because of the smaller section to reflect laser light, significant fine ship emission particles were still detected in fine size range. The explanation is that these particles are non-spherical fractal agglomerates, such as BC particles, having significantly larger cross sections to reflect laser light and thus be detected in SPAMS. The non-spherical
- 25 fractal shape of fresh Vanadium particles is consistent with the typical particle shapes from fresh combustion sources. Similar observations were observed were also reported in other studies using single particle mass spectrometer in ultra-fine size range (Ault et al., 2010).

The size distribution agrees with the sizes of ship emission particles measured in other studies (Gonzalez et al., 2011;Merico et al., 2016). It is reported that for ship exhaust their particle number contributions are more considerable than their mass contributions (Donateo et al., 2014;Jonsson et al., 2011;Merico et al., 2016). Size distribution of fresh particles from ship

exhaust shows that the number concentration mainly concentrated in UF mode (<100 nm) (Gonzalez et al., 2011;Moldanova et al., 2013). The less significant increase in PM_{2.5} mass concentration than NO_x and SO₂ in plumes are due to, in one respect, that the emission factors of PM are typically much lower than NO_x and SO₂ (Agrawal et al., 2010;Moldanova et al., 2013), and second, that the smaller size of fresh vanadium particles before secondary accumulation happens in the atmosphere

7

(Moldanova et al., 2013). Considering that the secondary processing occurred on every particle, the particle number concentrations of fresh shipping emissions is more significance than their initial mass concentrations. The study suggests that particle number quantification is more appropriate in an accurate evaluation of primary ship emissions.

3.2.2 Particle types in fresh ship emission plumes

- 5 After the separation of background ship emission particles, fresh Vanadium particles are isolated to represent fresh ship emission particles. Further analysis is performed to study their composition and emission characters, which will be helpful in particle source identification of SPAMS in future studies. In general, the fresh Vanadium particles could be grouped into 4 types based on their chemical composition: V-OC, V-EC, V-ECFe and V-Ash, and the average mass spectra are shown in Figure 5. The negative mass spectra of the four types are common in the dominating SO₄⁻ peak in addition to other negative
- 10 EC peaks in spectra, consistent with the high SO₂ concentration in ship emission-plumes. The major chemical differences of the four particle types are in the positive mass spectra as depicted in Figure 5. The V-OC type are characterized by the dominate OC peaks $C_2H_3^+, C_2H_5^+, C_2H_3O^+$, etc., with non or insignificant EC peaks. Considering the low ionization probability of OC in laser ionization, this particle type is deemed to be mainly composed of condensed organics in engine exhaust plume (Lack et al., 2009;Moldanova et al., 2013). The V-EC particles produce dominant EC peaks from C_1^+ to C_{13}^+
- 15 and metal peaks of V and Na, but without iron peaks Fe⁺. This type is also the most abundant type of all Vanadium particles. The V-ECFe type is similar as V-EC except for the addition of Fe⁺ and Ca⁺, Ni⁺ peaks of lower frequencies. The V-Ash particles produce minor or no EC peaks except some metal peaks of V, Fe and Ni in positive spectra. These metals are used as lubricant additives or inherently present in RFO, therefore their presence in ship emission particles are expected and commonly found (Becagli et al., 2012;Moldanova et al., 2013).
- 20 The chemical composition and size distribution of these four types suggest they have distinct physical properties (Fig. 6). Temporal trends of these 4 types show that their concentrations display daily fluctuations, with higher concentrations in daytime than nighttime. However, the inter-correlations among their concentrations are low ($r^2 < 0.4$), suggesting the different emission process among them (variant engine types, operation modes, etc.). The V-OC particles, although having low ionization probabilities, are found to concentrate in specific cases of plumes. Because the information of individual ships
- 25 is not yet available, it is therefore not attempt to directly link V-OC particle plumes to specific ship types in the present study. The V-OC particle type probably corresponds to the amorphous organic particles in TEM images, usually having inner metallic impurities (Moldanova et al., 2013), as indicated by the metal peaks in mass spectrum. In incomplete combustion (e.g., starting up phase) the organic vapors in fuels will condense onto particles (ash, EC particles) in the cooler ambient environment, resulting uniform size distribution compared with other types (lower right panel in Figure 6). The V-OC type
- 30 are more transient in that the peak width of its concentration peaks are normally narrower than other types (~1 hours contrasting to 3~5 hours of other types), suggesting they are principally emitted in specific phases of engine operations. The V-EC particles are the most abundant types found in ship plumes in this study. As mentioned earlier the enrichment of EC particles in UF size regions indicates the fractal agglomerates shapes typical of soot from engine, which is a common
 - 8

product of combustion of RFO (Moldanova et al., 2013). The V-Ash particles, which is most probably the ash spheres from combustion process of inorganic constituents in RFO and lubricants, are mainly detected in larger size range (> 0.5 um) (Fig. 6). It is reminded that the measured aerodynamic size by SPAMS is dependent both on particle physical diameter and particle densities. Therefore the larger size of V-Ash particles does not imply they are physically larger, because the ash spheres are mainly composed of metal oxides or salts of higher effective densities than EC. The origin of V-ECFe types are probably the result of internal mixing between V-EC and V-Ash particles. Their size distribution is more similar as V-Ash type, suggesting that the adhesion of fractal EC particles to ash spheres has not change the diameter of the latter significantly.

3.3 Contributions of ship emission to ambient pollutants in port area

5

- Quantification of ship emissions impact on regional and local air quality is an important issue to assess its impact on environment (Donateo et al., 2014;Aulinger et al., 2016;Merico et al., 2016). In East Asia, an earlier emission inventory in Shanghai area estimated that the ship emissions were 58160, 51180, 6960 tons/year for NO_x, SO₂ and PM respectively in 2003 (Yang et al., 2007). Over the last decade Shanghai port throughput of goods has dramatically increased. In 2010, the total ship emissions of NO_x, SO₂ and PM_{2.5} in YRD have grown to 7.1×10^5 , 3.8×10^5 and 5.1×10^4 tons/year, respectively (Fan et al., 2016). A more recent study estimated that the primary PM_{2.5} from ships ranged from 0.63 to 3.58 µg/m³,
- (15) accounting for 4.23 % of the total PM_{2.5} in Shanghai Port (Zhao et al., 2013), based on a marine port measurement off coast of Shanghai. Such information of port in coastal areas is needed since their closer distance to the urban of Shanghai city. In this study, the ambient measurement data in a coastal port is utilized to quantitatively assess the contribution of ship emissions to air quality, making current study more comparable with others.
- Due to its close proximity to urban region, the measurement site is under more important influences from land emissions than the marine port far from coastal area (Zhao et al., 2013). To give a more nuance illustration, the role of wind fields of portside on influencing the measured pollutants are shown for SO₂, O₃, NO, NO₂, PM_{2.5} and Vanadium particle numbers (Fig. 7); Wind roses of some pollutants (NO, NO₂ and PM_{2.5}) clearly showed that they are under the overwhelming influences of land emissions on the sampling site. On the contrary, the SO₂ concentrations and Vanadium particle numbers are dominant only when winds from port sectors. The hotspots in wind rose of Vanadium particle are most probably produced by
- 25 individual docks along the riverside. The wind dependence of Θ_{z} one concentrations is less apparent, except showing depletion in regions of high NO_x and SO₂ levels in wind roses, as previously explained. Because the air pollution in this two conditions are so different, it is appropriate to separate land-based emissions influences by limiting wind directions only to port directions. Therefore, in the present study two reference periods are considered in calculating shipping contributions: the entire study period (irrespective of wind) and only when the site is in downwind of port direction.
- 30 The calculations method of ship contributions is based on the extraction of ship emission plumes from background concentrations of pollutants (Merico et al., 2016):

$$\varepsilon_A = \frac{\Delta C_A F_{plm}}{C_A}$$

Where: ε_A , ship emission contributions of pollutants A; ΔC_A , the difference between average concentrations during plumes and non-plumes; F_{plm} , fraction of cases of plumes; C_A , the average concentration of pollutant A during reference period. Shipping emission contributions of respective pollutants in two reference periods are summarized in Table 2.

As shown in Table 2, shipping emission contribution for gaseous SO₂ (36.4 %) are much larger than for NO (0.7 %), NO₂

- 5 (5.1 %), and for PM_{2.5} (5.9 %), if not considering the factor of wind direction (entire period). The low contributions of NO_x from shipping are due to the inclusion of land emissions (mostly traffics) of stronger intensities. However, 49.5 % Vanadium particles numbers are contributed from shipping emissions despite the inclusion of land emissions, suggesting the importance of number concentration from shipping emissions. It is worth noting that in the entire period calculations, the frequency of winds at respective directions during the sampling period influences the calculated contributions. The calculations in this
- (10) way may somewhat overestimate ship emissions because of the prevailing southeast wind during summer in Shanghai.
 (10) During the period when southeast winds are prevalent, adjacent coastal regions other than port area may experience more ship emissions.

By limiting the sampling with time windows to periods when winds from port sector, the influences of land emissions could be largely eliminated, in which a less biased estimations of shipping emission are obtained. As shown in table 2, by

- 15 considering port sector wind, for all pollutants the ship emission contributions are magnified in amplitude. The most dramatic change occurs for gaseous NO_x , whose contributions from shipping upgraded to levels larger or comparable with SO₂. The larger contribution of primary NO than NO_2 and SO_2 indicates that the plumes are relatively fresh ones. Contributions obtained here can be compared with a very similar study carried out in a European port (Merico et al., 2016). Gaseous emissions of NO, NO_2 and SO_2 are similar between these two studies, which is an impressive result considering the
- 20 much larger throughput of goods in Shanghai port. However, the main differences between two studies are the larger contributions from the shipping for O₃ depletion (-16.6 % vs. -5 %) and PM_{2.5} concentrations (27.6 % vs. 9 %) in Shanghai port. In that study particle number contributions from shipping emissions (<50 %) are smaller than this study as measured by SPAMS (77 %).

In the marine port study at Yangshan island of Shanghai, the calculated PM_{2.5} contribution (~4 %) is smaller than present

- 25 study (5.9 %) (Zhao et al., 2013). In the Zhao et al study a different method is used to evaluate ship emissions, relying on Vanadium concentrations to indicate ship emissions. Considering the methodology differences, it is deemed that the results from the two studies are similar if land-based emission sources were included in Waigaoqiao port. A previous estimation in Shanghai area using inventories method showed that ship emissions contributed 9 % NO_x and 5.3 % PM_{2.5} in Shanghai area (Zhang et al., 2017), generally agrees with this study in the condition of including land-based emissions (Table 2). However,
- 30 for SO₂ the contribution in that estimation (12 %) is significantly smaller than the 36.4 % in this study. It is worth noting that the high SO₂ contributions in this study only represent the port site to emission sources. The detected pollutions in the port site are usually found to be fresh and with elevated concentrations. With transportations to the urban region their concentrations will dissipate and strength weakened. Quantifications of ship emission contributions in the urban regions will be the subject of future studies and beyond the scope of the present study.

4 Conclusions

The gaseous and particulate emissions from shipping emissions are measured in Shanghai port to study their emission eharacters. Pollutants from ship emission in port often result sharp peaks of concentrations during plumes passing the site. Of the measured pollutants (SO₂, NO, NO₂, O₃, PM_{2.5} and Vanadium particles), SO₂ and Vanadium particles show the best

- 5 responds to ship emission plumes. Using the SO_2 and Vanadium particles concentration as indicators, ship emission plumes can be successfully identified from background atmosphere. Statistics of pollutants during plumes show that the concentrations of SO_2 in plumes are about 3 times higher than the background concentrations. In non-plumes SO_2 concentrations in port share the trend of regional background level. Results show that the NO_x concentrations in port site are under much stronger influences from land emissions. The O_3 is depleted by primary NO_x and SO_2 emissions around the port,
- 10 resulting 11-33 % Ozone consumption compared with urban region of Shanghai. For particulate matters, shipping emitted PM is more embodied by the number concentrations rather than the mass concentrations (PM_{2.5}) during plumes. Individual particle size and chemical composition from shipping emission are online characterized in parallel with other measurement. It is found that, similar as SO₂, the ship emission particles in port could also be divided into fresh and background types. The mass spectra of fresh ship emission particles produce dominant peaks of EC, sulfate and trace metals
- 15 (V, Ni, Fe and Ca). Due to the low ionization efficiencies of organics, OC is only identified only in a small group of particles. Size distributions show that ship emission particles are more concentrated in fine size range (< 0.5 μm), which is mainly fractal EC agglomerates. Further classifying of ship emission particles indicates that trace metals are present in ash spheres of larger aerodynamic diameters, either externally or internally mixed with EC particles.</p>
- Based on the measured gaseous and particulate concentration, shipping emission contributions to the air quality in Shanghai 20 port area is quantitatively evaluated. The contributions of ship emissions are evaluated in two scenarios where the land emission sources are either included or excluded. With the inclusion of land-based emission sources, contributions of ship emission are mainly embodied in SO₂ (36.4 %) and Vanadium particle number concentrations (49.5 %). The ship emissions to PM_{2.5} is 5.9 % if include the land emission sources. NO_x contribution from shipping emissions is insignificant if include land traffic sources. However, if land sources are excluded, shipping contributions of NO_x became prominent comparable
- 25 with that of SO₂. 16.6 % of Ozone concentration is found to be depleted by ship emitted NO_x and SO₂ during wind from port direction. With the exclusion of land sources, ship emissions are found to contribute 27.6 % of PM_{2.5} and 77 % Vanadium particle numbers in port area.

References

Aman, A. A. and Bman, B. B.: The test article, J. Sci. Res., 12, 135-147, doi:10.1234/56789, 2015.

Aman, A. A., Cman, C., and Bman, B. B.: More test articles, J. Adv. Res., 35, 13–28, doi:10.2345/67890, 2014.
 References

Agrawal, H., Welch, W. A., Henningsen, S., Miller, J. W., and Cocker, D. R., III: Emissions from main propulsion engine on container ship at sea, Journal of Geophysical Research-Atmospheres, 115, 10.1029/2009jd013346, 2010.

Alföldy, B., Lööv, J. B., Lagler, F., Mellqvist, J., Berg, N., Beecken, J., Weststrate, H., Duyzer, J., Bencs, L., Horemans, B., Cavalli, F., Putaud, J. P., Janssens-Maenhout, G., Csordás, A. P., Van Grieken, R., Borowiak, A., and Hjorth, J.:

- 5 Measurements of air pollution emission factors for marine transportation in SECA, Atmos. Meas. Tech., 6, 1777-1791, 10.5194/amt-6-1777-2013, 2013.
 - Aulinger, A., Matthias, V., Zeretzke, M., Bieser, J., Quante, M., and Backes, A.: The impact of shipping emissions on air pollution in the greater North Sea region Part 1: Current emissions and concentrations, Atmospheric Chemistry and Physics, 16, 739-758, 10.5194/acp-16-739-2016, 2016.
- 10 Ault, A. P., Moore, M. J., Furutani, H., and Prather, K. A.: Impact of Emissions from the Los Angeles Port Region on San Diego Air Quality during Regional Transport Events, Environmental Science & Technology, 43, 3500-3506, 10.1021/es8018918, 2009.

Ault, A. P., Gaston, C. J., Wang, Y., Dominguez, G., Thiemens, M. H., and Prather, K. A.: Characterization of the Single Particle Mixing State of Individual Ship Plume Events Measured at the Port of Los Angeles, Environmental Science & Technology, 44, 1954-1961, 10.1021/es902985h, 2010.

- Becagli, S., Sferlazzo, D. M., Pace, G., di Sarra, A., Bommarito, C., Calzolai, G., Ghedini, C., Lucarelli, F., Meloni, D., Monteleone, F., Severi, M., Traversi, R., and Udisti, R.: Evidence for heavy fuel oil combustion aerosols from chemical analyses at the island of Lampedusa: a possible large role of ships emissions in the Mediterranean, Atmospheric Chemistry and Physics, 12, 3479-3492, 10.5194/acp-12-3479-2012, 2012.
- 20 Buffaloe, G. M., Lack, D. A., Williams, E. J., Coffman, D., Hayden, K. L., Lerner, B. M., Li, S. M., Nuaaman, I., Massoli, P., Onasch, T. B., Quinn, P. K., and Cappa, C. D.: Black carbon emissions from in-use ships: a California regional assessment, Atmospheric Chemistry and Physics, 14, 1881-1896, 10.5194/acp-14-1881-2014, 2014. Cappa, C. D., Williams, E. J., Lack, D. A., Buffaloe, G. M., Coffman, D., Hayden, K. L., Herndon, S. C., Lerner, B. M., Li,
- S. M., Massoli, P., McLaren, R., Nuaaman, I., Onasch, T. B., and Quinn, P. K.: A case study into the measurement of ship
 emissions from plume intercepts of the NOAA ship Miller Freeman, Atmospheric Chemistry and Physics, 14, 1337-1352, 10.5194/acp-14-1337-2014, 2014.

Chen, D., Zhao, N., Lang, J., Zhou, Y., Wang, X., Li, Y., Zhao, Y., and Guo, X.: Contribution of ship emissions to the concentration of PM2.5: A comprehensive study using AIS data and WRF/Chemmodel in Bohai Rim Region, China, Science of the Total Environment, 610, 1476-1486, 10.1016/j.scitotenv.2017.07.255, 2018.

30 Coggon, M. M., Sorooshian, A., Wang, Z., Metcalf, A. R., Frossard, A. A., Lin, J. J., Craven, J. S., Nenes, A., Jonsson, H. H., Russell, L. M., Flagan, R. C., and Seinfeld, J. H.: Ship impacts on the marine atmosphere: insights into the contribution of shipping emissions to the properties of marine aerosol and clouds, Atmospheric Chemistry and Physics, 12, 8439-8458, 10.5194/acp-12-8439-2012, 2012.

Corbett, J. J., Winebrake, J. J., Green, E. H., Kasibhatla, P., Eyring, V., and Lauer, A.: Mortality from ship emissions: A global assessment, Environmental Science & Technology, 41, 8512-8518, 10.1021/es071686z, 2007.

Donateo, A., Gregoris, E., Gambaro, A., Merico, E., Giua, R., Nocioni, A., and Contini, D.: Contribution of harbour activities and ship traffic to PM2.5, particle number concentrations and PAHs in a port city of the Mediterranean Sea (Italy), Environmental Science and Pollution Research, 21, 9415-9429, 10.1007/s11356-014-2849-0, 2014.

Fan, Q., Zhang, Y., Ma, W., Ma, H., Feng, J., Yu, Q., Yang, X., Ng, S. K. W., Fu, Q., and Chen, L.: Spatial and Seasonal Dynamics of Ship Emissions over the Yangtze River Delta and East China Sea and Their Potential Environmental Influence, Environmental Science & Technology, 50, 1322-1329, 10.1021/acs.est.5b03965, 2016.

5

- Fu, M., Liu, H., Jin, X., and He, K.: National- to port-level inventories of shipping emissions in China, Environmental
 Research Letters, 12, 10.1088/1748-9326/aa897a, 2017.
- Fuglestvedt, J., Berntsen, T., Eyring, V., Isaksen, I., Lee, D. S., and Sausen, R.: Shipping Emissions: From Cooling to Warming of Climate-and Reducing Impacts on Health, Environmental Science & Technology, 43, 9057-9062, 10.1021/es901944r, 2009.

Gonzalez, Y., Rodriguez, S., Guerra Garcia, J. C., Luis Trujillo, J., and Garcia, R.: Ultrafine particles pollution in urban coastal air due to ship emissions, Atmospheric Environment, 45, 4907-4914, 10.1016/j.atmosenv.2011.06.002, 2011.

- Healy, R. M., O'Connor, I. P., Hellebust, S., Allanic, A., Sodeau, J. R., and Wenger, J. C.: Characterisation of single particles from in-port ship emissions, Atmospheric Environment, 43, 6408-6414, 10.1016/j.atmosenv.2009.07.039, 2009.
 Johansson, L., Jalkanen, J.-P., and Kukkonen, J.: Global assessment of shipping emissions in 2015 on a high spatial and temporal resolution, Atmospheric Environment, 167, 403-415, 10.1016/j.atmosenv.2017.08.042, 2017.
- 20 Jonsson, A. M., Westerlund, J., and Hallquist, M.: Size-resolved particle emission factors for individual ships, Geophysical Research Letters, 38, 10.1029/2011gl047672, 2011.
- Lack, D. A., Corbett, J. J., Onasch, T., Lerner, B., Massoli, P., Quinn, P. K., Bates, T. S., Covert, D. S., Coffman, D., Sierau, B., Herndon, S., Allan, J., Baynard, T., Lovejoy, E., Ravishankara, A. R., and Williams, E.: Particulate emissions from commercial shipping: Chemical, physical, and optical properties, Journal of Geophysical Research-Atmospheres, 114, 10.1029/2008jd011300, 2009.
 - Li, L., Huang, Z., Dong, J., Li, M., Gao, W., Nian, H., Fu, Z., Zhang, G., Bi, X., Cheng, P., and Zhou, Z.: Real time bipolar time-of-flight mass spectrometer for analyzing single aerosol particles, International Journal of Mass Spectrometry, 303, 118-124, https://doi.org/10.1016/j.ijms.2011.01.017, 2011.

Liu, Z., Lu, X., Feng, J., Fan, Q., Zhang, Y., and Yang, X.: Influence of Ship Emissions on Urban Air Quality: A 30 Comprehensive Study Using Highly Time-Resolved Online Measurements and Numerical Simulation in Shanghai,

Environmental Science & Technology, 51, 202-211, 10.1021/acs.est.6b03834, 2017.
Merico, E., Donateo, A., Gambaro, A., Cesari, D., Gregoris, E., Barbaro, E., Dinoi, A., Giovanelli, G., Masieri, S., and Contini, D.: Influence of in-port ships emissions to gaseous atmospheric pollutants and to particulate matter of different sizes in a Mediterranean harbour in Italy, Atmospheric Environment, 139, 1-10, 10.1016/j.atmosenv.2016.05.024, 2016.

Moldanova, J., Fridell, E., Winnes, H., Holmin-Fridell, S., Boman, J., Jedynska, A., Tishkova, V., Demirdjian, B., Joulie, S., Bladt, H., Ivleva, N. P., and Niessner, R.: Physical and chemical characterisation of PM emissions from two ships operating in European Emission Control Areas, Atmospheric Measurement Techniques, *6*, 3577-3596, 10.5194/amt-6-3577-2013, 2013.

5 Murphy, S. M., Agrawal, H., Sorooshian, A., Padró, L. T., Gates, H., Hersey, S., Welch, W. A., Lung, H., and Miller, J. W.: Comprehensive simultaneous shipboard and airborne characterization of exhaust from a modern container ship at sea, Environmental Science & Technology, 43, 4626-4640, 2009.

Petzold, A., Hasselbach, J., Lauer, P., Baumann, R., Franke, K., Gurk, C., Schlager, H., and Weingartner, E.: Experimental studies on particle emissions from cruising ship, their characteristic properties, transformation and atmospheric lifetime in
the marine boundary layer, Atmospheric Chemistry and Physics, 8, 2387-2403, 10.5194/acp-8-2387-2008, 2008.

Spencer, M. T., Shields, L. G., Sodeman, D. A., Toner, S. M., and Prather, K. A.: Comparison of oil and fuel particle chemical signatures with particle emissions from heavy and light duty vehicles, Atmospheric Environment, 40, 5224-5235, 10.1016/j.atmosenv.2006.04.011, 2006.

Toner, S. M., Sodeman, D. A., and Prather, K. A.: Single particle characterization of ultrafine and accumulation mode

- particles from heavy duty diesel vehicles using aerosol time-of-flight mass spectrometry, Environmental Science & Technology, 40, 3912-3921, 10.1021/es051455x, 2006.
 Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, Atmospheric Chemistry and Physics, 9, 2891-2918,
- 20 Viana, M., Amato, F., Alastuey, A., Querol, X., Moreno, T., Garcia Dos Santos, S., Dolores Herce, M., and Fernandez-Patier, R.: Chemical Tracers of Particulate Emissions from Commercial Shipping, Environmental Science & Technology, 43, 7472-7477, 10.1021/es901558t, 2009.

Wenzel, R. J., Liu, D. Y., Edgerton, E. S., and Prather, K. A.: Aerosol time-of-flight mass spectrometry during the Atlanta Supersite Experiment: 2. Scaling procedures, Journal of Geophysical Research-Atmospheres, 108, 8427

25 10.1029/2001jd001563, 2003.

2009.

Xiao, Q., Li, M., Liu, H., Deng, F., Fu, M., Man, H., Jin, X., Liu, S., Lv, Z., and He, K.: Characteristics of marine shipping emissions at berth: profiles for PM and VOCs, Atmos. Chem. Phys. Discuss., 2018, 1-29, 10.5194/acp-2017-1132, 2018.
Yang, D.-q., Kwan, S. H., Lu, T., Fu, Q.-y., Cheng, J.-m., Streets, D. G., Wu, Y.-m., and Li, J.-j.: An Emission Inventory of Marine Vessels in Shanghai in 2003, Environmental Science & Technology, 41, 5183-5190, 10.1021/es061979c, 2007.

30 Zhang, Y., Yang, X., Brown, R., Yang, L., Morawska, L., Ristovski, Z., Fu, Q., and Huang, C.: Shipping emissions and their impacts on air quality in China, Science of the Total Environment, 581, 186-198, 10.1016/j.scitotenv.2016.12.098, 2017. Zhao, M., Zhang, Y., Ma, W., Fu, Q., Yang, X., Li, C., Zhou, B., Yu, Q., and Chen, L.: Characteristics and ship traffic source identification of air pollutants in China's largest port, Atmospheric Environment, 64, 277-286, 10.1016/j.atmosenv.2012.10.007, 2013.



Figure 1: Map of sampling site in Shanghai port and surrounding areas. Port region is indicated by shaded area. The insets are the satellite image of the port site and a photo taken on the roof of monitoring station seeing in port direction.



Figure 2: Temporal concentration of pollutants SO₂, NO, NO₂, O₃ and PM2.5 during 27-29 August 2016. Contemporary wind direction and speed, SO₂ concentration of Shanghai city and Vanadium particles number concentration as detected by SPAMS are included as a reference.



5

Figure 3: Mean mass spectra of fresh and background ship emission particles in port (a, c) and the wind rose of particle number concentration (in measure of particle number per hour) of these two particle types (b, d). Peaks in mass range of 70-150 in (a) and (c) are magnified by 10 times.



Figure 4: Particle number size distribution of fresh and background ship emission particles by SPAMS (Upper). Size distribution of these fresh and background types of ship emission particles normalized by total particles at each size (Lower).



5 Figure 5: Mean mass spectra of four major particle types from fresh ship emission.



Figure 6: Temporal trend of number concentrations of four fresh Vanadium particle types (Left panel); the number (upper right) and number fraction (lower right) of four Vanadium particle types as a function of particle size.



Figure 7: Pollution roses of SO₂, NO, NO₂, O₃, PM_{2.5} and Vanadium particles during the whole study period. Vanadium particles wind rose is based on number concentration as measured by SPAMS.

Tables.

Table 1: Statistics of pollutants concentration level during the whole sampling period. Numbers are average concentration followed by 25th and 75th quantiles in brackets. Average pollution levels in Shanghai city during the same period are included as a comparison.

5

10

	In plume		Non-plume		Non-plume (port sector)		Port average		Shanghai average	
$SO_2(ug/m^3)$	28.3	(17.6~31.8)	9.9	(8.1~11.6)	10.2	(8.2~12.1)	15.6	(8.7~16.8)	10.8	(9~12)
NO(<i>u</i> g/m3)	42.5	(7.6~47.5)	41.6	(7.1~59.1)	16.5	(1.8~18.1)	41.9	(7.3~55.3)	5.8	(3~6)
NO ₂ (<i>u</i> g/m3)	59.3	(36.1~72.4)	50.5	(27.8~60.8)	36.9	(22.1~46.1)	53.2	(30.3~65.0)	30.2	(18~38)
O₃(<i>u</i> g/m3)	53.1	(19.3~77.8)	54.6	(15.4~84.7)	71.3	(45.4~97.6)	54.1	(16.9~82.7)	81.1	(40~107)
PM _{2.5} (<i>u</i> g/m3)	30.2	(14.8~39.6)	25.1	(12.8~32.5)	19.6	(11.6~23.2)	26.7	(13.2~34.1)	31.4	(16~43)
Vanadium partilces (#/hour)	47.6	(31~55)	10.9	(5~17)	12.3	(7~19)	22.8	(7~29)		

Table 2: Contributions of ship emissions to ambient pollutants SO₂, NO, NO₂, O₃, PM_{2.5} and Vanadium particles in port area. Calculations are based on two situations: entire sampling period (all wind directions included) and only when site is in downwind direction of port emissions. Total lengths (in hours) of respective periods are given.

	SO ₂	NO	NO ₂	O ₃	PM _{2.5}	Vanadium particles*
Entire period (include land emissions):	36.4%	0.7%	5.1%	-0.9%	5.9%	49.5%
In port sector (exclude landemissions):	57.2%	71.9%	30.4%	-16.6%	27.6%	77.0%

Length of sampling (in hours): Entire period: 2256; Port sector: 1136; In plume: 694; Non-plume: 1563; Non-plume (port sector): 625.

* Number contribution