We are thankful to the three referees for their thoughtful and constructive comments which help improve the manuscript substantially. Following the reviewers' suggestions, we have revised the manuscript accordingly. Listed below are our point-by-point responses in blue to each comment that is repeated in italic.

Response to Reviewer #1

BrC was determined based on the optical property of aerosols, i.e. the difference between the total absorption and the absorption by BC at 370 nm. Then the absorption by BrC was apportioned to the OA factors from PMF analysis of AMS data. However, this paper did not discuss the compositions of the BrC. Since HR-AMS can measure the molecular fragments, the authors should also discuss the evidence of BrC from the molecular composition.

We thank the reviewer's comments. Molecular characterization of BrC is very important. Unfortunately, HR-AMS uses 70 eV electron impact for ionization, and the molecular information of organic compounds is difficult to retain. Therefore, we compared BrC with OA factors to investigate the potential sources of BrC in this study, and found that coal combustion and biomass burning were two major sources. Although we didn't have molecular information of BrC, previous studies have shown that nitrophenols and aromatic carbonyls are often the major BrC species (Desyaterik et al., 2013;Laskin et al., 2015).

Specific comments:

 The authors measured 50 vertical profiles of b_{sca} and b_{abs}, but just presented 10 profiles and discuss several ones. Did the other profiles have the same evolution with the discussed ones? The authors should present all the aerosol optical property profiles and the meteorological parameters, at least in the Supplementary materials.

We thank the reviewer's suggestions. In the revised manuscript, we added the rest profiles in Figure S5 in supplementary materials. As shown in Figure S5, most vertical profiles are overall similar to those discussed in the text.



Figure S5. Evolution of vertical profiles of b_{sca} , b_{abs} and SSA during 25-26 November (top) and 30 November – 2 December (bottom). Also shown are (a-d) meteorological variable of *T*, RH, WD, and WS, and (e) time series of b_{ext} at ground and MLH. M16-M25 refer to the number of vertical profiles.

Line 4 in page2: The abbreviation of non-refractory BC as rBC will cause confusion. Rewrite this sentence.

Revised.

Response to Reviewer #2

This work did a very nice job in measuring the aerosol optical properties at both ground and 260 m in winter Beijing. The quality of the measurement data were ensured by inter-comparisons across different instruments. The authors found substantial vertical differences in aerosol optical properties (e.g., SSA) at nighttime due to strong local emissions. They also clarified that the lensing effect played an important role in enhancing BC absorption in Beijing. Source apportionment of BrC absorption at 370 nm were conducted using two statistical tools, both suggesting that fossil fuel combustion is the dominant BrC contributor in urban Beijing. This work is very well designed and written, I would recommend this work to be published on Atmos. Chem. Phys. with only one comment to be addressed

We thank the reviewer's positive comments.

The comment is about the method for $b_{abs,370nm,BrC}$ estimation with Eqs (3) and (4). In Eq (3), the BC absorption at 370nm is estimated using a fitted power law, and BrC at 370 nm is then derived by subtracting the BC absorption from the total measured absorption. However, this work has demonstrated that the coating of organic aerosol on BC (or "lensing effect") can contribute to the light absorption enhancement of BC substantially (20~40%). In Eqs (3) and (4), such coating effect was not considered, which might lead to an overestimation of $b_{abs,370nm,BrC}$. So please provide an estimation on the uncertainty associated with $b_{abs,370nm,BrC}$ estimation due to OA coatings, or more discussions on the uncertainties.

This is a very good point. Previous studies by Liu et al. (2017) found that the absorption enhancement was small at M_R < 3. Assuming no absorption enhancement at M_R = 3 in this study, the increase in mass absorption cross-section (MAC) of BC due to coating materials was approximately 10 – 20%, and can be up to ~30% when M_R was above 6.5 (Fig. 9). Therefore, the upper limit of absorption enhancement of BC due to "lensing effect" in this study was ~30%, and mostly likely between 10% and 20%. With this, the *babs,370nm,BrC* could be overestimated by ~10 – 20%, but should be less than 30%.

Following the reviewer's comments, we added the uncertainties for estimation of BrC in the revised manuscript.

"It should be noted that we might overestimate $b_{abs,370nm,BrC}$ by approximately 10 – 20% if the contribution of "lensing effect" on BC absorption at M_R = 3 was negligible (Liu et al., 2017)."

Response to J. C. Corbin

This is a very brief comment to recommend that the authors use the terminology established in Petzold et al. (2013) when referring to aethalometer measurements. That is, use eBC rather than just BC.Additionally, I would suggest writing MAC(637nm) rather than simply MAC on the figures, as was done in the abstract.

Best regards,

Joel Corbin.

Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S.-M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, X.-Y.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365-8379, https://doi.org/10.5194/acp-13-8365-2013, 2013.

We thank Dr. Corbin's good suggestions. We changed "BC" to "eBC" in the revised manuscript, and included the wavelength for MAC.

References:

- Desyaterik, Y., Sun, Y., Shen, X., Lee, T., Wang, X., Wang, T., and Jeffrey Collett, J.: Speciation of "brown" carbon in cloud water impacted by agricultural biomass burning in Eastern China, J. Geophys. Res., 118, 7389–7399, doi:10.1002/jgrd.50561, 2013.
- Lack, D. A., and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, Atmospheric Chemistry and Physics, 10, 4207-4220, 10.5194/acp-10-4207-2010, 2010.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, Chem. Rev., 10.1021/cr5006167, 2015.
- Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S., Williams, P. I., Ting, Y.-C., Haslett, S., Taylor, J. W., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe, H., and Allan, J. D.: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, Nature Geosci, advance online publication, 10.1038/ngeo2901, 2017.

Vertical characterization of aerosol optical properties and brown

carbon in winter in urban Beijing, China

Conghui Xie^{1,2}, Weiqi Xu^{1,2}, Junfeng Wang⁴, Qingqing Wang¹, Dantong Liu⁵, Guiqian Tang¹, Ping Chen⁶, Wei Du^{1,2}, Jian Zhao^{1,2}, Yingjie Zhang¹, Wei Zhou^{1,2}, Tingting Han¹, Qingyun Bian^{2,7}, Jie Li¹, Pingqing Fu^{1,2}, Zifa Wang^{1,2,3}, Xinlei Ge⁴, James Allan^{5,8}, Hugh Coe⁵, Yele Sun^{1,2,3}

 ¹State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China
 ²College of Earth Sciences, University of Chinese Academy of Sciences, Beijing 100049, China
 ³Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China
 ⁴School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing 210044, China
 ⁵Centre for Atmospheric Science, School of Earth, Atmospheric and Environmental Science, University of Manchester, Manchester M13 9PL, UK
 ⁶Handix Scientific LLC, Boulder, CO 80301, USA
 ⁷CAS Key Laboratory of Regional Climate-Environment Research for Temperate East Asia, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

⁸National Centre for Atmospheric Science, The University of Manchester, Manchester, UK

20 Correspondence: Yele Sun (<u>sunyele@mail.iap.ac.cn</u>)

5

10

15

Abstract. Aerosol particles are of importance in the Earth's radiation budget since they scatter and absorb sunlight. While extensive studies of aerosol optical properties have been conducted at ground sites, vertical measurements and characterization are very limited in megacities. In this work, we present simultaneous real-time online measurements of aerosol optical properties at ground level

and at 260 m on a meteorological tower from 16 November to 13 December in 2016 in Beijing along with measurements of continuous vertical profiles during two haze episodes. The average $(\pm 1\sigma)$ scattering and absorption coefficients (b_{sca} and b_{abs} , $\lambda = 630$ nm) were 337.6 (± 356.0) and 36.6 (± 33.9) Mm⁻¹ at 260 m, which were 26.5% and 22.5% lower than those at ground level. Single scattering albedo (SSA), however, was comparable between the two heights with slightly higher values at ground level (0.89 ± 0.04). Although b_{sca} and b_{abs} showed overall similar temporal variations between ground and 260 m, the ratios of 260 m to ground varied substantially from less than 0.4 during the cleanest stages of haze episodes to > 0.8 in the late afternoon. A more detailed analysis indicates that vertical profiles of b_{sca} , b_{abs} , and SSA in the low atmosphere were closely related to the changes in meteorological conditions and mixing layer height. The mass absorption cross-section [MAC] of BC (equivalent black carbon (eBC, $\lambda = 630$ nm) varied substantially from 9.5 to 13.2 m² g⁻¹ in winter in Beijing, and it was strongly associated with the mass ratio of non-coating materials on refractory BC (*r*BC)-materials to *r*BC (M_R), and also the oxidation degree of organics in *r*BC-containing particles. Our results show that the increases in MAC of BC<u>eBC</u> in winter were mainly caused by photochemically produced secondary materials. Light absorption of organic carbon (brown carbon, BrC) was also important in winter, which on average accounted for 46 (±8.5) % and 48 (±9.3) % of the total absorption at 370 nm at ground level and 260 m, respectively. A Linear regression model combined with positive matrix factorization analysis was used to show that coal combustion was the dominant source contribution of BrC (48-55%) followed by biomass burning (17%) and photochemically processed secondary organic aerosol (~20%) in winter in Beijing.

1 Introduction

Light scattering and absorption of aerosols reduce visibility and affect the radiation and energy budget of the Earth (Rosenfeld, 1999;Kim and Ramanathan, 2008). Scattering aerosols cool atmosphere and exert a negative forcing while light absorbing materials
warm the atmosphere and, if they exist over a brighter underlying surface, contribute a positive forcing (Haywood and Boucher, 2000). While scattering particles mainly include ammonium sulfate, ammonium nitrate, and a majority of organics (Han et al., 2015), black carbon (BC) and brown carbon (BrC) are the two major light absorbing matters in fine particles (Bond et al., 2013). Although numerous studies have been conducted to investigate aerosol optical properties, accurate quantification of aerosol radiative forcing still remains challenging (Stocker et al., 2013). While the very complex optical properties affected by mixing states are one of the reasons (Peng et al., 2016;Cappa et al., 2012;Liu et al., 2017), our limited understanding of the vertical distributions of optical properties and their relationship to composition and mixing state is also important. For example, a recent study found that BC near the capping inversion is more effective in suppressing planetary boundary layer height and weakening turbulent mixing (Wang et the capping inversion is more effective in suppressing planetary boundary layer height and weakening turbulent mixing (Wang et the capping inversion is more effective in suppressing planetary boundary layer height and weakening turbulent mixing (Wang et the capping inversion is more effective in suppressing planetary boundary layer height and weakening turbulent mixing (Wang et the capping inversion is more effective in suppressing planetary boundary layer height and weakening turbulent mixing (Wang et the capping inversion is more effective in suppressing planetary boundary layer height and weakening turbulent mixing (Wang et the capping inversion is more effective in suppressing planetary boundary layer height

al., 2018; Ding et al., 2016).

5

10

Most previous measurements of aerosol particles in urban Beijing are conducted at ground level (Sun et al., 2013;Sun et al., 2014;Tao et al., 2015), which are subject to the influences of local emissions, such as biomass burning, cooking and traffic (Sun et al., 2014). Until recently, Sun et al. (2015) reported the first simultaneous measurements of PM₁ composition at 260 m and ground level in urban Beijing in winter 2013. Higher nitrate and much lower primary organic aerosol (POA) at 260 m than ground level was observed. This result is consistent with the subsequent observations in autumn 2015 (Zhao et al., 2017). The size distributions of aerosol at 260 m and ground level are also different especially for Aitken mode particles (Du et al., 2017). Such differences in concentration, composition and size distributions between 260 m and ground level may exert a potential influence on aerosol optical properties. However, the vertical distributions of aerosol optical properties are rarely characterized. Wang et al. (2017b) conducted the first continuous vertical measurements of particle extinction and BC from ground to 200-260 m during two severe haze episodes in winter. The results showed four distinct types of vertical profiles that were closely linked with boundary layer dynamics. However, this study is limited within a relatively short time, and the real-time measurements of aerosol optical properties at different heights in Beijing is yet to be performed.

Black carbon, as the major light absorbing matter in atmospheric aerosol (Bond et al., 2013), absorbs visible light strongly with an
absorption Angstrom exponent (AAE) of ~1 (Horvath, 1997). The light absorbing ability of BC is affected by the coating materials which may change substantially during the aging process. The enhancement of BC absorption due to non-BC materials in BC-containing particles is often referred to as lensing effect (Bond et al., 2006). Previous studies found that internally mixed particulate organic matter with BC could enhance absorption by up to 70% (Lack et al., 2012), and the absorption enhancement strongly depends on the BC coating amount and particle mixing state (Liu et al., 2017;Liu et al., 2015b). Although Peng et al. (2016) found that the
BC light absorption enhancement can be significantly increased in polluted urban environment by using a novel environmental chamber approach, the evaluation of such effects through field measurements are still limited in Beijing. In addition to BC, light-absorbing organic carbon (brown carbon, BrC) plays an important role in affecting radiative forcing at ultraviolet wavelength (Laskin et al., 2015) although its mass absorption cross-section (MAC, also known as mass absorption efficiency) is an order of

magnitude less than that of BC in the visible wavelength (Yang et al., 2009). The AAE of BrC is ubiquitously larger than 1, yet accurate quantification of absorption of BrC is challenging (Laskin et al., 2015), particularly in a megacity with complex source emissions and processes. For example, recent studies have found that coal combustion is also an important source of BrC in winter in Beijing in addition to biomass burning aerosol (Sun et al., 2017;Yan et al., 2017).

In this work, we conducted comprehensive measurements of aerosol optical properties including light extinction coefficients (b_{ext}), scattering coefficients (b_{sca}), and absorption coefficients (b_{abs}) at both ground level and 260 m on a meteorological tower in winter in 2016 along with two continuous vertical measurements of b_{sca} and b_{abs} between ground level and 260 m. The measurements of aerosol optical properties from different instruments are compared, and the vertical differences in b_{sca}, b_{abs}, and single scattering albedo (SSA) are characterized. Also, the evolution of vertical differences in aerosol optical properties and its relationship with meteorological conditions and aerosol composition are illustrated. In addition, the light absorption capability of BC is estimated, and the contribution of BrC to b_{abs} at 370 nm are quantified.

2. Experimental methods

15

20

2.1 Sampling site and measurements

All measurements at ground level and 260 m were conducted at the tower site of Institute of Atmospheric Physics (IAP), Chinese Academy of Sciences (39°58′28″N, 116°22′16″E) in Beijing from 16 November to 13 December 2016 as a part of APHH Beijing winter campaign. A detailed description of the sampling site is given in Zhou et al. (2018).

Table 1 lists a summary of measurements in this study. At the ground site, b_{ext} and b_{sca} (λ =630 nm) of dry particles (PM₁) were measured by a cavity attenuated phase shift single scattering albedo monitor (CAPS PM_{ssa}, Aerodyne Research Inc.) (Onasch et al., 2015) that was installed in a room on the rooftop of a two-storystorey building (~ 8 m). Note that the major uncertainty in scattering measurements caused by truncation effect is less than 2% (Han et al., 2017). The detailed description of the CAPS PM_{ssa} is given in Han et al. (2017). In the same room, a seven-wavelength (370, 470, 520, 590, 660, 880, and 950 nm) Aethalometer (AE33, Magee

Scientific Corp.) was used to measure BC at a time resolution of 1 min. This version of the Aethalometer uses a compensation algorithm based on "dual-spot" measurements to automatically correct the filter-based loading effects (Drinovec et al., 2015). According to Petzold et al. (2013), BC measured by AE33 was defined as equivalent BC (eBC) unless otherwise stated. In addition, a photoacoustic extinctioneter (PAX, Droplet Measurement Technologies) was used to measure b_{sca} and b_{abs} (λ =870 nm) of dry $PM_{2.5}$ in a container at ground level (~3 m). Other collocated measurements at the same ground site include non-refractory submicron (NR-PM₁) aerosol species (organics, sulfate, nitrate, ammonium, and chloride) by an Aerodyne high-resolution time-of-flight aerosol mass spectrometer (HR-AMS hereafter) and refractory black carbon (rBC) and BC-containing species by a soot particle aerosol mass spectrometer (SP-AMS)(Wang et al., 2017a;Onasch et al., 2012). In this study, the tungsten vaporizer of the SP-AMS was removed, thus only rBC-containing particles were measured. Positive matrix factorization (PMF) was performed onto the highresolution mass spectra of organic aerosol (OA) of the HR-AMS and the SP-AMS. Six OA factors were identified from HR-AMS 10 measurements including fossil-fuel-related OA (FFOA) dominantly from coal combustion, cooking OA (COA), biomass burning OA (BBOA), oxidized POA (OPOA), oxygenated OA (OOA), and aqueous-phase OOA (aq-OOA) (Xu et al., 2018). In comparison, four OA factors were identified from SP-AMS measurements including FFOA, BBOA, OOA1 and OOA2. More detailed descriptions of the operations and calibrations of the HR-AMS and SP-AMS, and subsequent PMF analysis can be found in Xu et al. (2015) and Wang et al. (2016).

5

15

20

At 260 m on the Beijing 325 m Meteorological Tower (BMT), b_{ext} (λ =630 nm) of PM_{2.5} was measured by a cavity attenuated phase shift extinction monitor (CAPS PMext Aerodyne Research Inc.), BCeBC was measured by AE33, and NR-PM1 aerosol species were measured by an Aerodyne aerosol chemical speciation monitor (ACSM) (Chen et al., 2015).

Continuous vertical measurements from ground to 240 m (nighttime)/260 m (daytime) were also conducted during 25-26 November and 30 November - 4 December, 2016. The AE33 and PAX were installed in a small container that was able to travel up and down the BMT. In total, 50 vertical profiles of BCeBC, b_{sca} and b_{abs} were obtained (Table S1). The meteorological variables of temperature (T), relative humidity (RH), wind speed (WS) and direction (WD) were measured at 15 heights (8, 15, 32, 47, 65, 80, 100, 120, 140, 160, 180, 200, 240, 280 and 320 m) on the BMT. In addition, mixing layer height (MLH) was retrieved from vertical attenuated backscatter coefficients measured by a single-lens ceilometer (CL51, Vaisala, Finland) (Tang et al., 2015;Tang et al., 2016). All data in this study are reported in Beijing local time (UTC + 8 hr).

2.2 Inter-comparisons

Aerosol optical properties and BC measurements from different instruments were evaluated through parallel sampling either before 5 or after the campaign. As shown in Fig. 1b, the one-week inter-comparison of the two AE33s shows an excellent agreement in **BCeBC** measurements (slope = 1.06, $R^2 = 0.94$). To be consistent, the **BCeBC** measured by AE33₂ was scaled by a factor of 1.06 according to that of AE33₁. b_{ext} measured by CAPS PM_{ext} was also highly correlated with that measured by CAPS PM_{ssa} (R^2 =0.99, slope =1.22) (Fig. 1a). Considering that the CAPS PM_{ssa} was not calibrated before this campaign, b_{ext} measured by the CAPS PM_{ssa} was multiplied by a factor of 1.22 in this study. We also compared the light extinction of PM_1 and $PM_{2.5}$ during the period of intercomparison. As indicated in Fig. 1c, the b_{ext} of PM_{2.5} is nearly twice that of PM₁ ($R^2 = 0.93$, slope =1.93), indicating that aerosol 10 particles between $1 - 2.5 \,\mu\text{m}$ are also important for light extinction in this study. In addition, b_{sca} measured by CAPS PM_{ssa} (630 nm) and PAX (870 nm) is highly correlated ($R^2=0.98$), and the slope of 2.2 suggests a scattering angstrom efficiency (SAE) of approximately 2.4 (Fig. 1d). Figure 2a shows a comparison of BCeBC measured by AE33 and SP-AMS for the entire study. It is clear that both independent measurements are highly correlated (R^2 =0.98). In our previous study, b_{abs} of BCeBC at 630 nm was derived using a mass absorption cross-section (MAC) of 7.4 m²g⁻¹, i.e., $b_{abs} = \frac{BCeBC}{X} \times 7.4$ (Han et al., 2017). The <u>BCeBC</u> derived 15 b_{abs} is highly correlated with that from the CAPS PM_{ssa} measurements ($R^2 = 0.89$, slope = 1.01), i.e., $b_{abs} = b_{ext} - b_{sca}$. All these results suggest that b_{abs} from different measurements agree reasonably well in this study. We then define the mass absorption crosssection (MAC) of BC at 630 nm as.

$$MAC = b_{abs} / rBC \tag{1}$$

20 Where b_{abs} is the absorption coefficient at 630 nm, and *r*BC is the refractory BC from SP-AMS measurements.

2.3 Calculations of single scattering albedo and brown carbon

Although single scattering albedo (SSA, $\lambda = 630$ nm) of PM₁ can be directly calculated from the CAPS PM_{ssa} measurements as SSA = b_{sca}/b_{ext} , the SSA of PM_{2.5} at ground level and 260 m are both derived as:

$$SSA = \frac{b_{\text{ext}} - b_{\text{abs}}}{b_{\text{ext}}} = \frac{b_{\text{ext}} - 7.4 \times BC}{b_{\text{ext}}}$$
(2)

5 Note that all SSA values discussed below refer to PM_{2.5} at 630 nm unless otherwise stated.

We also estimate the absorption of BrC assuming that BC is the only absorber at 950 nm with an AAE of 1 (Fig. S1). The absorption coefficient of BC at 370 nm is derived using a fitted power law relationship at seven wavelengths (Ran et al., 2016) (Eq. 3). BrC at 370 nm is then derived by subtracting the BC absorption from the total measured absorption (Eq. 4).

$$b_{\rm abs,370nm,BC} = b_{\rm abs,950nm} \left(\frac{370}{950}\right)^{-AAE}$$
 (3)

15

$$b_{\text{abs},370\text{nm},\text{BrC}} = b_{\text{abs},370\text{nm},\text{Total}} - b_{\text{abs},370\text{nm},\text{BC}}$$
(4)

Where $b_{abs,370nm,BC}$ is the BC absorption at 370 nm, and AAE is 1. It should be noted that we might overestimate $b_{abs,370nm,BrC}$ by approximately 10 – 20% if the contribution of "lensing effect" on BC absorption at $M_R = 3$ was negligible (Liu et al., 2017). Although the absorption of dust is much less than those of BC and BrC because of much lower MAC (Yang et al., 2009), the data with potential influences of dust as indicated by high Ca content (Fig. S2) were excluded in calculation of the absorption of BrC at 370 nm.

3. Results and discussion

3.1 General description

5

20

The average $b_{ext}(\pm 1\sigma)$ were 547.0 ± 555.9 and 387.8 ± 395.2 Mm⁻¹ at ground level and 260 m, respectively. Compared with previous measurements in winter at the same site (Han et al., 2017;Wang et al., 2017b), b_{ext} is much higher in this study due to more frequent occurrence of severe haze episodes, e.g., six haze episodes in Fig. 3. The<u>These</u> results suggest that the PM pollution in winter in Beijing was still severe although air quality has been continuously improved in recent years. We also note that the average b_{ext} at 260 m was on average 29.1% lower than that at ground level, suggesting a considerable vertical gradient in b_{ext} in winter.

As shown in Fig. 3, b_{sca} and b_{abs} varied similarly to b_{ext} , and the average b_{sca} and b_{abs} at 260 m were 337.6 (±356.0) and 36.6 (±33.9) Mm⁻¹, respectively, which were 26.5% and 22.5% lower than those at ground level (459.5 ±483.1 and 47.2 ±2.4 Mm⁻¹, respectively). Since the SSA is the ratio of b_{sca} and b_{ext} , the variations in SSA at ground level were similar to those at 260 m for most of the time, although several periods with much lower SSA at ground level were observed due to large emissions of BC at midnight, e.g., 17 November and 1 December. The average (±1 σ) SSA was 0.89 (±0.04) and 0.88 (±0.04) at ground level and 260 m, respectively, which was overall consistent with those (0.84 – 0.91) in previous studies (Wang et al., 2017b;Han et al., 2017;Han et al., 2015). Figure 3 also shows that high SSA values were typically associated with haze episodes while low values typically occur during clean periods. These results suggest an increasing role of scattering aerosols, e.g., secondary inorganic aerosol species, during haze episodes.

3.2 Vertical differences of aerosol optical properties.

All aerosol optical properties including b_{ext} , b_{sca} , and b_{abs} showed overall similar variations between ground level and 260 m with R^2 between ground and 260 m varying from 0.70 to 0.81. However, large vertical differences were also observed during specific periods, for example, the cleaning stages of haze episodes (26 November, 4 and 7 December), and several midnights with largely enhanced

BC emissions (17 and 29 November). As indicated in Figs. 4a and 4b, the periods with low values of $ratio_{260m/ground}$ were characterized by correspondingly low MLH (typically less than 300 m) and much higher RH at ground level than 260 m. Figure 5 further shows that the vertical $ratios_{260m/ground}$ of b_{sca} and b_{abs} varied substantially throughout the study with the values below 1 for most of the time (89% and 84%, respectively). These results suggest that the vertical differences of aerosol optical properties were complex in winter in Beijing, but overall higher heights showed lower values. We also observed different frequencies of the ratio_{260m/ground} between b_{sca} and b_{abs} (Fig. 5). While the ratio_{260m/ground} of b_{sca} varied mainly between 0.6 and 0.8, that of b_{abs} was between 0.8 and 1.0, highlighting the different contributions of scattering and absorbing aerosols at different heights. Although b_{abs} at 260 m was 21% lower than that at ground level (37 vs. 47 Mm⁻¹), the relative contribution of absorbing aerosols was relatively higher. One reason is that a large fraction of BC in Beijing is from regional transport (Sun et al., 2016), and the ratio of BC to scattering aerosol species in Beijing surrounding regions is high, likely due to the strict control of heavy-duty/diesel trucks and also much reduced coal combustion emissions in the city of Beijing.

5

10

Figure 6 shows the diurnal cycles of aerosol optical properties at 260 m and ground level after excluding the periods with cleaning processes, while those of entire study are presented in Fig. S3. The diurnal profiles of b_{sca} and b_{abs} were overall similar between ground level and 260 m, which were both characterized by higher values at night and lower values during daytime. Such diurnal patterns were opposite to those of MLH which showed gradual increases from 300 - 400 m at nighttime to ~600 - 700 m in the late afternoon (Fig. 5a). These results indicate that the variation of PBL height plays an important role in driving the diurnal variations of aerosol optical properties. This is further supported by the diurnal variations in ratios_{260m/ground}, and correlations between 260 m and ground (*R*²_{260m vs. ground}). As shown in Fig. 6, the ratio_{260m/ground} of *b*_{sca} increased from ~0.70 at nighttime to ~0.85 during daytime, while that of *b*_{abs} increased from 0.71 to 0.96 as a result of rising PBL. *R*² of *b*_{sca} and *b*_{abs} also increased up to > 0.90, suggesting that scattering and absorbing aerosols were relatively well mixed vertically. Because of the influences of enhanced local primary emissions and the reduced vertical mixing, both ratios and *R*² of *b*_{abs} between 260 m and ground level decreased substantially at nighttime. It should be noted that the ratio_{260m/ground} of *b*_{sca} did not reach 1 in the late afternoon even when the MLH was above 300 m. These results suggest that other factors also contributed the vertical differences of *b*_{sca} in addition to MLH. For example, *b*_{sca} is

mainly contributed by secondary aerosols (e.g., ammonium sulfate, ammonium nitrate, and SOA) (Han et al., 2015;Wang et al., 2015), while the secondary production at different heights could be different due to the different precursor concentrations and oxidants and also different phase partitioning since T falls and RH increases with height. Secondary aerosols are typically formed over a regional scale, which also supports the good correlations of b_{sca} between ground level and 260 m at nighttime.

SSA at ground level showed a clear diurnal trend with the values increasing from 0.85 at 8:00 to 0.89 at 12:00, and then remained at relatively high levels in the late afternoon. Such a diurnal profile is most likely related to photochemical production of secondary aerosol in the daytime, which is supported by the gradual increase in SSA as a function of photochemical age (-log(NO_x/NO_y, Fig. S4). Note that the diurnal variation of SSA is relatively smaller than those in autumn and summer at the same site (Han et al., 2015;Han et al., 2017). A possible explanation is that the enhancement of light scattering caused by photochemical production is weaker in winter. The diurnal profile of SSA at 260 m was relatively similar to that at ground level, but the variations were much smaller. We also noticed that SSA at ground level was lower than that at 260 m at nighttime due to enhanced light absorption of BC from local emissions. Indeed, such diurnal differences are closely related to composition differences between ground level and 260 m (Zhao et al., 2017).

3.3 Evolution of vertical profiles of aerosol optical properties.

15 50 vertical profiles of b_{sca} and b_{abs} from ground to 260 m (day) or 240 m (night) were obtained in this study (Table S1 and Fig. S5). As shown in Fig. 5, the vertical ratios of b_{sca} and b_{abs} between ground and 240/260 m agreed well with those determined from simultaneous real-time measurements at the two heights. Although the ratios_{260/ground} of b_{sca} and b_{abs} varied largely from ~0.1 to 1.6, those of SSA showed small vertical differences for most of the time, e.g., < 0.04. Figure 7 presents the evolution of vertical profiles during a severe haze episode on 2 – 4 December. Before the formation of haze episode, aerosol particles were relatively well mixed vertically, and the vertical differences of b_{sca} and b_{abs} (M16 and M17) were both less than 30%. This was consistent with the high MLH (~600 m) during this period of time. After M16 (16:00), b_{ext} increased rapidly from less than 100 Mm⁻¹ to > 300 Mm⁻¹ in four hours associated with a decrease of MLH from ~600 m to ~400 m and a change of air masses from the north to the south-west. We note that b_{sca} and b_{abs} increased simultaneously across different heights, suggesting that both regional transport and the decrease of MLH have played important roles in the formation of this haze episode. However, SSA showed a gradual increase from ~0.82 (M16) to ~0.90 (M19), consistent with the large increases in the contributions of scattering nitrate and SOA during this period (Xu et al., 2018).

5 The vertical profiles had significant changes at nighttime (M19 and M20) as the occurrence of T inversion. As shown in Fig. 7c, a strong T inversion by approximately 1 °C was observed between 150 – 320 m, and RH was decreased by ~5%. While b_{sca} and b_{abs} were relatively well vertically mixed below the T inversion, they had significant changes in both absolute values and relative contributions above the layer. In particular, SSA increased considerably from $\sim 0.81 - 0.82$ to $\sim 0.88 - 0.90$ above the T inversion, suggesting that scattering and absorbing aerosols differed significantly below and above T inversion. In fact, aerosol composition at 10 ground level during M19 and M20 was characterized by higher BC (10%) and organics (48 - 54%) than that at 260 m (8% and 44 -46%, respectively), and secondary nitrate was correspondingly low (16 - 17% vs. 23%). Because of the T inversion, the enhanced light absorbing aerosol from local emissions could not be well mixed to 260 m, leading to lower SSA at ground level. In addition, the higher mass fraction of scattering secondary inorganic aerosol (SIA) at 260 m could also increase the SSA independently (Fig. \$5\$6). Air pollution was the severest at the night of 3 December which was associated with a large increase in RH below 200 m, 15 mostly likely a light fog event, and consistently low MLH (≤ 200 m). As a result, large vertical differences were observed for both b_{sca} and b_{abs} . In particular, b_{sca} and b_{abs} start to decrease at ~120 m, and the ratios_{240m/ground} decrease rapidly from ~0.8 (M23) to <0.1 (M25). Such large vertical gradients were also due to the earlier cleaning of air pollutants at higher heights (Fig. 7d). Although the vertical changes in b_{sca} and b_{abs} were significant, those of SSA were small, which were due to the small changes in relative contributions of BC and non-BC aerosols at ground level and 260 m during this period of time (Xu et al., 2018). Our results suggest 20 that vertical profiles of aerosol optical properties can have significant changes during the formation and evolution of haze episodes depending on changes in meteorological conditions and source emissions.

3.3 MAC of BCeBC.

5

15

20

Figure 9a shows that the MAC of BCeBC increased substantially from 9.5 to 13.2 m² g⁻¹ as the increase of the mass ratio of nonrBC material to rBC in rBC-containing particles (M_R) from 3.2 to 6.7. Considering that the light absorption of BrC at 630 nm was relatively small, the increases in MAC could be mainly due to the lensing effect of BC-coated materials. The average M_R is 4.66 (±0.95) indicating that most rBC-containing particles contain a large amount of non-rBC constituents. According to SP-AMS measurements, the average composition of rBC-containing particles was dominated by organics (59%) and rBC (17%), of which 66% of organics are from primary emissions of coal combustion and biomass burning (Wang et al., 2018). This is also consistent with low O/C ratios (< 0.3) during this study (Fig. 10b). Although the rBC-coated materials were dominated by primary aerosol, the increases in MAC were more associated with secondary aerosol. As shown in Fig. 9b, the contributions of POA (BBOA+FFOA) 10 decreased from 67% to 16% as $M_{\rm R}$ increased from 3 to 7, in contrast, the contributions of secondary aerosol species (= SOA+SIA) increased from 29% to 80%. This was also supported by the correlations between MAC and $M_{\rm R}$ across different levels. As shown in Fig. 9a, the increases of MAC as a function of M_R were relatively small, typically less than 20% at low PM levels (< 100 μ g m⁻³) when the contribution of POA was much higher than that of secondary aerosol species (37% vs. 59%), while MAC increased significantly by nearly 40% during periods with high PM levels (> 150 μ g m⁻³) and higher contribution of secondary aerosol (Fig. <u>S6S7</u>). We also found that the increases in MAC and M_R were both associated with the corresponding increases in O/C ratios, indicating that photochemical processing plays an important role in changing the MAC and coating of BC in winter. Indeed, MAC showed a pronounced diurnal cycle with a clear daytime increase from ~ 11 to ~ 13 m² g⁻¹, (Fig. 10a), and such a diurnal pattern agreed well with that of O/C, an indicator of the extent of chemical aging of organic aerosol. Consistently, MAC was almost linearly correlated with O/C as indicated in Fig. 10b. As O/C increased from 0.1 to 0.25, MAC increased from ~10 to 14 m² g⁻¹. These results suggest that secondary production from photochemical processing can contribute to the light absorption enhancement of BC in daytime by approximately 20%, and up to 40% in this study.

3.4 Brown carbon.

5

10

As shown in Fig. S7SS, the average AAE derived from the wavelength-dependent absorption were 1.58 and 1.60 at the ground and at 260 m, respectively, indicating the importance of non-BC light absorbers. Indeed, the absorption of BrC ($b_{abs, BrC}$) on average account for 46% and 48% (Fig. S1) of the total absorption at 370 nm at ground and 260 m, supporting the importance of BrC absorption at ultraviolet wavelength. It should be noted that the extraction of $b_{abs, BrC}$ from total absorption may introduce uncertainty since the AAE of BC is also influenced by BC size and coatings (Liu et al., 2015a). It is interesting to note that the fraction of $b_{abs, BrC}$ related OA materials were more light absorbing than the total OA. In fact, the organic mass measured by the SP-AMS is dominated by coal combustion and biomass burning emissions (Wang et al., 2018), two important sources of BrC (Yan et al., 2015;Sun et al., 2017;Du et al., 2014). Further supporting evidence is that the mass absorption coefficient of BrC (MAC_{org. 370 nm}) showed a pronounced diurnal cycle with higher values at nighttime, consistent with the enhanced nighttime primary emissions in winter. Figure 11a also shows that the nighttime $b_{abs, BrC}$ at ground level was much higher than that at 260 m. One explanation is that a large amount of local primary emissions at nighttime is not well vertically transported to 260 m. The small late afternoon peak indicates that secondary aerosol from photochemical production can also contribute to the BrC absorption.

15 $b_{abs, BrC}$ correlated the best with BBOA and FFOA ($R^2 = 0.83$ and 0.81, respectively) and moderately correlated with photochemical SOA ($R^2 = 0.46$) at ground level. These results suggest that biomass burning, coal combustion, photochemical production are three major sources of BrC in winter. We then estimated the contributions of different OA sources to BrC using (1) a multiple linear regression model as indicated by Eq. (5) and PMF with OA factors and BrC absorption as input.

$$b_{\text{abs, BrC}} = m_1 \times [\text{FFOA}] + m_2 \times [\text{COA}] + m_3 \times [\text{BBOA}] + m_4 \times [\text{OOA}] + m_5 \times [\text{OPOA}] + m_6 \times [\text{aq-OOA}]$$
(5)

20 The results are very consistent between the two approaches which both show that coal combustion is the major source contribution of BrC at ground level, on average accounting for 48 – 55% (Fig. 12). The contributions of BBOA and OOA to BrC were comparable,

which were 17% and 19 - 20%, respectively. In comparison, cooking emissions and aqueous-phase processing tend to be minor contributions of BrC in winter in Beijing. Our results are consistent with a recent study which highlighted the importance of fossil source contribution to BrC in winter in Beijing (Yan et al., 2017). Also, large contributions of BBOA and OOA to BrC were also observed in Switzerland during winter (Moschos et al., 2018).

5 4 Conclusions

10

15

We conducted comprehensive measurements of aerosol optical properties at both ground level and 260 m in winter in Beijing using a suite of state-of-the-art instruments, e.g., two CAPSs, two AE33s, SP-AMS, and PAX. The inter-comparisons showed excellent agreements among different instruments. b_{ext} , b_{sca} , and b_{abs} varied dynamically throughout the campaign, but overall showed similar trends between ground level and 260 m. On average, b_{sca} and b_{abs} were 337.6 and 36.6 Mm⁻¹ at 260 m, which are 26.5% and 22.5% lower than those at ground level (459.5 and 47.2 Mm⁻¹, respectively). In fact, ratios_{260m/ground} of b_{sca} and b_{abs} varied substantially from <0.1 to ~1.5, indicating very complex vertical differences of aerosol optical properties in winter in Beijing. In particular, low ratios_{260m/ground} were frequently observed during cleaning stages of severe haze episodes and nighttime with strong local emissions and low MLH. The diurnal cycles of ratios_{260m/ground} and $R^2_{260m/ground}$ further indicated that the change of MLH played an important role in driving the vertical differences between daytime and nighttime. Although the vertical difference of SSA was not as much as those of b_{sca} and b_{abs} , we observed ubiquitously higher values at ground level than 260 m, suggesting relative more scattering aerosols at ground site. A case study further showed complex vertical changes of aerosol optical properties during the formation, evolution and cleaning stages of the haze episode, which were mainly due to the changes in meteorological conditions (e.g., *T* inversion), MLH, and source emissions.

The MAC of BCeBC showed a wide range from 9.5 to 13.2 m²g⁻¹. MAC was observed to increase substantially as a function of M_R during periods with high PM levels, and the increases were closely related with the increases in secondary aerosols species. The positive relationship between MAC and O/C, and their diurnal profiles further suggest that photochemical processing can increase MAC by approximately 20%, and up to 40%. It should be noted that the MAC measured in this study could be different from the MAC of ambient aerosol since the particles were dried and sampled at constant room temperature. BrC was a large contributor to the absorption at 370 nm in winter, on average accounting for 46% and 48% at ground level and 260 m, respectively. By linking with OA factors from PMF analysis, we found that BrC in winter is dominantly contributed by coal combustion (48 - 55%) followed by biomass burning (17%) and photochemical SOA (~20%).

5 Data availability. The data in this study are available from the authors upon request (sunyele@mail.iap.ac.cn).

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was supported by the National Natural Science Foundation of China (91744207, 41571130034, 41575120) and the National Key Project of Basic Research (2014CB447900). Qingqing Wang acknowledges the support from the General Financial Grant from the China Postdoctoral Science Foundation (2017M610972) and the National Postdoctoral Program for Innovative Talents (BX201600157).

References

10

15

- Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, Journal of Geophysical Research, 111, 10.1029/2006jd007315, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, Journal of Geophysical Research: Atmospheres, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., Jobson, B. T., Kolesar, K. R., Lack, D. A., Lerner, B. M., Li, S.-M., Mellon, D., Nuaaman, I., Olfert, J. S., Petäjä, T., Quinn, P. K., Song, C., Subramanian, R., Williams, E. J., and Zaveri, R. A.: Radiative Absorption Enhancements Due to the Mixing State of Atmospheric Black Carbon, Science, 337, 1078-1081, 10.1126/science.1223447, 2012.
- 25 Chen, C., Sun, Y. L., Xu, W. Q., Du, W., Zhou, L. B., Han, T. T., Wang, Q. Q., Fu, P. Q., Wang, Z. F., Gao, Z. Q., Zhang, Q., and Worsnop, D. R.: Characteristics and sources of submicron aerosols above the urban canopy (260 m) in Beijing, China, during the 2014 APEC summit, Atmospheric Chemistry and Physics, 15, 12879-12895, 10.5194/acp-15-12879-2015, 2015.

- Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V. M., Petäjä, T., Su, H., Cheng, Y. F., Yang, X. O., Wang, M. H., Chi, X. G., Wang, J. P., Virkkula, A., Guo, W. D., Yuan, J., Wang, S. Y., Zhang, R. J., Wu, Y. F., Song, Y., Zhu, T., Zilitinkevich, S., Kulmala, M., and Fu, C. B.: Enhanced haze pollution by black carbon in megacities in China, Geophysical Research Letters, 43, 2873-2879, 10.1002/2016gl067745, 2016.
- 5 Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Müller, T., Wiedensohler, A., and Hansen, A. D. A.: The "dual-spot" Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, Atmos. Meas. Tech., 8, 1965-1979, 10.5194/amt-8-1965-2015, 2015.
 - Du, W., Zhao, J., Wang, Y., Zhang, Y., Wang, Q., Xu, W., Chen, C., Han, T., Zhang, F., Li, Z., Fu, P., Li, J., Wang, Z., and Sun, Y.: Simultaneous measurements of particle number size distributions at ground level and 260 m on a meteorological
- 10 tower in urban Beijing, China, Atmospheric Chemistry and Physics Discussions, 1-24, 10.5194/acp-2016-1064, 2017. Du, Z., He, K., Cheng, Y., Duan, F., Ma, Y., Liu, J., Zhang, X., Zheng, M., and Weber, R.: A yearlong study of water-soluble organic carbon in Beijing II: Light absorption properties, Atmospheric Environment, 89, 235-241, 10.1016/j.atmosenv.2014.02.022, 2014.
 - Han, T., Xu, W., Chen, C., Liu, X., Wang, Q., Li, J., Zhao, X., Du, W., Wang, Z., and Sun, Y.: Chemical apportionment of aerosol optical properties during the Asia-Pacific Economic Cooperation summit in Beijing, China, Journal of Geophysical Research: Atmospheres, 120, 12,281-212,295, 10.1002/2015JD023918, 2015.
 - Han, T., Xu, W., Li, J., Freedman, A., Zhao, J., Wang, Q., Chen, C., Zhang, Y., Wang, Z., Fu, P., Liu, X., and Sun, Y.: Aerosol Optical Properties Measurements by a CAPS Single Scattering Albedo Monitor: Comparisons between Summer and Winter in Beijing, China, Journal of Geophysical Research: Atmospheres, 10.1002/2016jd025762, 2017.
- 20 Haywood, J., and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review, Reviews of Geophysics, 38, 513-543, 10.1029/1999rg000078, 2000.
 - Horvath, H.: Experimental calibration for aerosol light absorption measurements using the integrating plate method—Summary of the data, Journal of Aerosol Science, 28, 1149-1161, https://doi.org/10.1016/S0021-8502(97)00007-4, 1997.

Kim, D., and Ramanathan, V.: Solar radiation budget and radiative forcing due to aerosols and clouds, Journal of Geophysical

25

30

35

15

Research, 113, 10.1029/2007jd008434, 2008.

- Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, Proceedings of the National Academy of Sciences, 109, 14802-14807, 10.1073/pnas.1206575109, 2012.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, Chemical Reviews, 115, 4335-4382, 10.1021/cr5006167, 2015.
- Liu, D., Taylor, J. W., Young, D. E., Flynn, M. J., Coe, H., and Allan, J. D.: The effect of complex black carbon microphysics on the determination of the optical properties of brown carbon, Geophysical Research Letters, 42, 613-619, 10.1002/2014GL062443, 2015a.

Liu, D., Whitehead, J., Alfarra, M. R., Reves-Villegas, E., Spracklen, Dominick V., Reddington, Carly L., Kong, S., Williams, Paul I., Ting, Y.-C., Haslett, S., Taylor, Jonathan W., Flynn, Michael J., Morgan, William T., McFiggans, G., Coe, H., and Allan,

James D.: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, Nature Geoscience, 10, 184-188, 10.1038/ngeo2901, 2017.

- Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M. K., Cappa, C. D., Williams, L. R., Herndon, S. C., Massoli, P., Fortner, E. C., Chhabra, P. S., Brooks, W. A., Onasch, T. B., Jayne, J. T., Worsnop, D. R., China, S., Sharma, N., Mazzoleni, C., Xu, L., Ng,
- N. L., Liu, D., Allan, J. D., Lee, J. D., Fleming, Z. L., Mohr, C., Zotter, P., Szidat, S., and Prevot, A. S.: Enhanced light absorption by mixed source black and brown carbon particles in UK winter, Nat Commun, 6, 8435, 10.1038/ncomms9435, 2015b.

5

10

15

- Moschos, V., Kumar, N. K., Daellenbach, K. R., Baltensperger, U., Prévôt, A. S. H., and El Haddad, I.: Source Apportionment of Brown Carbon Absorption by Coupling Ultraviolet–Visible Spectroscopy with Aerosol Mass Spectrometry, Environmental Science & Technology Letters, 5, 302-308, 10.1021/acs.estlett.8b00118, 2018.
- Onasch, T. B., Trimborn, A., Fortner, E. C., Jayne, J. T., Kok, G. L., Williams, L. R., Davidovits, P., and Worsnop, D. R.: Soot Particle Aerosol Mass Spectrometer: Development, Validation, and Initial Application, Aerosol Science and Technology, 46, 804-817, 10.1080/02786826.2012.663948, 2012.
 - Onasch, T. B., Massoli, P., Kebabian, P. L., Hills, F. B., Bacon, F. W., and Freedman, A.: Single Scattering Albedo Monitor for Airborne Particulates, Aerosol Science and Technology, 49, 267-279, 10.1080/02786826.2015.1022248, 2015.
 - Peng, J., Hu, M., Guo, S., Du, Z., Zheng, J., Shang, D., Levy, Z. M., Zeng, L., Shao, M., and Wu, Y. S.: Markedly enhanced absorption and direct radiative forcing of black carbon under polluted urban environments, Proceedings of the National Academy of Sciences of the United States of America, 2016.
- Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S. M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto,
 - N., Wehrli, C., Wiedensohler, A., and Zhang, X. Y.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365-8379, 10.5194/acp-13-8365-2013, 2013.
 - Ran, L., Deng, Z. Z., Wang, P. C., and Xia, X. A.: Black carbon and wavelength-dependent aerosol absorption in the North China Plain based on two-year aethalometer measurements, Atmospheric Environment, 142, 132-144, <u>http://dx.doi.org/10.1016/j.atmosenv.2016.07.014</u>, 2016.
- 25 Rosenfeld, D.: TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall, Geophysical Research Letters, 26, 3105-3108, 10.1029/1999gl006066, 1999.
 - Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., and Bex, V.: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Computational Geometry, 18, 95-123, 2013.
- 30 Sun, J., Zhi, G., Hitzenberger, R., Chen, Y., Tian, C., Zhang, Y., Feng, Y., Cheng, M., Zhang, Y., Cai, J., Chen, F., Qiu, Y., Jiang, Z., Li, J., Zhang, G., and Mo, Y.: Emission factors and light absorption properties of brown carbon from household coal combustion in China, Atmospheric Chemistry and Physics, 17, 4769-4780, 10.5194/acp-17-4769-2017, 2017.
 - Sun, Y., Wang, Z., Fu, P., Jiang, Q., Yang, T., Li, J., and Ge, X.: The impact of relative humidity on aerosol composition and evolution processes during wintertime in Beijing, China, Atmospheric Environment, 77, 927-934, <u>http://dx.doi.org/10.1016/j.atmosenv.2013.06.019</u>, 2013.

- Sun, Y., Du, W., Wang, Q., Zhang, Q., Chen, C., Chen, Y., Chen, Z., Fu, P., Wang, Z., Gao, Z., and Worsnop, D. R.: Real-Time Characterization of Aerosol Particle Composition above the Urban Canopy in Beijing: Insights into the Interactions between the Atmospheric Boundary Layer and Aerosol Chemistry, Environ Sci Technol, 49, 11340-11347, 10.1021/acs.est.5b02373, 2015.
- 5 Sun, Y., Wang, Z., Wild, O., Xu, W., Chen, C., Fu, P., Du, W., Zhou, L., Zhang, Q., Han, T., Wang, Q., Pan, X., Zheng, H., Li, J., Guo, X., Liu, J., and Worsnop, D. R.: "APEC Blue": Secondary Aerosol Reductions from Emission Controls in Beijing, Sci Rep, 6, 20668, 10.1038/srep20668, 2016.
 - Sun, Y. L., Jiang, Q., Wang, Z. F., Fu, P. Q., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013, J. Geophys. Res.-Atmos., 119, 4380-4398, 10.1002/2014jd021641, 2014.
- 10 Tang, G., Zhu, X., Hu, B., Xin, J., Wang, L., Münkel, C., Mao, G., and Wang, Y.: Impact of emission controls on air quality in Beijing during APEC 2014: lidar ceilometer observations, Atmospheric Chemistry and Physics, 15, 12667-12680, 10.5194/acp-15-12667-2015, 2015.

20

- Tang, G., Zhang, J., Zhu, X., Song, T., Münkel, C., Hu, B., Schäfer, K., Liu, Z., Zhang, J., Wang, L., Xin, J., Suppan, P., and Wang, Y.: Mixing layer height and its implications for air pollution over Beijing, China, Atmospheric Chemistry and Physics, 16, 2459-2475, 10.5194/acp-16-2459-2016, 2016.
- Tao, J., Zhang, L., Gao, J., Wang, H., Chai, F., and Wang, S.: Aerosol chemical composition and light scattering during a winter season in Beijing, Atmospheric Environment, 110, 36-44, <u>http://dx.doi.org/10.1016/j.atmosenv.2015.03.037</u>, 2015.
- Wang, J., Onasch, T. B., Ge, X., Collier, S., Zhang, Q., Sun, Y., Yu, H., Chen, M., Prévôt, A. S. H., and Worsnop, D. R.: Observation of Fullerene Soot in Eastern China, Environmental Science & Technology Letters, 3, 121-126, 10.1021/acs.estlett.6b00044, 2016.
- Wang, J., Zhang, Q., Chen, M. D., Collier, S., Zhou, S., Ge, X., Xu, J., Shi, J., Xie, C., and Hu, J.: First chemical characterization of refractory black carbon aerosols and associated coatings over the Tibetan Plateau (4730 m a.s.l), Environmental Science & Technology, 51, 14072, 2017a.
- Wang, J., Liu, D., Ge, X., Wu, Y., Chen, M., Zhao, J., Xie, C., Wang, Q., Xu, W., Allan J., Joshi R., Fu P., Coe, H., and Sun, Y.: Characterization of black carbon-containing fine particles in Beijing during wintertime, 2018, submitted
- Wang, Q., Sun, Y., Jiang, Q., Du, W., Sun, C., Fu, P., and Wang, Z.: Chemical composition of aerosol particles and light extinction apportionment before and during heating season in Beijing, China, Journal of Geophysical Research Atmospheres, 120, 12708-12722, 2015.
- Wang, Q., Sun, Y., Xu, W., Du, W., Zhou, L., Tang, G., Chen, C., Cheng, X., Zhao, X., Ji, D., Han, T., Wang, Z., Li, J., and Wang,
 Z.: Vertically-resolved Characteristics of Air Pollution during Two Severe Winter Haze Episodes in Urban Beijing, China, Atmos. Chem. Phys. Discuss., 2017, 1-28, 10.5194/acp-2017-891, 2017b.
 - Wang, Q., Du, W., Zhou, W., Xie, C., Zhang, Y., Xu, W., Zhao, J., Marco, C., Nemitz, E., Fu, P., Q., Lee, J., Wang, Z., and Sun, Y, Real-Time Characterization of Vertical Distribution of Submicron Aerosol Chemical Species in Winter Based on the Beijing 325 m Meteorological Tower, 2018, inpreparation
- 35 Wang, Z., Huang, X., and Ding, A.: Dome effect of black carbon and its key influencing factors: a one-dimensional modelling study,

Atmospheric Chemistry and Physics, 18, 2821-2834, 10.5194/acp-18-2821-2018, 2018.

10

15

- Xu, W. Q., Sun, Y. L., Chen, C., Du, W., Han, T. T., Wang, Q. Q., Fu, P. Q., Wang, Z. F., Zhao, X. J., Zhou, L. B., Ji, D. S., Wang, P. C., and Worsnop, D. R.: Aerosol composition, oxidation properties, and sources in Beijing: results from the 2014 Asia-Pacific Economic Cooperation summit study, Atmos. Chem. Phys., 15, 13681-13698, 10.5194/acp-15-13681-2015, 2015.
- 5 Xu, W. Q., Sun, Y. L., Wang, Q. Q., Zhao J., Wang, J. F., Ge, X. L., Xie, C. H., Zhou, W., Du, W., Li J., Fu, P. Q., Wang, Z. F., Worsnop, D. R. and Coe, H.: Changes in aerosol chemistry from 2014 to 2016 in winter in Beijing: insights from high resolution aerosol mass spectrometry, 2018, submitted.
 - Yan, C., Zheng, M., Sullivan, A. P., Bosch, C., Desyaterik, Y., Andersson, A., Li, X., Guo, X., Zhou, T., Gustafsson, Ö., and Collett Jr, J. L.: Chemical characteristics and light-absorbing property of water-soluble organic carbon in Beijing: Biomass burning contributions, Atmospheric Environment, 121, 4-12, http://dx.doi.org/10.1016/j.atmosenv.2015.05.005, 2015.
 - Yan, C., Zheng, M., Bosch, C., Andersson, A., Desyaterik, Y., Sullivan, A. P., Collett, J. L., Zhao, B., Wang, S., He, K., and Gustafsson, O.: Important fossil source contribution to brown carbon in Beijing during winter, Sci Rep, 7, 43182, 10.1038/srep43182, 2017.
 - Yang, M., Howell, S. G., Zhuang, J., and Huebert, B. J.: Attribution of aerosol light absorption to black carbon, brown carbon, and dust in China – interpretations of atmospheric measurements during EAST-AIRE, Atmos. Chem. Phys., 9, 2035-2050, 10.5194/acp-9-2035-2009, 2009.
 - Zhao, J., Du, W., Zhang, Y., Wang, Q., Chen, C., Xu, W., Han, T., Wang, Y., Fu, P., Wang, Z., Li, Z., and Sun, Y.: Insights into aerosol chemistry during the 2015 China Victory Day parade: results from simultaneous measurements at ground level and 260 m in Beijing, Atmospheric Chemistry and Physics, 17, 3215-3232, 10.5194/acp-17-3215-2017, 2017.
- 20 Zhou, W., Zhao, J., Ouyang, B., Mehra, A., Xu, W., Wang, Y., Bannan, T. J., Worrall, S. D., Priestley, M., and Bacak, A.: Production of N2O5 and ClNO2 in summer in urban Beijing, China, Atmospheric Chemistry & Physics, 1-39, 2018.

 Table 1. A summary of measurements in this study

I

	Instruments	Manufacturer	Properties measured	Resolution
Ground	Cavity attenuated phase shift single scattering albedo monitor (CAPS PM _{ssa})	Aerodyne Research, Inc.	PM ₁ b_{ext} and b_{sca} (λ = 630 nm)	1 s
	Aethalometer (AE331)	Magee Scientific Corp.	BCeBC (370, 470, 520, 590, 660, 880, and 950 nm)	1 min
	Photoacoustic Extinctiometer (PAX)	Droplet Measurement Technologies	PM _{2.5} b_{ext} and b_{abs} (λ = 870 nm)	1 s
	High-Resolution Time-of- Flight Aerosol Mass Spectrometer (HR-AMS)	Aerodyne Research, Inc.	NR-PM ₁ species (organics, sulfate, nitrate, ammonium, and chloride)	5 min
	Soot Particle Aerosol Mass Spectrometer (SP-AMS)	Aerodyne Research, Inc.	BC-containing PM_1 species (<i>r</i> BC, organics, sulfate, nitrate, ammonium, and chloride)	5 min
	Cavity Attenuated Phase Shift Particulate Matter extinction monitor (CAPS PM _{ext}	Aerodyne Research, Inc.	PM _{2.5} b _{ext} (λ=630 nm)	1 s
260 m	Actuationiteter (AE351)Corp.and 950 nm)PhotoacousticDropletExtinctiometer (PAX)DropletHigh-ResolutionTime-of- TechnologiesPM2.5 b_{ext} and b_{abs} (λ = 870 nm)High-ResolutionTime-of- FlightAerodyne Research, Inc.NR-PM1 species (organics, nitrate, ammonium, and chlorideSoot Particle Aerosol Mass Spectrometer (SP-AMS)Aerodyne Research, Inc.BC-containingPM1 species organics, sulfate, nitrate, amm and chloride)Cavity Attenuated Phase Shift Particulate Matter extinction monitor (CAPS PMextAerodyne Research, Inc.BC-gBC (370, 470, 520, 590, 66 and 950 nm)Aethalometer (AE332)Magee Corp.Scientific Corp.BC-gBC (370, 470, 520, 590, 66 and 950 nm)Aerosol Chemical Species Monitor (ACSM, Aerodyne)Aerodyne Research, Inc.NR-PM1 species (organics, nitrate, ammonium, and chloride)Photoacoustic Extinctiometer (PAX)Droplet Measurement TechnologiesPM2.5 b_{sca} and b_{abs} (λ =870 nm)Aethalometer (AE331)Magee Magee ScientificPM2.5 b_{sca} and b_{abs} (λ =870 nm)	BCeBC (370, 470, 520, 590, 660, 880, and 950 nm)	1 min	
	Aerosol Chemical Species Monitor (ACSM, Aerodyne)	Aerodyne Research, Inc.	NR-PM ₁ species (organics, sulfate, nitrate, ammonium, and chloride)	5 min
Vertical	Photoacoustic Extinctiometer (PAX)	Droplet Measurement Technologies	PM _{2.5} $b_{\rm sca}$ and $b_{\rm abs}$ (λ =870 nm)	1 s
	Aethalometer (AE33 ₁)	Magee Scientific Corp.	BCeBC (370, 470, 520, 590, 660, 880, and 950 nm)	1 s

Table 2. A summary of average $(\pm 1\sigma)$ meteorological parameters and optical properties at ground level and 260 m during this study

	260 m	Ground	$R_{ m 260m/Ground}$		
Meteorological parameters					
<i>T</i> (°C)	2.3 ± 3.7	4.3 ± 3.7	0.53		
RH (%)	49.1 ± 23.5	48.5 ± 20.7	1.01		
WS (m s ⁻¹)	4.0 ± 2.4	1.4 ± 1.0	2.85		
Optical properties					
$b_{\rm ext} ({ m Mm^{-1}})$	387.8 ± 395.2	547.0 ± 555.9	0.71		
$b_{\rm sca}({\rm Mm}^{-1})$	337.6 ± 356.0	459.5 ± 483.1	0.73		
$b_{\rm abs}({ m Mm}^{-1})$	36.6 ± 33.9	47.2 ± 2.4	0.78		
SSA	0.88 ± 0.04	0.89 ± 0.04	0.98		
AAE	1.60 ± 0.18	1.58 ± 0.15	1.01		





Figure 1. Inter-comparisons of aerosol optical properties and BC: (a) *b*_{ext} of PM_{2.5} from CAPS PM_{ext} and CAPS PM_{ssa}, (b) **BC**<u>eBC</u> from two AE33s, (c) *b*_{ext} of PM2.5 from CAPS PM_{ext} vs. *b*_{ext} of PM1 from CAPS PM_{ssa}, (d) *b*_{sca} from CAPS PM_{ssa} vs. *b*_{sca} from PAX.



Figure 2. Comparisons of (a) BC measurements between AE33 and SP-AMS, (b) b_{abs} calculated from the difference of b_{ext} and b_{sca} of CAPS PM_{ssa} and that was derived from **BC**<u>eBC</u>.



Figure 3. Temporal variations of (a) temperature (T) and relative humidity (RH) at ground level and 280 m, (b) wind speed (WS) and wind direction (WD) at 280 m, and comparisons of the time series of optical properties of PM_{2.5} between ground level (black lines) and 260 m (color lines): (c) b_{ext} , (d) scattering coefficient, (e) absorption coefficient, (f) SSA.



Figure 4. Comparisons of (a) b_{sca} and (b) b_{abs} between 260 m and ground level. The scatter plot is color coded by mixed layer height, and the marker size is proportional to the RH difference between ground level and 280 m. (c) shows the frequencies of ratios of 260 m to ground level for b_{sca} and b_{abs} .



Figure 5. Time series of hourly aveage ratios_{260m/ground} for b_{sca} and b_{abs} during this study. The ratios are color coded by b_{ext} , and the black circles and squares are those from continuous vertical measurements with PAX.



Figure 6. Diurnal cycles of (a) b_{sca} , (b) b_{abs} , and (c) SSA at 260 m and ground level for the entire study. Right panel shows the diurnal cycles of correlation coefficients (R^2) and ratio_{260m/ground} for b_{sca} , b_{abs} , and SSA. Note that the diurnal cycles above excluded 5 periods with cleaning processes (November 19 1:00 to 9:00, November 26 1:00 to 12:00, November 26 19:00 to November 27 3:00, December 3 20:00 to December 4 16:00, and December 17 00:00 to 12:00) while those of entire study are shown in Fig. S3.



Figure 7. Evolution of vertical profiles of b_{sca} , b_{abs} and SSA during 2 – 4 December. Also shown are (a-d) meteorological variable of *T*, RH, WD, and WS, and (e) time series of b_{ext} at ground and MLH. M16-M25 refer to the number of vertical profiles.



Figure 8. Vertical profiles of b_{sca} , b_{abs} and SSA measured by PAX at 880 nm during (a) M19 and (b) M20. (c) and (d) show the average vertical profiles of T, RH, WS, and WD during M19 and M20, respectively. The pie charts show average chemical composition of PM₁ at 260 m (left) and ground level (right) during M19 and M20, respectively.



Figure 9. (a) Variations of mass absorption cross-section (MAC) as a function of $M_{\rm R}$ at different pollution levels and (b) variations of mass fraction of *r*BC-coated species as a function of $M_{\rm R}$.



Figure 10. (a) Diurnal cycles of <u>MACandMAC and</u> O/C ratio of SP-AMS (O/C_{SP-AMS}), and (b) variations of MAC as a function of O/C_{SP-AMS}. The data are binned according to O/C. The median (horizontal line), mean (triangle), 25th and 75th percentiles (lower and upper box), and 10th and 90th percentiles (lower and upper whiskers) are also shown.



Figure 11. (a) Diurnal cycles of *b*_{abs,BrC}/OA, i.e., MAC_{org, 370 nm} at ground level and 260 m, (b) variations of the ratio of BrC absorption to the total absorption at 370 nm as a function of the ratio of organics between SP-AMS and HR-AMS. The data

are binned according to O/C. The median (horizontal line), mean (triangle), 25th and 75th percentiles (lower and upper box), and 10th and 90th percentiles (lower and upper whiskers) are also shown.



Figure 12. Average contributions of OA sources to BrC absorption at ground level from (a) linear regression model-<u>analysis</u>, and (b) positive matrix factorization (Fig. <u>\$859</u>). The sources of OA include fossil-fuel-related OA (FFOA), cooking OA (COA), biomass burning OA (BBOA), oxygenated OA (OOA), oxidized primary OA (OPOA), aqueous-phase OOA (aq-OOA) (Xu et al. 2018).