

Reviewer 2 Comments

The simulations of particle number concentration and the corresponding source apportionment are very difficult in the model. We should encourage the relevant research. Still, more work should be included in the further study, especially for the model verification.

Comment 1. Maybe it is reasonable to set the first size bin as 7 nm in the model calculation, which will benefit the comparison.

Response: The reviewer makes a good suggestion to remove the uncertainty associated with comparing N7 measurements to N10 model predictions. We will incorporate this suggestion into future studies.

2. Page 12, line 289: I do not understand the Table 2. Particle number concentration and PM2.5 came from the different sources, so why you demonstrate the correlation (R2) ? Also, you use R in table 1, and R2 in table 2, why?

Response: The analysis showing correlation coefficients between particle number and PM2.5 was requested by a previous reviewer as another metric to test model performance. Predicted correlation coefficients tend to be higher than measured correlation coefficients suggesting that the model calculations do not capture all of the complexity in the real atmosphere. This has been clarified on line 294 of the revised manuscript. As noted in the current text, some of the higher correlation between particle number and PM2.5 in the model results may be caused by the artificial smoothing inherent in the 4km model grid cells.

The correlation coefficient in Table 2 has been changed to R to match with Table 1.

3. What is the Y-axis label for Figs. 10-12?

Response: The y-axis label should read $PM_{0.1}$ ($\mu\text{g m}^{-3}$). This has been corrected in the revised manuscript.

4. I think the figure label is wrong in Fig. 15, also why the scale of the x-axis in Figs. 15-17 is from 1 to 24?

Response: The figure label in Fig 15 has been corrected to label the panels a,b,c,d,e,f,g,h. The x-axis scale representing hours of the day as part of the diurnal profile has been revised to properly show hours 1-24 counting by 2.

5. There are several grammar mistakes in the text, the language and symbols should be checked carefully once more before publication.

Response: The text has been carefully proofread and necessary corrections have been made.

Regional Sources of Airborne Ultrafine Particle Number and Mass Concentrations in California

Xin Yu¹, Melissa Venecek², Anikender Kumar¹, Jianlin Hu³, Saffet Tanrikulu⁴, Su-Tzai Soon⁴, Cuong Tran⁴, David Fairley⁴, and Michael J. Kleeman^{1*}

¹Department of Civil and Environmental Engineering, University of California, Davis. One Shields Avenue, Davis CA. ²Department of Land, Air, and Water Resources, University of California, Davis. One Shields Avenue, Davis CA. ³School of Environmental Science and Engineering, Nanjing University of Information Science and Technology. ⁴Bay Area Air Quality Management District, San Francisco, CA.

*Corresponding author. Tel.: +1 530 752 8386; fax: +1 530 752 7872. E-mail address: mjkleeman@ucdavis.edu (M.J. Kleeman).

Abstract

Regional concentrations and source contributions are calculated for airborne particle number concentration (N_x) and ultrafine particle mass concentration ($PM_{0.1}$) in the San Francisco Bay Area (SFBA) and the South Coast Air Basin (SoCAB) surrounding Los Angeles with 4 km spatial resolution and daily time resolution for selected months in the years 2012, 2015, and 2016. Performance statistics for daily predictions of N_{10} concentrations meet the goals typically used for modeling of $PM_{2.5}$ (MFB < ± 0.5 and MFE < 0.75). The relative ranking and concentration range of source contributions to $PM_{0.1}$ predicted by regional calculations agree with results from receptor-based studies that use molecular markers for source apportionment at four locations in California. Different sources dominated regional concentrations of N_{10} and $PM_{0.1}$ because of the different emitted particle size distributions and different choices for heating fuels. Nucleation (24-57%) made the largest single contribution to N_{10} concentrations at the ten regional monitoring locations, followed by natural gas combustion (28-45%), aircraft (2-10%), mobile sources (1-5%), food cooking (1-2%), and wood smoke (0-1%). In contrast, natural gas combustion (22-52%) was the largest source of $PM_{0.1}$ followed by mobile sources (15-42%), food cooking (4%-14%), wood combustion (1-12%), and aircraft (2-6%). The study region encompassed in this project is home to more than 25M residents, which should provide sufficient power for future epidemiological studies on the

health effects of airborne ultrafine particles. All of the PM_{0.1} and N₁₀ outdoor exposure fields produced in the current study are available free of charge at http://webwolf.engr.ucdavis.edu/data/soa_v2/monthly_avg2.

35

1. Introduction

Numerous epidemiological studies have identified positive correlations between exposure to ambient particulate matter (PM) and increased risk of respiratory and cardiovascular diseases, premature mortality and hospitalization (Pope et al., 2002; Pope et al., 2004; Pope et al., 2009; Dockery and Stone, 2007; Ostro et al., 2015; Ostro et al., 2006; Ostro et al., 2010; Brunekreef and Forsberg, 2005; Fann et al., 2012; Gauderman et al., 2015; Miller et al., 2007). Most of these studies have not fully addressed ultrafine particles (UFPs; D_p<0.1μm) because these particles make a very small contribution to total ambient PM mass (Ogulei et al., 2007). Toxicity studies suggest that UFPs may be especially dangerous to human health since they have higher toxicity per unit mass (Li et al., 2003; Nel et al., 2006; Oberdorster et al., 2002) and can penetrate the lungs and enter the bloodstream and secondary organs (Sioutas et al., 2005). These toxicology results are suggestive but more epidemiological evidence is required before the threat to public health from UFPs can be fully assessed.

Most previous UFP epidemiology studies are based on particle number concentration (N_x – the number of particles with diameter less than X nm) measured at fixed sites using commercially-available instruments. These devices are expensive and they require regular maintenance which limits the number of measurement sites that can be deployed. Translating measured N_x into population exposure estimates is also difficult because UFP concentrations change more rapidly over shorter distances than PM_{2.5} (Hu et al., 2014b; Hu et al., 2015; Hu et al., 2014a). Land use regression (LUR) models could potentially be used to interpolate UFP concentrations between sparse measurement locations, but the atmospheric processes governing N_x concentrations are highly non-linear and (so far) sufficient training data is not generally available for LUR models to estimate N_x exposure over a large enough population to support a definitive epidemiology

study (Montagne et al., 2015). Previous attempts to use regional reactive chemical transport models to predict N_x in highly populated regions have focused on nucleation, yielding a wide range of predicted concentrations and only modest agreement with measurements when different nucleation algorithms were used (Elleman and Covert, 2009b; Zhang et al., 2010; Elleman and Covert, 2009a). Obtaining accurate exposure estimates to N_x in highly populated regions therefore remains a major challenge in UFP epidemiological studies.


Recent work has examined UFP mass ($PM_{0.1}$) as an alternative metric for UFP exposure, and demonstrated that $PM_{0.1}$ can be predicted with reasonable accuracy over large populations using regional reactive chemical transport models (Hu et al., 2014b; Hu et al., 2014a). The $PM_{0.1}$ exposure fields developed using this technique have been used in multiple epidemiological studies that revealed associations with mortality and pre-term birth (Ostro et al., 2015; Laurent et al., 2016). Despite the success of studies using $PM_{0.1}$, techniques that estimate N_x exposure are still needed because a large number of ongoing UFP studies are based on N_x and it is possible that $PM_{0.1}$ and N_x are associated with different types of health effects.

Here we extend the previous work using regional reactive chemical transport models for UFPs to include N_x in the San Francisco Bay Area (SFBA) and the South Coast Air Basin (SoCAB) region around Los Angeles which are the two most densely populated major metropolitan location in California. Source contributions to $PM_{0.1}$ and N_x are tracked using the University of California, Davis / California Institute of Technology (UCD/CIT) regional reactive chemical transport model with 4 km spatial resolution. Predicted concentrations during the year 2012 are compared to measurements available at ten regional monitoring sites. The spatial distribution fields of different particle metrics (N_x , $PM_{0.1}$, $PM_{2.5}$) are combined with population distributions to estimate exposure. To the best of our knowledge, this is the first integrated study of both UFP number and mass using a regional reactive chemical transport model in California.

2. Model Description

The UCD/CIT chemical transport model used in the current study has been successfully applied in several previous studies in the San Joaquin Valley (SJV) and the SoCAB (Ying et al., 2008b; Ying et al., 2008a; Hu et al., 2015; Hu et al., 2017; Chen et al., 2010; Held et al., 2004; Held et al., 2005; Hixson et al., 2010; Hixson et al., 2012; Hu et al., 2012; Kleeman and Cass, 2001; Kleeman et al., 2007; Kleeman et al., 1997; Mahmud, 2010; Mysliwiec and Kleeman, 2002; Rasmussen et al., 2013; Ying and Kleeman, 2006; Zhang and Ying, 2010). The model includes algorithms for emissions, transport, dry deposition, wet deposition, gas phase chemistry, gas-to-particle conversion, coagulation, and some condensed phase chemical reactions. Nucleation was added to the model for the first time in the current study using the ternary nucleation (TN) mechanism involving H₂SO₄-H₂O-ammonia (NH₃) (Napari et al., 2002). As was the case in previous studies using this algorithm, the resulting nucleation rate was adjusted using a tunable nucleation parameter set to 10⁻⁵ for new particle nucleation (Jung et al., 2010). The Kerminen and Kulmala (2002) parameterization was added in order to bridge the gap between the 1 nm particle nuclei and their appearance into the smallest size bin of the UCD/CIT model (~10 nm). The nuclei growth rate (GR) in the Kerminen and Kulmala (2002) parameterization is one of the factors that accounts for the competition between the condensation and nucleation of over-saturated compounds until the nucleated particles grow to the size of the smallest bin in the regional model, at which point this competition is represented explicitly by the model operators. In the current study, the GR for nucleated sulfate particles was calculated using the diffusion-limited condensation rate of sulfuric acid based on the recommendation of Kerminen and Kulmala. Once particles reach ~10nm, the full operators in the model calculations predict growth by condensation of sulfuric acid, nitric acid, ammonia, and secondary organic aerosol (SOA). Perturbation studies were conducted in the current analysis to test the effect of GR with a box model configured to represent a single grid cell using the full set of model operators. Initial conditions in the SAPRC11 gas-phase mechanism were 0.04 ppm O₃, 0.05 ppm NO, 0.0 ppm NO₂, 0.05 ppm HCHO, 0.1 ppm ISOPRENE, 0.1 ppm BENZENE, and 0.01 ppm ALK5. A nucleation event was initiated at 8am by setting H₂SO₄ concentrations to 10⁷ molecules cm⁻³ and NH₃ concentrations to 100 ppt. The

120 nominal GR was multiplied by a factor ranging from 0.5 to 2.0 to test the sensitivity of
the model results. Figure 1 illustrates the growth of nucleated particles between 5am and
12 noon for conditions representing July in California. The number concentration of
nucleated particles increases from zero to values between 2500 - 3000 # cm⁻³. SOA
condenses on the particles causing their size to increase above 100nm. Coagulation and
deposition processes remove particles over time. Three separate simulations are
125 illustrated in Figure 1 using the nominal GR along with perturbations of 0.5*GR and
2.0*GR. These model perturbations fall almost exactly on top of the basecase
simulations, suggesting that results are not overly sensitive to GR during the first few
seconds of nuclei growth before calculations are handed off to the regional model
algorithms.



130

Figure 1: Simulated particle nucleation event followed by growth due to SOA condensation under conditions representing July in California. Vertical axis displays the mean diameter of the nuclei mode while color represents the particle number concentration.


135 Several previous modeling studies have been conducted to evaluate the performance of the ternary nucleation mechanism on predicted N_x using global and regional models. Jung et al. (2010) found that a scaled version of the ternary H₂SO₄-NH₃-H₂O nucleation

theory (Napari et al., 2002 with a supplemental 10^{-5} nucleation tuning factor) added to the PMCAMx-UF model produced N_x predictions in reasonable agreement with
140 observations. The study of Westervelt et al. (2013) also showed that the ternary nucleation parameterization (with a supplemental 10^{-5} nucleation tuning factor) added to the Goddard Earth Observing System global chemical transport model (GEOS-Chem) produced reasonable N_x predictions on average when compared with measurements at five locations spanning various environments. Jung et al. (2008) considered multiple
145 nucleation parameterizations in the Dynamic Model for Aerosol Nucleation (DMAN) to predict the nucleation events and non-events observed during the Pittsburgh Pittsburgh Air Quality Study (PAQS) conducted between July 2001 and September 2002. Their results showed that the ternary nucleation mechanism ((Napari et al., 2002) with a supplemental 10^{-5} nucleation tuning factor) was a suitable nucleation scheme for 3-D
150 chemical transport models. Although there have been numerous significant efforts to incorporate nucleation algorithms into three-dimensional regional and global models (Jung et al., 2008; Jung et al., 2010; Westervelt et al., 2013; Zhang et al., 2010), nucleation modeling studies are still in the early stages of development and further efforts are needed to reduce the uncertainty in both the nucleation rate and growth mechanisms.

155 In the current study, emission, transport, deposition, and coagulation of UFPs were simulated using operators developed for the UCD/CIT model framework, leading to modification of the particle size distribution and the subsequent N_x concentrations. Dynamic condensation / evaporation is considered for all particle size bins with predicted UFP growth rates of 2-3 nm hr^{-1} or higher under favorable conditions. The regional
160 model operators are not well suited for the most extreme changes to the particle size distribution that occur within the first few seconds or minutes after emissions to the atmosphere (such as within 300 m of roadways). Dedicated simulations can predict the dynamic condensation/evaporation of particles at distances of 10's of meters downwind of the roadway (Zhang et al., 2005; Zhang et al., 2004) mostly due to the partitioning of
165 SOA (Anttila and Kerminen, 2003; Trostl et al., 2016), but these calculations are too expensive for domains spanning thousands of km. Regional calculations such as those illustrated in the current study rely on emissions characterization measurements that include a few minutes of aging to capture the “near-field” emissions of particle size and

composition that can then be used as the starting point for regional model calculations. In
170 some cases, evaporation of UFPs in the first few seconds after release to the atmosphere
is therefore represented by reducing the primary emissions of nano-particles based on
measurements conducted at high dilution factors (Xue et al., 2018a) or using
measurements of particle volatility to estimate the evaporation at high dilution factors
(May et al., 2013a; May et al., 2013b; Kuwayama et al., 2015). All of the results presented
175 in the current analysis focus on regional UFP concentrations with 4km resolution.

The model domains used in the study are shown in Figure 2. The parent domain with 24
km horizontal resolution covered the entire state of California (referred to as CA_24 km)
and the two nested domains with 4 km horizontal resolution covered the SFBA + SJV +
South Sacramento Valley air basins (referred as SJV_4 km) and the SoCAB surrounding
180 Los Angeles (referred as SoCAB_4 km). The UCD/CIT model was configured with 16
vertical layers up to a height of 5 km above ground level, with 10 layers in the first 1 km.
Previous studies have shown that this vertical configuration captures the air pollution
system above California (Hu et al., 2014a; Hu et al., 2014b; Hu et al., 2015). Particulate
number, mass, and composition are represented in 15 size bins, with particle diameters
185 being centered within equally spaced logarithmic size interval spanning the diameter
range from 0.01 to 10 μ m. Nucleated particles were initialized in a 16th size bin with
initial diameter of 0.007 μ m.



190 Figure 2: Modeling domains. Blue lines outline the CA_24 km domain, black lines
 outline the SoCAB_4 km (bottom) and SJV_4 km domains (top). Red crosses represent
 ten N_x sites (four sites operated by staff at the Bay Area Air Quality Management
 District (BAAQMD) and six sites from the Multiple Air Toxics Exposure Study IV
 (MATES IV)). Detailed location information for the N_x sites is listed in Table S3. Green
 195 dots represent BAAQMD $PM_{2.5}$ speciation network sites and the Interagency Monitoring
 of Protected Visual Environments (IMPROVE) sites; gray dots represent the $PM_{2.5}$
 federal reference method (FRM) sites.

2.1 Meteorological Fields

Hourly meteorological fields during the modeling period were generated by the Weather
 200 Research and Forecasting (WRF) model version 3.4 with three nested domains that had
 horizontal resolutions of 36 km, 12 km and 4 km, respectively. In the present simulations,
 the WRF model was configured with 50 vertical layers (up to 100 hpa) and four-

dimensional data assimilation (FDDA) nudging was utilized to improve the agreement between model predictions and observed meteorological patterns (Otte, 2008b, a). WRF
205 predictions for wind speed, temperature, and relative humidity were compared to
measurements for seven counties in the SFBA and two counties in SoCAB (see Table
S2). Temperature has mean bias (MB) within ~ 0.2 °C and root-mean-square errors
(RMSE) between 4-5 °C. Wind speed has mean fraction bias (MFB) within ± 0.20 and
RMSE generally < 2.0 m/s. This level of performance is consistent with performance of
210 WRF in previous studies conducted in California (Zhao et al., 2011; Hu et al., 2015).

2.2 Emissions

The emission inventories used in the SFBA were developed by the BAAQMD for the year 2012 based on the regulatory inventory provided by the California Air Resources Board for that same year. The SFBA inventory was processed using the Sparse Matrix
215 Operator Kernel Emissions (SMOKE) v3.7 software package provided by US EPA.
SMOKE was configured to separately tag emissions from on-road gasoline vehicles, off-road gasoline vehicles, on-road diesel vehicles, off-road diesel vehicles, food cooking, biomass burning, non-residential natural gas, and all other sources. The emission inventories used in South Sacramento Valley, SJV and SoCAB were provided by the
220 California Air Resources Board.

Measurements conducted in parallel with the current study found that particles emitted from natural gas combustion in home appliances were semi-volatile when diluted by a factor of 25 in clean air, but particles emitted from reciprocating engines did not evaporate under the same conditions (Xue et al., 2018a). Near-field emissions from all
225 natural gas sources combustion sources other than reciprocating engines were therefore set to 30% of their nominal levels. A map of the natural gas emissions distribution is shown in Supporting Information (Figure S3).

SMOKE results were transformed into size-resolved emissions of particle number, mass,
230 and composition using measured source profiles through an updated version of the emissions model described by Kleeman and Cass (1998). The PM profiles used for each

source type were specified as weighted averages from each of the detailed sources within each broad category as summarized in Table S1. Detailed PM source profiles for major sources of ultrafine particulate matter are based on measurements conducted during source tests (Li and Hopke, 1993;Kleeman et al., 1999, 2000;Robert et al., 2007a;Robert et al., 2007b;Mazaheri et al., 2009). In most cases, these emissions size distributions strongly influence the size distributions of particles in the ambient atmosphere (see Figures S1 and S4). A more detailed discussion of the emissions processing has been presented in a previous study (Hu et al., 2015).

240 3. Results

3.1 Statistical Evaluation

According to Taylor's Hypothesis (Shet et al., 2017), it is expected that the spatial distribution of model results is more important than the temporal distribution when evaluating performance. In the current study model performance evaluations are limited to the locations where measurements were made. Therefore, the temporal distribution is also considered by comparing predicted vs. measured daily average N_x , $PM_{2.5}$ and individual $PM_{2.5}$ species mass concentrations.

The evaluation data set was compiled from several measurement networks including the sites operated by staff at the Bay Area Air Quality Management District (BAAQMD), the IMPROVE sites, the MATES IV sites and FRM sites. In order to account for the uncertainty in predicted wind fields and spatial surrogates used to place emissions, "best-fit" model results were created by identifying the closest match within 3 grid cells of each measurement location. "Best-fit" model performance for $PM_{2.5}$ at routine monitoring sites (Figure 2) meets the performance criteria suggest by Boylan and Russell (Boylan and Russell, 2006) (mean fractional error (MFE) $\leq +0.75$ and mean fractional bias (MFB) $\leq \pm 0.5$) (Table S4). Table S5 shows the MFB and MFE values of gaseous species of O_3 , NO , NO_2 , CO and SO_2 using daily averages across all measurement sites during the entire simulated period. Gaseous species of O_3 , CO , NO , NO_2 and SO_2 have MFBs within ± 0.3 and MFE less than 0.5, indicating consistent behavior between predictions and measurement for these species. The ability of UCD/CIT predictions for key gas species,

mass and chemical component concentrations in the $PM_{0.1}$ and $PM_{2.5}$ size fractions was also evaluated in previous studies (Ying and Kleeman, 2006; Ying et al., 2008a; Ying et al., 2008b; Hu et al., 2012; Chen et al., 2010; Held et al., 2005; Hu et al., 2015; Hu et al., 2017; Venecek et al., 2018). The performance of the UCD/CIT air quality model in these studies generally meets standard model performance criteria. Of greatest interest in the current study, predicted “best-fit” N_{10} values were compared to measured N_7 values at four sites in the SFBA (Santa Rosa, San Pablo, Redwood City and Livermore) and six sites in SoCAB (Anaheim, Central Los Angeles, Compton, Huntington, Inland-valley and Rubidoux). N_7 measurements in the SFBA were made using an Environmental Particle Counter (EPC) Monitor Model 3783 (TSI Inc) while N_7 measurements in the SoCAB were made with EPC Model 3781 (TSI Inc). Both monitors can detect ultrafine particles down to 7 nm which is smaller than the first size bin of 10 nm used in model calculations. Previous studies conducted at Fresno, California, suggest that N_{7-10} accounts for approximately 8% of N_7 (Watson et al., 2011), and so some amount of negative bias is expected when comparing predicted N_{10} to measured N_7 . The evaluation results for “best-fit” N_{10} summarized in Table 1 follow this expected trend but mean fractional bias (MFB) and mean fractional error (MFE) at each comparison site still meet the $PM_{2.5}$ performance criteria suggested by Boylan and Russell (2006). This level of performance is comparable to the results from a previous UFP number simulation conducted in Northern California using a modified version of the WRF-Chem model (Lupascu et al., 2015). The level of agreement between predicted “best-fit” and measured $PM_{2.5}$, individual $PM_{2.5}$ species, key gas species and N_{10} builds confidence in the model skill for UFP predictions in the current study.

285 Table 1. Performance statistics for “best-fit” N₁₀ predictions vs. N₇ at individual monitoring sites. Threshold for PM modeling applications is typically MFB < ± 0.5 and MFE < 0.75.

	Ave Obs. Particles cm ⁻³	Ave Sim. Particles cm ⁻³	R	MFB	MFE	RMSE Particles cm ⁻³
Livermore	8219	9201	0.31	0.10	0.09	3615
Redwood city	11500	11325	0.97	0.02	0.08	1132
San Pablo	10481	15822	0.45	0.30	0.31	10302
Santa Rosa	8655	8967	0.78	0.05	0.15	2063
Anaheim	12850	14812	0.74	0.12	0.14	4239
Central LA	17378	25376	0.31	0.37	0.38	10328
Compton	16203	21036	0.36	0.24	0.26	8127
Huntington	23207	24103	0.77	0.04	0.08	3698
Inland-Valley	15028	16875	0.37	0.12	0.17	4290
Rubidoux	10728	11920	0.66	0.11	0.16	3069

290 Table 2 below summarizes the predicted correlations between daily-average particle number concentrations and PM_{2.5} along with the measured correlations for these metrics. Measured correlations (R²) are less than 0.525 at all locations except Santa Rosa where correlations are above 0.75. Model predictions for daily-average particle number concentrations and PM_{2.5} are more highly correlated, with R² ranging from 0.4722 to 0.8573. The higher correlation between particle number vs. PM_{2.5} in the predicted concentrations suggests that the model does not capture all of the complexity in the real atmosphere. Locations with high R² values such as central Los Angeles also have the highest MFB and MFE and so the high correlation between particle number and PM_{2.5} may reflect inaccuracies in the model inputs. At other locations where traditional model performance metrics suggest that predictions are more accurate, the high correlation between particle number and PM_{2.5} may be related to the model grid resolution. The 4km grid resolution used in the calculations smooths the sharp spatial gradients in the ultrafine particle concentration fields (see Figure 4 below). This same issue makes it difficult for point source measurements to accurately represent 4km average number concentrations. The particle number concentrations measured at a fixed monitoring location may not represent the variation in particle number concentrations a few km

295

300

305

away. PM_{2.5} concentration gradients are smoother, making model predictions and point measurements easier to compare.

Table 2. Daily-average correlation (R²) between PM_{2.5} mass and particle number concentration at 8 sites in California.

R ²	Livermore	Redwood City	San Pablo	Santa Rosa	Anaheim	Central LA	Compton	Rubidoux
Obs	0.200-04	0.100-01	0.400-16	0.760-58	0.280-08	0.370-14	0.390-15	0.470-22
Sim	0.530-28	0.700-49	0.740-55	0.470-22	0.710-51	0.850-73	0.780-64	0.710-50

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

310

3.2 PM_{0.1} and N₁₀ Source Apportionment in California

The UCD/CIT model uses a moving sectional approach to conserve particle number and mass while letting particle radius change due to condensation and evaporation (Kleeman et al., 1997). The method to calculate source contributions to number concentration is performed for each moving section individually. Number is explicitly conserved and correctly apportioned to sources in this algorithm. Each particle source type / moving size bin includes an artificial tracer equal to 1% of the primary particle mass. The mass of this tracer is related to the number of particles by the equation

$$\text{tracer_source_i} * 100 = N_source_i * 3.14159/6 * Dp_bin * \rho_i \text{ (eq1)}$$

where ρ_i is the density of primary particles emitted from source i . This equation can be easily rearranged to solve for N_source_i as a function of tracer_source_i in each size bin. Condensation/evaporation changes the particle diameter as semi-volatile components move on and off the particle but this does not change tracer_source_i or N_source_i . As a result, the moving sectional approach greatly simplifies the source apportionment of particle number compared to other models that use fixed particle size bins with condensation / evaporation transferring material between bins.

Coagulation complicates source apportionment calculations for particle number because coagulation events conserve particle mass but destroy particle number. The model calculations treat the most frequently occurring coagulation events between very small particles and very large particles in a manner analogous to condensation. When two

330

particles coagulate, the mass of the smaller particle is added to the mass of the larger particle. The number concentration of the smaller particle is discarded while the number concentration of the larger particle stays constant. This slightly reduces the accuracy of source apportionment calculations for particle number in the larger size bins because the tracer_source mass in the larger size bin is no longer proportional to the number concentration from that source. This issue is relatively minor since size bins larger than $1\mu\text{m}$ that act as the dominant sink during particle coagulation events typically account for less than 5% of the total number concentration.

Perturbation studies were conducted to test the accuracy of the source apportionment calculations by setting the UFP emissions for on-road gasoline vehicles to zero during the month August 2012. Emissions of gases and emissions of larger particles from on-road vehicles were not changed. The difference between this perturbation simulation vs. the basecase simulation was calculated to estimate the number concentration of particles associated with on-road gasoline vehicles. This “zero-out” concentration was then compared to the standard model source-apportionment calculations in Figure 3 below. The two methods for number source apportionment yield very similar spatial patterns and very similar maximum concentrations of ~ 0.5 kcounts cm^{-3} . The tracer source apportionment method accounts for all particle sizes, which produces slightly higher concentrations than the zero-out method that only considered particles smaller than 100 nm.




Figure 3: Particle number concentrations associated with on-road gasoline vehicles calculated using the zero-out method and the artificial tracer method in August 2012.

355 Many of the spatial patterns measured for airborne particle number concentrations in past
studies have focused on the gradients around roads (see for example (Zhu et al.,
2002a;Zhu et al., 2002b;Zhang et al., 2005;Zhang et al., 2004;Sowlat et al., 2016). These
gradients are impossible to resolve using a regional model with 4km resolution. A
limited set of additional simulations were conducted using the WRF/Chem model
360 configured with Large Eddy Simulation (LES) around Oakland California so that spatial
scales down to 250m could be examined. Maps of the predicted ultrafine particle mass
concentrations for gasoline, diesel, food cooking, wood combustion, and natural gas
combustion particles are shown in Figure 4 below. At 250m resolution, ultrafine
particles from diesel engines peak on major transportation corridors while ultrafine
365 particles from gasoline vehicles are more diffuse reflecting their increased activity on
adjacent surface streets. Ultrafine particles from natural gas combustion are even more
diffuse reflecting contributions from area sources across the region. As the spatial
resolution decreases to 1km and then 4km, the fine details around roadways are
artificially diluted in the larger grid cells. This process shifts the dominant source of
370 ultrafine particles over roadways from diesel engines at 250m resolution to natural gas
combustion at 4km resolution. These simulation results are consistent with
measurements of particle number in the proximity of roadways which show that the
traffic contribution to particle number concentration decays to background levels within
300 m (Zhu et al., 2002a;Zhu et al., 2002b). The measurements made by Zhu et al.
375 indicate that the traffic contribution to regional number concentration cannot be
distinguished from other sources on a regional scale using 4km grid cells which is the
focus of this study.




Figure 4: PM_{0.1} mass concentration associated with on-road diesel, on-road gasoline, and natural gas combustion at 250m, 1km, and 4km resolution over Oakland, California.

380

385 **3.2.1 UCD/CIT PM_{0.1} source contributions compared to Chemical Mass Balance (CMB) results**

A recently completed study measured the composition of PM_{0.1} at four sites in California and calculated source contributions using molecular markers (Xue et al., 2018b). Figures 5 and 6 compare the source contributions to PM_{0.1} OC concentrations predicted by the UCD/CIT model and “measured” using the molecular marker technique at San Pablo, 390 East Oakland, downtown Los Angeles and Fresno during a summer month (August 2015) and a winter month (February 2016). The “others” category in the molecular marker calculation represents unresolved sources, while in the UCD/CIT model “others” represents the sum of non-residential natural gas source combustion, aircraft emissions, and the sources that were not tagged in the current study. In general, the ranking and 395 concentration range of source contributions to PM_{0.1} OC from the molecular marker technique and the UCD/CIT model are consistent. Natural gas dominates PM_{0.1} OC in the summer of 2015 at San Pablo, East Oakland, downtown Los Angeles and Fresno, while wood smoke and aircraft are the major sources of PM_{0.1} OC in Fresno and East Oakland during the winter of 2016. The importance of ultrafine particles from natural gas 400 combustion has not previously been recognized because these particles lack a unique chemical signature, which causes them to be lumped into the “unresolved” category in receptor-based source apportionment studies. The source contribution results for the gasoline, diesel, wood burning, meat cooking and other source categories predicted by the UCD/CIT model and the molecular marker technique illustrated in Figures 5 and 6 405 build confidence in the accuracy of the UFP source predictions in the current study.




Figure 5: Source contribution to PM_{0.1} predicted by the CMB receptor model and the UCD/CIT model at four sites in California in August 2015. CMB results are calculated using 3-day average measurements composited for a full month.




Figure 6: Source contribution to PM_{0.1} predicted by the CMB receptor model and the UCD/CIT model at four sites in California in February 2016. CMB results are calculated using 3-day average measurements composited for a full month.

415 **3.2.2 PM_{0.1} and N₁₀₋₁₀₀₀ Source contributions in California**

Figures 7-9 and 10-12 show the seasonal variation of major source contributions to primary N₁₀ and PM_{0.1}, respectively. The black circles in Figure 7-9 represent the measured N₇₋₁₀₀₀ at four BAAQMD sites in SFBA and six MATES sites in Los Angeles and Riverside counties. Predicted “best-fit” N₁₀ follows the same trends as measured seasonal variations of N₇ at Livermore, Redwood City, Santa Rosa, Huntington Park, Inland Valley, and Rubidoux. The model over predicts N₇ at Anaheim, central Los Angeles, and Compton but overall model performance statistics for N₇ are within the target range for PM_{2.5} applications (see Table 1). Nucleation contributes to N₁₀ at all sites but makes negligible contributions to PM_{0.1} concentrations. Traffic sources including gasoline- and diesel-powered vehicles make significant contributions to PM_{0.1} concentrations at each measurement site depending on proximity to major freeways. Near-roadway effects on ultrafine particle concentrations are not apparent since these locations were chosen to be regional monitors and so they are more than 300 m from the nearest freeway. Predicted contributions from traffic sources are consistent with the molecular marker results illustrated in Figures 5-6. Traffic contributions to regional N₁₀ concentrations more than 300 m away from roadways are even smaller than PM_{0.1} contributions because the size distribution of particles emitted from motor vehicles peaks at 100 – 200 nm (Robert et al., 2007a; Robert et al., 2007b). Wood smoke makes strong contributions to regional PM_{0.1} concentrations in central California during winter but much smaller contributions in the SoCAB because wood burning is not typically used for home heating in this region. Wood burning contributions to N₁₀ are less dominant in central California because the size distribution of particles emitted from wood combustion peaks at 100-300 nm (Kleeman et al., 2008a). The largest primary source of N₁₀ in central California and N₁₀+PM_{0.1} in the SoCAB is natural gas combustion. Industrial processes and power generation that use natural gas do not follow strong seasonal cycles and so the strength of the natural gas source contributions is somewhat constant across seasons subject to variability caused by meteorological conditions.




Figure 7: Seasonal variation of measured N_7 (black circles) and major source contributions to “best-fit” N_{10} at Livermore, Redwood City, San Pablo and Santa Rosa, respectively. Results within each month have daily time resolution.




Figure 8: Seasonal variation of measured N_7 (black circles) and major source contributions to “best-fit” N_{10} at Anaheim, Central LA, and Compton, respectively. Results within each month have daily time resolution.




Figure 9: Seasonal variation of measured N_7 (black circles) and major source contributions to “best-fit” N_{10} at Huntington, Inland-Valley, and Rubidoux, respectively.

455 Results within each month have daily time resolution.




Figure 10: Seasonal variation of major source contributions to $PM_{0.1}$ at Livermore, Redwood City, San Pablo and Santa Rosa, respectively. Results within each month have daily time resolution.





Figure 11: Seasonal variation of major source contributions to PM_{0.1} at Anaheim, Central LA, and Compton, respectively. Results within each month have daily time resolution.



465 Figure 12: Seasonal variation of major source contributions to PM_{0.1} at Huntington, Inland-Valley, and Rubidoux, respectively. Results within each month have daily time resolution.

Figures 13 and Figures 14 show the source contributions to N₁₀ and PM_{0.1}, respectively, averaged over the days shown in Figures 7-9. Aside from nucleation, non-residential

470 natural gas combustion makes the largest predicted primary contribution to N_{10} at all the sites that were evaluated. Traditional sources that were tracked including meat cooking, wood smoke, and mobile (gasoline + diesel) accounted for approximately 5-15% of the predicted N_{10} at the sites selected for study. "Other" sources that were not tagged explicitly in the current study accounted for 5-31% of N_{10} across these sites. Nucleation
475 is a significant source for of N_{10} for both BAAQMD sites and MATES sites where sulfur emissions were highest, with contributions ranging from 24-57%.

The strong N_{10} contribution from natural gas combustion reflects the emitted particle size distribution combined with the ubiquitous use of this fuel in the SFBA and SoCAB regions. The chemical composition and size distribution information for non-residential
480 natural gas combustion emissions used in this study was measured by Hildemann (1991) and Li and Hopke (1993), respectively. Size distributions and volatility were further confirmed during on-going field studies conducted by the current authors (Xue et al., 2018a). The estimated non-residential natural gas combustion particle number and mass size distributions are shown in Figure S1 (left column). Clearly, the majority of particles
485 from non-residential natural gas combustion are typically found in diameters $<0.05 \mu\text{m}$, while particles emitted from other sources such as wood combustion tend to have slightly larger particle diameter (with lower number concentration per unit of emitted mass). These natural gas particles grow through the condensation of SOA once in the atmosphere, but they still contribute strongly to N_{10} concentrations.

490 Figure 14 shows that on-road vehicles (gasoline and diesel combined) are the largest $\text{PM}_{0.1}$ source at Anaheim (39%), central LA (31%), Huntington Park (33%), Inland Valley (39%), and Rubidoux (42%), while natural gas combustion still makes the largest contribution to $\text{PM}_{0.1}$ at other evaluation sites. Contributions from cooking and mobile sources are enhanced in $\text{PM}_{0.1}$ vs. N_{10} , with the cooking source accounting for 11% of
495 $\text{PM}_{0.1}$ at Santa Rosa. The different rankings of source contributions to N_{10} and $\text{PM}_{0.1}$ can be explained by the comparison of particle number-size distribution and particle mass-size distribution for the non-residential natural gas and wood burning sources at the four evaluated sites (Figure S1). Particles emitted from non-residential natural gas combustion and wood burning have number distributions that peak at particle diameters of 0.016-

500 0.025 μm and 0.025-0.04 μm , respectively. Non-residential natural gas combustion and wood burning mass distributions, however, peak at particle diameters of 0.025-0.04 μm and 0.10-0.16 μm , respectively.

Figure 15-17 show diurnal variations of measured N_{7-1000} and predicted “best-fit” N_{10} averaged over days in August and December 2012. These months span the temperature
505 range typically experienced across the year in California. Measured N_{7-1000} diurnal patterns in August generally peak in the afternoon hours between 12-3pm with an optional morning peak around 6am. The main afternoon peak appears to be related to nucleation events while the smaller early-morning peak appears to be related to early morning human activity including natural gas combustion. The predicted “best-fit” N_{10}
510 diurnal variations in August followed the same trends as measurements at six out of ten sites (Livermore, Anaheim, Compton, Huntington Park, Inland Valley, and Rubidoux). The model failed to capture the mid-day nucleation event at Redwood City and Santa Rosa possibly due to missing SO_2 sources in the emissions inventory upwind from these sites. The model overestimated mid-day peak values at Anaheim and central Los
515 Angeles. In December, the measured N_{7-1000} diurnal pattern were more distinctly bimodal with the first peak around 7:00-8:00am and the second peak in the evening at around 8pm. This pattern reflects both the emissions activity and the mixing status of the atmosphere throughout the day. The predicted “best-fit” N_{10} concentration follows this same pattern. Nucleation continues to play a role during winter but does not dominate to
520 the point that it produces a midday peak in N_{10} concentrations. Non-residential natural gas combustion is predicted to be the largest source of N_{10} during morning and evening peaks. The diurnal profiles of non-residential natural gas emissions are included in supplemental information (Figure S2) along with the regional distribution of those
525 emissions (Figure S3). These diurnal variations of the natural gas combustion emissions were obtained directly from the emissions inventory specified by the California Air Resources Board. Industrial natural gas combustion emissions peak during the daytime with lower values at night. Emissions from electricity generation powered by natural gas peak in the morning and evening. Commercial natural gas combustion emissions may either peak in the morning and evening or they may follow a uniform diurnal profile
530 depending on the specific source and location.




Figure 13: The relative source contributions to N₁₀ at Livermore, Redwood City, San Pablo, Santa Rosa, Anaheim, Central LA, Compton, Huntington, Inland-Valley and Rubidoux, respectively. Averaging time included all days shown in Figures 7-9. Values not displayed are $\leq 1\%$.

535




Figure 14: The relative source contributions to PM_{0.1} seasonally averaged at Livermore, Redwood City, San Pablo and Santa Rosa, Anaheim, Central LA, Compton, Huntington, Inland-Valley and Rubidoux, respectively. Averaging time included all days shown in Figures 10-12. Values not displayed are ≤ 1%.






Figure 15: Diurnal variations of measured N_7 and predicted "best-fit" N_{10} averaged for August 2012 (left column) and December 2012 (right column) at Livermore, Redwood City, San Pablo and Santa Rosa.



545

Figure 16: Diurnal variations of measured N_7 and predicted "best-fit" N_{10} averaged for August 2012 (left column) and December 2012 (right column) at Anaheim, Central LA, and Compton.



550

Figure 17: Diurnal variations of measured N_7 and predicted “best-fit” N_{10} averaged for August 2012 (left column) and December 2012 (right column) at Anaheim, Central LA, and Compton.

555 **3.2.3 Regional $N_{10-1000}$ Source contributions in California**

Figure 18 illustrates the predicted number concentration associated with primary emissions (Figures 18a-i) and nucleation (Figure 18j) in southern California averaged

over the months Aug-Dec 2012. Figure 18g shows that primary aircraft emissions in the plume downwind of the Los Angeles International Airport (LAX) are predicted to
560 account for 8 kcounts cm^{-3} and Figure 18j shows that nucleation of aircraft emissions in the LAX plume are predicted to account for 45 kcounts cm^{-3} yielding a total number concentration associated with LAX aircraft of approximately 53 kcounts cm^{-3} . Hudda et al. (2014) found that particle number concentrations increased by a factor of four to eight downwind of LAX based on measurements in June-July 2013. Total ground-level
565 number concentrations in the LAX plume reached 60-70 kcounts cm^{-3} . Given the 4km spatial resolution of the model calculations used in the current study, the predictions and measurements of particle number concentration downwind of LAX are consistent with one another.

It is noteworthy that military airbases in Figure 18g have significantly higher particle
570 number concentrations due to their use of aviation fuel with higher sulfur content, but nucleation plumes are not present downwind of these locations (Figure 18j). Particles emitted from military aircraft are represented as primary emissions in the current model calculations. Future measurements should compare particle number concentrations downwind of civilian and military airports to fully evaluate the impact of aviation fuel
575 sulfur content on ambient ultrafine particle concentrations.




Figure 18. Spatial distribution of particle number from major sources in Southern California (unit: kcount cm^{-3}).

580 Figure 19 illustrates the predicted particle number concentrations associated with primary sources and nucleation in northern California. The relative importance of sources and the prediction of nucleation downwind of major sulfur emissions are consistent in northern and southern California. Natural gas combustion is a notable strong source of ultrafine particles in both regions due to the widespread use of this fuel in numerous residential, commercial, and industrial applications. In many cases, the natural gas combustion 585 particles contribute strongly to the “urban background” concentrations over most California cities without the formation of individual plumes such as those found downwind of LAX. Future measurements could correlate ambient particle number concentrations and natural gas utilization across multiple cities to evaluate whether natural gas combustion is a significant source of particle number concentration.

590





Figure 19. Spatial distribution of particle number from major sources in Northern California (unit: kcount cm^{-3}).


The concentrations of nucleated particles in August, October, and December are shown in Figure 20 (Southern California) and Figure 21 (Northern California) below. Nucleation events occur in the regions where sulfur emissions are highest (typically airports, shipping ports and refining facilities). Concentrations of nucleated particles are higher in October and December than in August because colder temperatures increase nucleation rates if the precursor H_2SO_4 and NH_3 concentrations are relatively constant. A significant fraction of the H_2SO_4 in the current simulation is produced by the fast conversion of gas-phase SO_3 emissions to H_2SO_4 in the exhaust plume near the emissions source. SO_3 conversion does not depend on the presence of oxidants in the atmosphere and so the higher oxidant concentrations in the summer do not dominate the seasonal nucleation pattern.

Once H_2SO_4 forms in the exhaust plumes, it either condenses onto existing particles formed from lower volatility compounds in the plume, or it mixes with NH_3 in the background air and nucleates. This process is captured by dilution source sampling measurements that allow for a few minutes of aging time and so the size-resolved emissions profiles for many sources already account for the effects of nucleation within the “near-field” exhaust plume (within a few 10’s of meters after emission). SO_3 emissions from reciprocating internal combustion engines were therefore set to zero to avoid double counting the new particle formation downwind of these sources in the current study. Regular SO_2 emissions from these sources were not modified. Emissions from aircraft jet engines have high exit velocity which promotes rapid mixing with background air. SO_3 emissions were left at their nominal levels (3-4% of total SO_x) for jet engine aircraft in the current study. The consequence of these model treatments is that predicted concentrations of nucleated particles are highest downwind of LAX, which agrees with measurements of ambient particle number concentrations (Hudda et al., 2014).



620

Figure 20: Seasonal variation of nucleated particle concentrations in Southern California. Units are kcount cm⁻³.



625

Figure 21: Seasonal variation of nucleated particle concentrations in Northern California. Units are kcount cm⁻³.

4. Discussion

630 Previous researchers have used Positive Matrix Factorization (PMF) to calculate source contributions to N_7 (Sowlat et al., 2016; Morawska et al., 2008; Gu et al., 2011; Ogulei et al., 2007; Kasumba et al., 2009; Wang et al., 2013; Yue et al., 2008; Friend et al., 2013). The dominant factors resolved by these studies have been traffic, urban background, secondary aerosol, wood burning and nucleation (Sowlat et al., 2016; Morawska et al., 2008; Gu et al., 2011; Ogulei et al., 2007; Kasumba et al., 2009; Wang et al., 2013; Yue et al., 2008; Friend et al., 2013). Particles from natural gas combustion were not separately
635 identified by PMF because they do not contain a unique chemical tracer. It is very likely that natural gas combustion particles are artificially lumped into another source (e.g. traffic) or part of the “urban background” signal identified in previous studies. Natural gas combustion is used extensively in California for electric power, industrial,
640 commercial and residential use (Table S6), and so it seems plausible that this source contributes to ambient UFP concentrations.

The current UFP predictions rely on source profile measurements for wood burning, food cooking, mobile sources, and non-residential natural gas combustion (Cooper, 1989; Harley et al., 1992; Hildemann et al., 1991a; Hildemann et al., 1991b; Houck and L.
645 C., 1989; Kleeman et al., 2008b; Kleeman et al., 2000; Robert et al., 2007b; Robert et al., 2007a; Schauer et al., 1999b, a, 2001, 2002b, a; Taback, 1979). All of these size distributions were measured using appropriate instruments and methods by knowledgeable researchers, but some of these past studies were conducted more than a decade ago. Size distribution information for vehicles, natural gas, etc. have been added
650 to the supplemental information (Figure S4). Changes in fuel composition and emissions control technology in the interim years may have altered the emitted size distributions. New measurements of particle size distributions emitted from natural gas and biomethane combustion were made in parallel with the current project to confirm the source profile measurements from past studies (Xue et al., 2018a). The results of these measurements
655 are consistent with previous size distribution results (Li and Hopke, 1993).

California has tighter air pollution standards than many other regions in the United States due to the severe air quality problems that have historically occurred in the state.

California therefore has a unique combination of fuels and emissions control technology that may affect the mixture of sources that contribute to atmospheric ultrafine particle concentrations. Venecek et al. (2018) recently used the UCD/CIT air quality model with the 2011 National Emissions inventory to calculate source contributions to $PM_{0.1}$ in 39 major cities across the United States during peak summer photochemical smog episodes in the year 2010. The findings from this study show that natural gas combustion is a major source of ultrafine particles in the regional atmosphere over urban areas across the United States. The public health questions associated with ultrafine particles emitted by natural gas combustion have wide-ranging implications. Similar levels of ultrafine particle concentrations will likely occur in other regions across the world that extensively use natural gas as a fuel source, although other sources of ultrafine particles may also make strong contributions depending on the total mix of fuels in each region.

Recent theories suggest that primary particulate matter composed of semi-volatile organic compounds may evaporate after release to the atmosphere, which may reduce ambient N_x . Measurements conducted in parallel with the current study confirmed that particles emitted from natural gas combustion in home appliances partially evaporated when diluted by a factor of 25 in clean air, but particles emitted from reciprocating engines did not evaporate under the same conditions (Xue et al., 2018a). Future work should verify the accuracy of the size and composition distributions for all natural gas combustion sources given their apparent importance for predicted N_x .

Evidence from both toxicology and epidemiology will be required to assess the effect of UFPs on public health. It is essential to identify and quantify UFP sources based on both mass ($PM_{0.1}$) and N_x during this process (Friend et al., 2013). An accurate comparison of both $PM_{0.1}$ and N_x exposure could lay the groundwork for specific assessment of health effects of UFPs and potentially more efficient control strategies for PM emission from major sources (Yue et al., 2008). Ideally, spatial exposure patterns for N_x , $PM_{0.1}$, and $PM_{2.5}$ will be sufficiently unique to separate their individual effects in epidemiological studies. Regression statistics for different metrics were calculated by using all grid cells in the model domain of the current study. The correlations between the various particle metrics were: $R^2(PM_{2.5}$ vs. $N_{10})=0.35$, $R^2(PM_{2.5}$ vs. $PM_{0.1})=0.63$, $R^2(PM_{0.1}$ vs. $N_{10})=0.75$.

It seems likely that future epidemiological studies will be able to differentiate between the effects of PM_{2.5} and N_x based on the low R² value. The potential for comparisons between PM_{2.5} and PM_{0.1} is less clear cut, but previous work helps understand what may be possible. Ostro et al. (2015) compared the associations between IHD mortality and PM_{2.5} vs. PM_{0.1} in the California Teachers Study (CTS) cohort. Associations between IHD mortality and the sum of PM_{2.5} mass (p-value=0.001) were stronger than associations between IHD mortality and the sum of PM_{0.1} mass (p-value=0.01) but individual components of mass (EC, OC, Cu, etc) all had stronger associations with IHD mortality in the PM_{0.1} size fraction than the PM_{2.5} size fraction.

The current study focuses on outdoor exposure to UFPs that may be useful in future epidemiological studies. Indoor or in-vehicle exposure to UFPs can also be significant (Wallace and Ott, 2011; Rim et al., 2010; Bhangar et al., 2011; Weichenthal et al., 2015; Fruin et al., 2008) but characterizing these micro-environments is beyond the scope of the current manuscript.

5. Conclusions

The UCD/CIT regional chemical transport model has been updated with a nucleation algorithm and combined with the existing size-resolved source profiles of particulate matter emissions to predict regional source contributions to airborne particle number concentration (N₁₀) and airborne particulate ultrafine mass (PM_{0.1}). Predicted 24-hour average N₁₀ follows the same trend as measured N₇ at ten sites across California in summer (Aug) and winter (Dec). Predicted diurnal variation of N₁₀ follows the same trend as measured concentrations at the majority of the evaluation sites in August and December, but the results suggest that further refinement is needed for both primary emissions and nucleation algorithms. Predicted PM_{0.1} source contributions follow the same trends as PM_{0.1} source contributions measured in a molecular marker study at four sites across California in summer (August) and winter (December) months. Natural gas combustion is the largest primary source of regional N₁₀ at all locations outside of the immediate vicinity of other major combustion sources. Nucleation contributed strongly to particle number during both the summer and winter months. Traffic sources

contributed to N_{10} but did not dominate over regions more than 300 m away from freeways. Combustion sources such as wood burning, food cooking, and mobile sources made stronger contributions to $PM_{0.1}$ at heavily urbanized locations. Wood burning for
720 home heating had strong seasonal patterns with peak concentrations in winter while other sources contributed more consistently throughout the seasons. Nucleation made a negligible contribution to $PM_{0.1}$ over the urban areas at the focus of the current study.

The current study identifies natural gas combustion as an important source of ultrafine particle number and mass concentrations in urban regions throughout California. The
725 health implications of these natural gas combustion particles should be investigated in future epidemiology studies.

Data Availability: All of the $PM_{0.1}$ and N_x outdoor exposure fields produced in the current study are available free of charge at <http://faculty.engineering.ucdavis.edu/kleeman/> which provides a link to the most recent version of the dataset (currently
730 http://webwolf.engr.ucdavis.edu/data/soa_v2/monthly_avg2). Model source code and model input files are available to collaborators via direct email to the corresponding author at mikleeman@ucdavis.edu.

Acknowledgements: This research was supported by the Bay Area Air Quality
735 Management District under project #2013.218, the Coordinating Research Council under project #A-96, and the California Air Resources Board under project #14-314. None of the project sponsors nor any person acting on their behalf: (1) makes any warranty, express or implied, with respect to the use of any information, apparatus, method, or process disclosed in this report, or (2) assumes any liabilities with respect to use, or
740 damages resulting from the use or inability to use, any information, apparatus, method, or process disclosed in this report.

Reference

- Anttila, T., and Kerminen, V. M.: Condensational growth of atmospheric nuclei by organic vapours, *J Aerosol Sci*, 34, 41-61, 10.1016/s0021-8502(02)00155-6, 2003.
- 745 Bhangar, S., Mullen, N. A., Hering, S. V., Kreisberg, N. M., and Nazaroff, W. W.: Ultrafine particle concentrations and exposures in seven residences in northern California, *Indoor Air*, 21, 132-144, 10.1111/j.1600-0668.2010.00689.x, 2011.
- Boylan, J. W., and Russell, A. G.: PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models, *Atmos Environ*, 40, 4946-4959, 10.1016/j.atmosenv.2005.09.087, 2006.
- 750 Brunekreef, B., and Forsberg, B.: Epidemiological evidence of effects of coarse airborne particles on health, *Eur Respir J*, 26, 309-318, 2005.
- Chen, J. J., Ying, Q., and Kleeman, M. J.: Source apportionment of wintertime secondary organic aerosol during the California regional PM10/PM2.5 air quality study, *Atmos Environ*, 44, 1331-1340, 2010.
- 755 Cooper, J. A. E. A.: Dinal AppendixV-G, PM10 source composition
- library for the South Coast Air Basin, South Coast Air Quality Management District, Diamond Bar, California, 1989.
- 760 Dockery, D. W., and Stone, P. H.: Cardiovascular risks from fine particulate air pollution, *New Engl J Med*, 356, 511-513, 2007.
- Elleman, R. A., and Covert, D. S.: Aerosol size distribution modeling with the Community Multiscale Air Quality modeling system in the Pacific Northwest: 2. Parameterizations for ternary nucleation and nucleation mode processes, *J Geophys Res-Atmos*, 114, 2009a.
- 765 Elleman, R. A., and Covert, D. S.: Aerosol size distribution modeling with the Community Multiscale Air Quality modeling system in the Pacific Northwest: 1. Model comparison to observations, *J Geophys Res-Atmos*, 114, 2009b.
- Fann, N., Lamson, A. D., Anenberg, S. C., Wesson, K., Risley, D., and Hubbell, B. J.: Estimating the National Public Health Burden Associated with Exposure to Ambient PM2.5 and Ozone, *Risk Anal*, 32, 81-95, 2012.
- 770 Friend, A. J., Ayoko, G. A., Jager, D., Wust, M., Jayaratne, E. R., Jamriska, M., and Morawska, L.: Sources of ultrafine particles and chemical species along a traffic corridor: comparison of the results from two receptor models, *Environ Chem*, 10, 54-63, 2013.
- Fruin, S., Westerdahl, D., Sax, T., Sioutas, C., and Fine, P. M.: Measurements and predictors of on-road ultrafine particle concentrations and associated pollutants in Los Angeles, *Atmos Environ*, 42, 207-219, 10.1016/j.atmosenv.2007.09.057, 2008.
- 775 Gauderman, W. J., Urman, R., Avol, E., Berhane, K., McConnell, R., Rappaport, E., Chang, R., Lurmann, F., and Gilliland, F.: Association of Improved Air Quality with Lung Development in Children, *New Engl J Med*, 372, 905-913, 2015.
- 780 Gu, J. W., Pitz, M., Schnelle-Kreis, J., Diemer, J., Reller, A., Zimmermann, R., Soentgen, J., Stoelzel, M., Wichmann, H. E., Peters, A., and Cyrys, J.: Source apportionment of ambient particles: Comparison of positive matrix factorization analysis applied to particle size distribution and chemical composition data, *Atmos Environ*, 45, 1849-1857, 2011.
- 785 Harley, R. A., Hannigan, M. P., and Cass, G. R.: Respeciation of Organic Gas Emissions and the Detection of Excess Unburned Gasoline in the Atmosphere, *Environmental science & technology*, 26, 2395-2408, DOI 10.1021/es00036a010, 1992.

- Held, T., Ying, Q., Kaduwela, A., and Kleeman, M.: Modeling particulate matter in the San Joaquin Valley with a source-oriented externally mixed three-dimensional photochemical grid model, *Atmos Environ*, 38, 3689-3711, 2004.
- 790 Held, T., Ying, Q., Kleeman, M. J., Schauer, J. J., and Fraser, M. P.: A comparison of the UCD/CIT air quality model and the CMB source-receptor model for primary airborne particulate matter, *Atmos Environ*, 39, 2281-2297, 2005.
- Hildemann, L. M., Markowski, G. R., and Cass, G. R.: Chemical-Composition of Emissions from Urban Sources of Fine Organic Aerosol, *Environmental science & technology*, 25, 744-759, DOI 10.1021/es00016a021, 1991a.
- 795 Hildemann, L. M., Markowski, G. R., Jones, M. C., and Cass, G. R.: Submicrometer Aerosol Mass Distributions of Emissions from Boilers, Fireplaces, Automobiles, Diesel Trucks, and Meat-Cooking Operations, *Aerosol Sci Tech*, 14, 138-152, Doi 10.1080/02786829108959478, 1991b.
- 800 Hildemann, L. M., G.R. Markowski, and G.R. Cass: Chemical composition of emissions from urban sources of fine organic aerosol, *Environmental Science and Technology*, 25, 744-759, 1991.
- Hixson, M., Mahmud, A., Hu, J. L., Bai, S., Niemeier, D. A., Handy, S. L., Gao, S. Y., Lund, J. R., Sullivan, D. C., and Kleeman, M. J.: Influence of regional development policies and clean technology adoption on future air pollution exposure, *Atmos Environ*, 44, 552-562, 2010.
- 805 Hixson, M., Mahmud, A., Hu, J., and Kleeman, M. J.: Resolving the interactions between population density and air pollution emissions controls in the San Joaquin Valley, USA, *Journal of the Air & Waste Management Association*, 62, 566-575, 10.1080/10962247.2012.663325, 2012.
- Houck, J. E., Chow, J. C., Watson, J. G., Simons, C. A., Prichett, and L. C., G., J. M., Frazier, C. A.: Determination of particle size distribution and chemical composition of particulate matter from selected sources in California, California Air Resources Board, OMNI Environment Service Incorporate, Desert Research Institute, Beaverton, Oregon, 1989.
- 810 Hu, J., Howard, C. J., Mitloehner, F., Green, P. G., and Kleeman, M. J.: Mobile source and livestock feed contributions to regional ozone formation in Central California, *Environmental science & technology*, 46, 2781-2789, 10.1021/es203369p, 2012.
- Hu, J., Zhang, H., Chen, S., Ying, Q., Wiedinmyer, C., Vandenberghe, F., and Kleeman, M. J.: Identifying PM2.5 and PM0.1 sources for epidemiological studies in California, *Environmental science & technology*, 48, 4980-4990, 10.1021/es404810z, 2014a.
- Hu, J., Zhang, H., Chen, S. H., Wiedinmyer, C., Vandenberghe, F., Ying, Q., and Kleeman, M. J.: Predicting primary PM2.5 and PM0.1 trace composition for epidemiological studies in California, *Environmental science & technology*, 48, 4971-4979, 10.1021/es404809j, 2014b.
- 820 Hu, J., Zhang, H., Ying, Q., Chen, S. H., Vandenberghe, F., and Kleeman, M. J.: Long-term particulate matter modeling for health effect studies in California – Part 1: Model performance on temporal and spatial variations, *Atmospheric Chemistry and Physics*, 15, 3445-3461, 10.5194/acp-15-3445-2015, 2015.
- Hu, J. L., Jathar, S., Zhang, H. L., Ying, Q., Chen, S. H., Cappa, C. D., and Kleeman, M. J.: Long-term particulate matter modeling for health effect studies in California - Part 2: Concentrations and sources of ultrafine organic aerosols, *Atmospheric Chemistry and Physics*, 17, 5379-5391, 2017.
- Hudda, N., Gould, T., Hartin, K., Larson, T. V., and Fruin, S. A.: Emissions from an International Airport Increase Particle Number Concentrations 4-fold at 10 km Downwind, *Environmental science & technology*, 48, 6628-6635, 10.1021/es5001566, 2014.
- 830 Jung, J. G., Pandis, S. N., and Adams, P. J.: Evaluation of nucleation theories in a sulfur-rich environment, *Aerosol Sci Tech*, 42, 495-504, 2008.
- Jung, J. G., Fountoukis, C., Adams, P. J., and Pandis, S. N.: Simulation of in situ ultrafine particle formation in the eastern United States using PMCAMx-UF, *J Geophys Res-Atmos*, 115, 2010.

- 835 Kasumba, J., Hopke, P. K., Chalupa, D. C., and Utell, M. J.: Comparison of sources of submicron particle number concentrations measured at two sites in Rochester, NY, *Sci Total Environ*, 407, 5071-5084, 2009.
- Kerminen, V. M., and Kulmala, M.: Analytical formulae connecting the "real" and the "apparent" nucleation rate and the nuclei number concentration for atmospheric nucleation events, *J Aerosol Sci*, 33, 609-622, 2002.
- 840 Kleeman, M. J., Cass, G. R., and Eldering, A.: Modeling the airborne particle complex as a source-oriented external mixture, *J Geophys Res-Atmos*, 102, 21355-21372, 1997.
- Kleeman, M. J., and Cass, G. R.: Source contributions to the size and composition distribution of urban particulate air pollution, *Atmos Environ*, 32, 2803-2816, 1998.
- Kleeman, M. J., Schauer, J. J., and Cass, G. R.: Size and composition distribution of fine particulate matter emitted from wood burning, meat charbroiling, and cigarettes, *Environmental science & technology*, 33, 3516-3523, 10.1021/es981277q, 1999.
- 845 Kleeman, M. J., Schauer, J. J., and Cass, G. R.: Size and composition distribution of fine particulate matter emitted from motor vehicles, *Environmental science & technology*, 34, 1132-1142, 10.1021/es981276y, 2000.
- 850 Kleeman, M. J., and Cass, G. R.: A 3D Eulerian source-oriented model for an externally mixed aerosol, *Environmental science & technology*, 35, 4834-4848, 2001.
- Kleeman, M. J., Ying, Q., Lu, J., Mysliwiec, M. J., Griffin, R. J., Chen, J. J., and Clegg, S.: Source apportionment of secondary organic aerosol during a severe photochemical smog episode, *Atmos Environ*, 41, 576-591, 2007.
- 855 Kleeman, M. J., Robert, M. A., Riddle, S. G., Fine, P. M., Hays, M. D., Schauer, J. J., and Hannigan, M. P.: Size distribution of trace organic species emitted from biomass combustion and meat charbroiling (vol 42, pg 3059, 2008), *Atmos Environ*, 42, 6152-6154, 2008a.
- Kleeman, M. J., Robert, M. A., Riddle, S. G., Fine, P. M., Hays, M. D., Schauer, J. J., and Hannigan, M. P.: Size distribution of trace organic species emitted from biomass combustion and meat charbroiling, *Atmos Environ*, 42, 3059-3075, 10.1016/j.atmosenv.2007.12.044, 2008b.
- 860 Kuwayama, T., Collier, S., Forestieri, S., Brady, J. M., Bertram, T. H., Cappa, C. D., Zhang, Q., and Kleeman, M. J.: Volatility of Primary Organic Aerosol Emitted from Light Duty Gasoline Vehicles, *Environmental Science & Technology*, 49, 1569-1577, 2015.
- Laurent, O., Hu, J., Li, L., Kleeman, M. J., Bartell, S. M., Cockburn, M., Escobedo, L., and Wu, J.: A Statewide Nested Case-Control Study of Preterm Birth and Air Pollution by Source and Composition: California, 2001-2008, *Environ Health Perspect*, 124, 1479-1486, 10.1289/ehp.1510133, 2016.
- 865 Li, N., Sioutas, C., Cho, A., Schmitz, D., Misra, C., Sempf, J., Wang, M. Y., Oberley, T., Froines, J., and Nel, A.: Ultrafine particulate pollutants induce oxidative stress and mitochondrial damage, *Environ Health Persp*, 111, 455-460, 2003.
- 870 Li, W., and Hopke, P. K.: Initial Size Distributions and Hygroscopicity of Indoor Combustion Aerosol Particles, *Aerosol Sci Tech*, 19, 305-316, 10.1080/02786829308959638, 1993.
- Lupascu, A., Easter, R., Zaveri, R., Shrivastava, M., Pekour, M., Tomlinson, J., Yang, Q., Matsui, H., Hodzic, A., Zhang, Q., and Fast, J. D.: Modeling particle nucleation and growth over northern California during the 2010 CARES campaign, *Atmospheric Chemistry and Physics*, 15, 12283-12313, 2015.
- 875 Mahmud, A., Hixson, M., Hu, J., Zhao, Z., Chen, S. H., Kleeman, M. J.: Climate impact on airborne particulate matter concentrations in California using seven year analysis periods, *Atmospheric Chemistry and Physics*, 10, 11097-11114, 10.5194/acp-10-11097-2010, 2010.

- 880 May, A. A., Levin, E. J. T., Hennigan, C. J., Riipinen, I., Lee, T., Collett, J. L., Jimenez, J. L., Kreidenweis, S. M., and Robinson, A. L.: Gas-particle partitioning of primary organic aerosol emissions: 3. Biomass burning, *J Geophys Res-Atmos*, **118**, 11327-11338, 2013a.
- May, A. A., Presto, A. A., Hennigan, C. J., Nguyen, N. T., Gordon, T. D., and Robinson, A. L.: Gas-particle partitioning of primary organic aerosol emissions: (1) Gasoline vehicle exhaust, 885 *Atmospheric Environment*, **77**, 128-139, 2013b.
- Mazaheri, M., Johnson, G. R., and Morawska, L.: Particle and Gaseous Emissions from Commercial Aircraft at Each Stage of the Landing and Takeoff Cycle, *Environmental science & technology*, **43**, 441-446, 10.1021/es8013985, 2009.
- 890 Miller, K. A., Siscovick, D. S., Sheppard, L., Shepherd, K., Sullivan, J. H., Anderson, G. L., and Kaufman, J. D.: Long-term exposure to air pollution and incidence of cardiovascular events in women, *New Engl J Med*, **356**, 447-458, 2007.
- Montagne, D. R., Hoek, G., Klompmaker, J. O., Wang, M., Meliefste, K., and Brunekreef, B.: Land Use Regression Models for Ultrafine Particles and Black Carbon Based on Short-Term Monitoring Predict Past Spatial Variation, *Environmental science & technology*, **49**, 8712-8720, 2015.
- 895 Morawska, L., Ristovski, Z., Jayaratne, E. R., Keogh, D. U., and Ling, X.: Ambient nano and ultrafine particles from motor vehicle emissions: Characteristics, ambient processing and implications on human exposure, *Atmos Environ*, **42**, 8113-8138, 2008.
- Mysliwiec, M. J., and Kleeman, M. J.: Source apportionment of secondary airborne particulate matter in a polluted atmosphere, *Environmental science & technology*, **36**, 5376-5384, 2002.
- 900 Napari, I., Noppel, M., Vehkamäki, H., and Kulmala, M.: Parametrization of ternary nucleation rates for H₂SO₄-NH₃-H₂O vapors, *J Geophys Res-Atmos*, **107**, 2002.
- Nel, A., Xia, T., Madler, L., and Li, N.: Toxic potential of materials at the nanolevel, *Science*, **311**, 622-627, 2006.
- 905 Oberdorster, G., Sharp, Z., Atudorei, V., Elder, A., Gelein, R., Lunts, A., Kreyling, W., and Cox, C.: Extrapulmonary translocation of ultrafine carbon particles following whole-body inhalation exposure of rats, *J Toxicol Env Heal A*, **65**, 1531-1543, 2002.
- Ogulei, D., Hopke, P. K., Chalupa, D. C., and Utell, M. J.: Modeling source contributions to submicron particle number concentrations measured in Rochester, New York, *Aerosol Sci Tech*, **41**, 179-201, 2007.
- 910 Ostro, B., Broadwin, R., Green, S., Feng, W. Y., and Lipsett, M.: Fine particulate air pollution and mortality in nine California counties: Results from CALFINE, *Environ Health Persp*, **114**, 29-33, 2006.
- Ostro, B., Lipsett, M., Reynolds, P., Goldberg, D., Hertz, A., Garcia, C., Henderson, K. D., and Bernstein, L.: Long-Term Exposure to Constituents of Fine Particulate Air Pollution and Mortality: 915 Results from the California Teachers Study, *Environ Health Persp*, **118**, 363-369, 2010.
- Ostro, B., Hu, J., Goldberg, D., Reynolds, P., Hertz, A., Bernstein, L., and Kleeman, M. J.: Associations of mortality with long-term exposures to fine and ultrafine particles, species and sources: results from the California Teachers Study Cohort, *Environ Health Perspect*, **123**, 549-556, 10.1289/ehp.1408565, 2015.
- 920 Otte, T. L.: The impact of nudging in the meteorological model for retrospective air quality simulations. Part II: Evaluating collocated meteorological and air quality observations, *J Appl Meteorol Clim*, **47**, 1868-1887, 10.1175/2007jamc1791.1, 2008a.
- Otte, T. L.: The impact of nudging in the meteorological model for retrospective air quality simulations. Part I: Evaluation against national observation networks, *J Appl Meteorol Clim*, **47**, 925 1853-1867, 10.1175/2007jamc1790.1, 2008b.

- Pope, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and Thurston, G. D.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *Jama-J Am Med Assoc*, 287, 1132-1141, 2002.
- 930 Pope, C. A., Burnett, R. T., Thurston, G. D., Thun, M. J., Calle, E. E., Krewski, D., and Godleski, J. J.: Cardiovascular mortality and long-term exposure to particulate air pollution - Epidemiological evidence of general pathophysiological pathways of disease, *Circulation*, 109, 71-77, 2004.
- Pope, C. A., Ezzati, M., and Dockery, D. W.: Fine-Particulate Air Pollution and Life Expectancy in the United States., *New Engl J Med*, 360, 376-386, 2009.
- 935 Rasmussen, D. J., Hu, J. L., Mahmud, A., and Kleeman, M. J.: The Ozone-Climate Penalty: Past, Present, and Future, *Environmental science & technology*, 47, 14258-14266, 2013.
- Rim, D. H., Wallace, L., and Persily, A.: Infiltration of Outdoor Ultrafine Particles into a Test House, *Environmental science & technology*, 44, 5908-5913, 2010.
- 940 Robert, M. A., Kleeman, M. J., and Jakober, C. A.: Size and composition distributions of particulate matter emissions: Part 2- Heavy-duty diesel vehicles, *Journal of the Air & Waste Management Association*, 57, 1429-1438, 2007a.
- Robert, M. A., VanBergen, S., Kleeman, M. J., and Jakober, C. A.: Size and composition distributions of particulate matter emissions: Part 1 - Light-duty gasoline vehicles, *Journal of the Air & Waste Management Association*, 57, 1414-1428, 2007b.
- 945 Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 2. C-1 through C-30 organic compounds from medium duty diesel trucks, *Environmental science & technology*, 33, 1578-1587, DOI 10.1021/es980081n, 1999a.
- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 1. C-1 through C-29 organic compounds from meat charbroiling, *Environmental science & technology*, 33, 1566-1577, DOI 10.1021/es980076j, 1999b.
- 950 Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 3. C-1-C-29 organic compounds from fireplace combustion of wood, *Environmental science & technology*, 35, 1716-1728, DOI 10.1021/es001331e, 2001.
- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 5. C-1-C-32 organic compounds from gasoline-powered motor vehicles, 955 *Environmental science & technology*, 36, 1169-1180, 10.1021/es0108077, 2002a.
- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 4. C-1-C-27 organic compounds from cooking with seed oils, *Environmental science & technology*, 36, 567-575, 10.1021/es002053m, 2002b.
- 960 Shet, C. S., Cholevari, M. R., and Veeravalli, S. V.: Eulerian spatial and temporal autocorrelations: assessment of Taylor's hypothesis and a model, *Journal of Turbulence*, 1-15, 10.1080/14685248.2017.1357823, 2017.
- Sioutas, C., Delfino, R. J., and Singh, M.: Exposure assessment for atmospheric ultrafine particles (UFPs) and implications in epidemiologic research, *Environ Health Persp*, 113, 947-955, 2005.
- 965 Sowlat, M. H., Hasheminassab, S., and Sioutas, C.: Source apportionment of ambient particle number concentrations in central Los Angeles using positive matrix factorization (PMF), *Atmospheric Chemistry and Physics*, 16, 4849-4866, 10.5194/acp-16-4849-2016, 2016.
- Taback, H. J., Brienza, A. R., Macko, J., and Brunetz, N.: Fine particle emissions from stationary and miscellaneous sources in the South Coast Air Basin. KVB Report 5806-783., KVB Incorporated, Tustin, California, 1979.
- 970 Trostl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer, A. K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T.,

975 Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kurten, A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Mohler, O., Nieminen, T., Onnela, A., Petaja, T., Piel, F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipila, M., Smith, J. N., Steiner, G., Tome, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P. L., Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop, D. R., Donahue, N. M., and Baltensperger, U.: The role of low-volatility organic
 980 compounds in initial particle growth in the atmosphere, *Nature*, 533, 527-+, 10.1038/nature18271, 2016.

Venecek, M., Yu, X., and Kleeman, M.: Ultrafine Particulate Matter Source Contributions across the Continental United States, *Atmospheric Chemistry and Physics*, submitted for review, 2018.

985 Wallace, L., and Ott, W.: Personal exposure to ultrafine particles, *J Expo Sci Environ Epidemiol*, 21, 20-30, 10.1038/jes.2009.59, 2011.

Wang, Z. B., Hu, M., Wu, Z. J., Yue, D. L., He, L. Y., Huang, X. F., Liu, X. G., and Wiedensohler, A.: Long-term measurements of particle number size distributions and the relationships with air mass history and source apportionment in the summer of Beijing, *Atmospheric Chemistry and Physics*, 13, 10159-10170, 2013.

990 Watson, J. G., Chow, J. C., Sodeman, D. A., Lowenthal, D. H., Chang, M. C. O., Park, K., and Wang, X.: Comparison of four scanning mobility particle sizers at the Fresno Supersite, *Particuology*, 9, 204-209, 10.1016/j.partic.2011.03.002, 2011.

Weichenthal, S., Van Ryswyk, K., Kulka, R., Sun, L., Wallace, L., and Joseph, L.: In-vehicle exposures to particulate air pollution in Canadian metropolitan areas: the urban transportation
 995 exposure study, *Environmental science & technology*, 49, 597-605, 10.1021/es504043a, 2015.

Westervelt, D. M., Pierce, J. R., Riipinen, I., Trivitayanurak, W., Hamed, A., Kulmala, M., Laaksonen, A., Decesari, S., and Adams, P. J.: Formation and growth of nucleated particles into cloud condensation nuclei: model-measurement comparison, *Atmospheric Chemistry and Physics*, 13, 7645-7663, 2013.

1000 Xue, J., Li, Y., Peppers, J., Wan, C., Kado, N., Green, P. G., Young, T., and Kleeman, M.: Ultrafine particle emissions from natural gas, biogas and biomethane combustion, *Environmental Science and Technology*, in review, 2018a.

Xue, J., Xue, W., Sowlat, M., Sioutas, C., Lilinco, A., Hasson, A., and Kleeman, M.: Annual trends in ultrafine particulate matter (PM_{0.1}) source contributions in polluted California cities,
 1005 *Environmental Science and Technology*, in preparation, 2018b.

Ying, Q., and Kleeman, M. J.: Source contributions to the regional distribution of secondary particulate matter in California, *Atmos Environ*, 40, 736-752, 2006.

Ying, Q., Lu, J., Allen, P., Livingstone, P., Kaduwela, A., and Kleeman, M.: Modeling air quality during the California Regional PM₁₀/PM_{2.5} Air Quality Study (CRPAQS) using the UCD/CIT
 1010 source-oriented air quality model - Part I. Base case model results, *Atmos Environ*, 42, 8954-8966, 2008a.

Ying, Q., Lu, J., Kaduwela, A., and Kleeman, M.: Modeling air quality during the California Regional PM₁₀/PM_{2.5} Air Quality Study (CPRAQS) using the UCD/CIT Source Oriented Air
 1015 Quality Model - Part II. Regional source apportionment of primary airborne particulate matter, *Atmos Environ*, 42, 8967-8978, 2008b.

Yue, W., Stolzel, M., Cyrys, J., Pitz, M., Heinrich, J., Kreyling, W. G., Wichmann, H. E., Peters, A., Wang, S., and Hopke, P. K.: Source apportionment of ambient fine particle size distribution using positive matrix factorization in Erfurt, Germany, *Sci Total Environ*, 398, 133-144, 2008.

1020 Zhang, H. L., and Ying, Q.: Source apportionment of airborne particulate matter in Southeast Texas using a source-oriented 3D air quality model, *Atmos Environ*, 44, 3547-3557, 2010.

- Zhang, K. M., Wexler, A. S., Zhu, Y. F., Hinds, W. C., and Sioutas, C.: Evolution of particle number distribution near roadways. Part II: the 'road-to-ambient' process, *Atmos Environ*, 38, 6655-6665, 10.1016/j.atmosenv.2004.06.044, 2004.
- 1025 Zhang, K. M., Wexler, A. S., Niemeier, D. A., Zhu, Y. F., Hinds, W. C., and Sioutas, C.: Evolution of particle number distribution near roadways. Part III: Traffic analysis and on-road size resolved particulate emission factors, *Atmos Environ*, 39, 4155-4166, 10.1016/j.atmosenv.2005.04.003, 2005.
- Zhang, Y., Liu, P., Liu, X. H., Jacobson, M. Z., McMurry, P. H., Yu, F. Q., Yu, S. C., and Schere, K. L.: A comparative study of nucleation parameterizations: 2. Three-dimensional model application and evaluation, *J Geophys Res-Atmos*, 115, 10.1029/2010jd014151, 2010.
- 1030 Zhao, Z., Chen, S. H., Kleeman, M. J., Tyree, M., and Cayan, D.: The Impact of Climate Change on Air Quality-Related Meteorological Conditions in California. Part I: Present Time Simulation Analysis, *J Climate*, 24, 3344-3361, 2011.
- 1035 Zhu, Y. F., Hinds, W. C., Kim, S., Shen, S., and Sioutas, C.: Study of ultrafine particles near a major highway with heavy-duty diesel traffic, *Atmos Environ*, 36, 4323-4335, 10.1016/s1352-2310(02)00354-0, 2002a.
- Zhu, Y. F., Hinds, W. C., Kim, S., and Sioutas, C.: Concentration and size distribution of ultrafine particles near a major highway, *Journal of the Air & Waste Management Association*, 52, 1032-1042, 10.1080/10473289.2002.10470842, 2002b.
- 1040