

Interactive comment on “Climatology in Asian dust activation and transport based on MISR satellite observations and trajectory analysis” by Yan Yu et al.

Anonymous Referee #2

Received and published: 25 October 2018

The paper “Climatology in Asian dust activation and transport based on MISR satellite observations and trajectory analysis” presents and discusses the transport of dust aerosols, emitted from the arid and semiarid deserts of Taklamakan and Gobi, over the northern Pacific Ocean. The study falls within the scope of ACP. The study is based on MISR observations, MINX aerosol top height, and accordingly, forward HYSPLIT trajectory analysis. The manuscript is well-written/structured, the presentation clear, the language fluent. However, the submitted study is subject to major deficiencies and I would recommend publishing in ACP considering major revision.

Comments:

Printer-friendly version

Discussion paper



1) Regarding the “Asian dust activation climatology”. Dust aerosol classification is crucial in the scope of the study, since it is the initial point of the trajectories analysis. Therefore, I would recommend to the authors to describe briefly the dust aerosol classification in MISR/MINX (including the necessary references). The scientific methods and assumptions are not clearly outlined. How is a “dust plume” defined in the paper and how is a “dust event”? In addition, in case of air parcels containing dust aerosols originating from both the Taklimakan and Gobi desert, how is the discrimination performed to the different sources? Which are the uncertainties in the classification?

2) Regarding the “Asian dust transport climatology”. Although the paper presents an interesting approach to study dust transport the results are not sufficient to support the conclusions, due to the lack of observations provided on parallel with the trajectories. The study uses MISR observations-MINX provided top height to initiate HYSPLIT forward trajectories. Accordingly the climatology of trajectories is provided and not the Asian dust transport climatology. The difference is substantial. HYSPLIT computes the air parcel’s transport and dispersion from a source region (Taklamakan and Gobi here) and describes where the air parcel will go. In the framework of the study, the climatology of the trajectories is provided (spatial distribution - % of trajectory endpoints / Trajectory passage frequency - % of trajectories after a specific number of days), without providing any observation/evidence on the presence of dust (per trajectory, distance or area). Dust aerosols may already have been removed along the transport/trajectory due to dry (gravitational settling) or wet deposition, although the air parcel will reach further distances. The paper does not even provide quantitative information on the probability of dust to have been transported. The trajectory may extend over the Pacific Ocean, and even further, to the western coast of the United States, however this does not provide any guarantee that dust is present and has reached that distance. I would suggest the authors to do any necessary modifications to the manuscript. Either provide dust observations per trajectory or to focus on the trajectories analysis without giving the impression on the presence (and transport) of dust to the trajectories endpoint. Which are the uncertainties? Alternatively, the authors could implement ob-

[Printer-friendly version](#)[Discussion paper](#)

servations on the presence of dust to the western coast of USA (i.e. AERONET and AE, MODIS DT AOD and AE over ocean/ CALIOP volume/particle depolarization ratio) and use HYSPLIT back-trajectories. In addition, assuming a dust plume over an area, HYSPLIT initiated at different altitudes may provide different dust transport pathways. Therefore the study is representative only for the trajectories of the dust top-height and not for the dust plume (trajectories initiated at center of mass/scale height most probably would be more representative to discuss dust transport climatology).

3) The title does not reflect the contents of the paper and is misleading.

4) The figures are of high quality. I would suggest on Figure 2 to reverse the axes, time on horizontal axis and Injection height at the vertical height.

5) Regarding Figure 3 and the Gobi Desert the lack of continuity in wind speeds between 3 and 12 m/s is a strange feature. I would suggest the authors to describe this feature.

6) Regarding references, a brief list of references is provided. I would suggest the authors to expand the list of references in order to strengthen the manuscript and at the same time to give credit to related work. Indicatively, here a brief list of related studies is provided, describing features of dust aerosol transport emitted from the Taklamakan and Gobi deserts, based on synergies of passive and active ground-based and satellite-based instrumentation, models, campaigns and the meteorological and topographical mechanisms.

Bory, A. J. M., Biscaye, P. E. and Grousset, F. E.: Two distinct seasonal Asian source regions for mineral dust deposited in Greenland (NorthGRIP), *Geophys. Res. Lett.*, 30(4), 1167, doi:10.1029/2002GL016446, 2003.

Chen, S., Huang, J., Li, J., Jia, R., Jiang, N., Kang, L., Ma, X. and Xie, T.: Comparison of dust emissions, transport, and deposition between the Taklimakan Desert and Gobi Desert from 2007 to 2011, *Sci. China-Earth Sci.*, 60(7), 1338–1355,

[Printer-friendly version](#)[Discussion paper](#)

doi:10.1007/s11430-016-9051-0, 2017.

de Leeuw, G., Sogacheva, L., Rodriguez, E., Kourtidis, K., Georgoulas, A. K., Alexandri, G., Amiridis, V., Proestakis, E., Marinou, E., Xue, Y. and van der A, R.: Two decades of satellite observations of AOD over mainland China using ATSR-2, AATSR and MODIS/Terra: data set evaluation and large-scale patterns, *Atmos. Chem. Phys.*, 18(3), 1573–1592, doi:10.5194/acp-18-1573-2018, 2018.

Duce, R., Unni, C., Ray, B., Prospero, J. and Merrill, J.: Long-Range Atmospheric Transport of Soil Dust from Asia to the Tropical North Pacific - Temporal Variability, *Science*, 209(4464), 1522–1524, doi:10.1126/science.209.4464.1522, 1980.

Huebert, B. J., Bates, T., Russell, P. B., Shi, G. Y., Kim, Y. J., Kawamura, K., Carmichael, G. and Nakajima, T.: An overview of ACE-Asia: Strategies for quantifying the relationships between Asian aerosols and their climatic impacts, *J. Geophys. Res.-Atmos.*, 108(D23), 8633, doi:10.1029/2003JD003550, 2003.

Liu, Z., Liu, D., Huang, J., Vaughan, M., Uno, I., Sugimoto, N., Kittaka, C., Trepte, C., Wang, Z., Hostetler, C. and Winker, D.: Airborne dust distributions over the Tibetan Plateau and surrounding areas derived from the first year of CALIPSO lidar observations, *Atmos. Chem. Phys.*, 8(16), 5045–5060, 2008.

McKendry, I. G., Macdonald, A. M., Leaitch, W. R., van Donkelaar, A., Zhang, Q., Duck, T. and Martin, R. V.: Trans-Pacific dust events observed at Whistler, British Columbia during INTEX-B, *Atmos. Chem. Phys.*, 8(20), 6297–6307, 2008.

Proestakis, E., Amiridis, V., Marinou, E., Georgoulas, A. K., Solomos, S., Kazadzis, S., Chimot, J., Che, H., Alexandri, G., Biniotoglou, I., Daskalopoulou, V., Kourtidis, K. A., de Leeuw, G. and Ronald, J. van der A.: Nine-year spatial and temporal evolution of desert dust aerosols over South and East Asia as revealed by CALIOP, *Atmos. Chem. Phys.*, 18(2), 1337–1362, doi:10.5194/acp-18-1337-2018, 2018.

Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E. and Gill, T. E.: Environmental

Printer-friendly version

Discussion paper



characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40(1), 1002, doi:10.1029/2000RG000095, 2002.

Shaw, G.: Transport of Asian Desert Aerosol to the Hawaiian-Islands, *J. Appl. Meteorol.*, 19(11), 1254–1259, doi:10.1175/1520-0450(1980)019<1254:TOADAT>2.0.CO;2, 1980.

Sogacheva, L., Leeuw, G. de, Rodriguez, E., Kolmonen, P., Georgoulias, A. K., Alexandri, G., Kourtidis, K., Proestakis, E., Marinou, E., Amiridis, V., Xue, Y. and A. R. J. van der: Spatial and seasonal variations of aerosols over China from two decades of multi-satellite observations – Part 1: ATSR (1995–2011) and MODIS C6.1 (2000–2017), *Atmospheric Chemistry and Physics*, 18(15), 11389–11407, doi:https://doi.org/10.5194/acp-18-11389-2018, 2018.

Stith, J. L., Ramanathan, V., Cooper, W. A., Roberts, G. C., DeMott, P. J., Carmichael, G., Hatch, C. D., Adhikary, B., Twohy, C. H., Rogers, D. C., Baumgardner, D., Prenni, A. J., Campos, T., Gao, R., Anderson, J. and Feng, Y.: An overview of aircraft observations from the Pacific Dust Experiment campaign, *J. Geophys. Res.-Atmos.*, 114, D05207, doi:10.1029/2008JD010924, 2009.

Tan, S.-C., Li, J., Che, H., Chen, B. and Wang, H.: Transport of East Asian dust storms to the marginal seas of China and the southern North Pacific in spring 2010, *Atmos. Environ.*, 148, 316–328, doi:10.1016/j.atmosenv.2016.10.054, 2017.

Uno, I., Amano, H., Emori, S., Kinoshita, K., Matsui, I. and Sugimoto, N.: Trans-Pacific yellow sand transport observed in April 1998: A numerical simulation, *J. Geophys. Res.-Atmos.*, 106(D16), 18331–18344, doi:10.1029/2000JD900748, 2001.

Uno, I., Yumimoto, K., Shimizu, A., Hara, Y., Sugimoto, N., Wang, Z., Liu, Z. and Winker, D. M.: 3D structure of Asian dust transport revealed by CALIPSO lidar and a 4DVAR dust model, *Geophys. Res. Lett.*, 35(6), L06803, doi:10.1029/2007GL032329,

[Printer-friendly version](#)[Discussion paper](#)

2008.

Xu, H., Zheng, F. and Zhang, W.: Variability in Dust Observed over China Using A-Train CALIOP Instrument, *Adv. Meteorol.*, 1246590, doi:10.1155/2016/1246590, 2016.

Yumimoto, K., Eguchi, K., Uno, I., Takemura, T., Liu, Z., Shimizu, A. and Sugimoto, N.: An elevated large-scale dust veil from the Taklimakan Desert: Intercontinental transport and three-dimensional structure as captured by CALIPSO and regional and global models, *Atmos. Chem. Phys.*, 9(21), 8545–8558, 2009.

Zhang, X. Y., Arimoto, R. and An, Z. S.: Dust emission from Chinese desert sources linked to variations in atmospheric circulation, *J. Geophys. Res.-Atmos.*, 102(D23), 28041–28047, doi:10.1029/97JD02300, 1997.

Zhang, X. Y., Gong, S. L., Shen, Z. X., Mei, F. M., Xi, X. X., Liu, L. C., Zhou, Z. J., Wang, D., Wang, Y. Q. and Cheng, Y.: Characterization of soil dust aerosol in China and its transport and distribution during 2001 ACE-Asia: 1. Network observations, *J. Geophys. Res.-Atmos.*, 108(D9), 4261, doi:10.1029/2002JD002632, 2003.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2018-857>, 2018.

Printer-friendly version

Discussion paper

