

Interactive comment on “Near-surface and path-averaged mixing ratios of NO₂ derived from car DOAS zenith-sky and tower DOAS off-axis measurements in Vienna: a case study” by Stefan F. Schreier et al.

Anonymous Referee #1

Received and published: 10 November 2018

General Comments

This paper presents near-surface and path-averaged mixing ratios of NO₂ derived from car DOAS zenith-sky and tower DOAS off-axis, measurements performed in Vienna city during several days on 2015 and 2016. This paper provides an useful intercomparison between tower DOAS, mobile DOAS and in-situ observations.

Specific Comments

Section 2 - Instrument and car journeys, in this section you should add few info about

Printer-friendly version

Discussion paper



the in-situ instruments (type, error, etc.). Also please add a map (a new Figure) or include in Figure 1 the location of the in-situ monitoring stations and also the location of the DOAS tower instrument.

Please describe the tower DOAS instrument, I suggest you to introduce a Table with the technical characteristics of the two instruments (tower DOAS and mobile DOAS).

Figure 2. Could you explain the very low peak of intensity? Is it related to a tree, tunnel, or a bridge? Did you filter all the DSCDs function of RMS and O4?

Figure 3, please introduce the DSCD error. Also please introduce the error of each DSCD presented in the manuscript.

3.2.1 Temporal resolution and computation of horizontal NO₂ gradients- Could you specify the exposure time for the mobile DOAS instrument? (or this info could be included on the suggested Table for the two DOAS instrument).

3.2.2 Stratospheric NO₂ columns, Could you specify the error of Bremen 3d CTM (B3dCTM) model?

3.2.3 Conversion to tropospheric NO₂ vertical column densities SCD_{ref}, could you specify why you don't have a SCD_{ref} for each day? SCD_{ref} is quite important if you want to have qualitative data. I suggest to the authors to introduce more details about SCD_{ref} calculation, e.g. exact time of the selected SCD_{ref}. SCD_{ref} having 1.3×10^{15} , 1.1×10^{15} , and 2.2×10^{15} molecules/cm² as tropospheric contribution could be realistic. Considering that SCD_{ref} contain stratospheric and tropospheric contributions, did you cancel the stratospheric contribution? why do you refer to SCD_{ref} as having only tropospheric contributions?

A chapter to describe the AMF calculation (using NO₂ profiles, albedo, geometry, PBL, etc.) is mandatory for this study, I suggest to the authors to use a table. Figure 6 should be part of this section and should include the AMF calculations for several days which are presented in this study.

[Printer-friendly version](#)[Discussion paper](#)

The authors should give more details about the error calculation of tropospheric NO₂ VCD, or a section of errors would be more appropriate.

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2018-866>, 2018.

ACPD

Interactive
comment

Printer-friendly version

Discussion paper

