

# Sensitivity of Arctic sulfate aerosol and clouds to changes in future surface seawater dimethylsulfide concentrations

Rashed Mahmood<sup>1,2</sup>, Knut von Salzen<sup>1,2</sup>, Ann-Lise Norman<sup>3</sup>, Martí Galí<sup>4</sup> and Maurice Levasseur<sup>5</sup>

<sup>1</sup>School of Earth and Ocean Sciences, University of Victoria, Victoria, British Columbia, Canada

<sup>2</sup>Canadian Center for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, British Columbia, Canada

<sup>3</sup>Department of Physics and Astronomy, University of Calgary, Calgary, Alberta, Canada

<sup>4</sup>Takuvik Joint International Laboratory & Québec-Océan, Université Laval, Québec, Québec, Canada

<sup>5</sup>Département de biologie and Québec-Océan, Université Laval, Québec, Québec, Canada

*Correspondence to* : Knut von Salzen (knut.vonsalzen@canada.ca)

**Abstract.** Dimethylsulfide (DMS), outgassed from ocean waters, plays an important role in the climate system, as it oxidizes to methane sulfonic acid (MSA) and sulfur dioxide (SO<sub>2</sub>), which can lead to the formation of sulfate aerosol. Newly formed sulfate aerosol resulting from DMS oxidation may grow by condensation of gases, in-cloud oxidation, and coagulation to sizes where they may act as cloud condensation nuclei (CCN) and influence cloud properties. Under future global warming conditions, sea-ice in the Arctic region is expected to decline significantly, which may lead to increased emissions of DMS from the open ocean and changes in cloud regimes. In this study we evaluate impacts of DMS on Arctic sulfate aerosol budget, changes in cloud droplet number concentration (CDNC), and cloud radiative forcing in the Arctic region under current and future sea ice conditions using an atmospheric global climate model. Given that future DMS concentrations are highly uncertain, several simulations with different surface seawater DMS concentrations and spatial distributions in the Arctic were performed in order to determine the sensitivity of sulfate aerosol budgets, CDNC, and cloud radiative forcing to Arctic surface seawater DMS concentrations. For any given amount and distribution of Arctic surface seawater DMS, similar amounts of sulfate are produced by oxidation of DMS in 2000 and 2050 despite large increases in DMS emission in the latter period due to sea ice retreat in the simulations. This relatively low sensitivity of sulfate burden is related to enhanced sulfate wet removal by precipitation in 2050. However simulated aerosol nucleation rates are higher in 2050, which results in an overall increase in CDNC and substantially more negative cloud radiative forcing. Thus potential future reductions in sea ice extent may cause cloud albedos to increase, resulting in a negative climate feedback on radiative forcing in the Arctic associated with ocean DMS emissions.

## 1 Introduction

Dimethylsulfide is produced in the surface ocean by biological processes that involve phytoplankton, zooplankton, and bacteria (Simó 2001; Stefels et al. 2007). A fraction of the surface seawater DMS is vented to the atmosphere, depending on turbulence at the air-water interface (generally parameterized as a function of wind speed; Wanninkhof et al., 2009; see also Jahne et al., 1987; Merlivat and Memery, 1983; Memery and Merlivat, 1985; Monahan and Spillane, 1984) and

the depth of the oceanic upper mixed layer that exchanges with the atmosphere (Galí and Simó, 2015). Atmospheric DMS is subsequently oxidized to MSA and SO<sub>2</sub>. The latter is further oxidized to sulfuric acid (H<sub>2</sub>SO<sub>4</sub>), which can cause  
45 formation of new aerosols or condense on pre-existing aerosols. These aerosols may then act as CCN and affect cloud microphysical properties, especially in remote marine environments where concentrations of other types of CCN are low (Clarke et al., 1998; Leaitch et al., 2013; Dall'Osto et al., 2017, Collins et al., 2017). Willis et al. (2016) found that gaseous MSA may also play an important role in the initial growth of new particle formation.

50 According to the so-called CLAW hypothesis (Charlson et al., 1987), a negative feedback loop operates between ocean ecosystems and the Earth's climate. In particular, formation of new aerosol particles from ocean DMS emissions to the atmosphere leads to increased cloud albedo and reduced surface ocean temperature and/or incident irradiance, which then suppresses production of DMS in the ocean and emission to the atmosphere. Recent studies concluded that there is little evidence to support CLAW under present day climate conditions (Woodhouse et al., 2010; Quinn and Bates, 2011;  
55 Browse et al., 2014) since the sources of cloud condensation nuclei in marine boundary layer are numerous and the response of clouds to changes in aerosols are more complex than previously thought at the time of CLAW hypothesis (Quinn and Bates, 2011). However, marine DMS emission may still drive submicron aerosol populations over much of the remote marine atmosphere (Quinn et al., 2017). Important local impacts of DMS on climate may still exist in the Arctic where summertime aerosol and clouds are strongly influenced by DMS (Leaitch et al., 2013).

60 Considerable concentrations of ultrafine particles and DMS, which are likely involved in naturally occurring aerosol/climate interactions, have been observed in the Arctic in summer (Willis et al., 2016; Ghahremaninezhad et al., 2016; Burkart et al., 2017). Furthermore, Grandey and Wang (2015) found that artificially enhanced DMS emissions in different latitude bands could potentially offset greenhouse gas induced warming across most of the world and especially in the Arctic region. The loss of Arctic sea-ice allows further penetration of sunlight into surface ocean water (e.g.  
65 Nicolaus et al., 2012) that can increase net production of algae and phytoplankton (e.g. Arrigo and van Dijken, 2015). In addition, when sea-ice is melted the surface ocean water is more prone to wind stress (Rainville et al., 2011; Martin et al., 2014) that can enhance air-fluxes of DMS and other gases (e.g. Bates et al. 2006). Possible increases in DMS emissions and sulfate aerosol concentrations are particularly important in the Arctic where changes in aerosol radiative forcings are amplified by powerful Arctic feedback processes, including the surface albedo feedback (Pithan and Mauritsen, 2014;  
70 Gagné et al. 2015).

Arguably, uncertainties in surface seawater DMS concentrations and parameterizations of DMS emission fluxes limit scientific progress on climate effects associated with CLAW. For instance, the widely used climatology by Lana et al. (2011) is based on a compilation of data sets from different Arctic field campaigns that took place during the time period  
75 from 1985 to 2008. Only measurements from the warm season were used, mainly from the Atlantic portion of the Arctic. The small number of measurements from other locations in the Arctic is problematic, as recent research in the NETCARE network (Abbatt et al., 2019) has shown. Surface seawater DMS concentrations measured in the Canadian Arctic in July and August of 2014 and 2016 were substantially higher than those used by Lana et al. for July and August (e.g., NETCARE median concentrations of 4.4 nmol/L and 7.3 nmol/L, Martine Lizotte, personal communication;

80 median concentration range from 0.5 to 4.4 nmol/L for Lana et al., <https://saga.pmel.noaa.gov/dms/>). Furthermore, melt  
ponds on sea ice represent a yet missing source of DMS in studies of the Arctic (Mungall et al., 2016;  
Ghahremaninezhad et al., 2016; Gourdal et al. 2018; Abbatt et al., 2019). Galí et al. (2018) argue that biases in the  
climatology by Lana et al. arise from the application of objective interpolation procedures to a limited amount of  
measurements. Consequently, Arctic surface seawater DMS concentrations based on Lana et al. (2011) differ  
85 substantially from those of an earlier climatology (Kettle and Andreae, 2000), ocean biogeochemical models, and DMS  
parameterizations (Tesdal et al., 2016a), indicating large uncertainties in estimates of surface seawater DMS  
concentrations.

Over the last few decades, Arctic temperature has increased at a rate much faster than in other parts of the world (ACIA,  
90 2005; AMAP, 2017). Enhanced Arctic warming is largely caused by sea-ice-albedo feedbacks and it is expected that  
Arctic summer sea-ice may completely disappear well before the end of this century if warming continues at rates  
simulated by current climate models (Stroeve et al., 2012). Browse et al. (2014) found a weak response of CCN  
concentrations to enhanced Arctic DMS emissions from complete loss in summer sea-ice due to efficient scavenging of  
aerosol by drizzle associated with stratocumulus clouds. They did not find evidence for climate feedbacks through  
95 changes in cloud properties from enhanced aerosol sources in an ice-free summertime Arctic. However, Browse et al.  
used an atmospheric chemical transport model with specified meteorological conditions, which excludes responses of  
Arctic clouds and precipitation to changes in sea ice conditions. In a subsequent study, Ridley et al. (2016) performed  
fully interactive simulations of sea-ice, ocean biology (with DMS estimated using an embedded empirical algorithm;  
Simó and Dachs, 2002), aerosols, and clouds with the HadGEM2-ES model. They found a two- to five-fold increase in  
100 DMS emissions, an increase in sulfate CCN concentration, and an associated  $1 \text{ W m}^{-2}$  reduction in simulated summer  
cloud shortwave radiative forcing in the Arctic.

Despite substantial research activities (e.g. Gabric et al., 2005, Thomas et al., 2010, Woodhouse et al., 2010; Browse et  
al., 2014; Ridley et al., 2016) it is still very challenging to estimate DMS emissions and even more so how sea-ice  
105 reductions may affect DMS emissions in the Arctic in the future. Results of several modeling studies indicated that  
global warming related sea-ice loss would result in enhanced DMS emission fluxes in the Arctic region (Bopp et al.,  
2003; Gabric et al., 2005; Levasseur, 2013; Browse et al., 2014; Galí et al., *submitted*). Other studies suggest that the  
changes in DMS emission flux may be negative in sign due to potentially enhanced ocean acidification (e.g. Six et al.,  
2013; Schwinger et al., 2017). Phytoplankton species composition is a controlling factor for DMS concentrations given  
110 the wide range of cellular dimethylsulfoniopropionate (DMSP) quota among different phytoplankton species (Stefels  
et al. 2007). For example, the cellular DMSP quota of haptophytes is greater than that of diatoms by a factor of  $>10$  (Stefels  
et al. 2007). Long-term observational studies provide evidence that high DMSP-producing haptophytes are becoming  
more prevalent in the Arctic in the last decade (Winter et al., 2013; Nöthig et al., 2015; Soltwedel et al., 2016).  
Furthermore, Arrigo et al. (2008) suggest that primary productivity may increase more than 3 times compared to 1998-  
115 2002, if Arctic sea ice loss continues. A combination of a shift in the species composition and an increase in primary  
productivity (e.g. Yool et al., 2005; Vancoppenolle et al., 2013) could imply a multiplicative increase in surface seawater  
DMS concentrations in future climate.

120 Given large uncertainties in present-day surface seawater DMS concentrations and potentially large increases in future concentrations, the approach in the current study is to consider a wide range in concentrations in the Arctic for present-day and future conditions. The lower bound of the concentration range is provided by Lana et al. (2011), and the upper bound is obtained by scaling these concentrations by a factor of 10. A number of scenarios for DMS concentrations within this range are considered. These will be used to determine relationships between surface seawater DMS concentrations and climate variables in the Arctic. By selecting widely different scenarios, the robustness of the relationships can be tested without making a priori assumptions that only apply to specific DMS scenarios. Model sensitivity tests with similarly enhanced DMS concentrations have previously been performed (e.g. Grandy and Wang, 2015; Fiddes et al., 2018). Sensitivities of sulfate aerosols, CDNC and cloud radiative forcing to different DMS emission scenarios are investigated using a state-of-the-art atmospheric global climate model. Based on the sensitivity simulations we provide an assessment of Arctic annual mean changes in sulfate aerosol budget and cloud microphysical properties in relation to the mean DMS concentration in the Arctic. Note that an evaluation of surface seawater DMS data sets is outside the scope of the study.

## 2 Summary of model features

135 We used version 4.3 of the Canadian Atmospheric Model (CanAM4.3) which is an improved version of CanAM4 (von Salzen et al., 2013). The improvements to version 4.3 compared to version 4 include a higher vertical resolution, improved parameterizations for land surface and snow processes, DMS emissions, and clear-sky radiative transfer. CanAM4.3 has 49 vertical levels extending up to 1 hPa with a resolution of approximately 100 meters near the surface. Model simulations are performed using a spectral resolution of T63 which is equivalent to the horizontal resolution of approximately  $2.8 \times 2.8$  degrees. The model uses separate parameterizations for layer and convective clouds. Aerosol microphysical processes are based on the Piecewise Lognormal Approximation (von Salzen, 2006; Ma et al., 2008; Peng et al., 2012; Mahmood et al., 2016; AMAP, 2015). A detailed description of parameterizations of ocean DMS flux to atmosphere, oxidation and removal processes is provided in Tesdal et al. (2016a). Briefly, surface seawater DMS is ventilated to the atmosphere based on modeled wind speed and the piston velocity parameterization of Nightingale et al. (2000). There are no DMS emissions from sea ice. Furthermore, the model does not account for emissions of organic aerosol species from the ocean.

In the atmosphere, DMS is oxidized to MSA and  $\text{SO}_2$  by hydroxyl (OH) radicals during day time and nitrate radical ( $\text{NO}_3$ ) during night, with further oxidation of  $\text{SO}_2$  to sulfuric acid ( $\text{H}_2\text{SO}_4$ ) by OH in clear-sky conditions. MSA is treated as sulfuric acid in model for simplicity. Binary homogeneous nucleation of  $\text{H}_2\text{SO}_4$  and water vapour may cause formation of new aerosol particles, depending on temperature and relative humidity (Kulmala et al., 1998; von Salzen et al., 2000). In-cloud production of sulfate requires ozone ( $\text{O}_3$ ) and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) as oxidants (von Salzen et al., 2000), with oxidant (OH,  $\text{NO}_3$ ,  $\text{H}_2\text{O}_2$ ,  $\text{O}_3$ ) concentrations specified according as climatological results from the Model for Ozone and Related Chemical Tracers (MOZART, Brasseur et al., 1998). Dry deposition of aerosol depends on concentrations of aerosols in the near surface model layer (Zhang et al., 2001). Wet deposition includes in-cloud

scavenging in both convective clouds and layer clouds, and below-cloud scavenging. Emissions of non-DMS biogenic and anthropogenic aerosol precursors and primary aerosols for the time period up to year 2000 are specified according to Lamarque et al. (2010) and after that according to the IPCC RCP4.5 scenario (Moss et al., 2010).

160 Cloud droplet number concentrations are calculated based on the assumption of a parcel of air which ascends from the subcloud layer into the cloud layer with a characteristic vertical velocity (Peng et al., 2005), where the standard deviation of the subgrid-scale cloud vertical velocity probability distribution is parameterized using the approach by Ghan et al. (1997). Aerosol particles that are suspended in the parcel of air may activate and grow into cloud droplets by condensation of water vapour. A numerically efficient solution of the condensational droplet growth equation (e.g. 165 Seinfeld and Pandis, 1998) is employed for this purpose.

### 3 Sulfate concentrations in CanAM4.3

We performed two sets of historical model simulations, one with the full model with all natural and anthropogenic 170 aerosols and their precursors included (“hisCont”) and one with zero surface seawater DMS (“hisNoDMS”). Monthly mean surface seawater DMS concentrations in hisCont are specified according to the climatology of Lana et al. (2011) [hereafter referred to as L10]. The global annual DMS emission flux in hisCont is 24.96 TgS/yr, which is very close to 25.3 TgS/yr reported in Tesdal et al. (2016a) and well within previously reported ranges (e.g. Lana et al., 2011; Tesdal et al., 2016a). Both simulations were integrated for the time period 1991 to 2003 during which extensive observations of 175 sulfate are available. Wind and temperature in each simulation were nudged towards specified results from a common simulation with CanAM4.3 for this time period. According to Kooperman et al. (2012) nudging of meteorological model variables reduces the influence of natural variability and therefore improves estimates of differences in diagnosed aerosol indirect effects. Similarly, biases in simulated aerosol and CDNC concentrations between the simulations are also reduced according to a statistical analysis of CanAM model results (not shown). The contribution of DMS oxidation to 180 total sulfate concentrations is determined by calculating the difference in simulated sulfate concentrations between these two simulations (i.e. hisCont - hisNoDMS), which is interpreted as biogenic sulfate in the following.

A slightly different version of CanAM has previously been evaluated (Eckhardt et al., 2015; Tesdal et al., 2016b). In a 185 multi-model comparison, Eckhardt et al. (2015) compared model simulations of sulfate and black carbon aerosols with observations from different stations and aircraft campaigns and found that most models, except CanAM, significantly underestimated observed concentrations in the Arctic region. Mahmood et al. (2016) found that black carbon concentration differences in four models are related to differences in wet removal processes in the models.

For the current study we used observed data from various ship-based campaigns and observations from Alert in Canada 190 to further validate simulations of sulfate, with particular emphasis on the role of biogenic emissions. Shipboard data from the National Oceanic and Atmospheric Administration, Pacific Marine Environmental Laboratory (NOAA PMEL) was obtained from cruises that fell within the period 1992-2002. Only non-sea-salt sulfate ( $\text{nss-SO}_4^{2-}$ ) data was selected and

summed for all available bin sizes. The gridded model data was matched to the nearest location of the observations, shown in Fig. 1a.

195  
Figure 1 shows that simulated sulfate concentrations agree well with observations, especially in regions where modeled DMS contributions are relatively large. The mean value for all ship-based observations is  $3.416 \pm 4.018$  ( $\mu\text{g}/\text{m}^3$ ) and the model mean value is  $2.079 \pm 1.815$  ( $\mu\text{g}/\text{m}^3$ ), corresponding to a model underestimate of  $\sim 39\%$ . Most of the underestimates in simulated mean concentrations are associated with locations where the model simulates a large contribution of fossil-fuel (non-biogenic) sulfate to total sulfate concentrations.

200  
From the Alert data, with highly variable contributions of DMS, it is evident that the model overestimates the contribution of DMS to total sulfate concentrations at this location (Fig. 1b). Overall, CanAM4.3 is able to capture the sulfate annual cycle very well at Alert, with slight underestimation in winter and spring and overestimation in summer (Fig. 2). The correlation coefficient for the mean annual cycle between model and observations is 0.95. Mean observed and simulated sulfate concentrations at Alert are  $0.475 \pm 0.413$  and  $0.419 \pm 0.228$   $\mu\text{g}/\text{m}^3$  respectively, corresponding to a model underestimate of  $\sim 12\%$ .

210  
An analysis of isotopic data is available for Alert, which can be used to distinguish between contributions of biogenic and fossil-fuel sources to sulfate concentrations in the observations at this site (Norman et al., 1999). The ratio between sulfur isotopes  $^{34}\text{S}$  and  $^{32}\text{S}$  of an observed sample is compared with an international standard ratio based on sulfur isotopes in Vienna-Cañon Diablo Troilite (Beaudoin et al., 1994; Krouse and Grinenko, 1991; Norman et al., 1999). The results are expressed in parts per thousand (‰). The sulfate concentration at Alert consists of the sum of marine biogenic, anthropogenic, and sea salt sulfate with the delta isotopic ratios of  $+17.5\%$ ,  $+5\%$ ,  $+21\%$  respectively.

215  
A comparison with the isotopic data indicates that the model underestimates fossil-fuel sulfate and overestimates biogenic sulfate (Fig. 2b and 2c). This difference is particularly pronounced in spring and early summer, when observed biogenic sulfate concentrations are particularly high. An interesting feature is the double peak in biogenic sulfate concentrations during the annual cycle with one peak occurring in May and the other in October (Fig. 2c), which is essentially captured by the model. The correlation coefficient for the mean annual cycle between observed and simulated biogenic sulfate concentrations is 0.73.

220  
Another interesting feature is the relative contribution of biogenic sources to total sulfate concentrations (Fig. 2d). Although absolute sulfate aerosol concentrations in summer are much lower than during other seasons, both observations and model results indicate a much larger contribution of biogenic sources to total sulfate concentration in this season compared to other seasons (Fig. 2d).

#### 4 Sensitivity of sulfate, clouds, and radiation to changes in DMS



230 Five ensembles of five simulations each were performed, where ensemble members were generated by introducing random perturbations in radiative flux calculations (a total of 25 simulations). The five experiments differ in terms of specified surface seawater DMS concentrations in the Arctic region, defined here as the region from 62.78°-90°N. A wide range of different Arctic surface seawater DMS concentration patterns is considered in order to account for substantial uncertainties in surface seawater DMS concentrations. For present day, uncertainty in specified Arctic DMS  
235 concentration climatologies arises from a lack of observational data, and concentrations that are highly variable in space and time (Lana et al., 2011; Tesdal et al., 2016a; Galí et al., 2018). Furthermore, very little is known about how DMS concentration may evolve in the future. Outside the Arctic, monthly mean ocean DMS concentrations in the simulations are specified according to the L10 climatology.

240 For the first set of simulations, DMS concentrations were specified as zero in the Arctic, hereafter referred to as CNTRL. All subsequent simulations are compared to CNTRL in order to estimate the contribution of Arctic DMS emissions to simulated biogenic sulfate concentrations and climate. For the second set of simulations, CLIM, the L10 monthly climatology is used in the Arctic. In 10×CLIM, the L10 climatology was multiplied by a factor of 10 at each model grid point in the Arctic region. In order to analyze the impact of spatial variability in DMS concentrations, a single DMS  
245 concentration (i.e. 16.9 nM) was assigned to each grid cell in the Arctic region hereafter referred to as UNFM. Finally, as a further test of spatial variability of DMS, we used a satellite-based estimate of surface seawater DMS concentration (Galí et al., 2018) and multiplied it by a factor of 10 in 10×SAT. A value of 5 nM was applied in 10×SAT over the central Arctic region where satellite observations are not available, upon the observation that available sea-surface DMS measurements in the Arctic winter have an average of ~0.5 nM. Note this earlier version of Galí et al. (2018) satellite  
250 based DMS estimation had a small negative bias in magnitude, however, the spatial distribution remained largely unchanged after correction. Surface seawater DMS concentrations in all simulations are summarized in Fig. 3 and Table 1.

Simulated horizontal wind and temperature in each individual member of an ensemble (i.e. 5 separate simulations) were  
255 nudged towards specified results from a corresponding simulation (i.e. separate free running model simulation) with CanAM4.3 using a nudging time scale of 6 h. The model was integrated over four years for 1998-2001 and annual mean model results during the last three years of the simulations were analyzed (hereafter referred to as 2000). Present-day sea ice amounts and sea surface temperatures (SSTs) are specified according to reanalysis data from Climate Forecast System Version 2 (Saha et al., 2014). In addition, all of the above experiments were repeated for 2048-2051 (referred to  
260 as 2050). The projected sea ice amounts and SSTs from a large ensemble of simulations with CanESM2 for the RCP8.5 emission scenario were used in order to represent conditions in 2050 (Sigmond and Fyfe, 2016). In CanESM2, the Arctic is devoid of sea ice in September by 2050, consistent with results from other models for this relatively high emission scenario (Stroeve et al., 2012). The difference of sea ice extent in the simulation time periods is summarized in Fig. S1 (in supplementary data). The Arctic annual mean sea ice fractions are 75.6% (2000) and 50% (2050) for grid  
265 cells where the sea ice fraction is 0.15 or larger. Similar to simulations corresponding to year 2000, simulated horizontal winds and temperature were nudged towards specified results from 5 simulations with different meteorological conditions.

270 The total annual Arctic DMS fluxes for the two simulations time periods are summarized in Table 1. For 2000, DMS  
emission fluxes are approximately linearly proportional to the mean surface seawater DMS concentrations. For example,  
the total annual Arctic fluxes for the 10xCLIM are 84% higher than for 10xSAT, corresponding to 76% higher  
concentrations of Arctic-mean surface seawater DMS. This indicates that the sensitivity of Arctic-mean DMS fluxes to  
deviations of spatial distributions from the mean surface seawater concentrations is relatively low for 2000. Similar  
275 results for 2050 give evidence for relatively low sensitivity of fluxes to spatial distributions of surface seawater DMS  
concentrations.

Sea ice fraction in 2050 in summer and autumn is much lower than in 2000 (Fig. S1). For CLIM and 10xCLIM the total  
Arctic sulfur flux increases by 33% from the earlier to the later time period due to the reduction in future sea ice fraction.  
The difference is up to 47% for 10xSAT and 53% for UNFM (Table S1). Regionally, differences in fluxes are strongly  
280 correlated with changing sea ice fractions, with increases in regions with reduced sea ice fraction in 2050 and only minor  
changes over the open ocean (Fig. 4).

Owing to large DMS emission fluxes in the Atlantic region, the spatial pattern of the biogenic sulfate burden (relative to  
CNTRL) produces a maximum in this region for all surface seawater DMS data sets (Fig. 5). However, for CLIM  
285 diagnosed biogenic sulfate burdens are statistically significant over the Greenland Sea and nearby Baffin Bay for 2050  
but not for 2000. Differences in Arctic-mean biogenic sulfate burdens between 2000 and 2050 are relatively small for all  
of the scenarios, ranging from just -1% for UNFM to +21% for CLIM, despite the relatively large increases in DMS  
emissions between 2000 and 2050 (Table 1 and Fig. 5). The weak responses in biogenic sulfate burdens to DMS  
emissions are caused by increased precipitation and aerosol wet removal in the Arctic in 2050 (Tables S1 and S2; Figs.  
290 S2 and S3). Thus the wet deposition of biogenic sulfate from Arctic DMS emissions becomes more efficient in the future.  
Whereas emissions of Arctic DMS increase between 33.3% (CLIM) and 53.2% (UNFM), wet deposition of biogenic  
sulfate from Arctic DMS emissions increases more strongly, between 42.45% (CLIM) and 72.1% (UNFM) from 2000 to  
2050 (Table S2). The fraction of Arctic DMS emissions that is removed by wet deposition increases from between 55.3%  
(UNFM) and 76.5% (CLIM) in 2000 to between 62.1% (UNFM) and 81.8% (CLIM) in 2050 (Table S3).

295 On the other hand, projected reductions in anthropogenic sulfur emissions between 2000 and 2050 lead to reductions in  
total wet deposition of sulfate in the Arctic by -47.7% in CNTRL (Table S1). In the sensitivity experiments with  
increases in Arctic emissions of DMS between 2000 and 2050 reductions in total sulfate wet deposition in the Arctic  
between 2000 and 2050 are weaker, i.e. between -7.5% (10xCLIM) and -40.6% (CLIM). Considering the very wide  
300 range of surface seawater DMS concentrations applied here, a nearly complete compensation of aerosol production from  
oceanic DMS by increased wet deposition seems to be a robust feature of the future Arctic, largely independent of DMS  
emission patterns and amounts.

Changes in CDNC are important for radiative effects of sulfate aerosols. Impacts of climatological DMS emissions on  
305 CDNC are not statistically significant in CLIM, i.e. they are within the range of meteorological variability in the



ensemble of simulations (Fig. 6). The relatively weak simulated impact of present-day climatological DMS concentrations on CDNC and cloud microphysics is in agreement with previous studies (e.g. Browse et al., 2014; Ridley et al., 2016). Similarly, for 2050, few regions in the Arctic show significant impacts of present-day DMS emissions on CDNC although local increases are up to about 10%. On the other hand, the other sets of simulations (i.e. 10×CLIM, UNFM and 10×SAT) produce significant changes in CDNC, especially for 2050, with increases up to  $\sim 10^7 \text{ m}^{-3}$  for 10×CLIM. It is interesting to note that although the biogenic sulfate burdens are similar in 2000 and 2050, there are relatively large systematic increases in CDNC due to increased Arctic DMS emissions in 2050 for these simulations.

Increases in CDNC between 2000 and 2050 are related to increases in formation of new particles in the lower Arctic troposphere by between +128 and +269% (Table 2 and Fig. S3) for the range in surface seawater DMS concentration considered. This leads to large-scale increases in CCN concentrations near the surface in the Arctic (Fig. S3), which is in contrast to a more non-uniform response of CCN concentrations to reductions in sea ice fraction according to Browse et al. (2014), with relatively large simulated increases over the continental Arctic and small decreases over the central Arctic Ocean. Large-scale increases of CCN concentrations and nucleation rates in 2050 in simulations with CanAM4.3 can be attributed to several factors: First, global anthropogenic emissions of sulfur are considerably lower in 2050 compared to 2000, which causes a reduction in the burden of anthropogenic sulfate (-65% in CNTRL) and the associated condensation sink of sulfuric acid in the Arctic atmosphere, which can be expected to facilitate the formation of new particles (Wyslouzil et al., 1991). The condensation sink of sulfuric acid is further reduced by increased wet deposition of aerosols due to increased Arctic precipitation. Finally, increased evaporation of moisture from the ocean leads to increases in relative humidity in the Arctic, which also produces conditions that are more favorable to nucleation in 2050 than 2000. On the other hand, increases in the sulfuric acid condensation sink due to increased emissions of sea salt and organic aerosols from the open ocean are not accounted for in the current version of the model, which may lead to overestimates in nucleation rates in the simulations. According to Browse et al. (2014), the increase in the natural condensation sink due to increased production of sea salt and organic aerosol under ice-free conditions causes a substantial reduction in the near-surface nucleation rate. However, it is likely that increases in production of sea salt under ice-free conditions are accompanied by large increases in wet deposition of sea salt due to increased Arctic precipitation (Struthers et al., 2011), which has not been explicitly accounted for by Browse et al. (2014). In addition, recent observations indicate a dominant role of small particles in activation and formation of cloud condensation nuclei in clean Arctic conditions (Leitch et al., 2016) implying efficient nucleation of fine mode particles.

According to the first indirect effect of aerosols on climate, increases in CDNC may lead to smaller cloud droplets which are associated with more efficient scattering of incoming solar radiation and therefore stronger cloud radiative forcings, determined here as the difference in total-sky minus clear-sky shortwave radiative fluxes at top of the atmosphere (Soden et al., 2004). Subsequently, the cloud radiative forcing associated with biogenic DMS from Arctic DMS emissions is determined as difference in cloud radiative forcing between sensitivity experiments and CNTRL. As shown in Fig. 7 the cloud radiative forcing due to Arctic DMS emissions is small and not statistically significant for CLIM for both time periods. However, for 10×CLIM and 10×SAT the cloud radiative forcing in the Arctic due to Arctic DMS emissions is significant with maximum of up to  $-4 \text{ Wm}^{-2}$  for the Atlantic side of the Arctic for 10×CLIM in 2050, qualitatively in

agreement with differences in CDNC. Overall, the mean cloud radiative forcing in the Arctic due to Arctic DMS emissions increases by between 108% (CLIM) and 145% (UNFM) from 2000 to 2050 (Table 2). All DMS data sets produce similar patterns of changes, with systematically enhanced cloud radiative forcings for the Atlantic region of the Arctic where loss of sea ice leads to particularly large increases in DMS emissions in all cases.

On regional scales, differences in cloud radiative forcing due to Arctic DMS emissions in Fig. 7 are generally smaller than changes in cloud radiative forcing associated with changes in meteorological conditions and anthropogenic aerosol precursor emissions between 2000 and 2050 (Fig. S4). However, averaged over the Arctic, differences are similar. For instance, the mean cloud radiative forcing in the Arctic in CLIM is  $-0.13$  and  $-0.27$   $\text{Wm}^{-2}$  for 2000 and 2050 respectively (difference of  $-0.14$   $\text{Wm}^{-2}$ ). Similarly, CNTRL produces a difference in cloud radiative forcings of  $-0.65$   $\text{Wm}^{-2}$  in total cloud radiative forcing between 2000 and 2050. It is evident that the cloud radiative forcing from Arctic DMS (Fig. 7) acts to enhance negative cloud radiative forcings in the central Arctic and counteracts positive forcings in the Atlantic Arctic and north of Siberia (Fig. S4), especially for 10xCLIM (the Arctic mean difference for 10xCLIM is  $-0.84$   $\text{Wm}^{-2}$  between 2000 and 2050).

Mean results in the Arctic are summarized in Fig. 8 which provides an indication of robust aerosol and cloud responses at pan-Arctic scale, despite large differences in amount and spatial distribution of surface seawater DMS concentration in the different cases. For instance, Arctic-mean sulfate burdens due to Arctic DMS emissions are similar for present-day and future conditions (Fig. 8b) despite strongly increased DMS emissions in 2050 resulting from sea ice retreat. On the other hand, biogenic DMS emissions lead to more efficient formation of cloud droplets in the future in the Arctic. Therefore, cloud droplet number concentrations and cloud radiative forcing increase systematically as sea ice extent declines from 2000 to 2050 for each Arctic DMS, despite the low sensitivity of biogenic sulfate burdens to changes in sea ice. This provides evidence for a negative feedback of Arctic DMS emissions on Arctic radiative forcing, assuming that DMS concentrations in the ocean and atmospheric oxidant concentrations do not change between 2000 and 2050. To a good first approximation, the strength of the feedback is proportional to the mean surface seawater DMS concentration in the Arctic despite low sensitivity of sulfate burdens. The simulated responses of clouds and radiative forcing to changes in sea ice extent are found to be robust for a wide range of surface seawater DMS concentration scenarios.

## 5 Conclusions

Simulated sulfate concentrations from the Canadian Center for Climate Modeling and Analysis Atmospheric Model (CanAM4.3) were compared to observations from various shipboard campaigns and in-situ observations at Alert in Canada with a particular emphasis on the role of biogenic emissions. We found that the model reproduced seasonal variations in observed biogenic sulfur concentrations at Alert, although the model overestimates the biogenic contribution to total sulfate somewhat. Observed biogenic sulfur concentration maxima in May and in October are well reproduced by the model. Furthermore, comparisons with ship-based measurements from different field campaigns yield good agreement with simulated sulfate concentrations, especially in regions with large contributions of biogenic sulfate. However, it is plausible that the current Arctic surface seawater DMS concentrations are underestimated because models

and climatological data sets do not yet account for substantially enhanced concentrations in melt ponds, and near the ice edge (e.g. Mungall et al., 2016; Ghahremaninezhad et al., 2016; Hayashida et al., 2017; Gourdal et al. 2018). In addition, large uncertainties exist for nucleation parameterizations (e.g. Zhang et al., 2010).

385 We performed model simulations to understand the sensitivity of sulfate aerosols and cloud radiative forcing to projected changes in sea ice and climate conditions between 2000 and 2050. Several model experiments were performed using a wide range of different surface seawater DMS concentrations in order to account for uncertainties in present-day and future DMS and to explore the sensitivity of aerosol/climate interactions to differences in spatial patterns of DMS.  
390 Results of the simulations indicate that the enhanced wet removal efficiency from increased precipitation in 2050 largely counteracts the impact of the increase in DMS emissions on sulfate burden in the Arctic. Annual mean Arctic sulfate burden differences between 2000 and 2050 are small for any given scenario (differences ranging between -1 and 21%) despite large increases in DMS emission between 2000 and 2050 due to sea ice retreat (between +33 and +53%). The sensitivity of modeled DMS fluxes into the atmosphere and sulfate burdens to spatial variations in surface seawater DMS  
395 is relatively weak.

Similar to previous studies (e.g. Browse et al., 2014; Ridely et al., 2016) we found weak impacts of climatological DMS emissions on cloud radiative forcings for present day conditions (simulation CLIM). However, for 2050 simulations, biogenic DMS emissions lead to considerable impacts on simulated Arctic aerosol and cloud processes owing to  
400 conditions that are conducive to the formation of fine particles in the Arctic. In 2050, increased emissions of DMS from large ice-free regions of the Arctic ocean are associated with increased biogenic sulfate aerosol nucleation rates (between +128 and +269%) and cloud droplet number concentrations (between +35 and +133%) and thus enhanced cloud albedos, resulting in negative cloud radiative forcing of biogenic sulfate in the Arctic. The difference in cloud radiative forcing between years 2050 and 2000 based on simulations for four different Arctic surface seawater DMS data sets ranges from  
405 between  $-0.14 \text{ Wm}^{-2}$  (CLIM) to  $-0.84 \text{ Wm}^{-2}$  ( $10 \times \text{CLIM}$ ). Thus our model results provide evidence for a negative Arctic climate feedback. The essential ingredient of the feedback is a response of DMS emissions and cloud droplet number concentrations to sea ice retreat due to changes in radiative forcings in the climate system. This differs from CLAW which is rooted in the assumption of a change in biological production of DMS in the ocean in response to a change in radiative forcings. Furthermore, the strength of the Arctic climate feedback is proportional to the mean surface seawater  
410 DMS concentration in the Arctic. Consequently, potential future changes in primary productivity (Yool et al., 2005; Vancoppenolle et al., 2013), mixing and phytoplankton habitat (Harada, 2016) in the Arctic Ocean (Levasseur, 2013) may act to enhance the strength of the Arctic feedback.

The model simulations used in the current study are not interactively coupled with ocean and sea ice DMS and therefore  
415 rely on specified surface seawater DMS concentrations. The current model version does not account for increases in sulfuric acid condensation sink due to increased emissions of sea salt and organic aerosols from the open ocean, which may lead to overestimates in nucleation rates in the simulations. Additional uncertainty in the strength of the feedback arises from the fact that atmospheric oxidant concentrations are assumed to be steady in our study. More comprehensive assessments of the strength and impacts of DMS/climate feedbacks in the Arctic will become possible once a new

420 generation of Earth System Models with interactive ocean and sea ice DMS, chemistry, and climate processes becomes  
available.

### Acknowledgement

425 Shipping non-sea-salt sulfate data was obtained from NOAA PMEL Atmospheric Data Server (available at:  
*https://saga.pmel.noaa.gov/data/*). We thank Jim Christian for providing an internal review of the manuscript. We also  
thank Martine Lizotte and Hakase Hayashida for providing very helpful discussions of surface ocean DMS  
concentrations. Funding for this work was provided by the Natural Sciences and Engineering Research Council of  
Canada through the NETCARE project of the Climate Change and Atmospheric Research Program.

430

### References

- Abbatt, J. P. D., Leaitch, W. R., Aliabadi, A. A., et al.: Overview paper: New insights into aerosol and climate in the  
Arctic, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-2018-995>, 2019.
- 435 ACIA, 2005.: Arctic Climate Impact Assessment. Cambridge University Press, Cambridge, UK, 2005.
- AMAP, 2015.: Black carbon and ozone as Arctic climate forcers, Arctic Monitoring and Assessment Programme  
(AMAP), Oslo, Norway. vii + 116 pp., 2015.
- AMAP, 2017.: Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017. Arctic Monitoring and Assessment  
Programme (AMAP), Oslo, Norway. xiv + 269 pp., 2017.
- 440 Arrigo, K. R., van Dijken G., and Pabi S.: Impact of a shrinking Arctic ice cover on marine primary production,  
*Geophys. Res. Lett.*, 35, L19603, doi: 10.1029/2008GL035028, 2008.
- Arrigo, K. R., and van Dijken, G. L.: Continued increases in Arctic Ocean primary production. *Prog. Oceanogr.*, 136,  
60–70, 2015.
- Bates, N. R., Moran, S. B., Hansell, D. A., and Mathis, J. T.: An increasing CO<sub>2</sub> sink in the Arctic Ocean due to  
sea-ice loss. *Geophysical Research Letters*, 33(23), 2006.
- 445 Beaudoin, G., Taylor, B. E., Rumble III, D., and  
Thiemens, M.: Variations in the sulfur isotope composition of troilite from the Cañon Diablo iron meteorite,  
*Geochim. Cosmochim. Acta*, 58, 4253-4255, 1994.
- Bopp, L., Aumont, O., Belviso, S., and Monfray, P.: Potential impact of climate change on marine dimethyl sulfide  
emissions, *Tellus Ser. B*, 55, 11–22, doi:10.1034/j.1600-0889.2003.042.x, 2003.
- Brasseur, G. P., Hauglustaine, D. A., Walters, S., Rasch, P. J., Müller, J.-F., Granier, C., and Tie, X.: MOZART, a  
450 global chemical transport model for ozone and related chemical tracers: 1. Model description, *J. Geophys. Res.-  
Atmos.*, 103, 28265–28289, doi:10.1029/98JD02397, 1998.
- Browse, J., Carslaw, K. S., Mann, G. W., Birch, C. E., Arnold, S. R., and Leck, C.: The complex response of Arctic  
aerosol to sea-ice retreat, *Atmos. Chem. Phys.*, 14, 7543–7557, doi:10.5194/acp-14-7543-2014, 2014.
- 455 Burkart, J., Willis, M. D., Bozem, H., Thomas, J. L., Law, K., Hoor, P., Aliabadi, A. A., Köllner, F., Schneider, J.,  
Herber, A., Abbatt, J. P. D., and Leaitch, W. R.: Summertime observations of elevated levels of ultrafine  
particles in the high Arctic marine boundary layer, *Atmos. Chem. Phys.*, 17, 5515-5535,  
<https://doi.org/10.5194/acp-17-5515-2017>, 2017.

- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo, and climate, *Nature*, 326, 655–661, 1987.
- 460 Clarke, A. D., Varner, J. L., Eisele, F., Mauldin, R. L., Tanner, D., and Litchy, M.: Particle production in the remote marine atmosphere: cloud outflow and subsidence during ACE-1. *J. Geophys Res* 103 (D13): 16,397–16,409. doi: 10.1029/97JD02987, 1998.
- Collins, D. B., Burkart, J., Chang, R. Y.-W., Lizotte, M., Boivin-Rioux, A., Blais, M., Mungall, E. L., Boyer, M., Irish, V. E., Massé, G., Kunkel, D., Tremblay, J.-É., Papakyriakou, T., Bertram, A. K., Bozem, H., Gosselin, M.,  
465 Levasseur, M., and Abbatt, J. P. D.: Frequent ultrafine particle formation and growth in Canadian Arctic marine and coastal environments, *Atmos. Chem. Phys.*, 17, 13119-13138, <https://doi.org/10.5194/acp-17-13119-2017>, 2017.
- Dall’Osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Ström, J., Hansson, H.-C., Yoon, Y. J., Park, K.-T., Becagli, S., Udisti, R., Onasch, T., O’Dowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to  
470 atmospheric new particle formation, *Sci. Rep.*, 7, 3318, <http://doi.org/10.1038/s41598-017-03328-1>, 2017.
- Eckhardt, S., Quennehen, B., Olivie, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Lund, M. T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I. E., Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J.-C., Rumbold, S. T., Schulz, M., Sharma, S., Skeie, R. B., Skov, H., Uttal, T., von Salzen, K., and Stohl, A.: Current model capabilities for simulating black carbon and sulfate  
475 concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive measurement data set, *Atmos. Chem. Phys.*, 15, 9413-9433, <https://doi.org/10.5194/acp-15-9413-2015>, 2015.
- Fiddes, S. L., Woodhouse, M. T., Nicholls, Z., Lane, T. P., and Schofield, R.: Cloud, precipitation and radiation responses to large perturbations in global dimethyl sulfide, *Atmos. Chem. Phys.*, 18, 10177-10198, <https://doi.org/10.5194/acp-18-10177-2018>, 2018.
- 480 Fyfe, J. C., Derksen, C., Mudryk, L., Flato, G. M., Santer, B. D., Swart, N. C., Molotch, N. P., Zhang, X., Wan, H., Arora, V. K., Scinocca, J., and Jiao, Y.: Large near-term projected snowpack loss over the western United States, *Nat. Commun.*, 8, 14996, doi: 10.1038/ncomms14996, 2017.
- Gabric, A. J., Qu, B., Matrai, P., and Hirst, A. C.: The simulated response of dimethylsulphide production in the  
485 Arctic Ocean to global warming, *Tellus*, 57B, 391–403, 2005.
- Ghahremaninezhad, R., Norman, A.-L., Abbatt, J. P. D., Levasseur, M., and Thomas, J. L.: Biogenic, anthropogenic and sea salt sulfate size-segregated aerosols in the Arctic summer, *Atmos. Chem. Phys.*, 16, 5191–5202, <https://doi.org/10.5194/acp-16-5191-2016>, 2016.
- 490 Galí, M., Levasseur, M., Devred, E., Simó, R., and Babin, M.: Sea-surface dimethylsulfide (DMS) concentration from satellite data at global and regional scales, *Biogeosciences*, <https://doi.org/10.5194/bg-2018-18>, 2018.
- Galí, M., Devred, E., Babin, M., and Levasseur, M.: Decadal increase in Arctic dimethylsulfide emission, *submitted*, 2019
- Gagné, M.-É., Gillett, N. P., and Fyfe, J. C.: Impact of aerosol emission controls on future Arctic sea ice cover, *Geophys. Res. Lett.*, 42, 8481–8488, doi:10.1002/2015GL065504, 2015.

- 495 Ghan, S. J., Leung, L. R., Easter, R. C., and Abdul-Razzak, H.: Prediction of cloud droplet number in a general circulation model, *J. Geophys. Res.*, 102, D18, pp. 21777-21794, 1997.
- Gourdal, M., Lizotte, M., Massé, G., Gosselin, M., Poulin, M., Scarratt, M., Charette, J., and Levasseur, M.: Dimethyl sulfide dynamics in first-year sea ice melt ponds in the Canadian Arctic Archipelago, *Biogeosciences*, 15, 3169-3188, <https://doi.org/10.5194/bg-15-3169-2018>, 2018.
- 500 Grandey, B. S., and Wang, C.: Enhanced marine sulphur emissions offset global warming and impact rainfall, *Sci. Rep.*, 5, 13055, doi:10.1038/srep13055, 2015.
- Harada, N.: Review: potential catastrophic reduction of sea ice in the western Arctic Ocean: its impact on biogeochemical cycles and marine ecosystems, *Glob. Planet. Chang.*, 136, 1–17, 2016.
- Hayashida, H., Steiner, N., Monahan, A., Galindo, V., Lizotte, M., and Levasseur, M.: Implications of sea-ice biogeochemistry for oceanic production and emissions of dimethyl sulfide in the Arctic, *Biogeosciences*, 14, 3129-3155, <https://doi.org/10.5194/bg-14-3129-2017>, 2017.
- 505 Jahne, B., Munnich, K.O., Bosinger, R., Dutzi, A., Huber, W., and Libner, P.: On the parameters influencing air-water gas exchange, *J. Geophys. Res.*, 92, 1937-1949, 1987.
- Kettle, A. J., and Andreae, M. O.: Flux of dimethylsulfide from the oceans: A comparison of updated data sets and flux models, *J. Geophys. Res.-Atmos.*, 105, 26793–26808, doi:10.1029/2000JD900252, 2000.
- 510 Kooperman, G. J., Pritchard, M. S., Ghan, S. J., Wang, M., Somerville, R. C. J., and Russell, L. M.: Constraining the influence of natural variability to improve estimates of global aerosol indirect effects in a nudged version of the Community Atmosphere Model 5, *J. Geophys. Res.*, 117, D23204, doi:10.1029/2012JD018588, 2012.
- Krouse, H. R., and Grinenko, V. A.: Stable isotopes: natural and anthropogenic sulphur in the environment, Vol. 43, 515 Wiley, Chichester, UK, 1991.
- Kulmala, M., Laaksonen, A., and Pirjola, L.: Parameterizations for sulfuric acid/water nucleation rates, *J. Geophys. Res.*, 103, 8301-8307, 1998.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017-7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- 520 Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cycles*, 25, GB1004, doi:10.1029/2010GB003850, 2011.
- 525 Leaitch, W. R., Korolev, A., Aliabadi, A. A., Burkart, J., Willis, M. D., Abbatt, J. P. D., Bozem, H., Hoor, P., Köllner, F., Schneider, J., Herber, A., Konrad, C., and Brauner, R.: Effects of 20–100 nm particles on liquid clouds in the clean summertime Arctic, *Atmos. Chem. Phys.*, 16, 11107-11124, <https://doi.org/10.5194/acp-16-11107-2016>, 2016.
- 530 Leaitch, W. R., Sharma, S., Huang, L., Macdonald, A. M., Toom-Sauntry, D., Chivulescu, A., von Salzen, K., Pierce, J. R., Shantz, N. C., Bertram, A., Schroder, J., Norman, A.-L., and Chang R. Y.-W.: Dimethyl sulphide control



- of the clean summertime Arctic aerosol and cloud, *Elementa: Science of the Anthropocene*, 1, 000017, doi:10.12952/journal.elementa.000017, 2013.
- 535 Levasseur, M.: Impact of Arctic meltdown on the microbial cycling of sulphur. *Nature Geoscience* 6(9), 691-700, 2013
- Ma, X., von Salzen, K., and Cole, J.: Constraints on interactions between aerosols and clouds on a global scale from a combination of MODIS-CERES satellite data and climate simulations, *Atmos. Chem. Phys.*, 10, 9851-9861, 2010.
- 540 Martin, T., Steele, M., and Zhang, J.: Seasonality and long-term trend of Arctic Ocean surface stress in a model. *Journal of Geophysical Research: Oceans*, 119(3), 1723-1738, 2014.
- Mahmood, R., von Salzen, K., Flanner, M., Sand, M., Langner, J., Wang, H., and Huang, L.: Seasonality of global and Arctic black carbon processes in the Arctic Monitoring and Assessment Programme models, *J. Geophys. Res. Atmos.*, 121, doi:10.1002/2016JD024849, 2016.
- 545 Memery, L., and Merlivat L.: Modeling of gas flux through bubbles at the air-water interface, *Tellus Ser. B*, 37, 272-285, 1985.
- Merlivat, L., and Memery L.: Gas exchange across an air-water interface: Experimental results and modeling of bubble contribution to transfer, *J. Geophys. Res.*, 88, 707-724, 1983.
- 550 Monahan, E. C., and Spillane M. C.: The role of whitecaps in air-sea gas exchange, in *Gas Transfer at Water Surfaces*, edited by W. Brutsaert, and G.H. Jirka, pp. 495-504, D. Reidel, Norwell, Mass., 1984.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, <https://doi.org/10.1038/nature08823>, 2010.
- 555 Mungall, E. L., Croft, B., Lizotte, M., Thomas, J. L., Murphy, J. G., Levasseur, M., Martin, R. V., Wentzell, J. J. B., Liggio, J., and Abbatt, J. P. D.: Dimethyl sulfide in the summertime Arctic atmosphere: measurements and source sensitivity simulations, *Atmos. Chem. Phys.*, 16, 6665–6680, <https://doi.org/10.5194/acp-16-6665-2016>, 2016.
- 560 Nicolaus, M., Katlein, C., Maslanik, J., and Hendricks, S.: Changes in Arctic sea ice result in increasing light transmittance and absorption, *Geophys. Res. Lett.*, 39, L24501, <http://dx.doi.org/10.1029/2012GL053738>, 2012.
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cy.*, 14, 373y., 14doi:10.1029/1999GB900091, 2000.
- Norman, A. L., Barrie, L. A., Toom-Sauntry, D., Sirois, A., Krouse, H. R., Li, S. M., and Sharma, S.: Sources of aerosol sulphate at Alert: Apportionment using stable isotopes, *J. Geophys. Res.*, 104, 11619-11631, 1999.
- 565 Nöthig, E.-M., Bracher, A., Engel, A., Metfies, K., Niehoff, B., Peeken, I., Bauerfeind, E., Cherkasheva, A., Gäbler-Schwarz, S., Hardge, K., Kiliyas, E., Kraft, A., MebrahtomKidane, Y., Lalande, C., Piontek, J., Thomisch, K., and Wurst, M.: Summertime plankton ecology in Fram Strait—a compilation of long- and short-term observations, *Polar Research*, 34:1, DOI: 10.3402/polar.v34.23349, 2015.

- 570 Peng, Y., Lohmann, U., and Leaitch, W. R.: Importance of vertical velocity variations in the cloud droplet nucleation process of marine stratus clouds, *J. Geophys. Res.*, 110, D21213, doi:10.1029/2004JD004922, 2005.
- Peng, Y., von Salzen, K., and Li, J.: Simulation of mineral dust aerosol with Piecewise Log-normal Approximation (PLA) in CanAM4-PAM, *Atmos. Chem. Phys.*, 12, 6891-6914, doi:10.5194/acp-12-6891-2012, 2012.
- Pithan, F., and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models. *Nature Geosci.*, 7, 181–184, 2014.
- 575 Quinn, P. K., and Bates, T. S.: The case against climate regulation via oceanic phytoplankton sulphur emissions, *Nature*, 480, 51–56, doi:10.1038/nature10580, 2011.
- Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction of marine cloud condensation nuclei made up of sea spray aerosol, *Nature Geoscience*, 10(9), 674, 2017.
- 580 Rainville, L., Lee, C. M., and Woodgate, R. A.: Impact of wind-driven mixing in the Arctic Ocean, *Oceanography*, 24(3), 136-145, 2011. Ridley, J. K., Ringer, M. A., and Sheward, R. M.: The transformation of Arctic clouds with warming, *Climatic Change*, 139, 325-337, doi:10.1007/s10584-016-1772-4, 2016.
- Saha, S., and Coauthors .: The NCEP Climate Forecast System Version 2, *J. Climate*, 27, 2185-2208, doi: 10.1175/JCLI-D-12-00823.1, 2014.
- 585 Schwinger, J., Tjiputra, J., Goris, N., Six, K. D., Kirkevåg, A., Seland, Ø., Heinze, C., and Ilyina, T.: Amplification of global warming through pH dependence of DMS production simulated with a fully coupled Earth system model, *Biogeosciences*, 14, 3633-3648, <https://doi.org/10.5194/bg-14-3633-2017>, 2017.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: From air pollution to climate change*, Wiley, New York, 1998.
- 590 Sharma, S., Chan, E., Ishizawa, M., Toom-Saunty, D., Gong, S.L., Li, S.M., Leaitch, R., Norman, A.L., Quinn, P.K., Bates, T. S., Levasseur, M., and Barrie, L. A.: Influence of transport and ocean ice extent on biogenic aerosol sulfur in the Arctic atmosphere, *J. Geophys. Res. Atmos.*, 117 doi:10.1029/2011JD017074, 2012.
- Sigmond, M., and Fyfe, J. C.: Tropical Pacific impacts on cooling North American winters, *Nat. Clim. Change*, 6, 970-974, doi:10.1038/nclimate3069, 2016.
- 595 Simó, R.: Production of atmospheric sulfur by oceanic plankton: biogeochemical, ecological and evolutionary links. *Trends in Ecology & Evolution*, 16(6), 287-294, 2001.
- Six, K. D., Kloster, S., Ilyina, T., Archer, S. D., Zhang, K., and Maier-Reimer, E.: Global warming amplified by reduced sulphur fluxes as a result of ocean acidification, *Nat. Clim. Change*, 3, 975–978, doi:10.1038/nclimate1981, 2013.
- 600 Simó, R., and Dachs, J.: Global ocean emission of dimethylsulfide predicted from biogeophysical data, *Global Biogeochemical Cycles*, 16(4), 26-1, 2002.
- Soden, B. J., Broccoli, A. J., and Hemler, R. S.: On the use of cloud forcing to estimate cloud feedback, *J. Climate*, 17, 3661–3665, 2004.
- Soltwedel, T., Bauerfeind, E., Bergmann, M., Bracher, A., Budaeva, N., Busch, K., Cherkasheva, A., Fahl, K., Grzelak, K., Hasemann, C., Jacob, M., Kraft, A., Lalande, C., Metfies, K., Nöthig, E.-M., Meyer, K., Quéric, N.-V., Schewe, I., Włodarska-Kowalczyk M., Klages, M.: Natural variability or anthropogenically-induced
- 605

- variation? Insights from 15 years of multidisciplinary observations at the arctic marine LTER site HAUSGARTEN, *Ecological Indicators*, 65, 89-102, <https://doi.org/10.1016/j.ecolind.2015.10.001>, 2016.
- 610 Stefels, J., Steinke, M., Turner, S., Malin, G., and Belviso, S.: Environmental constraints on the production and removal of the climatically active gas dimethylsulphide (DMS) and implications for ecosystem modelling, *Biogeochemistry*, 83, 245-275, <https://doi.org/10.1007/s10533-007-9091-5>, 2007.
- Stroeve, J. C., Kattsov, V., Barrett, A., Serreze, M., Pavlova, T., Holland, M., and Meier, W. N.: Trends in Arctic sea ice extent from CMIP5, CMIP3 and observations, *Geophys. Res. Lett.*, 39, L16502, doi:10.1029/2012GL052676, 2012.
- 615 Struthers, H., Ekman, A. M. L., Glantz, P., Iversen, T., Kirkevåg, A., Mårtensson, E. M., Seland, Ø., and Nilsson, E. D.: The effect of sea ice loss on sea salt aerosol concentrations and the radiative balance in the Arctic, *Atmos. Chem. Phys.*, 11, 3459-3477, <https://doi.org/10.5194/acp-11-3459-2011>, 2011.
- Tesdal, J.-E., Christian, J. R., Monahan, A. H., and von Salzen, K.: Evaluation of diverse approaches for estimating sea-surface DMS concentration and air-sea exchange at global scale, *Environmental Chemistry*, doi:10.1071/EN14255, 2016a.
- 620 Tesdal, J.-E., Christian, J. R., Monahan, A. H., and von Salzen, K.: Sensitivity of modelled sulfate aerosol and its radiative effect on climate to ocean DMS concentration and air-sea flux, *Atmos. Chem. Phys.*, 16, 10847-10864, doi:10.5194/acp-16-10847-2016, 2016b.
- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Rast, S., Devasthale, A., Kloster, S., Feichter, J., and Lenton, T. M.: Quantification of DMS aerosol-cloud-climate interactions using the ECHAM5-HAMMOZ model in a current climate scenario, *Atmos. Chem. Phys.*, 10, 7425-7438, doi:10.5194/acp-10-7425-2010, 2010.
- 625 Vancoppenolle, M., Bopp L., Madec G., Dunne J., Ilyina T., Halloran P. R., and Steiner N.: Future Arctic Ocean primary productivity from CMIP5 simulations: Uncertain outcome, but consistent mechanisms, *Global Biogeochem. Cycles*, 27, 605-619, doi:10.1002/gbc.20055, 2013.
- 630 von Salzen, K., Leighton, H. G., Ariya, P. A., Barrie, L. A., Gong, S. L., Blanchet, J.-P., Spacek, L., Lohmann, U., and Kleinman, L. I.: Sensitivity of sulphate aerosol size distributions and CCN concentrations over North America to SO<sub>x</sub> emissions and H<sub>2</sub>O<sub>2</sub> concentrations, *J. Geophys. Res. Atmos.*, 105, 9741-9765, doi:10.1029/2000JD900027, 2000.
- von Salzen, K., Scinocca, J. F., McFarlane, N. A., Li, J., Cole, J. N. S., Plummer, D., Verseghy, D., Reader, M. C., Ma, X., Lazare, M., and Solheim, L.: The Canadian Fourth Generation Atmospheric Global Climate Model (CanAM4). Part I: Representation of physical processes, *Atmos. Ocean*, 51, 104-125, doi:10.1080/07055900.2012.755610, 2013.
- 635 von Salzen, K.: Piecewise log-normal approximation of size distributions for aerosol modelling, *Atmos. Chem. Phys.*, 6, 1351-1372, 2006.
- 640 Wanninkhof, R., Asher, W. E., Ho, D. T., Sweeney, C., and McGillis, W. R.: Advances in quantifying air-sea gas exchange and environmental forcing, *Annu. Rev. Mar. Sci.*, 1, 213-44, 2009. Willis, M. D., Burkart, J., Thomas, J. L., Köllner, F., Schneider, J., Bozem, H., Hoor, P. M., Aliabadi, A. A., Schulz, H., Herber, A. B., Leitch, W. R., and Abbatt, J. P. D.: Growth of nucleation mode particles in the summertime Arctic: a case study, *Atmos. Chem. Phys.*, 16, 7663-7679, doi:10.5194/acp-16-7663-2016, 2016.

645 Winter, A., Henderiks, J., Beaufort, L., Rickaby, R. E. M., and Brown, C. W.: Poleward expansion of the coccolithophore *Emiliana huxleyi*, *Journal of Plankton Research*, 36, 316–325, 2014.

Woodhouse, M. T., Carslaw, K. S., Mann, G. W., Vallina, S. M., Vogt, M., Halloran, P. R., and Boucher, O.: Low sensitivity of cloud condensation nuclei to changes in the sea-air flux of dimethyl-sulphide, *Atmos. Chem. Phys.*, 10, 7545–7559, doi:10.5194/acp-10-7545-2010, 2010.

650 Wyslouzil, B. E., Seinfeld, J. H., Flagan, R. C., and Okuyama, K.: Binary nucleation in acid water-system. 1. Methanesulfonic-acid water, *J. Chem. Phys.*, 94 (10), 6827-6841, 1991.

Yool, A., Popova, E. E., and Coward, A. C.: Future change in ocean productivity: Is the Arctic the new Atlantic?, *J. Geophys. Res. Oceans*, 120, 7771-7790, doi:10.1002/2015JC011167, 2015.

655 Zhang, L., Gong, S., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environment*, 35, 549-560, 2001.

Zhang, Y., McMurry, P. H., Yu, F., and Jacobson, M. Z.: A comparative study of nucleation parameterizations: 1. Examination and evaluation of the formulations, *J. Geophys. Res.*, 115, D20212, doi:10.1029/2010JD014150, 2010.

660

**Table 1.** Arctic mean surface seawater DMS concentration, total sulfur emission flux, and associated mean biogenic sulfate burden for 2000 (2050)


	DMS (nM)	Emissions (TgS/yr)	SO <sub>4</sub> <sup>2-</sup> Burden (kilotonnes)
CLIM	1.96	0.24 (0.32)	2.13 (2.58)
10×CLIM	19.58	2.41 (3.22)	20.05 (20.55)
UNFM	16.88	1.88 (2.87)	16.82 (16.64)
10×SAT	11.19	1.31 (1.92)	10.31 (11.63)

665

**Table 2.** Arctic mean aerosol nucleation rate, CDNC, and cloud radiative forcing in 2000 (2050) associated with emissions of DMS in the Arctic.

	Nucleation rate (×10 <sup>6</sup> m <sup>-2</sup> s <sup>-1</sup> )	CDNC (×10 <sup>6</sup> m <sup>-3</sup> )	Cloud radiative forcing (Wm <sup>-2</sup> )
CLIM	0.02517 (0.09294)	0.34716 (0.81023)	-0.13 (-0.27)
10×CLIM	0.41751 (0.98998)	2.9886 (4.1781)	-0.75 (-1.59)
UNFM	0.27358 (0.62366)	2.3019 (3.4893)	-0.40 (-0.98)
10×SAT	0.20618 (0.54223)	1.7853 (2.4045)	-0.55 (-1.18)

670



710 Figure 1: (a) Locations for Alert (red circle) and ship-based  $\text{nss-SO}_4^{2-}$  observations used for comparisons with CanAM4.3? model results. (b) Comparison of model and observed  $\text{nss-SO}_4^{2-}$ . For ship based observations, the size of the markers represent percentage of contribution of DMS to total  $\text{nss-SO}_4^{2-}$  derived from model results. For Alert, the percentage contribution to total  $\text{nss-SO}_4^{2-}$  is based on isotopic composition.

715





Figure 2: Multi-year mean monthly concentrations of total (a), fossil-fuel (b), biogenic (c) sulfate concentrations from simulations and observed nss-SO<sub>4</sub><sup>2-</sup> during 1994 to 2002 at Alert, (d) relative contribution of DMS source to total sulfate concentration at Alert. The whiskers represent minimum and maximum and the horizontal line inside the box represents the mean for the whole period. The box height represents the interquartile range of 25<sup>th</sup> and 75<sup>th</sup> percentiles. Unit: ng m<sup>-3</sup>



730

735



740


Figure 3: Annual mean distribution of surface seawater DMS concentration used in sensitivity simulations; (a) L10 DMS climatology, (b) L10 DMS climatology multiplied by 10 in the Arctic region, (c) uniform distribution of DMS, (d) satellite based DMS climatology multiplied by 10. The bottom panel shows zonal mean results.

745

750

755

760




765

Figure 4: DMS emission fluxes for the two different periods, similar to Fig. 3. Zonal mean results 2000 and 2050 are shown in the bottom panel.

770

775

780



785 **Figure 5: Biogenic sulfate burden due to Arctic DMS emissions for each of the 4 scenarios. The stippling represents regions where the burden difference relative to CNTRL is significant at 95% confidence level. Zonal mean results for 2000 and 2050 are shown in the bottom panel.**

790




Figure 6: Change in cloud droplet number concentration at first model level above the surface due to Arctic DMS emissions (relative to CNTRL). Stippling represents change significant at 95% confidence level. Zonal mean results are shown in the bottom panel.


800

805

810



815




820 **Figure 7: Cloud radiative forcing in the Arctic due to ocean DMS emissions. Stippling represents radiative forcing significant at 95% confidence level. Zonal mean results are shown in the bottom panel.**

825

830

835



840

**Figure 8: Relationship between annual Arctic mean DMS emission fluxes, sulfate burden, CDNC, and cloud forcing and the mean DMS concentration in the Arctic. The vertical lines represent 95% confidence interval based on two-tailed t-test. Dotted lines represent regression between four scenarios.**

845