

Figure S1. Spatial distribution of NH_3 column densities in mainland China observed by IASI sensor from 2008 to 2016. The area of interest in this study includes Beijing (A), Tianjin (B), Hebei (C), Shanxi (D), Shandong (E) and Henan (F) Provinces, as marked in the black rectangle. The small islands in South China Sea are not included in the figure.




Figure S2. Spatial distribution of satellite observations (a) and modelled tropospheric NH₃ column densities (b) for the year of 2016; (c) comparison of the regional mean NH₃ column densities between observations and simulations among different seasons in 2016.




Figure S3. Comparison of modelled PM_{2.5} sulfate (a), nitrate (b) and ammonium (c) concentrations with corresponding daily measurements of PM_{2.5} from 2014 to 2016 over North China Plain. The model results were extracted from the first model level (~25m aboveground) and averaged at one day. The Pearson correlation coefficients and normalized mean bias are shown inset.




Figure S4. The simulated annual-mean concentrations of PM_{2.5}-nitrate (µg/m³) over Northern China in recent years.

Table S1 Annual consumption of synthetic fertilizers, populations of major livestock animals and the proportions of the intensive system in livestock animal rearing in Northern China from 2008 to 2016.

	2008	2009	2010	2011	2012	2013	2014	2015	2016
Fertilizers (Ton)									
ammonium bicarbonate	146	128	99	108	107	99	61	54	42
Urea	477	489	520	506	507	509	538	532	523
Compound Fertilizer	120	127	135	140	145	149	160	163	168
Animals (Million)									
Beef	18.2	17.4	16.1	15.1	14.7	14.6	14.6	15.0	14.6
Dairy	3.5	3.6	4.3	4.7	4.9	4.8	5.1	5.0	4.8
Goat	49.2	45.4	43.4	42.8	42.1	42.1	43.0	44.2	37.8
Sheep	17.3	19.7	19.3	20.6	21.3	22.1	23.2	23.1	26.9
Pig	132	139	143	142	151	157	164	160	155
Poultry	3056	3141	3209	3369	3658	3605	3424	3504	3635
Proportion of the intensive system in livestock animal rearing (%)									
	28	32	35	37	39	41	43	46	48