

## Responses to Referee Comments

*The original referee comments are shown in italicized, black font.*

**Our replies are shown in bold, blue font.**

**We are very grateful to both reviewers for taking another look at our heavily revised manuscript. We respond below to additional, very helpful, suggestions from Referee #1.**

*Anonymous Referee #1*

*This paper has been significantly improved by the authors. It is shorter and more focused. The conclusions are supported by the work but it is still hampered by its use of models which are at least one generation behind the current state of the science. I don't see any obvious development in the intervening period that would completely invalidate the conclusions but it feels a little odd to be using results from these simulations.*

**Thank you for your guidance that helped us to focus and shorten the paper. We understand that these simulations are older than the reviewer would like. To our knowledge, this is the first application of the HTAP1 dataset to examine multi-model PAN source-receptor relationships, and this set of model simulations remains the most comprehensive available (i.e., a large set of sensitivity simulations performed consistently across 10+ models) for this type of analysis in search of observational constraints.**

*I would suggest that the conclusions are toned down a little bit. This feels like it is trying to prepare for grant funding to support PAN measurements into mountaintop sites. These observations are useful in the greater scheme of things but I'm not I would be as categorical about their value as the authors.*

**We have revised the conclusions and tried to “tone down” the recommendations for future measurements. We removed the final sentence of the conclusions, and moved the preceding sentences up to the second paragraph of the conclusions such that the paper now ends with the point about improving documentation of VOC emissions (including speciation) and their role as precursors to PAN and ozone in future multi-model efforts. We also deleted the sentences from the second paragraph that began to spell out “how-to” proceed with long-term PAN measurements. This second paragraph of the conclusions now reads:**

**“Establishing the strongest constraints possible on simulated SRRs for PAN and O<sub>3</sub> will require (1) measurements and simulations with chemical transport models that coincide, and (2) a sufficiently long measurement record to build a climatology suitable for evaluating chemistry-climate models that generate their own meteorology. Repeated sampling for the month of April may be sufficient to provide constraints on model responses to changes in anthropogenic emissions. PAN measurements over multiple seasons are necessary to evaluate model responses of PAN to climate change (e.g., by changing temperature and weather-sensitive precursor emissions) and the resulting influence on atmospheric O<sub>3</sub> and oxidizing capacity (e.g., Doherty et al., 2013). For example, changes in meteorology and biomass burning (Fischer et al., 2011; Zhu et al., 2015) such as those driven by ENSO (Koumoutsaris et al., 2008), as well as biogenic and lightning sources (Payne et al., 2017) vary from year to year and are expected to change as climate warms.”**

*There are a number of typos in the paper which reflect the author's corrections but it would be useful if these were corrected before publication.*


**We hope we have now caught all the typos, but will continue to read carefully for typos at the proofs stage as well.**

*The figures are at rather low resolution. Figure 2 especially looks very fuzzy in the version I have.*

**The resolution looks much better in the original eps files that we will be uploading directly for the final publication, so we hope that these issues will be resolved.**

Figure 3 would look better if the blue (multi-model mean) and the red (observations) were emphasised over the individual models. At the moment the most important symbols (blue and red) symbols are the least obvious.

**We have worked to improve this figure. We now increase the symbol size and change to solid filled circles for the multi-model mean (blue) and the observations (red, except for year 2001 at Schauinsland which we show in blue to emphasize that this point is the best comparison point for the multi-model mean):**



**Figure 3. April mean PAN abundances (ppt) simulated (black symbols, one per model as defined in Table 1; blue circles offset to the left show multi-model mean values) and measured at northern mid-latitude mountaintop sites (red circles offset to the right of the model values; the year 2001 value, which corresponds to the meteorological year used by most of the models, at Schauinsland is shown in blue).**

# Peroxy acetyl nitrate (PAN) measurements at northern mid-latitude mountain sites in April: A constraint on continental source-receptor relationships

Arlene M. Fiore<sup>1,2</sup>, Emily V. Fischer<sup>3</sup>, George P. Milly<sup>2</sup>, Shubha Pandey Deolal<sup>4</sup>, Oliver Wild<sup>5</sup>, Dan Jaffe<sup>6,7</sup>, Johannes Staehelin<sup>4</sup>, Olivia E. Clifton<sup>1,2</sup>, Dan Bergmann<sup>8</sup>, William Collins<sup>9</sup>, Frank Dentener<sup>10</sup>, Ruth M. Doherty<sup>11</sup>, Bryan N. Duncan<sup>12</sup>, Bernd Fischer<sup>13</sup>, Stefan Gilge<sup>14,15</sup>, Peter G. Hess<sup>16</sup>, Larry W. Horowitz<sup>17</sup>, Alexandru Lupu<sup>18,19</sup>, Ian MacKenzie<sup>11</sup>, Rokjin Park<sup>20</sup>, Ludwig Ries<sup>21</sup>, Michael G. Sanderson<sup>22</sup>, Martin G. Schultz<sup>23</sup>, Drew T. Shindell<sup>24</sup>, Martin Steinbacher<sup>25</sup>, David S. Stevenson<sup>11</sup>, Sophie Szopa<sup>26</sup>, Christoph Zellweger<sup>25</sup>, Guang Zeng<sup>27</sup>

<sup>1</sup>Department of Earth and Environmental Science, Columbia University, Palisades, NY, 10964, U.S.A.

<sup>2</sup>Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY, 10964, U.S.A.

<sup>3</sup>Department of Atmospheric Science, Colorado State University, Fort Collins, CO, 80521, U.S.A.

<sup>4</sup>Institute for Atmospheric and Climate Science, ETH Zürich, Switzerland

<sup>5</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

<sup>6</sup>School of STEM, University of Washington, Bothell, WA, 98011, U.S.A.

<sup>7</sup>Department of Atmospheric Science, University of Washington, Seattle, WA, 98195, U.S.A.

<sup>8</sup>Lawrence Livermore National Laboratory, Livermore, CA, 94550, U.S.A.

<sup>9</sup>Department of Meteorology, University of Reading, Reading, RG6 6BB, UK

<sup>10</sup>European Commission, Joint Research Centre, Ispra, I-21027, Italy

<sup>11</sup>School of GeoSciences, The University of Edinburgh, Edinburgh, EH9 3FF, UK

<sup>12</sup>Atmospheric Chemistry and Dynamics Laboratory, NASA GSFC, Greenbelt, MD 20720, U.S.A.

<sup>13</sup>Federal Environment Agency (UBA), Schaulinsland, 79254, Germany

<sup>14</sup>Meteorological Observatory Hohenpeissenberg, German Meteorological Service (DWD), Hohenpeissenberg, DE

<sup>15</sup>now at DWD, Research Center Human Biometeorology, Freiburg, DE

<sup>16</sup>Department of Biological and Environmental Engineering, Cornell University, Ithaca, NY, 14853, U.S.A.

<sup>17</sup>Geophysical Fluid Dynamics Laboratory, National Oceanic and Atmospheric Administration, Princeton, NJ, 08540, U.S.A.

<sup>18</sup>Centre for Research in Earth and Space Science, York University, Toronto, M3J 1P3, Canada

<sup>19</sup>now at Air Quality Research Division, Environment and Climate Change Canada, Toronto, M3H 5T4, Canada

<sup>20</sup>School of Earth and Environmental Sciences, Seoul National University, Seoul, 08826, Republic of Korea

<sup>21</sup>II4.5.7, German Environment Agency (UBA), Zugspitze, 82475, Germany

<sup>22</sup>Met Office, Exeter, EX1 3PB, UK.

<sup>23</sup>Jülich Supercomputing Centre, Forschungszentrum Jülich, 52425 Jülich, Germany

<sup>24</sup>Nicholas School of the Environment, Duke University, Durham, NC, 27708, U.S.A.

<sup>25</sup>Laboratory for Air Pollution / Environmental Technology, Empa – Swiss Federal Laboratories for Materials Science and Technology, Dübendorf, CH-8600, Switzerland

<sup>26</sup>Laboratoire des Sciences du Climat et de l'Environnement, Institut Pierre Simon Laplace, CEA/CNRS/UVSQ, Gif sur Yvette, France

<sup>27</sup>National Institute of Water and Atmospheric Research, Wellington, 6021, New Zealand

*Correspondence to:* Arlene M. Fiore (amfiore@ldeo.columbia.edu)

**Abstract.** Abundance-based model evaluations with observations provide critical tests for the simulated mean state in models of intercontinental pollution transport, and under certain conditions may also offer

constraints on model responses to emission changes. We compile multi-year measurements of peroxy acetyl nitrate (PAN) available from five mountaintop sites and apply them in a proof of concept approach that exploits an ensemble of global chemical transport models (HTAP1) to identify an observational “emergent constraint”. In April, when the signal from anthropogenic emissions on PAN is strongest, simulated PAN at northern mid-latitude mountaintops correlates strongly with PAN source-receptor relationships (the response to 20% reductions in precursor emissions within northern mid-latitude continents; hereafter, SRRs). This finding implies that PAN measurements can provide constraints on PAN SRRs by limiting the SRR range to that spanned by the subset of models simulating PAN within the observed range. In some cases, regional anthropogenic volatile organic compound (AVOC) emissions, tracers of transport from different source regions, and SRRs for ozone also correlate with PAN SRRs. Given the large observed interannual variability in the limited available datasets, establishing strong constraints will require matching meteorology in the models to the PAN measurements. Application of this evaluation approach to the chemistry-climate models used to project changes in atmospheric composition will require routine, long-term mountaintop PAN measurements to discern both the climatological SRR signal and its inter-annual variability.

## 1 Introduction

Peroxy acetyl nitrate (PAN) is produced alongside ozone ( $O_3$ ) from photochemical reactions involving precursor emissions of nitrogen oxides ( $NO_x$ ) and non-methane volatile organic compounds (VOC). Once ventilated from a source region to the free troposphere where it is more stable at colder temperatures, PAN can be efficiently transported throughout the hemisphere (Singh, 1987; Singh and Hanst, 1981). When a PAN-containing free tropospheric air mass subsides, PAN thermally decomposes to release  $NO_x$  and can thus facilitate  $O_3$  formation far downwind (Wild et al., 1996; Schultz et al., 1999; Jaeglé et al., 2003; Kotchenruther et al., 2001a; Hudman et al., 2004). Both PAN and  $O_3$  distributions over any northern mid-latitude region reflect the combined influence of production from sources within the region and transport from outside that region. At northern mid-latitudes, the intercontinental influence from anthropogenic emissions on surface  $O_3$  levels is largest during spring (e.g., HTAP 2010) and occurs via at least two pathways: (1)  $O_3$  can be produced within a polluted continental boundary layer, ventilated to the free troposphere and efficiently transported to other continents; and (2)  $O_3$  can be produced in transit from the export and subsequent chemical evolution of PAN and other precursors. Below, we examine the extent to which springtime PAN observations at northern mid-latitude mountaintop sites can be used to constrain the spread in multi-model estimates of source-receptor relationships (SRRs), where the sources are continental-scale regions and the receptors are the mountaintop sites, for both PAN and  $O_3$ .

Observations during several aircraft field campaigns in the Eastern Pacific and at mountain top sites in the Western U.S. and North Atlantic document efficient  $O_3$  production in the lower troposphere following subsidence of PAN-containing air masses (Fischer et al., 2010; Heald et al., 2003; Hudman et al., 2004;

Kotchenruther et al., 2001a,b; Val Martin et al., 2008; Zhang et al., 2008). When PAN decomposes in low- $\text{NO}_x$  regions of the atmosphere, the  $\text{NO}_x$  released can produce  $\text{O}_3$  up to eight times more efficiently than in polluted (high- $\text{NO}_x$ ) regions (Liang et al., 1998; Liu et al., 1987) and thus increase global  $\text{O}_3$  abundances (Moxim et al., 1996; Wang and Jacob, 1998), as  $\text{O}_3$  formation is  $\text{NO}_x$ -limited in most of the free troposphere (Chameides et al., 1992). The lifetime of PAN against thermal decomposition is about 1 hour at  $20^\circ\text{C}$ , and it approximately doubles for every  $4^\circ\text{C}$  decrease in temperature, leading to a lifetime of at least a month in the mid-troposphere during spring. This strong temperature dependence implies that a warmer climate will decrease PAN export from polluted continental boundary layers, although a rise in temperature-sensitive biogenic precursor emissions may temper this response (e.g. Doherty et al., 2013). Future projections of atmospheric composition under global change scenarios will thus benefit from a thorough understanding of the role PAN plays in transporting oxidized reactive nitrogen and thereby altering ozone production throughout the troposphere.

To better distinguish among disparate estimates for intercontinental  $\text{O}_3$  transport in the published literature, the Task Force on Hemispheric Transport of Air Pollution (HTAP) organized an international global modelling study, referred to here as HTAP1. The HTAP1 study identified a factor of two range across individual model estimates of surface  $\text{O}_3$  response to changes in anthropogenic precursor emissions from continental-scale, northern mid-latitude source regions (HTAP 2007; Fiore et al., 2009; HTAP 2010; Wild et al. 2012). The HTAP1 models do not distinguish between intercontinental  $\text{O}_3$  transport occurring due to  $\text{O}_3$  produced from PAN chemistry versus direct transport of  $\text{O}_3$  formed in a remote boundary layer, but other work indicates that both pathways contribute. Jaegle et al. (2003) find that 28% of the  $\text{O}_3$  in the Pacific Northwest free troposphere between 0-6 km is associated with PAN-to- $\text{NO}_x$  conversion, consistent with Jiang et al. (2016) who found that PAN produced from East Asian emissions and exported to the free troposphere contributes 35% and 25% in spring and summer, respectively, to the free tropospheric  $\text{O}_3$  abundance over western North America. Over East Asia, Lin et al. (2010) found that the export of PAN produced from European anthropogenic emission changes and subsequent downwind  $\text{O}_3$  formation contributed 20% of the spatially averaged response of surface  $\text{O}_3$  levels, and up to 50% of the  $\text{O}_3$  response at mountain sites.

In addition to the direct influence of PAN on intercontinental  $\text{O}_3$  transport, PAN may serve as a sensitive diagnostic of model uncertainties in  $\text{O}_3$  production chemistry and transport (Emmerson and Evans, 2009; Kuhn et al., 1998). Prior analysis of measurements and global model simulations suggests that PAN abundances at high altitude sites may be more sensitive than  $\text{O}_3$  itself to changes in precursor emissions (Fiore et al., 2011; Fischer et al., 2011; Jaffe et al., 2007). We interpret this stronger sensitivity of PAN than  $\text{O}_3$  to changes in precursor emissions as reflecting buffering of  $\text{O}_3$  by compensating changes to  $\text{O}_3$  losses, whereas PAN loss pathways are far less sensitive to changes in precursor emissions. PAN loss pathways include thermal decomposition (which dominates below approximately 7 km); photolysis in the

upper troposphere; and dry deposition within the boundary layer (Kirchner et al., 1999; Roberts, 2007; Turnipseed et al., 2006). All of the HTAP1 models include PAN formation, but the chemical mechanisms and kinetic rate coefficients differ, with likely implications for long-range transport (Emmerson and Evans, 2009; Knote et al., 2015). A prior multi-model study found that even with the same emissions, PAN differs widely across models, reflecting differences in simulated photochemistry (Emmons et al., 2015). While the absence of direct emissions and its low background make PAN a useful tracer of photochemistry, we note that O<sub>3</sub> typically responds more strongly to changes in NO<sub>x</sub> emissions, while PAN responds more strongly to changes in VOC emissions in many regions (Fischer et al., 2014; see their Figure 4).

A challenge in discriminating among model estimates of O<sub>3</sub> produced from different source regions is the lack of direct observational constraints on SRRs. For example, Fiore et al. (2009) did not find any relationship across models between their biases against surface O<sub>3</sub> observations and the strength of their response to emission changes. In the absence of an observable quantity to constrain these relationships, one approach is to identify an “emergent constraint” (Borodina et al., 2017), whereby a non-observable quantity correlates strongly across a multi-model ensemble with an observed variable. The inter-model range of the non-observable quantity is then narrowed by limiting it to the range encompassed by the models closest to the observed variable. This approach has gained traction for narrowing the spread across future climate projections (e.g., Hall and Qu, 2006; Cox et al. 2018). In light of its role as a proxy for ozone formation chemistry, its direct role in facilitating intercontinental ozone transport, and the large signature of PAN originating in the European boundary layer during spring found at Jungfraujoch (Pandey Deolal et al., 2013; 2014), we hypothesize that PAN measurements may offer much-needed constraints for discriminating across model estimates of intercontinental transport of PAN, and possibly O<sub>3</sub>. The number of models contributing to the HTAP1 study, which was designed to maximize comparability across individual model estimates of ozone responses to changes in precursor emissions within northern mid-latitude continental-scale source regions, offers an opportunity to evaluate this hypothesis.

We describe the HTAP1 model simulations, mountaintop measurements and our strategy to sample the models at these sites (Section 2) before illustrating our rationale for selecting the month of April to quantify the range of multi-model PAN distributions and PAN measurements at northern mid-latitude mountain sites (Section 3). We then borrow from the “emergent constraint” approach in climate science to show that correlations between simulated total PAN and SRRs for PAN are sufficiently strong as to permit PAN measurements at mountaintop sites (one in each of the three major mid-latitude source regions) to narrow the wide inter-model spread in estimates of PAN origin (Section 4). We further examine inter-model relationships between the simulated PAN SRRs at these three mountaintop sites and regional precursor emissions, and with a proxy for model transport (Section 5). Finally, we assess the relationship between PAN and O<sub>3</sub> SRRs (Section 6) and conclude with a summary and recommendations for future work based on our proof of concept analysis (Section 7).

## 2. Approach

### 2.1 HTAP1 model simulations

We use monthly mean PAN mixing ratios for the year 2001 simulated by fourteen global chemistry transport models (Table 1); the temporal resolution for three-dimensional chemical fields archived from the HTAP1 models is limited to monthly. We use four HTAP1 Source-Receptor (SR) simulations (Table 2): a base case (SR1) and three perturbation simulations in which anthropogenic O<sub>3</sub> precursor emissions (NO<sub>x</sub>, VOC, carbon monoxide and aerosols) are reduced simultaneously by 20% within East Asia (SR6EA), Europe and northern Africa (SR6EU), and North America (SR6NA). We calculate PAN SRRs by differencing the perturbation and base simulations (SR1-SR6XX), where XX refers to the region in which emissions of PAN precursors were decreased by 20%.

Of the models in Table 1, eleven used 2001 meteorological fields. Two models are chemistry-transport models coupled directly to a general circulation model forced by observed sea surface temperatures (STOC-HadAM3 and STOCHEM) and one model incorporates chemistry directly into a general circulation model (UM-CAM). We include these models as our evaluation compiles PAN measurements across several years (Section 2.2). The individual model specifications and emissions are described in Tables 1 and 2 of Fiore et al. (2009). For HTAP1, each model used its own emissions inventories (see Table A1 of Fiore et al., 2009); Fiore et al. (2009) provide emission totals within each HTAP1 source region for all (their Table A2) and anthropogenic (their Table A3) emissions of NO<sub>x</sub>, NMVOC, and CO. The relative inter-model spread in regional anthropogenic emissions is smallest for NO<sub>x</sub> emissions in EU and NA (<10%) and largest for VOC from EU (58%) (Fiore et al. 2009).

To separate the role of inter-model differences in transport from the combined impacts of inter-model differences in emissions and chemistry on simulated PAN at the mountaintop sites, we analyze an additional set of idealized tracer simulations available from eleven models (COfromXX in Table 1, where XX is the source region). In these simulations, a set of tagged carbon monoxide-like tracers are emitted, each from a single HTAP1 source region with a 50-day lifetime, and with identical emissions across models. Biomass burning emissions for the CO tracers are from GFED (van der Werf et al., 2006; 2010) and other emissions are from the RETRO project (Schultz et al., 2007; 2008). We refer to these tracers as “COfromEA”, “COfromNA”, and “COfromEU”, which denote the tracers emitted from EA, NA, and EU, respectively (Table 2; see also Doherty et al., 2013 and Shindell et al., 2008).

### 2.2 Multi-year PAN measurements at mountaintop sites and model sampling

To evaluate the HTAP1 models, we compiled April mean climatologies of lower tropospheric PAN measurements from northern mid-latitude mountain observatories (Table 3). Given the large interannual

variability in PAN abundances, we require at least two years of observations in April. PAN observations from Mount Bachelor (U.S.A.), Jungfrauoch (Switzerland), and Zugspitze (Schneefernerhaus), Hohenpeissenberg, and Schauinsland (all in Germany) meet these criteria. Taken together, these mountaintop measurements span 15 years, from 1995 to 2010 (Table 3), although only one site (Schauinsland) overlaps with the HTAP1 simulation year of 2001.

PAN was measured at all five mountain sites using gas chromatography with electron capture detection (ECD). A custom system using a Shimazu Mini-2 ECD was employed at Mount Bachelor (Fischer et al., 2010). The commercially available Meteorologie Consult (GmbH) system was used at the European sites (Zellweger et al., 2000). Calibrations generate PAN from the photolysis of excess acetone and NO in air (Warneck and Zerbach, 1992; Volz-Thomas et al., 2002). Reported detection limits are ~20 ppt for PAN measurements at Mount Bachelor, and ~50 ppt for the European sites, with total uncertainties of <10% (Fischer et al., 2010; Zellweger et al., 2003).

We include all available data at these sites without filtering for upslope winds or any other criteria. At Mount Bachelor, the cleanest of the 5 mountaintop sites (Supplemental Figure 1), Fischer et al. (2010) have shown that PAN mixing ratios are not primarily controlled by diurnal wind patterns, which lead to variations an order of magnitude smaller than the total observed range in measured PAN. When measurements fall below the detection limit, we include half of the detection limit. This assumption should not affect our conclusions as mountaintop sites generally sample free tropospheric air at night (e.g., Weiss-Penzias et al., 2004) but PAN values below the detection limit typically occur due to deposition in a shallow nocturnal boundary layer.

For comparison with the observations, we sample each model on its native grid (Table 1) at the horizontal grid cell containing the latitude and longitude of each mountain site. Orography at these mountain sites is poorly resolved at the relatively coarse HTAP1 model horizontal resolutions. This mismatch requires us to apply some approximations for vertical sampling. We convert the station altitude to an approximate pressure level by assuming a mean tropospheric temperature of 260 K, and a corresponding atmospheric scale height of 7.6 km. We then use monthly mean pressure fields from each model to linearly interpolate PAN based on the pressures of the two model grid cells that vertically bound the station pressure. While different sampling strategies may alter the exact value of simulated PAN and its comparison to observations, our primary interest is in the inter-model differences. Although the Zugspitze and Hohenpeissenberg sites fall within the same horizontal grid cell in the HTAP1 models, the station altitudes differ, so we consider the two sites separately.

Given that we seek constraints on intercontinental transport from the three major mid-latitude source regions, we conduct a more in-depth analysis at the highest altitude European site (Jungfrauoch), the most



likely of the available sites to measure PAN transported between continents in the free troposphere, as well as at Mount Bachelor in North America. At Jungfraujoch, we also evaluate SRRs in the models with an estimate of PAN originating in the European boundary layer based on an analysis of 20-day back trajectories (Pandey Deolal et al., 2013). We conduct a proof of concept analysis at Mount Waliguan in Asia (36.28°N, 100.90°E, 3816 m) to assess the potential for future PAN measurements at this site to narrow the inter-model range in SRRs. Short-term measurements have previously been collected at this site (Xue et al., 2011). While aircraft and satellite observations have advanced the understanding of the chemistry and dynamics of individual PAN plumes using models that archived higher temporal frequency chemical fields (e.g., Alvarado et al., 2010; Payne et al., 2014; Emmons et al., 2015), their limited temporal coverage is not well suited for comparison with the HTAP1 monthly mean PAN mixing ratios.

### **3. Modeled and measured lower tropospheric PAN at northern mid-latitudes in April**

Our goal is to assess the potential for mountaintop PAN measurements to discriminate among model estimates of PAN and O<sub>3</sub> produced by regional anthropogenic emissions and transported to the mountaintop sites. We thus focus our analysis on April when measured PAN reaches its seasonal maximum (Penkett and Brice, 1986; Singh and Salas, 1989; Bottenheim et al., 1994; Schmitt and Volz-Thomas, 2004; Supplemental Figure 1) and when the HTAP1 models indicate that the production of PAN from the EA, EU, and NA source regions dominates total simulated PAN (Figure 1). April thus offers the strongest possible signal of the influence of anthropogenic emissions from these three northern mid-latitude source regions in the mountaintop measurements.

Figure 2 shows the spatial distribution of the HTAP1 model ensemble mean PAN mixing ratios at 650 hPa (~3 km), the level sampled by the highest altitude sites on which we focus the majority of our analysis. PAN mixing ratios in April generally increase with latitude, as expected from the strong thermal dependence of the PAN lifetime, although some of the highest mixing ratios are simulated over the Asian source region. The multi-model spread in lower tropospheric PAN, represented by the coefficient of variation (standard deviation over the 14 models divided by the model ensemble mean) is within ±45% across much of the northern hemisphere (Figure 2). The large inter-model spread over much of Europe in Figure 2b implies that observational constraints in this region would be particularly valuable.

Observed and modelled PAN mixing ratios at the northern mid-latitude mountain sites are compared in Figure 3; see Supplemental Figure 1 for a comparison extended throughout the year). We consider the measured range across years to bound the “plausible” portion of the wide range in simulated total PAN across the models. The multi-model mean falls in the range of the measurements at four of the sites, but is higher than observed in any year at Mount Bachelor. The model rankings show some consistency across the different sites, suggesting systematic model differences that can be narrowed with a limited set of observational constraints, especially for models that rank similarly across the sites on all three continents

(Figure 3). For example, CAMCHEM and GEMAQ are consistently at the higher end of the range while GISS-PUCCINI and LLNL-IMPACT are at the low end. The two models falling closest to the observed 2001 value at Schauinsland (MOZECHE and MOZARTGFDL) fall into the observed range at either Mount Bachelor or Jungfraujoch; we analyze these two sites further in the following sections.

The longest observational dataset at Schauinsland varies by over a factor of three across years, consistent with large inter-annual variability found in prior analyses at mountaintop sites (Zellweger et al., 2003; Fischer et al., 2011; Pandey Deolal et al. 2013; 2014). All but one of the models (LLNL-IMPACT) fall within the wide range of observed inter-annual variability at Schauinsland, underscoring the tenuous nature of conclusions regarding model performance drawn from short observational records unless the modelled and observed meteorological years match. Future work to coordinate consistent time periods between measurements and models would provide tighter constraints than are possible with our proof-of-concept analysis described in the following sections.

#### 4. Exploring emergent constraints on model SRRs from measured total PAN

The range of the PAN SRRs across the HTAP1 models at Jungfraujoch, Mount Bachelor, and Mount Waliguan is wide for all three source regions, spanning a factor of five or more in several cases (Figure 4). The key to a successful emergent constraint analysis is for this range in inter-model PAN SRRs, our unobservable quantity, to correlate with the total PAN simulated at the mountaintop site, our observable variable. The strongest correlations emerge for PAN originating in the region where the mountain is located, but some intercontinental SRR pairs also show significant correlations ( $p \leq 0.05$ ) with total simulated PAN (Figure 4).

We illustrate here how PAN measurements can be used to narrow the inter-model range in the SRR pairs. For the sites with significant correlations, the range across years (i.e., red vertical lines in Figure 4) bound the April mean values observed at Jungfraujoch and Mount Bachelor. The models falling in this range are highlighted in red. We select these models to narrow the range in SRRs, indicated by the red horizontal dashed lines extending from the bounding models (red symbols) to the ordinate axis. Figure 4 shows that the constraint from total measured PAN narrows the inter-model range in SRRs for PAN by at least half, revealing some models as outliers. Other models simulate SRRs within the observationally constrained range (between the dashed red horizontal lines) despite falling outside the observed range for total PAN, possibly indicating a role for inter-model differences in non-anthropogenic sources of PAN or in the relative contributions from the individual mid-latitude source regions, which we investigate further in the next section. Given the year-to-year variability in total PAN, stronger constraints could be placed in future work where the model meteorology corresponds to the same year as the measurements.

At Jungfraujoch, we additionally consider PAN SRRs for the EU source region with those estimated previously by back-trajectory analysis (Pandey Deolal et al., 2013). While Pandey Deolal et al. (2013) also attribute trajectories to NA and EA, fewer than 15% and 4% of trajectories are attributed to those regions as compared to 25-50% from EU (range across years; see Figure 1 of Pandey Deolal et al. (2013)). Combining these low frequencies with the inevitable growth in uncertainty as trajectories lengthen, we have the most confidence in the Pandey Deolal et al. (2013) estimates for the EU region. The horizontal blue dashed lines indicate the bounds obtained from this trajectory-based approach to estimating PAN from EU. The models falling in these bounds overlap with those constrained by the total PAN measurements, lending some confidence that these two independent approaches (one using total PAN and the correlated inter-model spread in SRRs; the other using back-trajectories to estimate SRRs) yield useful constraints on the influence of the EU source region on PAN measured at Jungfraujoch.

We note that for consistency with the model SRRs in Figure 4, which are the responses to 20% emission reductions in the source region, we divide the Pandey Deolal et al. (2013) EU SRRs by five to scale back from their estimated “full contribution” (100%). This linear scaling of the PAN response between 20% and 100% may incur errors due to non-linear chemistry. With an additional simulation in which the FRSGCUCI model sets European anthropogenic emissions of NO<sub>x</sub>, CO and VOC to zero (a 100% perturbation), we estimate this error to be ~10%. For intercontinental regions, this error reduces to < 3%. Earlier work shows that the smaller non-linearity in PAN for intercontinental versus regional source-receptor pairs also holds for ozone (Fiore et al., 2009; Wu et al., 2009; Wild et al., 2012), and demonstrates approximate linearity between the simulated tropospheric ozone burden and ±50% of present-day global NO<sub>x</sub> emissions (Stevenson et al., 2006).

## 5. Factors contributing to the inter-model range in PAN SRRs

We investigate the role of inter-model differences in regional emissions of PAN precursors versus transport in contributing to inter-model differences in the PAN response to continental-scale emission changes at the three mountaintop sites shown in Figure 4. At each site, we examine the correlation across models between simulated PAN SRRs and regional anthropogenic emissions of VOC (AVOC; Figure 5) or NO<sub>x</sub> (ANO<sub>x</sub>). The relationships for the EA SRRs are not significant, even at Mount Waliguan. We find, however, that the inter-model range in regional AVOC emissions explains as much as 64% of the variation in PAN attributed to EU emissions, and at least 25% of the variance in PAN attributed to the NA region (Figure 5). In contrast to AVOC, we find little relationship between the range in simulated PAN SRRs at the mountain sites and the model spread in regional ANO<sub>x</sub> emissions. Fischer et al. (2014) have previously shown that PAN abundances respond more strongly to changes in emissions of VOC than of NO<sub>x</sub>. Our analysis supports that earlier finding and furthermore highlights a key role for model differences in regional AVOC emissions in contributing to the inter-model range in PAN SRRs.

Differences in model transport (e.g., Arnold et al., 2015; Orbe et al., 2017) may also contribute to the inter-model differences in PAN SRRs. Our analysis of the HTAP1 idealized CO tracers, however, reveals little correlation between inter-model differences in these idealized tracers (which have identical regional emissions and lifetimes applied in all of the models) and in the PAN SRRs sampled at these sites. Although we do not find any clear overall correlation, differences in the idealized CO tracers explain some of the scatter in Figure 5. For example, at Jungfraujoch for EU AVOC emissions of 22 Tg C a<sup>-1</sup>, the lowest model (GISS-PUCCINI) has one of the smallest values for the CO<sub>fromEU</sub> tracer, whereas the highest model (STOC-HadAM3) has the largest value of CO<sub>fromEU</sub>.

In light of the dependence of inter-model differences in PAN attributed to EU and NA during April and the corresponding regional AVOC emissions, we illustrate how one could extend our emergent constraints in Figure 4 (horizontal dashed red lines) to the regional AVOC emission estimates shown in Figure 5. A major caveat underlying this analysis is the mis-match between meteorological years for the models and measurements as discussed above, and the underlying assumption that the relationships in Figure 5 can exclusively be attributed to differences in the AVOC emissions (as opposed to chemistry or transport). The observationally-constrained SRRs between PAN from NA and total PAN measured at Jungfraujoch and Mount Bachelor can be used to narrow the range of NA AVOC emissions to 12-18 Tg C a<sup>-1</sup> (the low end is ruled out by the constraint imposed by PAN from NA at Jungfraujoch; the high end is ruled out by PAN from NA at Mount Bachelor). Similarly, the range for EU AVOC emissions would narrow to 16-25 Tg C a<sup>-1</sup>.

We consider next the importance that various models ascribe to a given source region relative to another source region. We first correlate the ratios of PAN from two different source regions with the total PAN simulated by the individual models in April. We find little relationship, with the exception of Mount Bachelor, where the observational constraint implies that more PAN originating from EA should be present at Mount Bachelor than PAN originating from NA (Figure 6a). We interpret this as indicating that models with higher total PAN at Mount Bachelor are overestimating North American influence at this mountain site (which samples free tropospheric air). This interpretation is supported by the idealized CO tracer simulations (with identical regional emissions and the same lifetime applied in all the models), which suggest that some of the variance in the ratio of PAN from NA versus EA at Mount Bachelor is due to differences in transport from the two regions (Figure 6b). We emphasize that these transport differences do not simply reflect the use of different meteorology to drive the CTMs (Figure 6b).

By comparing NA:EA at Mount Bachelor, EU:NA at Jungfraujoch, and EA:EU at Mount Waliguan, we examine the relative importance of emissions within the source region, where the measurement site is located, versus the upwind intercontinental source region on PAN (Supplemental Figure 2). At Mount Bachelor, the HTAP1 multi-model mean SRRs from NA, EA, and EU, are roughly equal in April (Figure

1). The differences across the HTAP models in the relative importance of the NA:EA source regions on PAN (which range from about 0.5 to 2.5) correlate roughly equally with the ratio of the NA:EA CO transport tracers and with the ratio of the NA:EA AVOC emissions (spearman rank correlation coefficient ( $r$ ) = 0.6 for both cases); we find no relationship with the ratio of the NA:EA ANO<sub>x</sub> emissions (Figure 6b and Supplemental Figure 2 left column). At Jungfraujoch, the HTAP1 multi-model mean attributes much of the PAN to emissions from the EU and NA source regions during April (Figure 1). The ratio of PAN attributed to EU versus NA at Jungfraujoch, however, varies from approximately 0.5 to 2 across the individual HTAP1 models (Supplemental Figure 2). In contrast to our findings at Mount Bachelor, this ratio at Jungfraujoch depends most strongly on the ratio of ANO<sub>x</sub> emissions in the EU to NA regions ( $r$  = 0.6), and more weakly on the ratio of EU:NA AVOC emissions ( $r$  = 0.5; Supplemental Figure 2). The correlation is even weaker between the ratio of PAN SRRs for these two regions with inter-model differences in transport as diagnosed with the CO tracers from EU versus NA ( $r$  = 0.4). At Mount Waliguan, the strongest relationship is found for the ratio of AVOC emissions ( $r$  = 0.5; Supplemental Figure 2).

We repeat this correlation analysis of inter-model differences in ratios of ANO<sub>x</sub> emissions, AVOC emissions, or the idealized CO tracers of transport from a region, but for the ratio of PAN SRRs from two intercontinental regions. At Mount Bachelor, the EU and EA source regions contribute similar amounts to multi-model mean PAN during April (Figure 1). Across the individual models, however, the ratio of the EU to EA source region on PAN at Mount Bachelor varies from less than half to a factor of two (Supplemental Figure 3). We find that the ratio of PAN attributed to the EU versus EA source regions at Mount Bachelor correlates strongly across the models with the ratio of the anthropogenic volatile organic compound (AVOC) emissions in the respective source regions ( $r$  = 0.8; Supplemental Figure 3). In contrast, the ratio of EU:EA anthropogenic emission influence on PAN at Mount Bachelor shows little correlation with the respective regional NO<sub>x</sub> emissions used in the models, or with the differences in the simulated transport tracers ( $r$ =0.3 for both cases). As at Mount Bachelor, the model spread in the contribution to total simulated PAN from the EA versus NA source regions at both Jungfraujoch and Mount Waliguan depend most on the regional AVOC ratios ( $r$  = 0.8 and 0.6, respectively; Supplemental Figure 3), with little correlation with inter-model differences in NA:EA ANO<sub>x</sub> emissions. Some correlation also emerges between the NA:EA source-receptor relationships for PAN and the NA:EA transport tracers ( $r$  = 0.6 at both sites; Supplemental Figure 3). Finally, we do not find any obvious link between PAN SRRs and the choice of meteorological fields (the individual symbols in Supplemental Figures 2 and 3).

## 6. Linking PAN and O<sub>3</sub> SRRs

We address here the extent to which observational constraints on PAN SRRs might also serve to narrow the range of uncertainty in the inter-model spread in intercontinental SRRs for O<sub>3</sub> (e.g., Fiore et al., 2009). We expect some commonality between the sensitivity of PAN and O<sub>3</sub> to changes in precursor emissions

because (1) both species are produced from chemical reactions involving  $\text{NO}_x$  and VOC, and (2) PAN serves as a  $\text{NO}_x$  reservoir, which upon decomposition releases  $\text{NO}_x$  that can then produce  $\text{O}_3$  far downwind of the region where the PAN (and  $\text{O}_3$ ) precursors were originally emitted. Furthermore, earlier analysis of HTAP1 ozone continental-scale SRRs also identified a correlation with the model AVOC emissions, particularly over EU (Fiore et al., 2009).

We assess the extent to which the inter-model range in source region influence on mountaintop PAN levels in April is relevant for interpreting  $\text{O}_3$  SRRs by correlating PAN and  $\text{O}_3$  SRRs at the three mountaintop sites (Figure 7). Relationships vary across the individual source-receptor pairs, with the inter-model variability in PAN explaining 16-60% of the inter-model differences in  $\text{O}_3$  at the mountain sites. The strongest relationships occur for the influence of regional sources at Mount Bachelor (from NA) and Jungfrauoch (from EU). At Mount Waliguan, the EU and EA source-receptor relationships for PAN and  $\text{O}_3$  are of similar strength ( $r = 0.7$ ). Intercontinental source-receptor pairs for  $\text{O}_3$  and PAN at Mount Bachelor and Mount Waliguan are also significant to within 90%, with variability in the PAN attributed to intercontinental source regions explaining 25-35% and 30-45%, respectively, of the variability in the corresponding  $\text{O}_3$  SRRs.

We expand the correlation analysis of ozone and PAN SRRs from the free troposphere sampled at the mountaintop sites to large-scale SRRs in surface air over the HTAP1 continental regions. Of the significant relationships in Figure 7 ( $p < 0.10$ ), 6 out of 7 also emerge as significant in Figure 8. We infer that conclusions drawn from a limited number of mountaintop sites regarding PAN SRRs and their relationship to ozone SRRs are relevant, at least according to the models, on much broader scales.

We repeat the analysis in Figure 5 but for  $\text{O}_3$  to consider the influence of the three source regions on the three mountaintop sites (nine total source-receptor pairs), but find little relationship between the model spread in the simulated  $\text{O}_3$  SRRs and in the magnitude of the regional AVOC or  $\text{ANO}_x$  emissions. Model differences in transport as diagnosed by the idealized regional CO tracers correlates more with  $\text{O}_3$  SRRs than for PAN for all source-receptor pairs, though the correlations remains weak except for CO from EU with  $\text{O}_3$  SRRs at Jungfrauoch. Overall, this analysis supports earlier findings that PAN is more sensitive to changes in emissions (and subsequent chemistry), particularly for VOC precursors, than  $\text{O}_3$ .

The correlations between SRRs for PAN and  $\text{O}_3$  could reflect a role for PAN transport in contributing to  $\text{O}_3$  production over the receptor region, or may instead reflect co-production of PAN and  $\text{O}_3$  from oxidation of regional precursor emissions followed by transport in the same air mass. In the latter case, PAN is serving as a proxy for  $\text{O}_3$  transport whereas in the former case, PAN is serving as the actual pathway by which  $\text{O}_3$  is transported. We do not have model diagnostics that allow us to distinguish between these two roles for PAN. The correlations between PAN and  $\text{O}_3$  SRRs, however, suggest that long-term PAN measurements

contain signals relevant for constraining the relative importance of regional vs. intercontinental emissions on both PAN and O<sub>3</sub>. We examine the strength of these signals by correlating the O<sub>3</sub> SRRs at each site with total PAN as simulated at each site. Relationships are far weaker than for the PAN SRRs and total PAN shown in Figure 4, but **correlations are significant** between total PAN at Jungfraujoch for O<sub>3</sub> from EU (r=0.67; p=0.03) and at Mount Bachelor for O<sub>3</sub> from NA (r=0.61; p=0.04; Supplemental Figure 4).

## 7. Conclusions and recommendations

Our proof of concept approach applies the HTAP1 multi-model ensemble to identify a strong inter-model correlation between PAN source-receptor relationships (SRRs; defined as the difference in simulations with 20% emission reductions separately within each of the northern mid-latitude continents) and simulated total PAN at mountaintop sites during April. Our findings imply promise for developing “emergent constraints” (e.g., Hall and Qu, 2006; Borodina et al., 2017; Cox et al., 2018) from **more routine** PAN measurements to narrow uncertainty in wide-ranging model estimates of PAN SRRs, quantities that are not directly observable yet relevant to air quality policy (e.g., HTAP 2010). Inter-model correlations of the responses of PAN versus O<sub>3</sub> to perturbations in regional anthropogenic emissions (Figures 7 and 8) imply that constraints on PAN SRRs are relevant for lowering uncertainty in **O<sub>3</sub> SRR estimates. This connection** between PAN and O<sub>3</sub> likely reflects the dual role of PAN as both a pathway for O<sub>3</sub> transport (by producing O<sub>3</sub> upon its decomposition following transport), and as a proxy for O<sub>3</sub> transport (as it is produced alongside O<sub>3</sub> in the polluted continental boundary layer).

Establishing the strongest constraints possible on simulated SRRs for PAN and O<sub>3</sub> will require (1) measurements and simulations with chemical transport models that coincide, and (2) a sufficiently long measurement record to build a climatology suitable for evaluating chemistry-climate models that generate their own meteorology. **Repeated sampling for the month of April may be sufficient to provide constraints on model responses to changes in anthropogenic emissions. PAN measurements over multiple seasons are necessary to evaluate model responses of PAN to climate change (e.g., by changing temperature and weather-sensitive precursor emissions) and the resulting influence on atmospheric O<sub>3</sub> and oxidizing capacity (e.g., Doherty et al., 2013). For example, changes in meteorology and biomass burning (Fischer et al., 2011; Zhu et al., 2015) such as those driven by ENSO (Koumoutsaris et al., 2008), as well as biogenic and lightning sources (Payne et al., 2017) vary from year to year and are expected to change as climate warms.**

We identified only five multi-year datasets at mountain sites, four of which are located near each other in Europe, and only one of which continues at present (Schauinsland). Our analysis suggests that **future** measurements at Mount Waliguan would provide constraints on PAN SRRs, particularly for PAN originating in EA (Figure 4). Additional work could systematically examine over 60 stations at altitudes above 2500 m in the Tropospheric Ozone Assessment Report (TOAR) database (Schultz et al., 2017).



We recommend archival of daily model fields for future applications of this multi-model emergent constraint approach to SRRs. Access to daily model fields permits (1) a more rigorous process-oriented evaluation of specific events (e.g., Fischer et al., 2010; Alvarado et al., 2010; Arnold et al., 2015), and (2) comparison with satellite-derived tropospheric PAN columns, which show promise for documenting PAN distributions, particularly in the upper troposphere, and their temporal variability and spatial patterns across the globe (e.g., Fadnavis et al., 2014; Jiang et al., 2016; Payne et al., 2014; 2017; Zhu et al., 2015; 2017). We also suggest archiving daily tracers tagged by emission region to isolate the role of model differences in transport during individual events. In addition, Lin et al. (2017) have demonstrated that applying a filtering technique based on daily idealized CO regional tracers can better isolate free tropospheric air from surface air masses when comparing coarse resolution models with high altitude measurements.

By focusing on April, our analysis largely minimizes complexities introduced by inter-model differences in biogenic, fire, and lightning sources that further complicate disentangling summertime discrepancies in simulated PAN and O<sub>3</sub> (e.g., Arnold et al., 2015; Emmons et al., 2015) and restricts inter-model differences to those associated with anthropogenic emissions and the subsequent chemistry and transport. Nevertheless, we find a wide range in inter-model SRR relationships that reflects uncertainties in emissions and different model representations of VOC chemistry, including PAN yields from VOCs (Figure 5; Emmerson and Evans, 2009; Fischer et al., 2014; Arnold et al., 2015; Emmons et al., 2015; Knote et al., 2015). Future multi-model efforts could seek to parse separately the influence of differences in total anthropogenic VOC emissions, the mix of emitted VOC species and their reactivity, and the chemical production of PAN and O<sub>3</sub>. Documenting these aspects of model configuration would help to establish benchmarks for inter-model differences in simulated total PAN, O<sub>3</sub>, and their SRRs, against which future model simulations (and multi-model ensembles) can be assessed.

## Acknowledgments

We thank Mathew Evans (York University, UK) and Terry Keating (U.S. EPA) for useful discussions, and two anonymous referees for their constructive comments. AMF and BND acknowledge NASA MAP (NNX14AM38G). DSS acknowledges NERC (grants NE/K001329/1 and NE/N003411/1) and the ARCHER UK National Supercomputing Service (<http://www.archer.ac.uk>). Data for the Mt. Bachelor Observatory are archived and available at the University of Washington data archive: <https://digital.lib.washington.edu/researchworks/browse?type=subject&value=Mt.+Bachelor+Observatory>. The PAN data for the European Mountain sites is archived by the World Data Centre for Greenhouse Gases (<http://ds.data.jma.go.jp/gmd/wdcgg/>). Upon publication, the data used to generate the figures will be placed in a CSU digital repository that we have already established for this manuscript (<https://hdl.handle.net/10217/185610>). This is Lamont contribution number 8251.



## References

- Alvarado, M. J., and 35 others: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmos. Chem. Phys.*, 10, 9739–9760, doi:10.5194/acp-10-9739-2010, 2010.
- Arnold, S. R., Emmons, L. K., Monks, S. A., Law, K. S., Ridley, D. A., Turquety, S., Tilmes, S., Thomas, J. L., Bouarar, I., Flemming, J., Huijnen, V., Mao, J., Duncan, B. N., Steenrod, S., Yoshida, Y., Langner, J., and Long, Y.: Biomass burning influence on high-latitude tropospheric ozone and reactive nitrogen in summer 2008: a multi-model analysis based on POLMIP simulations, *Atmos. Chem. Phys.*, 15, 6047–6068, <https://doi.org/10.5194/acp-15-6047-2015>, 2015.
- Balzani Lööv, J. M., S. Henne, G. Legreid, J. Staehelin, S. Reimann, A. S. H. Prévôt, M. Steinbacher, and M. K. Vollmer: Estimation of background concentrations of trace gases at the Swiss Alpine site Jungfraujoch (3580 m asl), *J. Geophys. Res.*, 113, D22305, doi:[10.1029/2007JD009751](https://doi.org/10.1029/2007JD009751), 2008.
- Borodina, A., Fischer, E. M., R. Knutti, Emergent Constraints in Climate Projections: A Case Study of Changes in High-Latitude Temperature Variability, *Journal of Climate*, <https://doi.org/10.1175/JCLI-D-16-0662.1>, 2017.
- Bottenheim, J. W., Sirois, A., Brice, K. A., and Gallant, A. J.: Five years of continuous observations of PAN and ozone at a rural location in eastern Canada, *Journal of Geophysical Research: Atmospheres*, 99, 5333–5352, 10.1029/93JD02716, 1994.
- Carpenter, L. J., Green, T. J., Mills, G. P., Bauguutte, S., Penkett, S. A., Zanis, P., Schuepbach, E., Schmidbauer, N., Monks, P. S., Zellweger, C: Oxidized nitrogen and ozone production efficiencies in the springtime free troposphere over the Alps, *J. Geophys. Res.*, 105, D11, 14,547–14,559, 2000.
- Chameides, W. L., Fehsenfeld, F. C., Rodgers, M. O., Cardelino, C., Martinez, J., Parrish, D., Lonnerman, W., Lawson, D. R., Rasmussen, R. A., Zimmerman, P., Greenberg, J., Middleton, P., and Wang, T.: Ozone precursor relationships in the ambient atmosphere, *Journal of Geophysical Research*, 97, 6037–6055, 1992.
- Cox, P. M., Huntingford, C., and Williamson, M. S., Emergent constraint on equilibrium climate sensitivity from global temperature variability: *Nature*, 553, 319, 10.1038/nature25450, 2018.
- Doherty, R. M., et al.: Impacts of climate change on surface ozone and intercontinental ozone pollution: A multi-model study, *J. Geophys. Res. Atmos.*, 118, doi:10.1002/jgrd.50266, 2013.
- Emmerson, K. M., and Evans, M. J.: Comparison of tropospheric gas-phase chemistry schemes for use within global models, *Atmos. Chem. Phys.*, 9, 1831–1845, 10.5194/acp-9-1831-2009, 2009.
- Emmons, L. K., Arnold, S. R., Monks, S. A., Huijnen, V., Tilmes, S., Law, K. S., Thomas, J. L., Raut, J.-C., Bouarar, I., Turquety, S., Long, Y., Duncan, B., Steenrod, S., Strode, S., Flemming, J., Mao, J., Langner, J., Thompson, A. M., Tarasick, D., Apel, E. C., Blake, D. R., Cohen, R. C., Dibb, J., Diskin, G. S., Fried, A., Hall, S. R., Huey, L. G., Weinheimer, A. J., Wisthaler, A., Mikoviny, T., Nowak, J., Peischl, J., Roberts, J. M., Ryerson, T., Warneke, C., and Helmig, D.: The POLARCAT Model

Intercomparison Project (POLMIP): overview and evaluation with observations, *Atmos. Chem. Phys.*, 15, 6721-6744, <https://doi.org/10.5194/acp-15-6721-2015>, 2015.

Fadnavis, S., Schultz, M. G., Semeniuk, K., Mahajan, A. S., Pozzoli, L., Sonbawne, S., Ghude, S. D., Kiefer, M., and Eckert, E.: Trends in peroxyacetyl nitrate (PAN) in the upper troposphere and lower stratosphere over southern Asia during the summer monsoon season: regional impacts, *Atmos. Chem. Phys.*, 14, 12725-12743, <https://doi.org/10.5194/acp-14-12725-2014>, 2014.

Fiore, A. M., Dentener, F. J., Wild, O., Cuvelier, C., Schultz, M. G., Hess, P., Textor, C., Schulz, M., Doherty, R. M., Horowitz, L. W., MacKenzie, I. A., Sanderson, M. G., Shindell, D. T., Stevenson, D. S., Szopa, S., Van Dingenen, R., Zeng, G., Atherton, C., Bergmann, D., Bey, I., Carmichael, G., Collins, W. J., Duncan, B. N., Faluvegi, G., Folberth, G., Gauss, M., Gong, S., Hauglustaine, D., Holloway, T., Isaksen, I. S. A., Jacob, D. J., Jonson, J. E., Kaminski, J. W., Keating, T. J., Lupu, A., Marmer, E., Montanaro, V., Park, R. J., Pitari, G., Pringle, K. J., Pyle, J. A., Schroeder, S., Vivanco, M. G., Wind, P., Wojcik, G., Wu, S., and Zuber, A.: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *J. Geophys. Res.*, 114, D04301, [10.1029/2008jd010816](https://doi.org/10.1029/2008jd010816), 2009.

Fiore, A. M., Levy II, H., and Jaffe, D. A.: North American isoprene influence on intercontinental ozone pollution, *Atmos. Chem. Phys.*, 11, 1697-1710, [10.5194/acp-11-1697-2011](https://doi.org/10.5194/acp-11-1697-2011), 2011.

Fischer, E. V., Jaffe, D. A., Reidmiller, D. R., and Jaegle, L.: Meteorological controls on observed peroxyacetyl nitrate (PAN) at Mount Bachelor during the spring of 2008, *Journal of Geophysical Research*, 115 doi: [10.1029/2009JD0012776](https://doi.org/10.1029/2009JD0012776), 2010.

Fischer, E. V., Jaffe, D. A., and Weatherhead, E. C.: Free tropospheric peroxyacetyl nitrate (PAN) and ozone at Mount Bachelor: potential causes of variability and timescale for trend detection, *Atmos. Chem. Phys.*, 11, 5641-5654, [10.5194/acp-11-5641-2011](https://doi.org/10.5194/acp-11-5641-2011), 2011.

Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot, F., Singh, H. B., Roiger, A., Ries, L., Talbot, R. W., Dzepina, K., and Pandey Deolal, S.: Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos. Chem. Phys.*, 14, 2679-2698, [10.5194/acp-14-2679-2014](https://doi.org/10.5194/acp-14-2679-2014), 2014.

Gilge, S., Plass-Dülmer, C., Fricke, W., Kaiser, A., Ries, L., Buchmann, B., and Steinbacher, M.: Ozone, carbon monoxide and nitrogen oxides time series at four alpine GAW mountain stations in central Europe, *Atmos. Chem. Phys.*, 10, 12295-12316, [10.5194/acp-10-12295-2010](https://doi.org/10.5194/acp-10-12295-2010), 2010.

Hall, A., and Qu, X.: Using the current seasonal cycle to constrain snow albedo feedback in future climate change: *Geophys. Res. Lett.*, 33, L03502, doi:[10.1029/2005GL025127](https://doi.org/10.1029/2005GL025127), 2006.

Heald, C. L., Jacob, D. J., Fiore, A. M., Emmons, L. K., Gille, J. C., Deeter, M. N., Warner, J., Edwards, D. P., Crawford, J. H., Hamlin, A. J., Sachse, G. W., Browell, E. V., Avery, M. A., Vay, S. A., Westberg, D. J., Blake, D. R., Singh, H. B., Sandholm, S. T., Talbot, R. W., and Fuelberg, H. E.: Asian outflow and trans-Pacific transport of carbon monoxide and ozone pollution: An integrated satellite, aircraft, and model perspective, *J. Geophys. Res.*, 108, 4804, [10.1029/2003jd003507](https://doi.org/10.1029/2003jd003507), 2003.

HTAP: Task Force on Hemispheric Transport of Air Pollution 2007 Interim Report, United Nations Economic Commission for Europe, New York and Geneva, 2007.

HTAP: HEMISPHERIC TRANSPORT OF AIR POLLUTION 2010 PART A: OZONE AND PARTICULATE MATTER, Air Pollution Studies No. 17, UNITED NATIONS, New York, 2010.

Hudman, R. C., Jacob, D. J., Cooper, O. R., Evans, M. J., Heald, C. L., Park, R. J., Fehsenfeld, F., Flocke, F., Holloway, J., Hübler, G., Kita, K., Koike, M., Kondo, Y., Neuman, A., Nowak, J., Oltmans, S., Parrish, D., Roberts, J. M., and Ryerson, T.: Ozone production in transpacific Asian pollution plumes and implications for ozone air quality in California, *J. Geophys. Res.*, 109, D23S10, 10.1029/2004jd004974, 2004.

Jaeglé, L., Jaffe, D. A., Price, H. U., Weiss-Penzias, P., Palmer, P. I., Evans, M. J., Jacob, D. J., and Bey, I.: Sources and budgets for CO and O<sub>3</sub> in the northeastern Pacific during the spring of 2001: Results from the PHOBEA-II Experiment, *Journal of Geophysical Research*, 108, doi:10.1029/2002JD003121, 2003.

Jaffe, D., Thornton, J., Wolfe, G., Reidmiller, D., Fischer, E. V., Jacob, D. J., Cohen, R., Singh, H., Weinheimer, A., and Flocke, F.: Can we Detect an Influence over North America From Increasing Asian NO<sub>x</sub> Emissions?, *EOS Trans. AGU*, 88(52), Fall Meet. Suppl., Abstract A51E-04, 2007.

Jiang, Z., Worden, J. R., Payne, V. H., Zhu, L., Fischer, E., Walker, T., and Jones, D. B. A.: Ozone export from East Asia: The role of PAN, *Journal of Geophysical Research: Atmospheres*, 121, 6555-6563, 10.1002/2016JD024952, 2016.

Kirchner, F., Mayer-Figge, A., Zabel, F., and Becker, K. H.: Thermal stability of peroxy nitrates, *International Journal of Chemical Kinetics*, 31, 127-144, 1999.

Knote, C., Tuccella, P., Curci, G., Emmons, L., Orlando, J. J., Madronich, S., Baro, R., Jimenez-Guerrero, P., Luecken, D., Hogrefe, C., Forkel, R., Werhahn, J., Hirtl, M., Perez, J., San Jose, R., Giordano, L., Brunner, D., Yahya, K., Zhang, Y.: Influence of the choice of gas-phase mechanism on predictions of key gaseous pollutants during the AQMEII phase-2 intercomparison, *Atmos. Environ.*, 115, 553-568, 2015.

Kotchenruther, R. A., Jaffe, D. A., and Jaeglé, L.: Ozone photochemistry and the role of PAN in the springtime northeastern Pacific Troposphere: Results from the PHOBEA Campaign. *J. Geophys. Res.*, 106, 28731-28741, 2001a.

Kotchenruther, R. A., Jaffe, D. A., Beine, H. J., Anderson, T., Bottenheim, J. W., Harris, J. M., Blake, D., and Schmitt, R.: Observations of ozone and related species in the Northeast Pacific during the PHOBEA Campaigns: 2. Airborne observations. *J. Geophys. Res.*, 106, 7463-7483, 2001b (corrected Table 1 published in Vol. 106 (D17), p.20507).

Koumoutsaris, S., Bey, I., Generoso, S., and Thouret, V.: Influence of El Niño-Southern Oscillation on the interannual variability of tropospheric ozone in the northern midlatitudes, *J. Geophys. Res.*, 113, D19301, 10.1029/2007jd009753, 2008.

Kuhn, M., Builtjes, P. J. H., Poppe, D., Simpson, D., Stockwell, W. R., Andersson-Sköld, Y., Baart, A., Das, M., Fiedler, F., Hov, Ø., Kirchner, F., Makar, P. A., Milford, J. B., Roemer, M. G. M., Ruhnke, R., Strand, A., Vogel, B., and Vogel, H.: Intercomparison of the gas-phase chemistry in several chemistry and transport models, *Atmospheric Environment*, 32, 693-709, 1998.

Liang, J., Horowitz, L. W., Jacob, D. J., Wang, Y., Fiore, A. M., Logan, J. A., Gardner, G. M., and Munger, J. W.: Seasonal variations of reactive nitrogen species and ozone over the United States and export fluxes to the global atmosphere, *Journal of Geophysical Research*, 103, 13,435-413,450, 1998.

Lin, M., Horowitz, L. W., Payton, R., Fiore, A. M., and Tonnesen, G.: US surface ozone trends and extremes from 1980 to 2014: quantifying the roles of rising Asian emissions, domestic controls, wildfires, and climate, *Atmos. Chem. Phys.*, 17, 2943-2970, <https://doi.org/10.5194/acp-17-2943-2017>, 2017.

Lin, M., Holloway, T., Carmichael, G. R., and Fiore, A. M.: Quantifying pollution inflow and outflow over East Asia in spring with regional and global models, *Atmos. Chem. Phys.*, 10, 4221-4239, 10.5194/acp-10-4221-2010, 2010.

Liu, S. C., Trainer, M., Fehsenfeld, F. C., Parrish, D. D., Williams, E. J., Fahey, D. W., Hubler, G., and Murphy, P. C.: Ozone production in the Rural Troposphere and the Implications for Regional and Global Ozone Distributions, *Journal of Geophysical Research*, 92, 4191-4207, 1987.

Moxim, W. J., Levy, H., II, and Kasibhatla, P. S.: Simulated global tropospheric PAN: Its transport and impact on NO<sub>x</sub>, *J. Geophys. Res.*, 101, 12621-12638, 10.1029/96jd00338, 1996.

Orbe, C., D. W. Waugh, H. Yang, J.-F. Lamarque, S. Tilmes, and D. E. Kinnison (2017), Tropospheric transport differences between models using the same large-scale meteorological fields, *Geophys. Res. Lett.*, 44, 1068–1078, doi:10.1002/2016GL071339.

Pandey Deolal, S., Henne, S., Ries, L., Gilge, S., Weers, U., Steinbacher, M., Staehelin, J., and Peter, T.: Analysis of elevated springtime levels of Peroxyacetyl nitrate (PAN) at the high Alpine research sites Jungfraujoch and Zugspitze, *Atmos. Chem. Phys.*, 14, 12553-12571, 10.5194/acp-14-12553-2014, 2014.

Pandey Deolal, S., Staehelin, J., Brunner, D., Cui, J., Steinbacher, M., Zellweger, C., Henne, S., and Vollmer, M. K.: Transport of PAN and NO<sub>y</sub> from different source regions to the Swiss high alpine site Jungfraujoch, *Atmospheric Environment*, 64, 103-115, <https://doi.org/10.1016/j.atmosenv.2012.08.021>, 2013.

Payne, V. H., Alvarado, M. J., Cady-Pereira, K. E., Worden, J. R., Kulawik, S. S., and Fischer, E. V.: Satellite observations of peroxyacetyl nitrate from the Aura Tropospheric Emission Spectrometer, *Atmos. Meas. Tech.*, 7, 3737-3749, 10.5194/amt-7-3737-2014, 2014.

Payne, V. H., Fischer, E. V., Worden, J. R., Jiang, Z., Zhu, L., Kurosu, T. P., and Kulawik, S. S.: Spatial variability in tropospheric peroxyacetyl nitrate in the tropics from infrared satellite observations in 2005 and 2006, *Atmos. Chem. Phys.*, 17, 6341-6351, 10.5194/acp-17-6341-2017, 2017.

- Penkett, S. A., and Brice, K. A.: The spring maximum in photo-oxidants in the Northern Hemisphere troposphere, *Nature*, 319, 655-657, 1986.
- Roberts, J. M.: PAN and Related Compounds, in: *Volatile Organic Compounds in the Atmosphere*, edited by: Koppmann, R., Blackwell Publishing, 500, 2007.
- Schmitt, R., and Volz-Thomas, A.: Climatology of Ozone, PAN, CO, and NMHC in the Free Troposphere Over the Southern North Atlantic, *Journal of Atmospheric Chemistry*, 28, 245-262, 10.1023/A:1005801515531, 1997.
- Schultz, M. G. et al.: On the origin of tropospheric ozone and NO<sub>x</sub> over the tropical South Pacific, *J. Geophys. Res.*, 104, D5, 5829-5843, 1999.
- Schultz, M., Rast, S., van het Bolscher, M., Pulles, T., Brand, R., Pereira, J., Mota, B., Spessa, A., Dalsoren, S., van Noije, T., and Szopa, S.: Emission data sets and methodologies for estimating emissions, Hamburg, 2007.
- Schultz, M. G., Heil, A., Hoelzemann, J. J., Spessa, A., Thonicke, K., Goldammer, J. G., Held, A. C., Pereira, J. M. C., and van het Bolscher, M.: Global wildland fire emissions from 1960 to 2000, *Global Biogeochemical Cycles*, 22, GB2002, 10.1029/2007GB003031, 2008.
- Schultz MG, Schröder S, Lyapina O, Cooper O, Galbally I, Petropavlovskikh I, et al.: Tropospheric Ozone Assessment Report: Database and Metrics Data of Global Surface Ozone Observations, *Elem Sci Anth.*, 5, 58, DOI: <http://doi.org/10.1525/elementa.244>, 2017.
- Shindell, D. T., Chin, M., Dentener, F., Doherty, R. M., Faluvegi, G., Fiore, A. M., Hess, P., Koch, D. M., MacKenzie, I. A., Sanderson, M. G., Schultz, M. G., Schulz, M., Stevenson, D. S., Teich, H., Textor, C., Wild, O., Bergmann, D. J., Bey, I., Bian, H., Cuvelier, C., Duncan, B. N., Folberth, G., Horowitz, L. W., Jonson, J., Kaminski, J. W., Marmer, E., Park, R., Pringle, K. J., Schroeder, S., Szopa, S., Takemura, T., Zeng, G., Keating, T. J., and Zuber, A.: A multi-model assessment of pollution transport to the Arctic, *Atmos. Chem. Phys.*, 8, 5353-5372, 10.5194/acp-8-5353-2008, 2008.
- Singh, H. B.: Reactive nitrogen in the troposphere, *Environmental Science and Technology*, 21, 320-327, 1987.
- Singh, H. B., and Hanst, P. L.: Peroxyacetyl nitrate (PAN) in the unpolluted atmosphere: An important reservoir for nitrogen oxides, *Geophysical Research Letters*, 8, 941-944, 1981.
- Singh, H. B., and Salas, L. J.: Measurements of peroxyacetyl nitrate (pan) and peroxypropionyl nitrate (ppn) at selected urban, rural and remote sites, *Atmospheric Environment* (1967), 23, 231-238, [https://doi.org/10.1016/0004-6981\(89\)90115-7](https://doi.org/10.1016/0004-6981(89)90115-7), 1989.
- Stevenson, D.S., et al.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, 111, D08301, doi:10.1029/2005JD006338, 2006.
- Turnipseed, A. A., Huey, L. G., Nemitz, E., Stickel, R., Higgs, J., Tanner, D. J., Slusher, D. L., Sparks, J. P., Flocke, F., and Guenther, A.: Eddy covariance fluxes of peroxyacetyl nitrates (PANs) and NO<sub>y</sub> to a coniferous forest, *Journal of Geophysical Research: Atmospheres*, 111, D09304, 10.1029/2005jd006631, 2006.

- Val Martin, M., Honrath, R. E., Owen, R. C., and Lapina, K.: Large-scale impacts of anthropogenic pollution and boreal wildfires on the nitrogen oxide levels over the central North Atlantic region, *Journal of Geophysical Research*, 113, doi:10.1029/2007JD009689, 2008.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr, A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423-3441, 10.5194/acp-6-3423-2006, 2006.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707-11735, 10.5194/acp-10-11707-2010, 2010.
- Volz-Thomas, A., Xueref, I., and Schmitt, R.: An automatic gas chromatograph and calibration system for ambient measurements of PAN and PPN, *Environmental Science and Pollution Resources*, Special Issue 4, 72-76, 2002.
- Wang, Y., and Jacob, D. J.: Anthropogenic forcing on tropospheric ozone and OH since preindustrial times, *Journal of Geophysical Research*, 103, 31,123-131,135, 1998.
- Warneck, P., and Zerbach, T.: Synthesis of peroxyacetyl nitrate by acetone photolysis, *Environmental Science and Technology*, 26, 74-79, 1992.
- Weiss-Penzias, P., Jaffe, D. A., Jaeglé, L., and Liang, Q.: Influence of long-range-transported pollution on the annual and diurnal cycles of carbon monoxide and ozone at Cheeka Peak Observatory, *Journal of Geophysical Research: Atmospheres*, 109, n/a-n/a, 10.1029/2004JD004505, 2004.
- Wild, O., Law, K., McKenna, D., Bandy, B., Penkett, S., Pyle, J.: Photochemical trajectory modeling studies of the North Atlantic region during August 1993, *Journal of Geophysical Research: Atmospheres*, 101, D22, p. 29269-29288, 1996.
- Wild, O., Fiore, A. M., Shindell, D. T., Doherty, R. M., Collins, W. J., Dentener, F. J., Schultz, M. G., Gong, S., MacKenzie, I. A., Zeng, G., Hess, P., Duncan, B. N., Bergmann, D. J., Szopa, S., Jonson, J. E., Keating, T. J., and Zuber, A.: Modelling future changes in surface ozone: a parameterized approach, *Atmos. Chem. Phys.*, 12, 2037-2054, 10.5194/acp-12-2037-2012, 2012.
- Wu, S., Duncan, B. N., Jacob, D. J., Fiore, A. M., and Wild, O.: Chemical nonlinearities in relating intercontinental ozone pollution to anthropogenic emissions, *Geophys. Res. Lett.*, 36, L05806, 10.1029/2008gl036607, 2009.
- Xue, L. K., Wang, T., Zhang, J. M., Zhang, X. C., Deliger, Poon, C. N., Ding, A. J., Zhou, X. H., Wu, W. S., Tang, J., Zhang, Q. Z., and Wang, W. X.: Source of surface ozone and reactive nitrogen speciation at Mount Waliguan in western China: New insights from the 2006 summer study, *J. Geophys. Res.*, 116, D07306, doi:10.1029/2010JD014735, 2011.
- Zellweger, C., Ammann, M., Buchmann, B., Hofer, P., Lugauer, M., Rüttimann, R., Streit, N., Weingartner, E., and Baltensperger, U.: Summertime NO<sub>y</sub> speciation at the Jungfrauoch, 3580 m asl, Switzerland, *Journal of Geophysical Reserach*, 105, 2000.

- Zellweger, C., Forrer, J., Hofer, P., Nyeki, S., Schwarzenbach, B., Weingartner, E., Ammann, M., and Baltensperger, U.: Partitioning of reactive nitrogen ( $\text{NO}_y$ ) and dependence on meteorological conditions in the lower free troposphere, *Atmospheric Chemistry and Physics*, 3, 779-796, 2003.
- Zhang, L., Jacob, D. J., Boersma, K. F., Jaffe, D. A., Olson, J. R., Bowman, K. W., Worden, J. R., Thompson, A. M., Avery, M. A., Cohen, R. C., Dibb, J. E., Flocke, F. M., Fuelberg, H. E., Huey, L. G., McMillian, W. W., Singh, H. B., and Weinheimer, A. J.: Transpacific transport of ozone pollution and the effect of recent Asian emission increases on air quality in North America: an integrated analysis using satellite, aircraft, ozonesonde, and surface observations, *Atmospheric Chemistry and Physics*, 8, 6117-6136, 2008.
- Zhu, L., V. H. Payne, T. W. Walker, J. R. Worden, Z. Jiang, S. S. Kulawik, and E. V. Fischer (2017), PAN in the eastern Pacific free troposphere: A satellite view of the sources, seasonality, interannual variability, and timeline for trend detection, *J. Geophys. Res. Atmos.*, 122, 3614–3629, doi:10.1002/2016JD025868.
- Zhu, L., Fischer, E. V., Payne, V. H., Worden, J. R., and Jiang, Z.: TES observations of the interannual variability of PAN over Northern Eurasia and the relationship to springtime fires, *Geophysical Research Letters*, 42, 7230-7237, 10.1002/2015GL065328, 2015.

**Table 1: Models contributing to the HTAP1 simulations (SR1, SR6xx, and COfromXX) used in this study.**

Model	Resolution (lat-lon-layers)	Institute	Model contact	SR1	SR6xx	COfromxx	Plotting symbol
CAMCHEM-3311m13	2.5°x2°x30	NCAR, USA	Peter Hess	X	X	X	Filled circle
FRSGCUCI-v01	2.81°x2.81°x37	Lancaster Univ., UK	Oliver Wild	X	X	X	Filled upward triangle
GEMAQ-v1p0	2°x2°x28	York Univ., Canada	Alex Lupu	X	X	X	Filled downward triangle
GEOSChem-v07	2.5°x2°x30	Harvard Univ., USA	Rokjin Park	X	X	X	Filled diamond
GISS-PUCCINI-modelE	5°x4°x23	NASA GISS, USA	Drew Shindell	X	X	X	Filled square
GMI-v02f	2.5°x2°x42	NASA GSFC, USA	Bryan Duncan	X	X	X	Open circle
LMDZ3-INCA1	3.75° x 2° x 19	CEA, France	Sophie Szopa	X	X		Open upward triangle
LLNL-IMPACT-T5a	2.5° x 2° x 48	LLNL, USA	Dan Bergmann	X			Open downward triangle
MOZARTGFDL-v2	1.88° x 1.88° x 28	NOAA GFDL, USA	Arlene Fiore	X	X	X	Open diamond
MOZECHE-v16	1.88° x 1.88° x 28	FZ Julich, Germany	Martin Schultz	X	X	X	Open square
STOC-HadAM3-v01	5° x 5° x 19	University of Edinburg, UK	Ruth Doherty, David Stevenson	X	X	X	Plus sign
STOCHEM-v02	3.75° x 2.5° x 20	Met Office, Hadley Center, UK	Bill Collins, Michael	X			X



			Sanderson				
TM5-JRC-cy2- ipcc-v1	1° x 1° x 25	JRC, Italy	Frank Dentener	X	X	X	Filled right facing triangle
UM-CAM-v01	3.75° x 2.5° x 19	University of Cambridge, UK	Guang Zeng	X	X	X	Filled left facing triangle

757

758

759

760

761

762

763

764

765

766

767

768

769

**Table 2: Simulations from HTAP1 used in this study.**


Simulation	Description
SR1	Base case (see Section 2.1 for details)
SR6EA	SR1 but with anthropogenic emissions of all O <sub>3</sub> precursors (NO <sub>x</sub> +CO+NMVOC) and aerosols within EA decreased by 20%
SR6EU	SR1 but with 20% emissions reductions within the EU region
SR6NA	SR1 but with 20% emissions reductions within the NA region.
COfromEA	Idealized tracer simulation in which all models use identical CO emissions, emitted within the EA region, with a 50-day e-folding lifetime.
COfromEU	Same as COfromEA but for the EU region.
COfromNA	Same as COfromEA but for the NA region.

776 **Table 3: Mountaintop sites with multiple years of PAN observations used in this study.**


Site	Location	Elevation	Measurement Period (s)	Reference (s)
Mount Bachelor	43.979° N, 121.687° W	2763m	3 April – 18 June 2008, 30 August – 7 October 2008, 26 March – 20 May 2009, 23 March – 25 May 2010	(Fischer et al., 2010;Fischer et al., 2011)
Hohenpeissenberg	47.80° N, 11.02° E	985 m	January 2003 – December 2008	<a href="http://www.dwd.de/de/GAW">http://www.dwd.de/de/GAW</a> (Gilge et al., 2010)
Jungfraujoch	46.55°N, 7.98°E	3580 m	April 1997 – May 1998, Aug 30 2005 – Sept 16 2005, Throughout 2005, but not continuous	(Balzani Lööv et al., 2008;Carpenter et al., 2000;Zellweger et al., 2000;Zellweger et al., 2003)
Zugspitze	47.42° N, 10.98° E	2960 m	May 2004 – December 2008	<a href="http://gaw.kishou.go.jp">http://gaw.kishou.go.jp</a>
Schauinsland	47.92° N, 7.92° E	1205m	January 1995 – December 2010	www.umweltbundesamt.de

777


778




**Figure 1: Multi-model monthly mean total PAN mixing ratios (black circles and solid lines) at Mount Bachelor (left) and Jungfraujoch (right). We take the difference between the base simulation (SR1) and one in which emissions are decreased by 20% and then multiply the difference by 5 to estimate a 100% contribution associated with anthropogenic precursor emissions from Europe (green), North America (red), East Asia (blue). The sum of the anthropogenic contribution from these three regions is shown (dashed black) for comparison with total simulated PAN.**




**Figure 2: Multi-model ensemble ( $n=14$ ; Table 1) average PAN mixing ratios (ppt; left panels) and relative standard deviation (the absolute standard deviation across the models divided by the ensemble mean; right panels) at 650 hPa in April; relative standard deviations are masked out (white) for regions where multi-model mean PAN falls below 100 ppt. The models were sampled at 650 hPa by vertically interpolating between the bounding grid cells and then re-gridded horizontally to a common  $1^\circ \times 1^\circ$  grid. White lines denote the HTAP1 source regions: North America (NA), Europe and North Africa (EU), and East Asia (EA) from left to right. White circles indicate the five mountain sites with multi-year PAN observations used in our analysis (note: Zugspitze and Hohenpeissenberg are too close to differentiate on the map; see Table 3). Mount Waliguan in Asia, where we lack multi-year measurements but conduct model analysis, is denoted by the white square.**



**Figure 3. April mean PAN abundances (ppt) simulated (black symbols, one per model as defined in Table 1; blue circles offset to the left show multi-model mean values) and measured (red circles offset to the right of the model values) at northern mid-latitude mountaintop sites: Mount Bachelor (MBO), Zugspitze (ZUG), Jungfraujoch (JFJ), Hohenpeissenberg (HOH), Schauinsland (SCH) and Mount Waliguan (MTW). The observed year 2001 April mean, which corresponds to the meteorological year used by most of the models, at Schauinsland is shown in blue to the right of the models.**



**Figure 4. Simulated total PAN versus source-receptor relationships (SRRs) at each of three northern mid-latitude sites. For Jungfraujoch and Mount Bachelor, vertical red lines bound the observed range in total PAN. For source-receptor pairs with significant correlations ( $p \leq 0.05$ ), models falling within the observed range (across years) are colored red, and horizontal red dashed lines extend to the ordinate, representing the emergent constraint (narrower range resulting from selecting only those models falling in the observed range of total PAN). At Jungfraujoch, the range (across years) in PAN attributed to the EU source region by back-trajectory analysis (Pandey Deolal *et al.*, 2013) is indicated by horizontal dashed blue lines. Individual models are denoted by the symbols defined in Table 1.**



**Figure 5: SRRs diagnosed as the difference between the SR1 and SR6xx simulations in Table 1 for PAN (ppt) at Jungfraujoch (top), Mount Bachelor (middle), and Mount Waliguan (bottom) in each HTAP1 model (see Table 1 for symbol assigned to each model) versus the annual emission of anthropogenic VOC (AVOC; Tg C a<sup>-1</sup>) within the NA (left), EU (middle) and EA (right) source regions. The Spearman rank correlation coefficient (more robust to outliers than the traditional Pearson coefficient) and associated p-value are shown in each panel. The horizontal red lines correspond to the values identified with the red symbols in Figure 4.**






Figure 6: Ratio of the PAN response to 20% emission reductions within NA versus EA plotted against (a) total PAN and (b) the ratio of idealized tracers of model transport emitted from NA versus EA (COfromNA/COfromEA; see Table 2) at Mount Bachelor as simulated by the HTAP1 models. Each symbol in (a) represents a model as defined in Table 1; the range of observed total PAN at Mount Bachelor is indicated by the black vertical lines. The colored symbols in (b) represent the meteorological fields used in the simulation: blue triangles for GEOS winds; red circles for NCEP; black diamonds for ECMWF; cyan upside-down triangles for CMC; green squares for general circulation models forced by observed sea surface temperatures and sea ice. Both panels show Spearman rank correlation coefficients and p-values, as well as a black dashed horizontal line at 1 to separate the models suggesting a higher NA influence (above) versus higher EA influence (below) on PAN SRRs.





**Figure 7: SRRs for  $O_3$  versus PAN at Jungfraujoch (top), Mount Bachelor (middle), and Mount Waliguan (bottom), obtained by subtracting the SR6XX from the SR1 simulations (Table 2) available from 12 models, where XX denotes the NA (left), EU (middle) or EA (right) source region. Each model thus contributes one point (symbols defined in Table 1) in each panel. Spearman (rank) correlation coefficient and p-values are also shown.**



**Figure 8: SRRs for O<sub>3</sub> versus PAN in surface air over each of the HTAP1 northern mid-latitude continent regions: EU (top), NA (middle), and EA (bottom), obtained by subtracting the SR6XX from the SI simulations, where XX denotes the NA (left), EU (middle) or EA (right) source region. Spearman (rank) correlation coefficient and p-values are also shown. Symbols denote individual models as defined in Table 1. STOC-HadAM3-v01 is excluded here as an outlier that artificially raised the correlation significance.**