

Supplementary Information

Particle Measurements at Alert, Nunavut

The experimental methods for the particle microphysical and filter-based chemical measurements in Fig. 7 are discussed by Leitch et al. (2013, 2018).

Monthly-averaged values of $N_{>50}$ (number concentration > 50 nm diameter), $N_{>100}$, N_{15-30} (number concentration in the size interval 15 nm to 30 nm), N_{30-50} , and N_{50-100} for the inclusive period of 2012–2014 are shown in Fig. 7a.

One estimate of the effect of growth of newly-formed particles on the concentrations of particles in the range of 15–100 nm is demonstrated using Fig. 2. Baseline concentrations are estimated by linearly interpolating each of the N_{15-30} , N_{30-50} and N_{50-100} from May to October. These are shown as the dashed lines in Fig. 7a. The increases in N_{15-30} , N_{30-50} and N_{50-100} during the summer months, relative to the estimated baseline concentrations, are shown in Fig. 7b. The progressive decrease in N_{15-30} to N_{30-50} to N_{50-100} is indicative of the average of growth from smaller to larger sizes. There is no obvious increase in particles larger than 100 nm during the summer, suggesting growth to such sizes is mostly undetectable by this approach. Despite the high frequency of NPF, the growth to 50 nm or larger during August is almost identical to that in July, suggesting that the levels of precursors for condensational growth are on average lower during August than July. The monthly-average OM/SO_4^{2-} , also shown in Figure 7b (from Leitch et al. (2018)), increases substantially and coincidentally with the increases in 15–50 nm particle concentrations, suggesting that organic components are important for the growth of newly formed particles in the summer Arctic.

References

Leitch, W. R., Sharma, S., Huang, L., Toom-Sauntry, D., Chivulescu, A., Macdonald, A. M., von Salzen, K., Pierce, J. R., Betram, A. K., Schroder, J. C., Shantz, N. C., Chang, R. and Norman, A. L.: Dimethyl sulfide control of the clean summertime Arctic aerosol and cloud, *Elementa*, 1, 1–000017, doi:10.12952/journal.elementa.000017, 2013.

Leitch, W. R., Russell, L. M., Liu, J., Kolonjari, F., Toom, D., Huang, L., Sharma, S., Chivulescu, A., Veber, D. and Zhang, W.: Organic functional groups in the submicron aerosol at 82.5° N, 62.5° W from 2012 to 2014, *Atmos. Chem. Phys.*, 18(5), 3269–3287, doi:10.5194/acp-18-3269-2018, 2018.