

We would like to thank the reviewers for the valuable and constructive comments, which help us to improve the manuscript. Listed below are our point-by-point responses to the comments, including the corresponding changes made to the revised manuscript. The reviewers' comments are marked in black and our answers are marked in blue, and the revision in the manuscript is further formatted as '*Italics*'.

#### **Referee #1:**

This paper shows a new and innovative method of trying to represent observed changes in the 'coating' of black carbon, as detected as changes in the  $D_p/D_c$  metric produced by the DMT SP2. This works by combining an emission field with a Lagrangian transport model and an empirical parameterisation for the evolution of black carbon in the atmosphere. While this 'k' parameter is tuned to the measurements, the model shows an impressive correlation with the observations, giving confidence that there is value in the technique. The authors then go on to make estimates of the contributions from inside and outside the city to the measured concentrations and properties.

While not an all-encompassing process model, this does present an intriguing new way of looking at the data that could conceivably bridge the gap between a highly detailed process model such as PARTMC-MOSAIC and Eulerian chemical transport models like WRF-CHEM. It is also good that it provides another new perspective on the phenomenon of pollution 'building up' in Beijing, offering further evidence that this is driven by regional transport and transformation rather than local sources and processes. While the main innovation with this paper is technical, the authors do use it to interpret air quality sources and phenomena, so I would say this is in-scope for ACP (as opposed to AMT or GMD). While I am not completely convinced of the results (see below), I think that the novelty of the technique alone means that it deserves to be discussed within the scientific literature. Overall, the quality of English is good, but I do have certain issues regarding how the results are interpreted and presented. As such, I recommend publication after revisions.

#### **1. General comments:**

While the earlier sections of the paper were well written, I was not impressed with how the paper interpreted the implications of the results. Sections 4 and 5 in particular seem to just largely restate what was already said in section 3 in different ways, so this entire portion of the paper could do with rewriting and sharpening up. But more generally, I would question what the key implications of this paper are; the conclusions seem to work off the reduction of Babs being a regulatory motivation, however most evidence supports BC being a more important metric for human health, which while related, is more directly related to emissions rather than processing. While the evolution of the mixing state of BC is important to consider for wet removal and radiative transfer, I would say it is debatable as to whether reducing the coatings (as opposed to overall BC) should be considered an objective for policymaking (while BC is recognized as a

global climate forcing agent, the majority of this is likely from biomass burning). It is my opinion that the conclusions of this paper are more applicable to process-level atmospheric science, except for the part where the contributions to bulk BC are discussed. I would recommend the discussion and conclusions be reframed accordingly.

**Response:** Thanks for the suggestions. Following the reviewer's suggestion, we have rewritten and sharpened the Sections 4 (Discussion) and 5 (Conclusion). In terms of the Discussion section in the revised manuscript, we specifically focused on:

- presenting the simplified scheme of our model and its dependence on the model parameters,
- discussing the robustness of our model and how to extend our model calculation on BC mixing state at other seasons (e.g. summer) except for autumn/winter,
- exploring the possible mechanisms for BC aging during atmospheric transport in north china plain based on the retrieved aging rate coefficient, which is comparing with observations and those used in other models,
- implication in process-level atmospheric science.

To sharpen the conclusion of this work, we reorganized the Concluding Remarks section, as “*The effect of BC-containing particles on air quality and climate is not only dominated by BC mass concentration but also controlled by their mixing state. To better understand the mixing state of atmospheric BC in China, we developed a new approach to simulate the BC aging process during atmospheric transport based on the BC emission inventory and back-trajectory analysis. Our models track the BC mixing state (i.e.,  $D_p/D_c$  ratio) from an emitted source origin (e.g., a  $0.25^\circ \times 0.25^\circ$  grid) to a receptor (i.e., Tsinghua site). The model calculation can quantify the mass-averaged  $D_p/D_c$  ratio of overall BC particles over the receptor site from various origins, which agreed well with observed ones. The simulations can provide information on the BC mixing state with fine temporal and spatial resolutions.*

*Based on the simulations of BC mixing state during atmospheric transport, we find a strong dependence of BC mixing state on emissions during atmospheric transport. BC particles with higher aging degrees at our site were mainly from more intensive emission origins (e.g., southern Hebei) due to higher aging rates. On the other hand, when BC particles emitted from clean origins passed through polluted regions, they were also characterized by a greater  $D_p/D_c$  ratio by speeding up the aging process in polluted air. Our simulations demonstrated the importance of regional transport in BC light absorption in Beijing under polluted conditions. This provides a new perspective on the phenomenon of pollution building up in Beijing, further demonstrating that this is driven by regional transport and transformation rather than local sources and processes..”*

2. Specific comments:

(1) I found this paper extremely abstract and hard to read in places regarding certain quantities and it took me multiple reads before I think I began to understand what was going on. In particular, I would have appreciated a more intuitive explanation of what things like 'k' and 'EEI' physically represent (along with many others).

**Response:** Thanks for the comment. Following the reviewer's suggestion, we have made an explanation on a more intuitive explanation of what things like 'k' and 'EEI' physically represent. We have added the descriptions regarding the terms ('k', 'EEI', "TE" and " $(\frac{D_p}{D_c})_{ini}$ ") in the revised manuscript, as shown below:

*"The parameter k characterizes the rate coefficient of coating materials produced on the surface of BC by atmospheric aging such as condensation, coagulation and cloud process, which is influenced by meteorological factors, chemistry, aerosol phase state as well as other parameters (e.g., particle size)."*

*"The EEI represents the effective BC amount transported to the receptor site from the emission origins, taking into account the magnitude of BC emission from origin regions, the transport, hydrophobic-to-hydrophilic transformation, as well as dry and wet depositions during atmospheric transport."*

*"The TE is defined to quantify the transport ability of BC from origin regions to the receptor site based on transformation (i.e., hydrophobic-to-hydrophilic BC) and removal processes of BC (i.e., dry and wet depositions) in the atmospheric."*

*"The  $(\frac{D_p}{D_c})_{ini}$  represents the initial the initial value of the  $D_p/D_c$  ratio of BC particles before atmospheric transport, which characterizes the BC aging degree near emissions."*

(2) More discussion should be given to the possible mechanisms for the increase in  $D_p/D_c$ , i.e. coagulation and secondary aerosol formation. Given that there are already other models out there that consider these processes, is it possible to compare the value of the 'k' parameter to equivalent timescales in other models?

**Response:** Thanks. Following the reviewer's suggestion, we have added the statement on the possible mechanisms for the increase in  $D_p/D_c$ . In our work, the aging rate coefficient (i.e.,  $k_{aging}$ ) is determined to be  $\sim 17\% \text{ h}^{-1}$ , corresponding to the timescale of BC aging process about  $\sim 6 \text{ h}^{-1}$ . Riemer et al. (2004) revealed that the time scale of BC aging process by condensation (2-8 h) was significantly shorter than that by coagulation (10-40 h). Higher aging rate coefficient obtained in our work indicates the BC aging in north china plain is most likely dominated by condensation process.

To the second question, the  $k_{\text{aging}}$  used in our models ( $\sim 17\% \text{ h}^{-1}$ ) is comparable with the observed values in north china plain (i.e.,  $\sim 20\% \text{ h}^{-1}$  at Yufa site (Cheng et al., 2012) and  $\sim 21\% \text{ h}^{-1}$  at Xianghe site (Zhang et al., 2018)), but significantly higher than those ( $1\text{-}5\% \text{ h}^{-1}$ ) derived in developed countries and then used in models (Moteki et al., 2007; Shiraiwa et al., 2007; Cooke and Wilson, 1996; Lohmann et al., 2000; Jacobson, 2001; Koch, 2001). The difference between  $k_{\text{aging}}$  used in our model and other models is consistence with the distinction between the timescale of BC aging measured in Beijing ( $\sim 4.6 \text{ h}$  corresponding to  $k_{\text{aging}}$  of  $\sim 22\% \text{ h}^{-1}$ ) and Houston ( $\sim 20 \text{ h}$  corresponding to  $k_{\text{aging}}$  of  $\sim 5\% \text{ h}^{-1}$ ) using an environment chamber.

Correspondingly, the related discussion has been added in the revised manuscript, as “*The retrieved aging rate coefficient  $k_{\text{aging}}$  ( $\sim 17\% \text{ h}^{-1}$ , corresponding to the timescale of BC aging process about  $\sim 6 \text{ h}^{-1}$ ) indicated BC particles under polluted environment at autumn/winter in NCP underwent fast aging during atmospheric transport. The retrieved  $k_{\text{aging}}$  with a value of  $\sim 17\% \text{ h}^{-1}$  used in our models is comparable with the observed values at other suburban sites in NCP, namely up to  $\sim 20\% \text{ h}^{-1}$  and  $\sim 21\% \text{ h}^{-1}$  at Yufa site and Xianghe site, respectively (Cheng et al., 2012; Zhang et al., 2018). However, the aging rate coefficient commonly used in other models ( $1\text{-}5\% \text{ h}^{-1}$ ) was significantly smaller than that used in our model (Cooke and Wilson, 1996; Jacobson, 2001; Koch, 2001; Lohmann et al., 2000;), indicating that the values used in previous models derived in developed countries (Moteki et al., 2007; Shiraiwa et al., 2007) could not represent the characteristic of BC aging process in China. A similar conclusion also found by Peng et al., (2007) using an environment chamber, namely the timescale of BC aging with significant distinction between urban cities in Beijing ( $\sim 5 \text{ h}$  corresponding to  $k_{\text{aging}}$  of  $\sim 20\% \text{ h}^{-1}$ ) and Houston ( $\sim 18 \text{ h}$  corresponding to  $k_{\text{aging}}$  of  $\sim 5.6\% \text{ h}^{-1}$ ). Higher aging rate coefficient suggested the BC aging under polluted environment in NCP is most likely dominated by condensation process during atmospheric transport, taking into account a clear difference between timescale of BC aging by condensation ( $2\text{-}8 \text{ h}$ ) and coagulation ( $10\text{-}40 \text{ h}$ ) (Riemer et al., 2004). ”*

- (3) Ultimately, while good correlation between measurement and model is reached, it is possible that other yet-to-be-identified factors may be responsible for this agreement rather than the 'k' parameter being a good representation of ageing, which is the working hypothesis. The authors should spend more time discussing how best to further test the robustness of this model. In particular, a major limitation of this work is that it is restricted to autumn/winter datasets. During the summer in Beijing, there is typically much more photochemistry (reflected in high ozone concentrations) but BC concentrations and coating thicknesses are both reduced according to Liu et al. (<https://www.atmos-chem-phys-discuss.net/acp-2018-1142/>). Can this result be reconciled with this model? Would it mean that the 'k' value would need to have a seasonal dependence if this model were to be extended to other

months?

**Response:** Thanks to the reviewer to point this out. Following reviewer's suggestion, we have discussed how best to further test the robustness of our model. The uncertainties of our model calculation are strongly related to the retrieved parameters of aging rate coefficient ( $k_{\text{aging}}$ ) and the initial value of  $D_p/D_c$  ratio ( $(\frac{D_p}{D_c})_{\text{ini}}$ ). It is noted that the  $k_{\text{aging}}$  value ( $\sim 17\% \text{ h}^{-1}$ ) in this work is retrieved by the measurements under polluted environment at autumn/winter in Beijing. The retrieved  $k_{\text{aging}}$  in this work can represent the characteristics of BC aging in polluted NCP at autumn/winter. In terms of BC aging at summer, the observed aging rate coefficient at other two site (xianghe and Yufa) in NCP showed pronounced diurnal cycle with a maximum ( $\sim 20\% \text{ h}^{-1}$ ) at noon time due to stronger photochemistry (Cheng et al., 2012; Zhang et al., 2018). Figure R1 shows the aging rate coefficient of BC in the range of 0.2-21%  $\text{h}^{-1}$  at summer in NCP. The maximum of  $k_{\text{aging}}$  at summer noontime was comparable with that at autumn/winter. However, other time especially nighttime is significantly smaller, which can explain thinner coatings of BC at summer compared with those at autumn/winter (Liu et al., 2019).

On the other hand, the initial values of  $D_p/D_c$  ratio ( $(\frac{D_p}{D_c})_{\text{ini}}$ ) used in our model at autumn/winter are also different with that at summer. The BC transported to Beijing at autumn/winter are dominated by industrial and residential emissions with higher  $D_p/D_c$  ratio near sources; however, the sources of BC in Beijing at summer are controlled by traffic and industrial emission and with smaller  $(\frac{D_p}{D_c})_{\text{ini}}$ , which can also explain thinner coatings of BC at summer compared with those at autumn/winter (Liu et al., 2019).

In summary, our model can be used to calculate BC mixing state during atmospheric transport at different seasons with distinct  $k_{\text{aging}}$  and  $(\frac{D_p}{D_c})_{\text{ini}}$  values. Both  $k_{\text{aging}}$  and  $(\frac{D_p}{D_c})_{\text{ini}}$  values show a seasonal dependence. Quantification of BC mixing state at different seasons using our model will be the subject of future work. In this work, we focused on the development of model.




Fig. S1 Diurnal cycle of BC aging rate coefficient ( $k_{\text{aging}}$ ) at summer in Xianghe site (data from Zhang et al., (2018)).

To make this point clear, the related discussion has been added in the revised manuscript, as “Although our model was developed and validated with autumn/winter datasets, it can be applied in other seasons with distinct model parameters (i.e. aging rate coefficient and the initial value of  $D_p/D_c$  ratio). The two parameters exhibit a seasonal dependence. In terms of the aging rate coefficient at summer (Cheng et al., 2012; Zhang et al., 2018), the measurements at two suburban site (xianghe and Yufa) in NCP showed pronounced diurnal cycle in the range of 0.2-20%  $\text{h}^{-1}$  with a maximum at noon time due to stronger photochemistry. The  $k_{\text{aging}}$  around noontime at summer was comparable with that ( $\sim 17\% \text{ h}^{-1}$ ) at autumn/winter. However, other time especially nighttime at summer is significantly smaller. For the  $(\frac{D_p}{D_c})_{\text{ini}}$ , the seasonal dependence is due to different dominated sources of BC. The BC particles in Beijing at summer are controlled by traffic and industrial emission with smaller higher  $D_p/D_c$  ratio near sources, but the dominated sources of BC at autumn/winter are industrial and residential emissions with higher  $(\frac{D_p}{D_c})_{\text{ini}}$  (D. Liu et al., 2019; H. Liu et al., 2019). In summary, lower values both of aging rate coefficient and the initial value of  $D_p/D_c$  ratio would lead to thinner coatings of BC at summer compared with those at autumn/winter in NCP, which was consistence with observations in Beijing (D. Liu et al., 2019).”

## References

Cheng, Y. F., Su, H., Rose, D., Gunthe, S. S., Berghof, M., Wehner, B., Achtert, P., Nowak, A., Takegawa, N., Kondo, Y., Shiraiwa, M., Gong, Y. G., Shao, M., Hu, M., Zhu, T., Zhang, Y. H., Carmichael, G. R., Wiedensohler, A., Andreae, M. O., and Pöschl, U.: Size-resolved measurement of the



1 mixing state of soot in the megacity Beijing, China: diurnal cycle, aging and parameterization, *Atmos.*  
2 *Chem. Phys.*, 12, 4477-4491, 2012.

3 Cooke, W. F. and Wilson, J. J. N.: A global black carbon aerosol model, *J. Geophys. Res.*, 101, 19395–  
4 19409, 1996.

5 Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols,  
6 *Nature*, 409, 695-697, 2001.

7 Koch, K.: Transport and direct radiative forcing of carbonaceous and sulphate aerosols in the GISS GCM,  
8 *J. Geophys. Res.*, 106, 20311–20332, 2001.

9 Liu, D., Joshi, R., Wang, J., Yu, C., Allan, J. D., Coe, H., Flynn, M. J., Xie, C., Lee, J., Squires, F.,  
10 Kotthaus, S., Grimmond, S., Ge, X., Sun, Y., and Fu, P.: Contrasting physical properties of black carbon  
11 in urban Beijing between winter and summer, *Atmos. Chem. Phys.*, 19, 6749-6769, 10.5194/acp-19-6749-  
12 2019, 2019.

13 Liu, H., Pan, X., Liu, D., Liu, X., Chen, X., Tian, Y., Sun, Y., Fu, P., and Wang, Z.: Mixing characteristics  
14 of refractory black carbon aerosols determined by a tandem CPMA-SP2 system at an urban site in Beijing,  
15 *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-244>, in review, 2019.

16 Lohmann, U., Feichter, J., Penner, J., and Leaitch, R.: Indirect effect of sulfate and carbonaceous aerosols:  
17 a mechanistic treatment, *J. Geophys. Res.*, 105, 12193–12206, 2000.

18 Moteki, N., Kondo, Y., Miyazaki, Y., Takegawa, N., Komazaki, Y., Kurata, G., Shirai, T., Blake, D. R.,  
19 Miyakawa, T., and Koike, M.: Evolution of mixing state of black carbon particles: Aircraft measurements  
20 over the western Pacific in March 2004, *Geophys. Res. Lett.*, 34, 2007.

21 Peng, J., Hu, M., Guo, S., Du, Z., Zheng, J., Shang, D., Levy Zamora, M., Zeng, L., Shao, M., Wu, Y.-  
22 S., Zheng, J., Wang, Y., Glen, C. R., Collins, D. R., Molina, M. J., and Zhang, R.: Markedly enhanced  
23 absorption and direct radiative forcing of black carbon under polluted urban environments, *Proc. Natl.*  
24 *Acad. Sci. USA*, 113, 4266-4271, 2016.

25 Riemer, N., Vogel, H., and Vogel, B.: Soot aging time scales in polluted regions during day and night,  
26 *Atmos. Chem. Phys.*, 4, 1885-1893, 10.5194/acp-4-1885-2004, 2004.

27 Shiraiwa, M., Kondo, Y., Moteki, N., Takegawa, N., Miyazaki, Y., and Blake, D. R.: Evolution of mixing  
28 state of black carbon in polluted air from Tokyo, *Geophys. Res. Lett.*, 34, L16803,  
29 doi:10.1029/2007GL029819, 2007.

30 Zhang, Y., Su, H., Ma, N., Li, G., Kecorius, S., Wang, Z., Hu, M., Zhu, T., He, K., Wiedensohler, A.,  
31 Zhang, Q., and Cheng, Y.: Sizing of Ambient Particles From a Single-Particle Soot Photometer  
32 Measurement to Retrieve Mixing State of Black Carbon at a Regional Site of the North China Plain, *J.*  
33 *Geophys. Res.-Atmos.*, 123, 12,778-712,795, doi:10.1029/2018JD028810, 2018b.

**Referee #2:**

The work developed a new approach to simulate the BC mixing state based on an emissions inventory and back-trajectory analysis. They quantified the mass-averaged aging degree of total BC particles and found the aging process of BC during atmospheric transport showed that it strongly dependent on emission levels. In general, from the modelling development to the modelling implication, I think that the result is worth to publish in the ACP after one minor revision.

1. Introduction P2L16-17 need references such as Li et al., JGR, 121(22), 13,784-713,798.

**Response:** Thanks. We have added some related references (Li et al., 2016; Jacobson, 2001; Moffet et al., 2009; Liu et al., 2017).

2. P2L39 The author review several method to determine the BC particles. Also, Wang et al., ESTL, 2017 measure the Df in polluted air following soot particle aging. Does the model possibly consider the dimension fraction of BC? That would be more interesting.

**Response:** Thanks for the comment. We have added some statement on change in BC morphology (quantified by dimension fraction) associated with BC mixing state under polluted conditions, as *“Moreover, Li et al., (2016) and Wang et al., (2017) revealed that the significant change in BC morphology (e.g. increase of fraction dimension) associated with their mixing state (i.e., from bare-like or partly coated to embedded BC) in polluted air, which could enhance BC light absorption (Liu et al., 2015; Cappa et al., 2012).”*

3. P7L24 deleted were P7L25 identified to indicated P7L27 import?

**Response:** Thanks. We have revised.

4. Change the word P9L2-L3 Here it is possible to consider how RH influence BC coating? If the authors want to get the conclusion, firstly you need to exclude other possible factors. Therefore, as you try to do it.

**Response:** Thanks. In our model calculation, we just considered the effect of emissions on BC mixing state, and excluded the influences of other possible factors. A rather simplified scheme was adopted where the aging rate is assumed proportional to the emissions. Actually, the aging rate coefficient ( $k$ ) in our models was related to meteorological factors (e.g., temperature, relative humidity (RH)), chemistry (e.g., oxidation), aerosol phase state (e.g., heterogeneous reaction) as well as other parameter (e.g., particle size). The aging rate coefficient was presumed as a constant in the model calculation, and therefore the change in BC aging degree obtained from our simulations is only caused by distinct emissions.



To make this point clear, the related statement in the revised manuscript was “*In this study, a rather simplified scheme was adopted where the aging rate is assumed proportional to the emissions without detailed consideration of the effects of temperature, particle sizes, phase state, hygroscopicity and chemistry (Riemer et al. 2009; Cheng et al. 2008, 2012, 2015; Mu et al. 2018). Actually, these factors can influence the model parameter of aging rate coefficient. Our simulations strongly depends on the emission, initial value of  $D_p/D_c$  ratio and aging rate coefficient.*”

5. P9L6-13 Here some logical problem. if the aerosol in clean regions pass through the polluted area, how you know these BC don't deposit. How do you know BC at the sampling site were from clean region instead of the close polluted areas. Seemly, these are difficult questions about the mixture of air masses during the transport. How the authors separate them here?

**Response:** We thank the reviewer for raising these questions. To the first question, our model calculations consider the dry and wet deposit. In Eq. (7) in the manuscript, the  $TE$  represents the transport efficiency, taking into account transformation, transport, and removal processes of BC in the atmosphere. The  $TE$  is be treated as a weighting factor to quantify effective amount of BC from the emission origin to receptor site.

To the second question, we can separate BC from various emission origins (i.e.,  $0.25^\circ \times 0.25^\circ$  grid). Here, we discussed the gridded  $D_p/D_c$  ratio, representing the aging degree of BC particles that are transported to the receptor site from a certain origin (i.e., a grid  $h$ ) rather than all origins. Our model calculation can track the evolution of BC aging degree from a certain origin (i.e., a grid  $h$ ) to the receptor site throughout different regions.

6. P10-P11, the implications and discussion should be combined? The implications should not be in Result section. The authors need to consider the structure in the two section.

**Response:** Thanks. We have changed “3.3 Implication for BC light absorption” into “BC light absorption in the atmosphere”

7. P12L10-11, capability should be individual BC or total here? My question is that the statement could be one problem. If you think total capability of BC in upper layer is higher that of BC in the lower. You need to know the BC distribution in the column within PBL. Normally the BC concentration is much higher than them in upper layer. Even mixing state is higher in upper layer but their concentration is lower. Therefore, the total capacity need be questioned.

**Response:** Thanks. Sorry for the misleading and here the capability of BC light absorption represents the light absorption per unit BC mass, namely mass cross section (MAC) of BC. The MAC of BC strongly depends on their mixing state due to lens effect of coating materials on BC surface. The

MAC and BC mass concentration ( $C_{BC}$ ) are two factors to determine BC light absorption ( $\sigma_{ab}$ ), namely  $\sigma_{ab} = \text{MAC} \times C_{BC}$ .

To make it clear, we have changed the “*capability of BC light absorption*” into “*capability of BC light absorption (i.e., mass cross section of BC-containing particles including coating materials on BC surface)*”.

8. P12 section 5 concluding remarks. Most of them repeat the results or others. Is that possible to shorten it? The details should not be concluded here such as P13L2-5 and so on. The reference should be removed in the section.

**Response:** Thanks. Following the reviewer’s suggestion, we have sharpened the Concluding Remarks section, as “*The effect of BC-containing particles on air quality and climate is not only dominated by BC mass concentration but also controlled by their mixing state. To better understand the mixing state of atmospheric BC in China, we developed a new approach to simulate the BC aging process during atmospheric transport based on the BC emission inventory and back-trajectory analysis. Our models track the BC mixing state (i.e.,  $D_p/D_c$  ratio) from an emitted source origin (e.g., a  $0.25^\circ \times 0.25^\circ$  grid) to a receptor (i.e., Tsinghua site). The model calculation can quantify the mass-averaged  $D_p/D_c$  ratio of overall BC particles over the receptor site from various origins, which agreed well with observed ones. The simulations can provide information on the BC mixing state with fine temporal and spatial resolutions.*

*Based on the simulations of BC mixing state during atmospheric transport, we find a strong dependence of BC mixing state on emissions during atmospheric transport. BC particles with higher aging degrees at our site were mainly from more intensive emission origins (e.g., southern Hebei) due to higher aging rates. On the other hand, when BC particles emitted from clean origins passed through polluted regions, they were also characterized by a greater  $D_p/D_c$  ratio by speeding up the aging process in polluted air. Our simulations demonstrated the importance of regional transport in BC light absorption in Beijing under polluted conditions. This provides a new perspective on the phenomenon of pollution building up in Beijing, further demonstrating that this is driven by regional transport and transformation rather than local sources and processes.”*

## References

Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., Jobson, B. T., Kolesar, K. R., Lack, D. A., Lerner, B. M., Li, S.-M., Mellon, D., Nuaaman, I., Olfert, J. S., Petäjä, T., Quinn, P. K., Song, C., Subramanian, R., Williams, E. J., and

1 Zaveri, R. A.: Radiative Absorption Enhancements Due to the Mixing State of Atmospheric Black  
2 Carbon, *Science*, 337, 1078-1081, 2012.

3 Cheng, Y. F., Wiedensohler, A., Eichler, H., Su, H., Gnauk, T., Brüggemann, E., Herrmann, H.,  
4 Heintzenberg, J., Slanina, J., Tuch, T., Hu, M., and Zhang, Y. H.: Aerosol optical properties and related  
5 chemical apportionment at Xinken in Pearl River Delta of China, *Atmos. Environ.*, 42, 6351-6372,  
6 <http://dx.doi.org/10.1016/j.atmosenv.2008.02.034>, 2008.

7 Cheng, Y. F., Su, H., Rose, D., Gunthe, S. S., Berghof, M., Wehner, B., Achtert, P., Nowak, A.,  
8 Takegawa, N., Kondo, Y., Shiraiwa, M., Gong, Y. G., Shao, M., Hu, M., Zhu, T., Zhang, Y. H.,  
9 Carmichael, G. R., Wiedensohler, A., Andreae, M. O., and Pöschl, U.: Size-resolved measurement of the  
10 mixing state of soot in the megacity Beijing, China: diurnal cycle, aging and parameterization, *Atmos.*  
11 *Chem. Phys.*, 12, 4477-4491, 2012.

12 Cheng, Y., Su, H., Koop, T., Mikhailov, E., and Pöschl, U.: Size dependence of phase transitions in  
13 aerosol nanoparticles, *Nat Commun.*, 6, 5923, 2015.

14 Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols,  
15 *Nature*, 409, 695-697, 2001.

16 Li, W., Sun, J., Xu, L., Shi, Z., Riemer, N., Sun, Y., Fu, P., Zhang, J., Lin, Y., Wang, X., Shao, L., Chen,  
17 J., Zhang, X., Wang, Z., and Wang, W.: A conceptual framework for mixing structures in individual  
18 aerosol particles, *J. Geophys. Res.-Atmos.*, 121, 13,784-713,798, 10.1002/2016jd025252, 2016.

19 Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S.,  
20 Williams, P. I., Ting, Y.-C., Haslett, S., Taylor, J. W., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe,  
21 H., and Allan, J. D.: Black-carbon absorption enhancement in the atmosphere determined by particle  
22 mixing state, *Nature Geosci.*, 10, 184-188, 2017.

23 Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M. K., Cappa, C. D., Williams, L. R., Herndon, S. C.,  
24 Massoli, P., Fortner, E. C., Chhabra, P. S., Brooks, W. A., Onasch, T. B., Jayne, J. T., Worsnop, D. R.,  
25 China, S., Sharma, N., Mazzoleni, C., Xu, L., Ng, N. L., Liu, D., Allan, J. D., Lee, J. D., Fleming, Z. L.,  
26 Mohr, C., Zotter, P., Szidat, S., and Prévôt, A. S. H.: Enhanced light absorption by mixed source black  
27 and brown carbon particles in UK winter, *Nat Commun.*, 6, 8435, 2015.

28 Moffet, R. C., and Prather, K. A.: In-situ measurements of the mixing state and optical properties of soot  
29 with implications for radiative forcing estimates, *Proc. Natl. Acad. Sci. USA*, 106, 11872-11877,  
30 10.1073/pnas.0900040106, 2009.

31 Mu, Q., Shiraiwa, M., Octaviani, M., Ma, N., Ding, A., Su, H., Lammel, G., Pöschl, U., and Cheng, Y.:  
32 Temperature effect on phase state and reactivity controls atmospheric multiphase chemistry and transport  
33 of PAHs, *Sci. Adv.*, 4, 2018.

- 1 Riemer, N., West, M., Zaveri, R. A., and Easter, R. C.: Simulating the evolution of soot mixing state with  
2 a particle-resolved aerosol model, *J. Geophys. Res.-Atmos.*, 114, 10.1029/2008jd011073, 2009.
- 3 Wang, Y., Liu, F., He, C., Bi, L., Cheng, T., Wang, Z., Zhang, H., Zhang, X., Shi, Z., and Li, W.: Fractal  
4 Dimensions and Mixing Structures of Soot Particles during Atmospheric Processing, *Environmental*  
5 *Science & Technology Letters*, 4, 487-493, 10.1021/acs.estlett.7b00418, 2017.

# Modeling the aging process of black carbon during atmospheric transport using a new approach: a case study in Beijing

Yuxuan Zhang<sup>1,2</sup>, Meng Li<sup>2</sup>, Yafang Cheng<sup>2</sup>, Guannan Geng<sup>3</sup>, Chaopeng Hong<sup>4</sup>, Haiyan Li<sup>1</sup>, Xin Li<sup>1</sup>, Dan Tong<sup>4</sup>, Nana Wu<sup>1</sup>, Xin Zhang<sup>1</sup>, Bo Zheng<sup>5</sup>, Yixuan Zheng<sup>6</sup>, Yu Bo<sup>1,7</sup>, Hang Su<sup>2</sup>, and Qiang Zhang<sup>1</sup>

<sup>1</sup>Department of Earth System Science, Tsinghua University, Beijing 100084, China

<sup>2</sup>Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz 55128, Germany

<sup>3</sup>Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, Georgia 30322, USA

<sup>3</sup>Department of Earth System Science, University of California, Irvine, California 92697, USA

<sup>3</sup>Laboratoire des Sciences du Climate et de l'Environnement LSCE, Batiment 706, Pte 25, Orme de Merisiers 91191 Gif-sur-Yvette, France

<sup>6</sup>Department of Global Ecology, Carnegie Institution for Science, CA, 94305, USA

<sup>7</sup>RCE-TEA, Institute of Atmospheric Physics, Chinese Academy of Science, Beijing, 100029, China

*Correspondence to:* Qiang Zhang ([qiangzhang@tsinghua.edu.cn](mailto:qiangzhang@tsinghua.edu.cn)) and Yu Bo ([boyu@mail.iap.ac.cn](mailto:boyu@mail.iap.ac.cn))

**Abstract.** The effect of black carbon (BC) on air quality and the climate is still unclear, which is partly because of the poor understanding regarding the BC aging process in the atmosphere. In this work, we developed a new approach to simulate the BC mixing state (i.e., other species coated on the BC surface) based on an emissions inventory and back-trajectory analysis. The model tracks the evolution of the BC aging degree (characterized by the ratio of the whole particle size and BC core) during atmospheric transport. Using the models, we quantified the mass-averaged aging degree of total BC particles transported to a receptor (e.g., an observation site) from various emission origins (i.e.,  $0.25^\circ \times 0.25^\circ$  grids). The simulations showed good agreement with the field measurements, which validated our model calculation. Modeling the aging process of BC during atmospheric transport showed that it strongly dependent on emission levels. BC particles from extensive emission origins (i.e., polluted regions) were characterized by a higher aging degree during atmospheric transport due to more co-emitted coating precursors. On the other hand, high-emission regions also controlled the aging process of BC particles that were emitted from cleaner regions and passed through these polluted regions during atmospheric transport. The simulations identified the important roles of extensive emission regions in the BC aging process during atmospheric transport, implying the enhanced contributions of extensive emission regions to BC light absorption. This revealed that emission reductions in polluted regions could achieve more benefits for improving air pollution and climate change. Emission reductions in polluted regions not only decreased the aging degree of BC emitted from these regions but also reduced the aging process of BC emitted from other origins during atmospheric transport. Moreover, emissions reduction in polluted regions may be more efficient in counteract the suppression of planetary boundary layer (PBL) by BC particles (e.g., Zdunkowski et al. 1976; Jacobson 1998; Wendisch et al. 2008; Barbaro et al. 2013; Ding et al. 2016; Wang et al., 2018), because a greater decrease in the BC aging degree during atmospheric transport would weaken the light absorption capability of BC in the upper PBL. The simulation of the BC aging degree during atmospheric transport provided more clues for improving air pollution and climate change.

## 1 Introduction

Black carbon (BC) plays an important role in the global warming and deterioration of air quality (Bond et al. 2013). The effects of BC aerosols on air quality and climate strongly depend on their light absorption. Accurately assessing the radiative effects

1 of BC aerosols continues to be a major challenge in atmospheric/climate sciences, partly due to unclear light absorption  
2 capability of ambient BC particles. Estimates of the light absorption enhancement for BC-containing aerosols caused by  
3 coating materials on BC surface differ by a factor of  $\sim 3.5$ , spanning over a wide range from 1.05-3.5 (Cappa et al., 2012;  
4 Jacobson, 2001; Moffet et al., 2009; Peng et al., 2016). Generally, the aged BC during atmospheric transport exhibit a stronger  
5 absorption capability compared to near-source BC aerosols (Dahmkötter et al., 2014; Gustafsson and Ramanathan, 2016). In  
6 some climate model studies, the light absorption properties obtained from near-source BC aerosols are taken to estimate the  
7 direction radiative forcing (DRF) of BC (X. Wang et al., 2014; Schulz et al., 2006; Myhre et al., 2009). However, the climate  
8 effects of BC aerosols are on the regional to even global scales. Meanwhile, the effect of BC on air quality by the suppression  
9 on planetary boundary layer (PBL) is associated with atmospheric aging process of BC particles (Ding et al., 2016; Wang et  
10 al., 2018). Therefore, better understanding light absorption properties during atmospheric transport can improve the model  
11 prediction of BC effects on climate and air quality.

12 The change of light absorption of BC-containing particles during atmospheric transport is associated with the evolution of  
13 their mixing state. During atmospheric transport from emission sources, BC can internally mix with other atmospheric species  
14 (e.g., sulfate, nitrate, secondary organic matter and named coating materials) by condensation and coagulation processes  
15 (Jacobson, 2001; Li et al., 2016; Liu et al., 2017; Moffet et al., 2009). The interaction between BC and other aerosol components  
16 is defined as the BC aging process. Jacobson, (2001), Cheng et al., (2006), Lack and Cappa, (2010), Liu, et al., (2015) and  
17 Zhang et al., (2016) pointed out that aged BC can exhibit light absorption amplification by 2-3 times due to the lensing effect  
18 of the coating material on the BC surface, which then also influences the DRF and impact on the development of PBL (Cheng  
19 et al., 2008, 2009; Chung et al., 2012; Moffet et al., 2009; Ramanathan and Carmichael, 2008; Wendisch et al., 2008; Nordmann  
20 et al., 2014; Ding et al., 2016). However, the light absorption capability of BC during the aging process is still under debate  
21 (Cappa et al., 2012; Jacobson, 2001; Liu et al., 2017) partly due to a lack of understanding of the BC mixing state during  
22 atmospheric transport.

23 The mixing state of atmospheric BC-containing particles can be quantified by field observations, aircraft measurements  
24 and model simulations. Field measurements obtain the mixing state of BC particles as they are transported to an observation  
25 site (Cheng et al., 2006 and 2012; Moffet et al., 2009; Sedlacek et al., 2012; Zhang, et al., 2018a and 2018c). These observations  
26 characterize the average mixing state of BC over the observation site and cannot distinguish the mixing state of BC particles  
27 from different source origins. Moreover, field observations cannot be used to understand the temporal and spatial variations in  
28 the BC mixing state during atmospheric transport. Aircraft measurements are applied to explore the evolution of the BC mixing  
29 state during atmospheric transport from emission sources (Dahmkötter et al., 2014; Ditas et al., 2018; McMeeking, et al., 2011;  
30 Moteki et al., 2007). However, aircraft measurements are currently limited due to high costs, especially in developing countries  
31 with high BC emissions (e.g., China and India). On the other hand, some models have been developed to simulate the mixing  
32 state of BC-containing aerosols based on the mass or volume concentrations of BC and non-BC components (Jacobson, 2001;  
33 Matsui, et al., 2013; Oshima, et al., 2009). The major challenge of these models is how to treat non-BC components as coating  
34 materials of BC and BC-free particles. Moreover, when using these models, the computational cost is high when simulating  
35 the BC mixing state during atmospheric transport (Matsui, et al., 2013). A lack of information on the BC mixing state during  
36 the transport process will prevent a good understanding of the light absorption of BC particles in the atmosphere.

37 Considering the important contribution of BC from polluted regions to the BC amount present in the regional atmosphere  
38 (Lu et al., 2012; Zhang, et al., 2018c), studies on BC aging during atmospheric transport should pay more attention to BC  
39 particles from polluted regions. Based on in situ measurements, Zhang et al., (2018c) found that the light absorption capability  
40 of BC increased with increasing levels of air pollution due to more coating materials of BC under more polluted conditions.  
41 Cheng et al. (2012) showed that the aging process in polluted areas, such as Beijing, was much faster than that in clean or less  
42 polluted regions (Moteki et al., 2007; Shiraiwa et al., 2007) and in modeling studies (Cooke and Wilson, 1996; Jacobson 2001;  
43 Koch, 2001; Lohmann et al., 2000). Peng et al., (2016) also pointed out a higher aging rate of BC particles under more polluted



environments. Moreover, Li et al. (2016) and Wang et al. (2017) revealed that the significant change in BC morphology (e.g. increase of fraction dimension) associated with their mixing state (i.e., from bare-like or partly coated to embedded BC) in polluted air, which could enhance BC light absorption (Liu et al., 2015; Cappa et al., 2012). These studies identified the importance of understanding the mixing state of BC during atmospheric transport from polluted regions.

In this work, we developed a new approach to simulate the evolution of the BC mixing state during atmospheric transport based on BC emission inventory and back-trajectory analyses. First, the model calculation was used to simulate the mass-weighted mixing state of overall BC particles at a Beijing site with a fine temporal resolution of one hour, which was compared with the in situ measurements to evaluate our models. We then used our model to separate the mixing state of BC-containing particles from various spatial origins ( $0.25^\circ \times 0.25^\circ$  resolution) as they were transported to a receptor site in Beijing during a pollution period in late autumn. Based on the simulations, we focused on the mixing state of BC from polluted regions and discussed the dependence of the BC mixing state during atmospheric transport on emission levels. Finally, we explored the implication on the BC light absorption during atmospheric transport, especially for BC from polluted regions.

## 2 Data and methods

### 2.1 Data

#### 2.1.1 Measured aging degree of BC

In this study, the aging degree of BC particles was characterized by the  $D_p/D_c$  ratio (i.e., the ratio of the whole particle (including coatings and BC core) and the BC core). The observed  $D_p/D_c$  ratio of BC-containing particles is measured using a single-particle soot photometer (SP2) in this work. The SP2 technique (Droplet Measurement Technology, Boulder, CO, USA) has been described in detail elsewhere (Sedlacek et al., 2012; Moteki and Kondo, 2010; Zhang et al., 2016 and 2018b). In brief, SP2 uses incandescence and scattering signals induced by a Nd:YAG intracavity laser beam at 1064 nm to quantify the refractory BC (rBC) mass and the scattering cross section of individual BC-containing particles. The rBC mass determined from incandescence signal of SP2 was calibrated with Aquadag particles of known masses. Details on SP2 calibration was shown in previous studies (Zhang et al., 2018c). To retrieve the scattering cross section of BC-containing particles from scattering signals of SP2, a leading edge only (LEO) fit method was used (Gao et al., 2007).  $D_p$  and  $D_c$  were derived from the SP2 measurement and Mie theory, as given by Zhang et al. (2016, 2018c). In Mie calculation, the refractive indices of rBC core and coating materials was prescribed the values of 2.26-1.26i (Taylor et al., 2015; Zhang et al., 2018b) and 1.50-0i (Cappa et al., 2012; Zhang et al., 2018c); the density of rBC core was used as the value of  $1.8 \text{ g cm}^{-3}$  (Cappa et al., 2012; Taylor et al., 2015). The hourly mass-average  $D_p/D_c$  ratio of BC-containing particles is used in this work.

Table 1 lists the field observations: BJNOV2014 (performed on 17-30 November 2014), BJOC2014 (performed from 28 October to 2 November 2014), BJSEP2015 (performed on 12-19 September 2015) and BJAUG2015 (performed on 17-23 August 2015). The observation site ( $40^\circ 00' 17'' \text{ N}$ ,  $116^\circ 19' 34'' \text{ E}$ , shown in Fig. 2a) is located at Tsinghua University in the downtown area of Beijing and can be representative of the urban environment (Zhang et al., 2018a and 2018c). In this study, the Tsinghua site was taken as the receptor of BC particles from emission origins (e.g., Hebei, Tianjin, Shandong, Shanxi, Shaanxi and Inner Mongolia, Fig. 2a) during atmospheric transport. The BJNOV2014 measurement contained several pollution episodes (Fig. 1a), which featured the evolution of BC aging degree (i.e.,  $D_p/D_c$  ratio) associated with air pollution in the range of 1.4-2.3 (Fig. 1b). During the BJNOV2014 campaign period, BC amount transported to the Tsinghua site was dominated by the emission of Beijing and its surrounding areas (i.e., Hebei, Tianjin, Shanxi, Shaanxi and Inner Mongolia) (Fig. 3). Moreover, Figure 3 shows that the spatial origins of BC over the observation site during the four the pollution episodes for the BJNOV2014 measurement were different (e.g., urban, rural and industrial sources) based on back-trajectory analysis (Lu et al., 2012). Thereby, the BC aging degree (i.e.,  $D_p/D_c$  ratio) obtained from BJNOV2014 measurement was representative and

were used to establish the model in this study. The other measurement periods (i.e., the BJOC2014, BJSEP2015 and BJAUG2015 cases) were characterized by the evolution of pollution episodes (i.e., from clean hours to slight pollution and then reaching to pollution period and finally retuning to clean hours), which were used to identify that the models could simulate BC aging degree with a high time resolution (i.e., an hour) to characterize the change of BC mixing state associate with air pollution.

### 2.1.2 Back-trajectory analysis and BC emissions

During atmospheric transport, information on the location, height and transport time of BC-containing particles is obtained from back-trajectory analysis using the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model, with the meteorological fields from the National Centers for Environmental Prediction (NCEP) Global Data Assimilation System (GDAS). Back-trajectories with an hourly temporal resolution were calculated. The arrival height was set as 100 m. We ran the trajectories backwards for 3, 5 and 7 days. The effective amount of BC transported to the observation site derived from the 5 day back-trajectories was similar to that from the 7 day back-trajectories, which was significantly larger than that from the 3 day back-trajectories (Fig. 1c). This result indicated that the atmospheric lifetime of BC during the campaign period was ~5 days. Therefore, the 5 day back-trajectories were used in the following model calculation.

The gridded BC emissions for the years 2014 and 2015 were obtained from the MIX inventory, with a resolution of  $0.25^\circ \times 0.25^\circ$  (<http://www.meicmodel.org/dataset-mix>). The MIX inventory includes the emission data of anthropogenic sources in Asia (Li et al., 2017). Figure 2b shows the gridded BC emissions at  $0.25^\circ \times 0.25^\circ$  in Beijing and its surrounding regions (i.e., Hebei, Tianjin, Shandong, Shanxi, Shaanxi and Inner Mongolia).

### 2.2 Model development of BC aging during atmospheric transport

The BC aging process during atmospheric transport depends on the formation of coating materials (i.e., other species (e.g., sulfate, nitrate and organics) on the BC surface by condensation and coagulation). The more coating materials there are on the BC surface, the more aged BC during atmospheric transport. The quantity of coating materials during atmospheric transport strongly depends on the pollutant emission levels and BC transport time. In this study, the rate of change in coating mass ( $m_{coating}$ ) on BC is defined as:

$$\frac{dm_{coating}}{dt} = k_{aging}E \quad (1)$$

where  $t$  represents the transport time;  $k_{aging}$  represents the aging rate coefficient; and  $E$  represents the emissions level of coating precursors, which are co-emitted with BC. To simplify the calculation,  $E$  is quantified by BC emissions from the MIX inventory. The variable  $m_{coating}$  is calculated by Eq. (2):

$$m_{coating} = \frac{1}{6}\pi D_p^3 \rho_p - \frac{1}{6}\pi D_c^3 \rho_c \quad (2)$$

where  $\rho_p$  and  $\rho_c$  represent the densities of the total BC-containing particles (including coating materials and BC core) and the BC cores only, respectively.

Combining Eqs. (1) and (2), the BC mixing state (i.e., the  $D_p/D_c$  ratio) during atmospheric transport can be calculated as:

$$\left(\frac{D_p}{D_c}\right)^3 = kE_{aver}t + \left(\frac{D_p}{D_c}\right)_{t=0}^3 \quad (3)$$

where  $E_{aver}$  is the average BC emissions during transport and  $k$  represents the normalized aging rate coefficient, which is expressed as:

$$k = \frac{6k_{aging}}{\rho_p \pi D_c^3} \quad (4)$$

Following Eq. (3), we calculated the  $D_p/D_c$  ratio of BC aerosols transported to the receptor site from different source origins with a  $0.25^\circ \times 0.25^\circ$  resolution (i.e., the origin-resolved mixing state of BC). The conceptual scheme of the evolution of the BC

mixing state (i.e., the  $D_p/D_c$  ratio) during atmospheric transport is shown in Fig. 4. When BC aerosols emitted in a grid,  $h$ , were transported to the receptor site following a trajectory,  $l$  (Fig. 4), the  $D_p/D_c$  ratio (i.e.,  $(\frac{D_p}{D_c})_{h,l}$ ) is given, as shown in Eq. (5):

$$(\frac{D_p}{D_c})_{h,l}^3 = kE_{aver,h,l}t_{h,l} + (\frac{D_p}{D_c})_{ini}^3 \quad (5)$$

where  $E_{aver,h,l}$  and  $t_{h,l}$  represent the average BC emissions (unit of t/grid/year) and BC transport time (unit of h) from the grid  $h$  to the receptor site following the trajectory  $l$ , respectively, and  $(\frac{D_p}{D_c})_{ini}$  represents the initial value of the  $D_p/D_c$  ratio of BC before transport (Fig. 4), which characterizes BC aging from the emission source to the top of the planetary boundary layer (PBL) of the source origin (i.e., grid  $h$ ).

For a trajectory,  $l$ , BC particles pass through a series of grids (i.e.,  $h_1, h_2, h_3, \dots$ ) to the receptor (i.e.,  $hn$ ), the average  $D_p/D_c$  ratio (i.e.,  $(\frac{D_p}{D_c})_l$ ) of overall BC particles transported to the receptor site from various source origins (i.e.,  $h_1, h_2, h_3, \dots, h_i, \dots, h_n$ ) is determined by the mass-weighted  $(\frac{D_p}{D_c})_{h,l}$ , which is expressed by Eq. (6):

$$(\frac{D_p}{D_c})_l = \sum_{h=1}^{TNG_l} [(\frac{D_p}{D_c})_{h,l} \times W_{h,l}] \quad (6)$$

where  $TNG_l$  represents the total number of contributing grids to BC over the receptor site following the trajectory  $l$ ;  $W_{h,l}$  represents the weighting factor of BC from the grid  $h$ , which is determined by the effective emission intensity (EEI; defined by Lu et al. (2012)). The EEI represents the effective BC amount transported to the receptor site from the emission origins, taking into account the magnitude of BC emission from origin regions, the transport, hydrophobic-to-hydrophilic transformation, as well as dry and wet depositions during atmospheric transport.  $W_{h,l}$  is calculated by Eq. (7):

$$W_{h,l} = \frac{EEI_{h,l}}{\sum_{h=1}^{TNG_l} EEI_{h,l}} \quad (7).$$

Following the algorithm developed by Lu et al. (2012), the EEI of BC transported the receptor site from the surface grid  $h$  following a trajectory  $l$  (i.e.,  $EEI_{h,l}$  in Eq. (7)) can be determined by Eq. (8):

$$EEI_{h,l} = E_h \times TE_{h,l}, \quad (8)$$

where  $TE_{h,l}$  represents the BC transport efficiency following the trajectory  $l$ , which is calculated following Eqs. (1)-(4) shown in Lu et al. (2012). The TE is defined to quantify the transport ability of BC from origin regions to the receptor site based on transformation (i.e., hydrophobic-to-hydrophilic BC) and removal processes of BC (i.e., dry and wet depositions) in the atmospheric.

Combining Eqs. (5), (6) and (7), the  $D_p/D_c$  ratio of BC aerosols transported to the receptor site following trajectory  $l$  (i.e.,  $(\frac{D_p}{D_c})_l$ ) can be calculated by Eq. (9), which quantifies the aging degree of the overall BC transported to the receptor site, with a temporal resolution of 1 hour.

$$(\frac{D_p}{D_c})_l = \sum_{h=1}^{TNG_l} \left[ \left( kE_{aver,h,l}t_{h,l} + (\frac{D_p}{D_c})_{ini}^3 \right)^{\frac{1}{3}} \times \frac{EEI_{h,l}}{\sum_{h=1}^{TNG_l} EEI_{h,l}} \right] \quad (9)$$

Two model parameters, namely  $(\frac{D_p}{D_c})_{ini}$  and  $k$ , need to be retrieved to determine BC aging degree during atmospheric transport. The  $(\frac{D_p}{D_c})_{ini}$  represents the initial value of the  $D_p/D_c$  ratio of BC particles before atmospheric transport, which characterizes the BC aging degree near emissions. The parameter  $k$  characterizes the rate coefficient of coating materials produced on the surface of BC by atmospheric aging such as condensation, coagulation and cloud process, which is influenced by meteorological factors, chemistry, aerosol phase state as well as other parameters (e.g., particle size).

The  $(\frac{D_p}{D_c})_{ini}$  value is estimated by the BC aging near different emission sources. In this work, the  $D_p/D_c$  values of BC near the industrial, residential and traffic sources were prescribed as 1.4, 1.6 and 1.2 (Liu et al., 2014; Liu et al., 2017; Liu et al.,

2015; Laborde et al., 2013; Healy et al., 2015; Kondo et al., 2011; Morgan et al., 2019; Pan et al., 2017; Ramnarine et al., 2018; Willis et al., 2016; Schwarz et al., 2008; Shi et al., 2019; Wang et al. 2018). The  $(\frac{D_p}{D_c})_{ini}$  at the source origin (i.e., grid  $h$ ) was taken as the mass-weighted values of different source types. Figure S1 shows that the  $(\frac{D_p}{D_c})_{ini}$  values in Beijing and its surrounding areas were dominated by 1.45-1.55, which was agreed with the lowest 5th percentile of the observed  $D_p/D_c$  ratio of ambient BC-containing particles at a Beijing site (Table 1). This validated the  $(\frac{D_p}{D_c})_{ini}$  value used in the model calculation.

The aging rate coefficient  $k$  is retrieved from in situ measurements. The  $k$  value can be determined with an assumption that the simulated  $D_p/D_c$  ratio of BC-containing particles was equal to the measured one. It was noted that the simulated  $(\frac{D_p}{D_c})_t$  values, with an hourly temporal resolution, from Eq. (9) might not be equal to the observed values at a certain hour because some BC particles transported were not transported out of the observed site within one hour. To reduce the influence of the incomplete dispersion of ambient aerosols, the experimentally determined  $k$  was calculated based on the observed  $D_p/D_c$  ratio of BC-containing particles during a period  $((\frac{D_p}{D_c})_{obs,p})$ , such as the mass-average values during a pollution episode and the whole campaign. In this work, the aging rate coefficient  $k$  was calculated with the assumption that the  $(\frac{D_p}{D_c})_{obs,p}$  values was equal to the simulated  $D_p/D_c$  ratio of BC-containing particles over the receptors from various source origins during a period following various trajectories (i.e.,  $(\frac{D_p}{D_c})_p$ ).

Following Eqs. (5)-(7),  $(\frac{D_p}{D_c})_p$  is calculated using Eqs. (10)-(12):

$$\left\{ \begin{aligned} (\frac{D_p}{D_c})_p &= \sum_{h=1}^{TNG_p} [(\frac{D_p}{D_c})_{h,p} \times W_{h,p}] \end{aligned} \right. \quad (10)$$

$$\left\{ \begin{aligned} (\frac{D_p}{D_c})_{h,p} &= \left( k \left( \frac{\sum_{l=1}^{TNT_{h,p}} E_{aver,h,l} t_{h,l}}{TNT_{h,p}} \right) + (\frac{D_p}{D_c})_{ini}^3 \right)^{\frac{1}{3}} \end{aligned} \right. \quad (11)$$

$$\left\{ \begin{aligned} W_{h,p} &= \frac{EEI_{h,p}}{\sum_{h=1}^{TNG_p} EEI_{h,p}} \end{aligned} \right. \quad (12)$$

where  $(\frac{D_p}{D_c})_{h,p}$  represents the  $D_p/D_c$  ratio of BC-containing aerosols transported to our observation site from a grid,  $h$ , during the period;  $W_{h,p}$  represents the weighting factor of BC from the grid  $h$  to the receptor site during the period;  $TNG_p$  represents the total number of contributing grids to BC over the receptor site during the period;  $TNT_{h,p}$  represents the total number of trajectories passing through the grid  $h$  during the period; and  $EEI_{h,p}$  represents the  $EEI$  in the surface grid  $h$  during our a period, which is calculated by Eq. (13) (Lu et al., 2012):

$$EEI_{h,p} = E_h \times [(\sum_{l=1}^{TNT_{h,p}} TE_{h,l}) / TNT_p] \quad (13)$$

where  $TNT_{h,p}$  represents the total number of trajectories passing through the grid  $h$  and  $TNT_p$  represents the total number of back-trajectories originating at our site during the whole period. Fig. 3 shows the TE density (TED) (defined as  $(\sum_{l=1}^{TNT_{h,p}} TE_{h,l}) / TNT_p$  in Eq. (13), Lu et al., 2012) and  $EEI$  on a  $0.25^\circ \times 0.25^\circ$  grid during the episodes 1-4 (Fig. 1a). The gridded  $EEI$  analysis revealed that BC particles over the Tsinghua site during the four episodes are from different source types, namely industrial and rural sources, urban and rural sources, urban and industrial sources as well as industrial source during episodes 1-4, respectively. This indicated that the model parameters obtained from the BJNOV2014 measurement are representative.

With the assumption that  $(\frac{D_p}{D_c})_p = (\frac{D_p}{D_c})_{obs,p}$ , the experimentally determined  $k$  was calculated by Eq. (14):

$$k = \left[ \left( \frac{D_p}{D_c} \right)_{obs,p}^3 - \left( \frac{D_p}{D_c} \right)_{ini}^3 \right] \times \frac{\sum_{h=1}^{TNG_p} EEI_{h,p}}{\sum_{h=1}^{TNG_p} \left[ \left( \frac{\sum_{l=1}^{TNT_{h,p}} E_{aver,h,l} t_{h,l}}{TNT_{h,p}} \right) \times EEI_{h,p} \right]} \quad (14).$$

In this work, the  $k$  value was calculated based on the measured  $D_p/D_c$  ratios of BC-containing particles during BJNOV2014 campaign period at the Tsinghua site (40°00'17" N, 116°19'34" E) in Beijing. In order to obtain the  $k$  value and then valid our models, the BJNOV2014 measurements were divided into two parts: the first two episodes (i.e., 17-24 November 2014) used to calculate the  $k$  value and the last two episodes (i.e., 24-30 November 2014) used for model validation. Following the Eq. (14), the normalized aging rate coefficient  $k$  was determined to have a value of  $1.8 \times 10^{-4} \text{ t}^{-1} \text{ h}^{-1}$ . Correspondingly, the value of  $k_{\text{aging}}$  is  $\sim 17\% \text{ h}^{-1}$ , which was comparable with the observation ones (Cheng et al., 2012; Zhang et al., 2018).

To evaluate our model calculation of the mixing state of BC-containing particles over the site, the simulated  $(\frac{D_p}{D_c})_l$  values with an hourly temporal resolution are compared with the observed hourly  $D_p/D_c$  ratio during 24-30 November 2014 (i.e., the last two episodes during the BJNOV2014 measurement). It is noted that the observed  $D_p/D_c$  ratio at a certain hour is not only dominated by the aging degree of BC transported to the site at this time but also impacted by the aging degree of BC over the site several hours beforehand due to the incomplete dispersion of ambient aerosols within one hour. In this work, we averaged the simulated  $(\frac{D_p}{D_c})_l$  at a certain hour and within  $n$  ( $n=0, 2, 4, 6, \dots$ ) hours beforehand ( $(\frac{D_p}{D_c})_{\text{aver},t}$ ) to compare with the observed hourly  $D_p/D_c$  ratio.  $(\frac{D_p}{D_c})_{\text{aver},t}$  is calculated by Eq. (15):

$$(\frac{D_p}{D_c})_{\text{aver},t} = \frac{\sum_{l=t-n}^t (\frac{D_p}{D_c})_l}{n+1} \quad (15).$$

Figure 5 shows an excellent agreement between the simulated and observed hourly  $D_p/D_c$  ratio with a difference of  $\sim 3\%$ . The simulated  $(\frac{D_p}{D_c})_{\text{aver},t}$  values, with  $n=2, 4$ , and  $6$ , exhibit a better correlation with the observed hourly  $D_p/D_c$  ratio than those without considering the effect of BC transported over our observation site within a few hours beforehand (i.e.,  $n=0$ ). When  $n=4$ , the linear relationship between simulated and observed  $D_p/D_c$  ratio exhibited a highest correlation coefficient (i.e.,  $R^2=0.86$ ), indicating that the residence time of BC-containing particles over the Tsinghua site during the campaign period (i.e., polluted periods) is  $\sim 5$  hours.

### 3 Results and discussion

#### 3.1 Simulating BC mixing state with a high time resolution

Following Eqs. (9) and (15), we calculated the hourly  $D_p/D_c$  ratios of BC-containing particles during the BJNOV2014, BJSEP2015 and BJAUG2015 measurement periods. Fig. 6 shows that the simulated  $D_p/D_c$  values of BC-containing particles exhibit significant changes with pollution levels (i.e., the  $\text{PM}_{2.5}$  and rBC concentrations), revealing that our model with a high time resolution can simulate the evolution of the BC aging degree during a pollution episode. The simulation results showed that under pollution conditions, not only the BC mass concentrations increased, but also the aging degree of BC-containing particles enhanced. This indicated that the amplification of BC light absorption associated with air pollution due to more coating material on BC surface (Zhang et al., 2018c).

It is important for simulation of BC mixing state with a high time resolution to evaluate effect on air quality. The effect of BC on air quality depends on both the mass concentration and aging degree of BC. Simultaneous increase in the mass concentration and aging degree of BC associated with air pollution could suppress PBL by the dome effect (Ding et al., 2016). In China, the air pollution often start rapidly within several hours (Zheng et al., 2016). It is necessary to understand hourly aging degree of BC-containing particles for better exploring the effect of BC on air quality. Our model calculation can provide BC aging degree (i.e.,  $D_p/D_c$  ratio) with a fine temporal resolution of an hour.

The simulated BC mixing state with high time resolution during different measurement periods was used to revealed that our models could be generally applied. Although the model is established based on the BJNOV2014 measurement (17-30

November 2014, Beijing), the retrieved model parameters are applicable to other measurement periods (i.e., October 2014, September 2015, and August 2015). Fig. 6 reveals an excellent agreement between the simulated and observed hourly  $D_p/D_c$  ratio for the BJOCT2014, BJSEP2015 and BJAUG2015 measurement. During these measurement periods, the linear relationship between the simulated  $D_p/D_c$  ratio and the observed values shows slopes of 0.98-1.01, with a correlation coefficient ( $R^2$ ) of 0.53-0.85. A good correlation further demonstrated the validity of our model for calculating the BC mixing state during atmospheric transport. Accurate simulations for the different measurement periods identify the generalness of our model, especially under polluted environments.

### 3.2 BC mixing state during the atmospheric transport process

#### 3.2.1 Aging degree of BC from emission origins to receptor sites

Using our models, we investigated the aging process of BC-containing particles transported from different spatial origins to the receptor (i.e., the Tsinghua site, Beijing) during the BJNOV2014 measurement period. The origin-resolved ( $0.25^\circ \times 0.25^\circ$ )  $D_p/D_c$  ratio of BC-containing particles over the receptor site calculated using Eq. (11), shown in Fig. 7. The BC-containing particles from various spatial origins exhibited significant difference in their mixing state with  $D_p/D_c$  ratio in the range of 1.4-2.8 as they reached the Tsinghua site. The  $D_p/D_c$  ratio of BC from the polluted regions (i.e., southern Heibei, northeastern Hebei and Tianjin) could up to ~2.3-2.8, which characterized the mixing state of fully aged BC in the North China Plain. These fully aged BC particles played an important role in regional light absorption (Gustafsson and Ramanathan, 2016; Peng et al., 2016). The high  $D_p/D_c$  ratio (~2.3-2.8) of BC transported to our site was consisted with the mixing state of fully aged BC particles ( $D_p/D_c$  ratio ~2.5) in Beijing reported by Peng et al., (2016), implying reliable values of gridded  $D_p/D_c$  ratio from our model calculation.

In this work, we classified the spatial emission sources of BC-containing particles over the receptor site into local (or Beijing (i.e., BJ origin), shown in Fig. 8a) and non-local (i.e., non-Beijing) origins based on political boundaries. The non-local origins were further divided into Hebei, Tianjin and other origins (i.e., HB, TJ or OT origins, respectively, as shown in Fig. 8a). Based on the gridded  $D_p/D_c$  ratio shown in Fig.7, the  $D_p/D_c$  ratios of BC-containing particles from Beijing, Hebei, Tianjin and other regions during the campaign period were estimated to be 1.8, 2.2, 2.1 and 2.0, respectively (Fig. 8b). Fig. 2b reveals that more intensive emission regions of BC are located in southern Beijing, southern Hebei, northeastern Hebei and Tianjin, which dominated BC amount over the receptor site during the campaign period (Fig. 3). The EEI analysis (Fig. 3) shows that BC-containing particles transported to the site during the campaign period mainly originate from Beijing, Hebei, Tianjin, Inner Mongolia, Shanxi and Shaanxi (the political boundaries of these regions are shown in Fig. 2a).

The contributions of different emission origins to the BC amount over the receptor site during the BJNOV2014 measurement period were estimated by EEI analysis. BC transported to the site was dominated by Hebei and Beijing as the major source regions, accounting for ~40.2% and ~40.0% of the total amount transported, respectively. In terms of non-local (i.e., non-Beijing) origins, the contribution of BC from Hebei was significantly higher than that from Tianjin (~4.2%) and others (~15.5%) due to more intensive emissions and a larger region in Hebei Province. Approximately 60% of BC from non-local origins (i.e., Hebei, Tianjin and others) indicated the importance of atmospheric transport to BC concentrations during polluted environments. During the campaign period, the average BC mass concentration was ~4.0  $\mu\text{g m}^{-3}$  (Fig. 1b). Based on EEI analysis, the mass concentrations of local (i.e., Beijing) and non-local (i.e., non-Beijing) BC at the observed site were estimated to be ~1.6  $\mu\text{g m}^{-3}$  and ~2.4  $\mu\text{g m}^{-3}$ , respectively.

Atmospheric transport not only played an important role in BC mass concentration in Beijing under polluted environments but also controlled BC aging. During the investigation period, the  $D_p/D_c$  ratio of non-local (i.e., non-Beijing) BC over the site was ~2.1, which was greater than that of local (i.e., Beijing) BC (~1.8). The higher aging degree of non-local BC could be attributed to the longer transport time and larger emissions from non-local origins. As more intensive emission sources (i.e.,



polluted regions), Hebei and Tianjin were identified as the two largest contributing regions for the BC mixing state at the site (Fig.7 and Fig. 8b). When BC particles were emitted from Hebei and Tianjin and then transported to the receptor site, their  $D_p/D_c$  ratios could reach up to 2.2 and 2.1, respectively, which were larger than that from other non-local origins (i.e., cleaner regions). This could be due to more pollutant emission from southern Hebei, northeastern Hebei and Tianjin than that from other regions (i.e., Inner Mongolia). The results revealed that BC particles emitted from polluted regions would exhibit a higher aging degree during atmospheric transport, which is most likely attributed to more rapid aging due to more co-emitted coating precursors (Peng et al., 2016).

On the other hand, high emission origins (e.g., southern Hebei, northeastern Hebei and Tianjin) also affect the aging process of BC particles that pass through these regions. When BC particles are emitted from a clean origin and then pass through high emission regions (e.g., southern Hebei) to the receptor site (cluster 3 shown in Fig. 8c), their  $D_p/D_c$  ratio could reach ~2.2 (Fig. 8d). However, the emitted BC particles from the clean origin passing through a series of clean origins (cluster 1 shown in Fig. 8c) show a lower  $D_p/D_c$  ratio (~1.6, Fig. 8d) as they reach the site. This difference identifies the important role of extensive emission regions (e.g., southern Hebei) in the atmospheric aging process of BC particles emitted from other clean regions (e.g., Inner Mongolia). When BC particles passed through these polluted regions, their aging degree could be accelerated due to high pollutant emission.

### 3.2.2 Emission dependence

As discussed above, the aging process of BC particles during atmospheric transport was closely associated with emissions from regions through which they pass. To investigate the dependence of the BC mixing state during atmospheric transport, we normalized the current emissions obtained from the MIX inventory as a unit (i.e., normalized emission of 1 shown in Fig. 9) and set the scenarios of BC emissions reduced by 20%, 40%, 50%, 60%, and 80% (corresponding to normalized emissions of 0.8, 0.6, 0.5, 0.4, and 0.2, respectively, in Fig. 9).

Fig. 9a1 shows that the  $D_p/D_c$  ratios of local (i.e., Beijing) BC, non-local (i.e., non-Beijing) BC and total BC (including both local and non-local BC) transported to the site were proportional to the normalized emissions (i.e., the sum of both local and non-local emissions), with slopes of 0.55, 0.26 and 0.44, respectively. The slope values revealed that the relative increase or decrease in the  $D_p/D_c$  ratio with emission change for non-local BC was ~2 times that of local BC. This result revealed that the mixing state of non-local BC was more sensitive to emission change than that of local BC. Therefore, emission reduction was more effective in lowering the aging degree of non-local BC compared with that of local BC. For example, when BC emissions were reduced by 50% (i.e., from normalized emissions of 1 to 0.5 in Fig. 9a1), the  $D_p/D_c$  ratios of non-local BC, local BC and total BC at our site decreased by 14%, 7% and 11%, respectively. A greater reduction in the aging degree of non-BC particles could be attributed to most of the non-local BC (~75%) from high emission origins (i.e., the southern Hebei, northeastern Hebei and Tianjin regions). The origin-resolved  $D_p/D_c$  ratio of BC transported to the site during the campaign period shown in Fig. 7 indicated that a greater reduction in aging degree would be found for BC from higher emission regions, such as southern Hebei, northeastern Hebei and Tianjin, which revealed the benefits of emission controls in extensive emission regions. The results identified the dependence of the BC aging process during atmospheric transport on emissions, especially for non-local BC from high emission origins.

To further evaluate the change in the mixing state of BC at the site from local (i.e., Beijing) and non-local (i.e., non-Beijing) emissions, we simulated the origin-resolved  $D_p/D_c$  ratio of BC over the receptor site only considering the changes in local (Fig. 9a2) or non-local (Fig. 9a3) emissions. It is noted that the mixing state of total BC at the receptor site not only depended on the respective  $D_p/D_c$  ratio of local and non-local BC but also their contributions to the total BC amount. When only altering local or non-local emissions in the simulations, the contributions of various source origins to the BC amount at our site would be changed.

When the changes in only local emissions were included in our simulations, the mixing state of total BC (including both

local (i.e., Beijing) and non-local BC (i.e., non-Beijing)) at the observed site was slightly sensitive to emissions. Fig. 9a2 presents the linear change in the  $D_p/D_c$  ratio of total BC with changing local emissions (i.e., normalized emissions of 0.2-1), with a slope of 0.07, which is significantly smaller than that (0.44) for the cases of changing both local and non-local emissions (Fig. 8a1). Taking 50% of the reduction in local emissions as an example, the  $D_p/D_c$  ratio of total BC at our site only decreased by 2%. Meanwhile, 50% of the local emission reduction resulted in ~5.0% and ~4.6% of the decreases in the  $D_p/D_c$  ratio for local and non-local BC, respectively. The reductions in the  $D_p/D_c$  ratio of local and non-local BC were larger than that in total BC (~2%), which was due to the increase in the contributions of non-local BC (characterized by a larger  $D_p/D_c$  ratio compared to local BC) with a local emission reduction. The results showed that altering local emissions had a slight effect on the BC mixing state at our site during the investigated period (i.e., under a polluted environment) due to slight change in the aging degree of non-local BC.

In terms of non-local (i.e., non-Beijing) emission change, the response of the mixing state of total BC at our site to emission changes was more significant than that for the case of local (i.e., Beijing) emission change. The linear correlation between the  $D_p/D_c$  ratio of total BC and normalized non-local emission (i.e., 0.2-1) shown in Fig. 9a3 exhibited a slope of 0.28, which was markedly greater than that (0.07) in the case of local emission change (Fig. 9a2). Fig. 9a3 shows that the slope for total BC (i.e., 0.28) was similar to that (i.e., 0.31) for non-local BC, indicating that the variation in the BC mixing state at our site during the investigated period (i.e., under a polluted environment) was controlled by the emission change of non-local BC. This result could be attributed to more aged BC particles were mainly from non-local regions (the southern Hebei, northeastern Hebei and Tianjin). These results indicated that the BC mixing state at our site was dominated by non-local emissions (in particular polluted regions with intensive emission), identifying the importance of atmospheric transport in the BC mixing state in Beijing during polluted periods.

### 3.3 BC light absorption in the atmosphere

BC light absorption depends on both the mass concentration and mixing state of BC. The light absorption of BC can be characterized by multiplying EEI by the  $D_p/D_c$  ratio (i.e.,  $EEI \cdot D_p/D_c$ ). In this study, the origin-resolved  $EEI \cdot D_p/D_c$  values represent the light absorption levels of BC particles as they were transported to the receptor site from various source origins ( $0.25^\circ \times 0.25^\circ$ ). Fig. 10a displays the origin-resolved  $EEI \cdot D_p/D_c$  values during the campaign period. High light-absorption levels of BC were mainly from the local Beijing area, southern Hebei, and northeastern Hebei and Tianjin, resulting from high BC emissions and strong BC aging in these regions.

The origin-resolved  $EEI \cdot (D_p/D_c)$  values revealed the contributions of BC from different source regions to light absorption at the site (Fig. 10b). During the investigated period (i.e., polluted period) in Beijing, Hebei Province was the largest contributing region, accounting for ~44% of BC light absorption transported to the observed site during the campaign period. Local Beijing was responsible for ~36% of the light absorption of BC at the site, which was lower than the contribution from Hebei. In total, ~64% of BC light absorption at the receptor site was contributed by non-local (i.e., non-Beijing) BC source origins, reflecting the importance of atmospheric transport on the light absorption of BC in Beijing during polluted periods. The contribution of non-local origins to BC light absorption at our site was larger than that to BC mass concentration (~60%, as quantified by the EEI analysis), which was due to the higher aging degree ( $D_p/D_c$  ratio of ~2.12) of non-local BC compared to that of local BC ( $D_p/D_c$  ratio of ~1.78). If the difference between the mixing states of local and non-local BC is not considered, the effect of atmospheric transport on BC light absorption in Beijing during the polluted period would be underestimated. The results revealed that the BC aging process during atmospheric transport strengthens the importance of emissions from surrounding areas (e.g., Hebei and Tianjin) to BC light absorption in Beijing during polluted periods.

The strong dependence of both the mass concentrations and mixing states of BC on emissions indicates that emission reduction could significantly lower light absorption and, thus, weaken the effect of BC on air quality and climate, especially during polluted periods. A linear decrease in BC light absorption with emission reductions was found (Fig. 9b). When both

local (i.e., Beijing) and non-local (i.e., non-Beijing) BC emissions were reduced (Fig. 9b1), the change in light absorption for non-local BC was much more significant than that for local BC. For the case of a 50% reduction in both local and non-local BC emissions, the light absorption of total BC, local BC and non-local BC decreased by 55%, 53% and 56%, respectively (Fig. 9b1). An extra decrease of ~6% in light absorption for non-local BC was greater than that (~3%) for local BC, which could be attributed to a much greater reduction in the aging degree (i.e.,  $D_p/D_c$  ratio) of non-local BC (Fig. 9a1). Compared with local BC, a much greater decrease in the light absorption of non-local BC implied that the emission reduction from non-Beijing sources rather than Beijing sources achieved many more benefits in terms of the BC effect in Beijing. This result also revealed that the emission reduction of surrounding areas of Beijing not only brought less BC but also lowered the BC mixing state, which enhanced the decrease in BC light absorption in Beijing.

The analyses of only reducing local emissions or non-local emissions further identified the importance of emission reduction of non-Beijing sources to weaken BC light absorption in Beijing during polluted periods. For the case of local (i.e., Beijing) emissions reduction (Fig. 9b2), the reduction in BC light absorption was significantly smaller than that for the case of non-local (i.e., non-Beijing) emissions reduction (Fig. 9b3), revealing that emissions reduction in non-Beijing sources played a more important role in decreasing BC light absorption in Beijing. For example, when emissions were reduced by 50% for local and non-local origins, the BC light absorption at the site decreased by 21% and 35%, respectively. A greater decrease in BC light absorption in Beijing with non-local emissions reduction resulted from larger contributions of non-Beijing source emissions (especially for polluted regions, such as southern Hebei, northeastern Hebei and Tianjin) to both the mass concentration and mixing state of BC over the site. Moreover, the extra reduction in BC light absorption in Beijing caused by weakening of the BC aging degree was also greater under non-Beijing emissions reduction compared with that under Beijing emissions reduction (e.g., ~5% and 1% extra reductions for the cases of 50% reductions in non-local and local emissions, respectively). The import contribution of emission reduction of non-Beijing sources to decrease of BC light absorption in Beijing under polluted environment was mainly due to significant decrease in both BC amount and its aging degree in Beijing caused by emission reduction of polluted regions (e.g., southern Hebei, northeastern Hebei and Tianjin). This suggested that the efforts to weaken the influence of BC on air pollution and climate change should pay more attention in emissions reduction in polluted regions.

## 4 Discussion

In this study, a rather simplified scheme was adopted where the aging rate is assumed proportional to the emissions without detailed consideration of the effects of temperature, particle sizes, phase state, hygroscopicity and chemistry (Riemer et al. 2009; Cheng et al. 2008, 2012, 2015; Mu et al. 2018). **Actually, these factors can influence the model parameter of aging rate coefficient. Our simulations strongly depends on the emission, initial value of  $D_p/D_c$  ratio and aging rate coefficient. Accurate  $(\frac{D_p}{D_c})_{ini}$  values requires more measurements of the mixing state of freshly emitted BC-containing particles.** In this work, the experimentally determined  $k$  value was derived from the observations during the pollution periods **at autumn/winter**. In terms of the pollution periods with low  $O_3$  concentrations and high RH, the BC mass-weighted  $O_3$  and RH levels at the observed site were 1-3 ppb and 50-63%, respectively.

**Although our model was developed and validated with autumn/winter datasets, it can be applied in other seasons with distinct model parameters (i.e. aging rate coefficient and the initial value of  $D_p/D_c$  ratio). The two parameters exhibit a seasonal dependence. In terms of the aging rate coefficient at summer (Cheng et al., 2012; Zhang et al., 2018), the measurements at two suburban site (xianghe and Yufa) in NCP showed pronounced diurnal cycle in the range of 0.2-20%  $h^{-1}$  with a maximum at noon time due to stronger photochemistry. The  $k_{aging}$  around noontime at summer was comparable with that (~17%  $h^{-1}$ ) at autumn/winter. However, other time especially nighttime at summer is significantly smaller. For the  $(\frac{D_p}{D_c})_{ini}$ , the seasonal**

dependence is due to different dominated sources of BC. The BC particles in Beijing at summer are controlled by traffic and industrial emission with smaller higher  $D_p/D_c$  ratio near sources, but the dominated sources of BC at autumn/winter are industrial and residential emissions with higher  $(\frac{D_p}{D_c})_{ini}$  (D. Liu et al., 2019; H. Liu et al., 2019). In summary, lower values both of aging rate coefficient and the initial value of  $D_p/D_c$  ratio would lead to thinner coatings of BC at summer compared with those at autumn/winter in NCP, which was consistence with observations in Beijing (D. Liu et al., 2019).

The retrieved aging rate coefficient  $k_{aging}$  ( $\sim 17\% \text{ h}^{-1}$ , corresponding to the timescale of BC aging process about  $\sim 6 \text{ h}$ ) indicated BC particles under polluted environment at autumn/winter in NCP underwent fast aging during atmospheric transport. The retrieved  $k_{aging}$  with a value of  $\sim 17\% \text{ h}^{-1}$  used in our models is comparable with the observed values at other suburban sites in NCP, namely up to  $\sim 20\% \text{ h}^{-1}$  and  $\sim 21\% \text{ h}^{-1}$  at Yufa site and Xianghe site, respectively (Cheng et al., 2012; Zhang et al., 2018). However, the aging rate coefficient commonly used in other models ( $1\text{-}5\% \text{ h}^{-1}$ ) was significantly smaller than that used in our model (Cooke and Wilson, 1996; Jacobson, 2001; Koch, 2001; Lohmann et al., 2000;), indicating that the values used in previous models derived in developed countries (Moteki et al., 2007; Shiraiwa et al., 2007) could not represent the characteristic of BC aging process in China. A similar conclusion also found by Peng et al., (2007) using an environment chamber, namely the timescale of BC aging with significant distinction between urban cities in Beijing ( $\sim 5 \text{ h}$  corresponding to  $k_{aging}$  of  $\sim 20\% \text{ h}^{-1}$ ) and Houston ( $\sim 18 \text{ h}$  corresponding to  $k_{aging}$  of  $\sim 5.6\% \text{ h}^{-1}$ ). Higher aging rate coefficient suggested the BC aging under polluted environment in NCP is most likely dominated by condensation process during atmospheric transport, taking into account a clear difference between timescale of BC aging by condensation (2-8 h) and coagulation (10-40 h) (Riemer et al., 2004).

More aged BC aloft increases atmospheric stability, which suppressed the development boundary layer and suppress or enhance the formation of different types of clouds (Zdunkowski et al. 1976; Jacobson 1998; Wendisch et al. 2008; Barbaro et al. 2013; Bond et al. 2013; Ding et al., 2016; Wang et al., 2018). Our simulations showed that the aging degree of BC from atmospheric transport along with polluted air masses was significantly higher than that of local BC under polluted conditions, implying that the light absorption capability (i.e., mass cross section of BC-containing particles including coating materials on BC surface) of BC in the upper PBL would be higher than that of BC in the lower PBL. This characteristic favored the formation of the inversion layer due to more heating in the upper PBL under polluted conditions (Ding et al., 2016; Wang et al., 2018), which consequently depressed PBL development. In China, air pollution generally occurs at the regional scale due to atmospheric transport (Sun et al., 2014; L. Wang et al., 2014; Yang et al., 2015; Zheng et al., 2015). Regional pollution would bring more aged BC due to faster aging processes during atmospheric transport. The enhanced effect caused by more aged BC from atmospheric transport would further suppress the development of boundary layer (e.g., Zdunkowski et al. 1976; Jacobson 1998; Wendisch et al. 2008; Barbaro et al. 2013; Ding et al. 2016) and together with the special haze chemistry (Cheng et al. 2016) strongly strengthens regional pollution.

## 5 Concluding remarks

The effect of BC-containing particles on air quality and climate is not only dominated by BC mass concentration but also controlled by their mixing state. To better understand the mixing state of atmospheric BC in China, we developed a new approach to simulate the BC aging process during atmospheric transport based on the BC emission inventory and back-trajectory analysis. Our models track the BC mixing state (i.e.,  $D_p/D_c$  ratio) from an emitted source origin (e.g., a  $0.25^\circ \times 0.25^\circ$  grid) to a receptor (i.e., Tsinghua site). The model calculation can quantify the mass-averaged  $D_p/D_c$  ratio of overall BC particles over the receptor site from various origins, which agreed well with observed ones. The simulations can provide information on the BC mixing state with fine temporal and spatial resolutions.

Based on the simulations of BC mixing state during atmospheric transport, we find a strong dependence of BC mixing state on emissions during atmospheric transport. BC particles with higher aging degrees at our site were mainly from more

intensive emission origins (e.g., southern Hebei) due to higher aging rates. On the other hand, when BC particles emitted from clean origins passed through polluted regions, they were also characterized by a greater  $D_p/D_c$  ratio by speeding up the aging process in polluted air. Our simulations demonstrated the importance of regional transport in BC light absorption in Beijing under polluted conditions. This provides a new perspective on the phenomenon of pollution building up in Beijing, further demonstrating that this is driven by regional transport and transformation rather than local sources and processes.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (41571130032, 41571130035, 41625020, and 91744310).

## References

- Barbaro, E., Vilà-Guerau de Arellano, J., Krol, M. C., and Holtslag, A. A. M.: Impacts of Aerosol Shortwave Radiation Absorption on the Dynamics of an Idealized Convective Atmospheric Boundary Layer, *Boundary-Layer Meteorology*, 148, 31-49, 10.1007/s10546-013-9800-7, 2013.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., Jobson, B. T., Kolesar, K. R., Lack, D. A., Lerner, B. M., Li, S.-M., Mellon, D., Nuaaman, I., Olfert, J. S., Petäjä, T., Quinn, P. K., Song, C., Subramanian, R., Williams, E. J., and Zaveri, R. A.: Radiative Absorption Enhancements Due to the Mixing State of Atmospheric Black Carbon, *Science*, 337, 1078-1081, 2012.
- Cheng, Y. F., Eichler, H., Wiedensohler, A., Heintzenberg, J., Zhang, Y. H., Hu, M., Herrmann, H., Zeng, L. M., Liu, S., Gnauk, T., Brüggemann, E., and He, L. Y.: Mixing state of elemental carbon and non-light-absorbing aerosol components derived from in situ particle optical properties at Xinken in Pearl River Delta of China, *J. Geophys. Res.-Atmos.*, 111, 2006.
- Cheng, Y. F., Wiedensohler, A., Eichler, H., Su, H., Gnauk, T., Brüggemann, E., Herrmann, H., Heintzenberg, J., Slanina, J., Tuch, T., Hu, M., and Zhang, Y. H.: Aerosol optical properties and related chemical apportionment at Xinken in Pearl River Delta of China, *Atmos. Environ.*, 42, 6351-6372, <http://dx.doi.org/10.1016/j.atmosenv.2008.02.034>, 2008.
- Cheng, Y. F., Berghof, M., Garland, R. M., Wiedensohler, A., Wehner, B., Müller, T., Su, H., Zhang, Y. H., Achtert, P., Nowak, A., Pöschl, U., Zhu, T., Hu, M., and Zeng, L. M.: Influence of soot mixing state on aerosol light absorption and single scattering albedo during air mass aging at a polluted regional site in northeastern China, *J. Geophys. Res.-Atmos.*, 114, 10.1029/2008JD010883, 2009.
- Cheng, Y. F., Su, H., Rose, D., Gunthe, S. S., Berghof, M., Wehner, B., Achtert, P., Nowak, A., Takegawa, N., Kondo, Y., Shiraiwa, M., Gong, Y. G., Shao, M., Hu, M., Zhu, T., Zhang, Y. H., Carmichael, G. R., Wiedensohler, A., Andreae, M. O., and Pöschl, U.: Size-resolved measurement of the mixing state of soot in the megacity Beijing, China: diurnal cycle, aging and parameterization, *Atmos. Chem. Phys.*, 12, 4477-4491, 2012.
- Cheng, Y., Su, H., Koop, T., Mikhailov, E., and Pöschl, U.: Size dependence of phase transitions in aerosol nanoparticles, *Nat Commun.*, 6, 5923, 2015.
- Cheng, Y., Zheng, G., Wei, C., Mu, Q., Zheng, B., Wang, Z., Gao, M., Zhang, Q., He, K., and Carmichael, G.: Reactive nitrogen chemistry in aerosol water as a source of sulfate during haze events in China, *Science advances*, 2, e1601530, 2016.

1 Chung, C. E., Ramanathan, V., and Decremer, D.: Observationally constrained estimates of carbonaceous aerosol radiative  
2 forcing, *Proc. Natl. Acad. Sci. USA*, 109, 11624-11629, 10.1073/pnas.1203707109, 2012.

3 Cooke, W. F. and Wilson, J. J. N.: A global black carbon aerosol model, *J. Geophys. Res.*, 101, 19395–19409, 1996.

4 Dahlkötter, F., Gysel, M., Sauer, D., Minikin, A., Baumann, R., Seifert, P., Ansmann, A., Fromm, M., Voigt, C., and Weinzierl,  
5 B.: The Pagami Creek smoke plume after long-range transport to the upper troposphere over Europe &ndash; aerosol properties  
6 and black carbon mixing state, *Atmos. Chem. Phys.*, 14, 6111-6137, 10.5194/acp-14-6111-2014, 2014.

7 Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V. M., Petäjä, T., Su, H., Cheng, Y. F., Yang, X. Q., Wang, M. H., Chi,  
8 X. G., Wang, J. P., Virkkula, A., Guo, W. D., Yuan, J., Wang, S. Y., Zhang, R. J., Wu, Y. F., Song, Y., Zhu, T., Zilitinkevich, S.,  
9 Kulmala, M., and Fu, C. B.: Enhanced haze pollution by black carbon in megacities in China, *Geophys. Res. Lett.*, 43, 2873-  
10 2879, doi:10.1002/2016GL067745, 2016.

11 Ditas, J., Ma, N., Zhang, Y., Assmann, D., Neumaier, M., Riede, H., Karu, E., Williams, J., Scharffe, D., Wang, Q., Saturno,  
12 J., Schwarz, J. P., Katich, J. M., McMeeking, G. R., Zahn, A., Hermann, M., Brenninkmeijer, C. A. M., Andreae, M. O., Pöschl,  
13 U., Su, H., and Cheng, Y.: Strong impact of wildfires on the abundance and aging of black carbon in the lowermost stratosphere,  
14 *Proc. Natl. Acad. Sci. USA*, 115, E11595-E11603, 10.1073/pnas.1806868115, 2018.

15 Gao, R. S., Schwarz, J. P., Kelly, K. K., Fahey, D. W., Watts, L. A., Thompson, T. L., Spackman, J. R., Slowik, J. G., Cross, E.  
16 S., Han, J. H., Davidovits, P., Onasch, T. B., and Worsnop, D. R.: A Novel Method for Estimating Light-Scattering Properties  
17 of Soot Aerosols Using a Modified Single-Particle Soot Photometer, *Aerosol Sci. Technol.*, 41, 125-135,  
18 10.1080/02786820601118398, 2007.

19 Gustafsson, Ö., and Ramanathan, V.: Convergence on climate warming by black carbon aerosols, *Proc. Natl. Acad. Sci. USA*,  
20 113, 4243-4245, 2016.

21 Healy, R. M., Wang, J. M., Jeong, C.-H., Lee, A. K. Y., Willis, M. D., Jaroudi, E., Zimmerman, N., Hilker, N., Murphy, M.,  
22 Eckhardt, S., Stohl, A., Abbatt, J. P. D., Wenger, J. C., and Evans, G. J.: Light-absorbing properties of ambient black carbon  
23 and brown carbon from fossil fuel and biomass burning sources, *J. Geophys. Res.-Atmos.* 120, 6619-6633,  
24 10.1002/2015jd023382, 2015.

25 Jacobson, M. Z.: Studying the effects of aerosols on vertical photolysis rate coefficient and temperature profiles over an urban  
26 airshed, *J. Geophys. Res.-Atmos.*, 103, 10593-10604, 1998.

27 Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols, *Nature*, 409, 695-  
28 697, 2001.

29 Koch, K.: Transport and direct radiative forcing of carbonaceous and sulphate aerosols in the GISS GCM, *J. Geophys. Res.*,  
30 106, 20311–20332, 2001.

31 Kondo, Y., Matsui, H., Moteki, N., Sahu, L., Takegawa, N., Kajino, M., Zhao, Y., Cubison, M. J., Jimenez, J. L., Vay, S.,  
32 Diskin, G. S., Anderson, B., Wisthaler, A., Mikoviny, T., Fuelberg, H. E., Blake, D. R., Huey, G., Weinheimer, A. J.,  
33 Knapp, D. J., and Brune, W. H.: Emissions of black carbon, organic, and inorganic aerosols from biomass burning in North  
34 America and Asia in 2008, *J. Geophys. Res.*, 116, D08204, <https://doi.org/10.1029/2010jd015152>, 2011.

35 Lack, D. A., and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and  
36 absorption wavelength dependence of black carbon, *Atmos. Chem. Phys.*, 10, 4207-4220, 2010.

37 Laborde, M., Crippa, M., Tritscher, T., Jurányi, Z., Decarlo, P. F., Temime-Roussel, B., Marchand, N., Eckhardt, S., Stohl, A.,  
38 Baltensperger, U., Prévôt, A. S. H., Weingartner, E., and Gysel, M.: Black carbon physical properties and mixing state in the  
39 European megacity Paris, *Atmos. Chem. Phys.*, 13, 5831–5856, <https://doi.org/10.5194/acp-13-5831-2013>, 2013.

40 Li, M., Zhang, Q., Kurokawa, J. I., Woo, J. H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng,  
41 Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission  
42 inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963,  
43 2017.



1 Li, W., Sun, J., Xu, L., Shi, Z., Riemer, N., Sun, Y., Fu, P., Zhang, J., Lin, Y., Wang, X., Shao, L., Chen, J., Zhang, X., Wang,  
2 Z., and Wang, W.: A conceptual framework for mixing structures in individual aerosol particles, *J. Geophys. Res.-Atmos.*, 121,  
3 13,784-713,798, 10.1002/2016jd025252, 2016.

4 Liu, D., Allan, J. D., Young, D. E., Coe, H., Beddows, D., Fleming, Z. L., Flynn, M. J., Gallagher, M. W., Harrison, R. M.,  
5 Lee, J., Prevot, A. S. H., Taylor, J. W., Yin, J., Williams, P. I., and Zotter, P.: Size distribution, mixing state and source  
6 apportionment of black carbon aerosol in London during wintertime, *Atmos. Chem. Phys.*, 14, 10061–10084,  
7 <https://doi.org/10.5194/acp-14-10061-2014>, 2014.

8 Liu, D., Joshi, R., Wang, J., Yu, C., Allan, J. D., Coe, H., Flynn, M. J., Xie, C., Lee, J., Squires, F., Kotthaus, S., Grimmond,  
9 S., Ge, X., Sun, Y., and Fu, P.: Contrasting physical properties of black carbon in urban Beijing between winter and summer,  
10 *Atmos. Chem. Phys.*, 19, 6749-6769, 10.5194/acp-19-6749-2019, 2019.

11 Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S., Williams, P. I., Ting,  
12 Y.-C., Haslett, S., Taylor, J. W., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe, H., and Allan, J. D.: Black-carbon absorption  
13 enhancement in the atmosphere determined by particle mixing state, *Nature Geosci.*, 10, 184-188, 2017.

14 Liu, H., Pan, X., Liu, D., Liu, X., Chen, X., Tian, Y., Sun, Y., Fu, P., and Wang, Z.: Mixing characteristics of refractory black  
15 carbon aerosols determined by a tandem CPMA-SP2 system at an urban site in Beijing, *Atmos. Chem. Phys. Discuss.*,  
16 <https://doi.org/10.5194/acp-2019-244>, in review, 2019.

17 Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M. K., Cappa, C. D., Williams, L. R., Herndon, S. C., Massoli, P., Fortner, E. C.,  
18 Chhabra, P. S., Brooks, W. A., Onasch, T. B., Jayne, J. T., Worsnop, D. R., China, S., Sharma, N., Mazzoleni, C., Xu, L., Ng,  
19 N. L., Liu, D., Allan, J. D., Lee, J. D., Fleming, Z. L., Mohr, C., Zotter, P., Szidat, S., and Prévôt, A. S. H.: Enhanced light  
20 absorption by mixed source black and brown carbon particles in UK winter, *Nat Commun.*, 6, 8435, 2015.

21 Lohmann, U., Feichter, J., Penner, J., and Leaitch, R.: Indirect effect of sulfate and carbonaceous aerosols: a mechanistic  
22 treatment, *J. Geophys. Res.*, 105, 12193–12206, 2000

23 Lu, Z., Streets, D. G., Zhang, Q., and Wang, S.: A novel back-trajectory analysis of the origin of black carbon transported to  
24 the Himalayas and Tibetan Plateau during 1996–2010, *Geophys. Res. Lett.*, 39, 2012.

25 Matsui, H., Koike, M., Kondo, Y., Moteki, N., Fast, J. D., and Zaveri, R. A.: Development and validation of a black carbon  
26 mixing state resolved three-dimensional model: Aging processes and radiative impact, *J. Geophys. Res.-Atmos.*, 118, 2304-  
27 2326, 10.1029/2012JD018446, 2013.

28 McMeeking, G. R., Morgan, W. T., Flynn, M., Highwood, E. J., Turnbull, K., Haywood, J., and Coe, H.: Black carbon aerosol  
29 mixing state, organic aerosols and aerosol optical properties over the United Kingdom, *Atmos. Chem. Phys.*, 11, 9037-9052,  
30 2011.

31 Moffet, R. C., and Prather, K. A.: In-situ measurements of the mixing state and optical properties of soot with implications for  
32 radiative forcing estimates, *Proc. Natl. Acad. Sci. USA*, 106, 11872-11877, 10.1073/pnas.0900040106, 2009.

33 Moteki, N., and Kondo, Y.: Dependence of Laser-Induced Incandescence on Physical Properties of Black Carbon Aerosols:  
34 Measurements and Theoretical Interpretation, *Aerosol Sci. Technol.*, 44, 663-675, 2010.

35 Moteki, N., Kondo, Y., Miyazaki, Y., Takegawa, N., Komazaki, Y., Kurata, G., Shirai, T., Blake, D. R., Miyakawa, T., and  
36 Koike, M.: Evolution of mixing state of black carbon particles: Aircraft measurements over the western Pacific in March 2004,  
37 *Geophys. Res. Lett.*, 34, 2007.

38 Morgan, W. T., Allan, J. D., Bauguutte, S., Darbyshire, E., Flynn, M. J., Lee, J., Liu, D., Johnson, B., Haywood, J., Longo, K.  
39 M., Artaxo, P. E., and Coe, H.: Transformation and aging of biomass burning carbonaceous aerosol over tropical South America  
40 from aircraft in-situ measurements during SAMBBA, *Atmos. Chem. Phys. Discuss.*, 2019, 1-32, 10.5194/acp-2019-157, 2019.

41 Mu, Q., Shiraiwa, M., Octaviani, M., Ma, N., Ding, A., Su, H., Lammel, G., Pöschl, U., and Cheng, Y.: Temperature effect on  
42 phase state and reactivity controls atmospheric multiphase chemistry and transport of PAHs, *Sci. Adv.*, 4, 2018.

1 Myhre, G., Berglen, T. F., Johnsrud, M., Hoyle, C. R., Berntsen, T. K., Christopher, S. A., Fahey, D. W., Isaksen, I. S. A., Jones,  
2 T. A., Kahn, R. A., Loeb, N., Quinn, P., Remer, L., Schwarz, J. P., and Yttri, K. E.: Modelled radiative forcing of the direct  
3 aerosol effect with multi-observation evaluation, *Atmos. Chem. Phys.*, 9, 1365-1392, 2009.

4 Nordmann, S., Cheng, Y. F., Carmichael, G. R., Yu, M., Denier van der Gon, H. A. C., Zhang, Q., Saide, P. E., Pöschl, U., Su,  
5 H., Birmili, W., and Wiedensohler, A.: Atmospheric black carbon and warming effects influenced by the source and absorption  
6 enhancement in central Europe, *Atmos. Chem. Phys.*, 14, 12683-12699, 10.5194/acp-14-12683-2014, 2014.

7 Oshima, N., Koike, M., Zhang, Y., Kondo, Y., Moteki, N., Takegawa, N., and Miyazaki, Y.: Aging of black carbon in outflow  
8 from anthropogenic sources using a mixing state resolved model: Model development and evaluation, *J. Geophys. Res.-Atmos.*,  
9 114, 10.1029/2008JD010680, 2009.

10 Peng, J., Hu, M., Guo, S., Du, Z., Zheng, J., Shang, D., Levy Zamora, M., Zeng, L., Shao, M., Wu, Y.-S., Zheng, J., Wang, Y.,  
11 Glen, C. R., Collins, D. R., Molina, M. J., and Zhang, R.: Markedly enhanced absorption and direct radiative forcing of black  
12 carbon under polluted urban environments, *Proc. Natl. Acad. Sci. USA*, 113, 4266-4271, 2016.

13 Ramanathan, V., and Carmichael, G.: Global and regional climate changes due to black carbon, *Nature Geosci.*, 1, 221-227,  
14 2008.

15 Ramnarine, E., Kodros, J. K., Hodshire, A. L., Lonsdale, C. R., Alvarado, M. J., and Pierce, J. R.: Effects of Near-Source  
16 Coagulation of Biomass Burning Aerosols on Global Predictions of Aerosol Size Distributions and Implications for Aerosol  
17 Radiative Effects, *Atmos. Chem. Phys. Discuss.*, 2018, 1-28, 10.5194/acp-2018-1084, 2018.

18 Riemer, N., Vogel, H., and Vogel, B.: Soot aging time scales in polluted regions during day and night, *Atmos. Chem. Phys.*, 4,  
19 1885-1893, 10.5194/acp-4-1885-2004, 2004.

20 Riemer, N., West, M., Zaveri, R. A., and Easter, R. C.: Simulating the evolution of soot mixing state with a particle-resolved  
21 aerosol model, *J. Geophys. Res.-Atmos.*, 114, 10.1029/2008jd011073, 2009.

22 Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S.,  
23 Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S.,  
24 Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-  
25 industrial simulations, *Atmos. Chem. Phys.*, 6, 5225-5246, 2006.

26 Sedlacek, A. J., Lewis, E. R., Kleinman, L., Xu, J., and Zhang, Q.: Determination of and evidence for non-core-shell structure  
27 of particles containing black carbon using the Single-Particle Soot Photometer (SP2), *Geophys. Res. Lett.*, 39, 2012.

28 Shi, S., Cheng, T., Gu, X., Guo, H., Wu, Y., and Wang, Y.: Biomass burning aerosol characteristics for different vegetation  
29 types in different aging periods, *Environment International*, 126, 504-511, <https://doi.org/10.1016/j.envint.2019.02.073>, 2019.

30 Shiraiwa, M., Kondo, Y., Moteki, N., Takegawa, N., Miyazaki, Y., and Blake, D. R.: Evolution of mixing state of black carbon  
31 in polluted air from Tokyo, *Geophys. Res. Lett.*, 34, L16803, doi:10.1029/2007GL029819, 2007.

32 Sun, Y., Jiang, Q., Wang, Z., Fu, P., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe  
33 haze pollution in Beijing in January 2013, *J. Geophys. Res.-Atmos.*, 119, 4380-4398, 2014.

34 Schwarz, J. P., Gao, R. S., Spackman, J. R., Watts, L. A., Thomson, D. S., Fahey, D. W., Ryerson, T. B., Peischl, J., Holloway,  
35 J. S., and Trainer, M.: Measurement of the mixing state, mass, and optical size of individual black carbon particles in urban  
36 and biomass burning emissions, *Geophys. Res. Lett.*, 35, L13810, <https://doi.org/10.1029/2008GL033968>, 2008.

37 Taylor, J. W., Allan, J. D., Liu, D., Flynn, M., Weber, R., Zhang, X., Lefer, B. L., Grossberg, N., Flynn, J., and Coe, H.:  
38 Assessment of the sensitivity of core / shell parameters derived using the single-particle soot photometer to density and  
39 refractive index, *Atmos. Meas. Tech.*, 8, 1701-1718, 10.5194/amt-8-1701-2015, 2015.

40 Wang, L. T., Wei, Z., Yang, J., Zhang, Y., Zhang, F. F., Su, J., Meng, C. C., and Zhang, Q.: The 2013 severe haze over southern  
41 Hebei, China: model evaluation, source apportionment, and policy implications, *Atmos. Chem. Phys.*, 14, 3151-3173, 2014.

1 Wang, X., Heald, C. L., Ridley, D. A., Schwarz, J. P., Spackman, J. R., Perring, A. E., Coe, H., Liu, D., and Clarke, A. D.:  
 2 Exploiting simultaneous observational constraints on mass and absorption to estimate the global direct radiative forcing of  
 3 black carbon and brown carbon, *Atmos. Chem. Phys.*, 14, 10989-11010, 2014.

4 Wang, Y., Liu, F., He, C., Bi, L., Cheng, T., Wang, Z., Zhang, H., Zhang, X., Shi, Z., and Li, W.: Fractal Dimensions and  
 5 Mixing Structures of Soot Particles during Atmospheric Processing, *Environmental Science & Technology Letters*, 4, 487-  
 6 493, 10.1021/acs.estlett.7b00418, 2017.

7 Wang, Y., Ma, P.-L., Peng, J., Zhang, R., Jiang, J. H., Easter, R. C., and Yung, Y. L.: Constraining Aging Processes of Black  
 8 Carbon in the Community Atmosphere Model Using Environmental Chamber Measurements, *Journal of Advances in*  
 9 *Modeling Earth Systems*, 10, 2514-2526, 10.1029/2018ms001387, 2018.

10 Wang, Z., Huang, X., and Ding, A.: Dome effect of black carbon and its key influencing factors: a one-dimensional modelling  
 11 study, *Atmos. Chem. Phys.*, 18, 2821-2834, 2018.

12 Wendisch, M., Hellmuth, O., Ansmann, A., Heintzenberg, J., Engelmann, R., Althausen, D., Eichler, H., Müller, D., Hu, M.,  
 13 Zhang, Y., and Mao, J.: Radiative and dynamic effects of absorbing aerosol particles over the Pearl River Delta, China, *Atmos.*  
 14 *Environ.*, 42, 6405-6416, 2008.

15 Willis, M. D., Healy, R. M., Riemer, N., West, M., Wang, J. M., Jeong, C. H., Wenger, J. C., Evans, G. J., Abbatt, J. P. D., and  
 16 Lee, A. K. Y.: Quantification of black carbon mixing state from traffic: implications for aerosol optical properties, *Atmos.*  
 17 *Chem. Phys.*, 16, 4693-4706, 10.5194/acp-16-4693-2016, 2016.

18 Yang, Y. R., Liu, X. G., Qu, Y., An, J. L., Jiang, R., Zhang, Y. H., Sun, Y. L., Wu, Z. J., Zhang, F., Xu, W. Q., and Ma, Q. X.:  
 19 Characteristics and formation mechanism of continuous hazes in China: a case study during the autumn of 2014 in the North  
 20 China Plain, *Atmos. Chem. Phys.*, 15, 8165-8178, 2015.

21 Zdunkowski, W. G., Welch, R. M., and Paegle, J.: One-Dimensional Numerical Simulation of the Effects of Air Pollution on  
 22 the Planetary Boundary Layer, *J. Atmos. Sci.*, 33, 2399-2414, 1976.

23 Zhang, Y., Li, X., Li, M., Zheng, Y., Geng, G., Hong, C., Li, H., Tong, D., Zhang, X., Cheng, Y., Su, H., He, K., and Zhang,  
 24 Q.: Reduction in black carbon light absorption due to multi-pollutant emission control during APEC China 2014, *Atmos. Chem.*  
 25 *Phys.*, 18, 10275-10287, 2018a.

26 Zhang, Y., Su, H., Ma, N., Li, G., Kecorius, S., Wang, Z., Hu, M., Zhu, T., He, K., Wiedensohler, A., Zhang, Q., and Cheng,  
 27 Y.: Sizing of Ambient Particles From a Single-Particle Soot Photometer Measurement to Retrieve Mixing State of Black  
 28 Carbon at a Regional Site of the North China Plain, *J. Geophys. Res.-Atmos.*, 123, 7778-7795, doi:10.1029/2018JD028810, 2018b.

29 Zhang, Y., Zhang, Q., Cheng, Y., Su, H., Kecorius, S., Wang, Z., Wu, Z., Hu, M., Zhu, T., Wiedensohler, A., and He, K.:  
 30 Measuring the morphology and density of internally mixed black carbon with SP2 and VTDMA: new insight into the  
 31 absorption enhancement of black carbon in the atmosphere, *Atmos. Meas. Tech.*, 9, 1833-1843, 2016.

32 Zhang, Y., Zhang, Q., Cheng, Y., Su, H., Li, H., Li, M., Zhang, X., Ding, A., and He, K.: Amplification of light absorption of  
 33 black carbon associated with air pollution, *Atmos. Chem. Phys.*, 18, 9879-9896, 2018c.

34 Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T., Chang, D., Pöschl, U.,  
 35 Cheng, Y. F., and He, K. B.: Exploring the severe winter haze in Beijing: the impact of synoptic weather, regional transport  
 36 and heterogeneous reactions, *Atmos. Chem. Phys.*, 15, 2969-2983, 10.5194/acp-15-2969-2015, 2015.


37 Zheng, G., Duan, F., Ma, Y., Zhang, Q., Huang, T., Kimoto, T., Cheng, Y., Su, H., and He, K.: Episode-Based Evolution  
 38 Pattern Analysis of Haze Pollution: Method Development and Results from Beijing, China, *Environ. Sci. Technol.*, 50, 4632-  
 39 4641, 10.1021/acs.est.5b05593, 2016.

40


1 Table 1 The in situ measurement in this work.  $(D_p/D_c)_{5\%}$  represents the average value of the lowest 5th percentile of the hourly  
2  $D_p/D_c$  ratio for ambient BC-containing particles during the campaign period.

Measurement	Period	Location	Hourly $PM_{2.5}$ ( $\mu g\ m^{-3}$ )	Hourly rBC ( $\mu g\ m^{-3}$ )	Hourly $D_p/D_c$	$(D_p/D_c)_{5\%}$
BJNOV2014	17-30 November 2014	Tsinghua site, Beijing	~10-440	~0.2-14	~1.4-2.3	1.54
BJOCT2014	28 October-2 November 2014	Tsinghua site, Beijing	~5-200	~0.1-7	~1.4-2.2	1.50
BJSEP2015	12-19 September 2015	Tsinghua site, Beijing	~3-190	~0.1-4	~1.5-2.2	1.55
BJAUG2015	16-23 August 2015	Tsinghua site, Beijing	~3-110	~0.1-3	~1.5-2.0	1.55

3  
4




**Figure 1.** Time series of the PM<sub>2.5</sub> concentration, rBC mass concentration,  $D_p/D_c$  ratio of BC-containing particles and the hourly EEI for BC transported to the observation site for 3 and 5 days.




**Figure 2.** (a) The location of the receptor site (Tsinghua site ( $40^{\circ}00'17''$  N,  $116^{\circ}19'34''$  E), marked as red star) during the BJNOV2014 measurement period. (b) Spatial distribution ( $0.25^{\circ} \times 0.25^{\circ}$ ) of BC emission inventory from MIX.







**Figure 3.** Spatial distribution ( $0.25^{\circ} \times 0.25^{\circ}$ ) of the TED and EEI for BC transported to the observation site ( $40^{\circ}00'17''$  N,  $116^{\circ}19'34''$  E) during episodes 1-4 of the BJNOV2014 measurement period. The TED and EEI were derived from BC emission and black-trajectory analysis (Lu et al., 2012). The TED represents effective trajectory density transported to the receptor site. The EEI quantifies the effective emission amount transported to the receptor site from emission origins.




**Figure 4.** Conceptual scheme of the evolution of the mixing state (i.e.,  $D_p/D_c$  ratio) of BC from the source origin (i.e., grid  $h$ ) to the receptor site following trajectory  $l$ .




**Figure 5.** Comparison of the hourly  $D_p/D_c$  ratio of BC between the model simulations and observations at our site with different  $n$  values in Eq. (15) on 24-30 November 2014 at Tsinghua site. The  $n$  values represent the simulated hourly  $D_p/D_c$  ratio considering the effects of 0, 2, 4, and 6 hour backward transport using Eq. (15).



**Figure 6.** Time series of the observed  $D_p/D_c$  ratio, simulated  $D_p/D_c$  ratio, and  $PM_{2.5}$  and rBC concentrations during the (a) BJOCT2014, (b) BJSEP2015 and (c) BJAUG2015 measurement period.




1  
2 **Figure 7.** Aging degree (i.e.,  $D_p/D_c$ ) of BC particles as they are transported to the receptor site (40°00'17'' N, 116°19'34'' E)  
3 from emission origins (0.25°×0.25°) during the BJNOV2014 campaign period. The gridded  $D_p/D_c$  ratio was calculated by  
4 Eq. (11).




**Figure 8.** BC aging from emission origins to the receptor site (40°00'17" N, 116°19'34" E; marked with a red star) during the BJNOV2014 measurement period: (a) classification of the BC emission source regions based on political boundaries (i.e., the Beijing source regions (BJ origins), Hebei source regions (HB origins), Tianjin source regions (TJ origins) and other source regions (OT origins)); (b) aging degree (i.e.,  $D_p/D_c$ ) of BC particles emitted from the four source regions as they are transported to the receptor site; (c) conceptual scheme of trajectories that pass through low emission regions (cluster 1), medium emission regions (cluster 2) and high emission regions (cluster 3) as they transport from clean origins (i.e., Inner Mongolia) to the receptor site; and (d) the degree of aging of BC reaching the receptor site through different trajectories (i.e., Clusters 1, 2 and 3 shown in (c)).





**Figure 9.** Variations in the (a)  $D_p/D_c$  ratio and (b) light absorption coefficient for total, local and non-local BC over the site with normalized emissions. The current emissions obtained from the MIX inventory (Fig. 2b) were normalized as a unit (i.e., normalized emission of 1), and the emissions reductions of 20%, 40%, 50%, 60% and 80% corresponded to the normalized emissions of 0.8, 0.6, 0.5, 0.4, and 0.2, respectively. a1 and b1 represent the simulations for the case of both local and non-local emission variations, respectively; a2 and b2 represent the simulations for the case of only local emissions variations; and a3 and b3 represent the simulations for the case of only non-local emissions variations. The light absorption coefficient of BC was estimated by the BC mass concentrations, the mass absorption cross section of BC ( $7.5 \text{ m}^2 \text{ g}^{-1}$  at 550 nm) and the  $D_p/D_c$  ratio. When the normalized emissions of total BC were equal to 1, the average mass concentration of total BC was  $\sim 4.0 \mu\text{g m}^{-3}$ , which was obtained by measurements during the campaign period. The mass concentration of local and non-local BC can be further calculated based on their EEI contributions (i.e.,  $\sim 1.6 \mu\text{g m}^{-3}$  for local BC and  $\sim 2.4 \mu\text{g m}^{-3}$  for non-local BC). Based on the linear decrease in BC mass concentration with emission reduction, the mass concentrations of total, local and non-local BC for different emission cases were calculated.





1  
2 Figure 10. (a) Spatial distribution ( $0.25^\circ \times 0.25^\circ$  resolution) for light absorption levels (i.e.,  $EEI*(D_p/D_c)$ ) of BC as it is  
3 transported to the receptor site ( $40^\circ 00' 17''$  N,  $116^\circ 19' 34''$  E) from various source origins. (b) Contributions of different source  
4 regions to BC light absorption at the receptor site. The classification of source regions based on political boundaries is shown  
5 in Fig. 8a. The BJ, HB, TJ and OT origins represent spatial sources of emitted BC transported to the receptor site from Beijing,  
6 Hebei, Tianjin and other regions, respectively.

7

1 *Supplement of*

2 **Modeling the aging process of black carbon during atmospheric**  
3 **transport using a new approach: a case study in Beijing**

4 Yuxuan Zhang<sup>1,2</sup>, Meng Li<sup>2</sup>, Yafang Cheng<sup>2</sup>, Guannan Geng<sup>3</sup>, Chaopeng Hong<sup>4</sup>, Haiyan Li<sup>1</sup>, Xin Li<sup>1</sup>, Dan  
5 Tong<sup>4</sup>, Nana Wu<sup>1</sup>, Xin Zhang<sup>1</sup>, Bo Zheng<sup>5</sup>, Yixuan Zheng<sup>6</sup>, Yu Bo<sup>1,7</sup>, Hang Su<sup>2</sup>, and Qiang Zhang<sup>1</sup>

6 <sup>1</sup>Department of Earth System Science, Tsinghua University, Beijing 100084, China

7 <sup>2</sup>Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz 55128, Germany

8 <sup>3</sup>Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, Georgia 30322, USA


9 <sup>3</sup>Department of Earth System Science, University of California, Irvine, California 92697, USA

10 <sup>3</sup>Laboratoire des Sciences du Climate et de l'Environnement LSCE, Batiment 706, Pte 25, Orme de Merisiers 91191 Gif-sur-  
11 Yvette, France

12 <sup>6</sup>Department of Global Ecology, Carnegie Institution for Science, CA, 94305, USA

13 <sup>7</sup>RCE-TEA, Institute of Atmospheric Physics, Chinese Academy of Science, Beijing, 100029, China

14 *Correspondence to:* Qiang Zhang ([qiangzhang@tsinghua.edu.cn](mailto:qiangzhang@tsinghua.edu.cn)) and Yu Bo ([boyu@mail.iap.ac.cn](mailto:boyu@mail.iap.ac.cn))



1  
 2 Figure S1. The  $(D_p/D_c)_{ini}$  of BC particles at emission origins ( $0.25^\circ \times 0.25^\circ$ ) during the BJNOV2014  
 3 campaign period.