

1 *Supplement of*

2 **Modeling the aging process of black carbon during**
3 **atmospheric transport using a new approach: a case**
4 **study in Beijing**

5 Yuxuan Zhang^{1,2}, Meng Li², Yafang Cheng², Guannan Geng³, Chaopeng Hong⁴,
6 Haiyan Li¹, Xin Li¹, Dan Tong⁴, Nana Wu¹, Xin Zhang¹, Bo Zheng⁵, Yixuan Zheng⁶,
7 Yu Bo^{1,7}, Hang Su², and Qiang Zhang¹

8 ¹Department of Earth System Science, Tsinghua University, Beijing 100084, China

9 ²Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz 55128, Germany

10 ³Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta,
11 Georgia 30322, USA


12 ³Department of Earth System Science, University of California, Irvine, California 92697, USA

13 ³Laboratoire des Sciences du Climat et de l'Environnement LSCE, Batiment 706, Pte 25, Orme de
14 Merisiers 91191 Gif-sur-Yvette, France

15 ⁶Department of Global Ecology, Carnegie Institution for Science, CA, 94305, USA

16 ⁷RCE-TEA, Institute of Atmospheric Physics, Chinese Academy of Science, Beijing, 100029, China

17 *Correspondence to:* Qiang Zhang (qiangzhang@tsinghua.edu.cn) and Yu Bo (boyu@mail.iap.ac.cn)



1

2 Figure S1. The $(D_p/D_c)_{ini}$ of BC particles at emission origins ($0.25^\circ \times 0.25^\circ$) during the
 3 BJNOV2014 campaign period.