

1

2 **Boreal forest fire CO and CH₄ emission factors derived from tower** 3 **observations in Alaska during the extreme fire season of 2015**

4 Elizabeth B. Wiggins^{1*}, Arlyn Andrews², Colm Sweeney², John B. Miller², Charles E. Miller³, Sander
5 Veraverbeke⁴, Roisin Commane⁵, Steven Wofsy⁶, John M. Henderson⁷, and James T. Randerson¹

6 ¹Department of Earth System Science, University of California, Irvine, California, USA, ²National Oceanic and Atmospheric
7 Administration, Boulder, Colorado, USA, ³Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California,
8 USA, ⁴Vrije University Amsterdam, Netherlands, ⁵Department of Earth and Environmental Sciences, Columbia University,
9 Palisades, New York, USA, ⁶School of Engineering and Applied Sciences, Harvard, Cambridge, Massachusetts, USA,
10 ⁷Atmospheric and Environmental Research, Inc., Lexington, Massachusetts, USA,*Now at NASA Langley Research Center,
11 Hampton, VA, USA

12

13 *Correspondence to:* Elizabeth B. Wiggins (Elizabeth.b.wiggins@nasa.gov)

14 **Abstract.** Recent increases in boreal forest burned area, which have been linked with climate warming, highlight the need to better
15 understand the composition of wildfire emissions and their atmospheric impacts. Here we quantified emission factors for CO and
16 CH₄ from a massive regional fire complex in interior Alaska during the summer of 2015 using continuous high-resolution trace
17 gas observations from the Carbon in Arctic Reservoirs Vulnerability Experiment (CRV) tower in Fox, Alaska. Averaged over the
18 2015 fire season, the mean CO/CO₂ emission ratio was 0.142 ± 0.051 and the mean CO emission factor was 127 ± 40 g kg⁻¹ dry
19 biomass burned. The CO/CO₂ emission ratio was about 39% higher than the mean of previous estimates derived from aircraft
20 sampling of wildfires from boreal North America. The mean CH₄/CO₂ emission ratio was 0.010 ± 0.004 and the CH₄ emission
21 factor was 5.3 ± 1.8 g kg⁻¹ dry biomass burned, which are consistent with the mean of previous reports. CO and CH₄ emission
22 ratios varied in synchrony, with higher CH₄ emission factors observed during periods with lower modified combustion efficiency
23 (MCE). By coupling a fire emissions inventory with an atmospheric model, we identified at least 34 individual fires that contributed
24 to trace gas variations measured at the CRV tower, representing a sample size that is nearly the same as the total number of boreal
25 fires measured in all previous field campaigns. The model also indicated that typical mean transit times between trace gas emission
26 within a fire perimeter and tower measurement were 1-3 days, indicating that the time series sampled combustion across day and
27 night burning phases. The high CO emission ratio estimates reported here provide evidence for a prominent role of smoldering
28 combustion, and illustrate the importance of continuously sampling fires across time-varying environmental conditions that are
29 representative of a fire season.

30 **1 Introduction**

31 Boreal forest fires influence the global carbon cycle and climate system through a variety of pathways. These fires initiate
32 succession, influence landscape patterns of carbon accumulation, and directly release carbon dioxide and other trace gases and
33 aerosols into the atmosphere [Johnson, 1996]. One of the largest reservoirs of global terrestrial carbon resides in organic soils
34 underlying boreal forests [Apps *et al.*, 1993; Rapalee *et al.*, 1998; Tarnocai *et al.*, 2009], and fires in the boreal forest can consume
35 significant amounts of aboveground and belowground biomass [Harden *et al.*, 2000; French *et al.*, 2004; Boby *et al.*, 2010; Walker
36 *et al.*, 2018]. Many boreal forest fires are stand replacing and high energy [Johnstone *et al.*, 2011; Rogers *et al.*, 2015], with enough

1 convective power to inject aerosols into the upper troposphere and lower stratosphere where they can be widely dispersed across
2 the Northern Hemisphere [Fromm *et al.*, 2000; Forster *et al.*, 2001; Turquety *et al.*, 2007; Peterson *et al.*, 2018].

3 Emissions from boreal fires are known to considerably influence atmospheric composition in downwind areas. Fire
4 plumes from regional fire complexes in Alaska and western Canada, for example, have been shown to influence air quality over
5 Nova Scotia [Duck *et al.*, 2007], the south-central United States [Wotawa *et al.*, 2001; Kasischke *et al.*, 2005] and Europe [Forster
6 *et al.*, 2001]. Similarly, emissions from boreal forest fires in Russia have caused unhealthy air quality in Moscow [Konovalov *et al.*,
7 2011] and have affected ozone and other trace gases concentrations across the western United States [Jaffe *et al.*, 2004]. Over
8 the past few decades, annual burned area in several regions in boreal North America has increased [Gillett *et al.*, 2004; Kasischke
9 and Turetsky, 2006; Veraverbeke *et al.*, 2017], and projections suggest further increases may occur in response to changes in fire
10 weather and a lengthening of the fire season [Flannigan *et al.*, 2001; de Groot *et al.*, 2013; Young *et al.*, 2017]. As a consequence,
11 fires are likely to play an increasingly important role in regulating air quality and climate during the remainder of the 21st century.

12 Emission factors provide a straightforward way to convert fire consumption of dry biomass into emissions of specific
13 trace gas species, such as CO, CH₄, and CO₂. This technique is commonly used to model emissions of select species in fire
14 inventories, allowing for comparison of atmospheric model simulations with in-situ or remotely sensed mole fraction or
15 concentration observations. The most frequently used boreal forest fire emission factors are derived from meta-analyses that
16 average together information from individual field campaigns [Andreae and Merlet, 2001; Akagi *et al.*, 2011; Andreae, 2019].
17 These syntheses often include in situ airborne and ground based measurements along with laboratory measurements of combusted
18 fuels. There is no consensus on how to combine information from different studies, and in past work individual studies have
19 sometimes been given equal weight when estimating biome-level means, even when the number of fires and duration of sampling
20 has varied considerably from one field campaign to another.

21 In past work, the most common approach for measuring emission factors from boreal fires is to fly aircraft through smoke
22 plumes, measuring trace gases using gas analyzers mounted in the aircraft or by collecting flasks of air that are measured later in
23 the laboratory. Over a period of more than 25 years, a total of at least 42 boreal fires have been sampled by aircraft, including 19
24 wildfires and 14 prescribed land management fires from boreal North America and 9 prescribed fires in Siberia (Table 1). Aircraft
25 sampling is a highly effective approach for sampling large and remote wildfires, especially for characterizing non-conserved trace
26 gas and particulate emissions that have lifetimes of hours to days. It is also important to recognize potential limits associated with
27 sampling fires in this way. Aircraft observations are mostly confined to periods with good visibility, often sampling well-developed
28 fire plumes during mid-day and during periods with relatively low cloud cover. These conditions represent a subset of the
29 environmental variability that a large wildland fire may experience in boreal forest ecosystems as it burns over a period of weeks
30 to months. An alternative approach for measuring in situ emission factors involves using a fixed surface site that continuously
31 samples trace gas concentrations in an area downwind of a fire. This approach has been used to estimate CO emission ratios during
32 a moderate fire season in Alaska [Wiggins *et al.*, 2016] and to estimate emission factors in other biomes [Collier *et al.*, 2016;
33 Benedict *et al.*, 2017; Selimovic *et al.*, 2019; Selimovic *et al.*, 2020]. Surface sampling near or within fire perimeters may have an
34 advantage with respect to providing measurements during intervals when aircraft are unable to fly, but are also more likely to under
35 sample emissions injected above the boundary layer by fire plumes and pyro-cumulus clouds [Selimovic *et al.*, 2019].

36 Environmental conditions, including weather, vegetation, and edaphic conditions are known to influence the composition
37 of emissions, in part by regulating the prevalence of flaming and smoldering combustion processes [Ward and Radke, 1993;
38 Yokelson *et al.*, 1997; Akagi *et al.*, 2011; Urbanski, 2014]. The relative amounts of smoldering and flaming combustion are difficult
39 to measure, but can be estimated using the modified combustion efficiency (MCE), defined as $\Delta\text{CO}_2/(\Delta\text{CO}_2 + \Delta\text{CO})$, where the Δ
40 notation denotes the fire-associated dry air mole fraction of a sample gas after background levels have been removed. Fire

1 emissions dominated by flaming combustion have an MCE from 0.92 – 1.0, while emissions dominated by smoldering combustion
2 have an MCE that often ranges between 0.65 and 0.85 [Akagi *et al.*, 2011; Urbanski, 2014]. MCE can be used to understand the
3 relative contribution of flaming and smoldering combustion processes to the composition of trace gases and aerosols in air
4 measured downwind of a fire. Smoldering combustion converts solid biomass to gases and aerosols, while flaming oxidizes some
5 emissions [Yokelson *et al.*, 1996, 1997]. As a consequence, smoldering combustion produces more CO, CH₄, and organic carbon
6 aerosol relative to CO₂ [Ward and Radke, 1993; Urbanski *et al.*, 2008]. Flaming combustion requires the presence of organic
7 material that burns efficiently [Ryan *et al.*, 2002], and often occurs in boreal forests when fires consume dry aboveground fuels,
8 including vegetation components with low moisture content, litter, and fine woody debris [French *et al.*, 2002]. Smoldering, in
9 contrast, is a dominant combustion process for burning of belowground biomass and larger coarse woody debris. Residual
10 smoldering combustion in boreal forests can continue to occur for weeks after a flaming fire front has passed through, especially
11 in peatland areas with carbon-rich organic soils [Harden *et al.*, 2000; Bertschi *et al.*, 2003]. Over the lifetime of a large fire,
12 smoldering combustion is more likely to occur during periods with lower temperatures and higher atmospheric humidity that
13 increase the moisture content of fine fuels [Stocks *et al.*, 2001; Ryan, 2002].

14 Here we used trace gas observations of CO, CH₄, and CO₂ from the CRV tower to estimate emission factors from boreal
15 forest fires that burned during the near-record high Alaska fire season of 2015. The summer of 2015 was the second largest fire
16 season in terms of burned area since records began in 1940 with about 2.1 million hectares burned [Hayasaka *et al.*, 2016; Partain
17 *et al.*, 2016]. An unseasonably warm spring and early snowmelt allowed fuels to dry early in the season [Partain *et al.*, 2016]. In
18 mid-June, thunderstorms caused an unprecedented number of lightning strikes (over 65,000) that ignited over 270 individual fires
19 on anomalously dry fuel beds over the course of a week [Hayasaka *et al.*, 2016; Veraverbeke *et al.*, 2017]. Fires expanded rapidly
20 during several hot and dry periods through mid-July, and then slowed down as multiple precipitation events and cool, damp weather
21 minimized fire growth for the rest of the summer fire season.

22 The CRV tower captured an integrated signal of trace gas emissions from multiple fires across interior Alaska during the
23 2015 fire season [Karion *et al.*, 2016]. The data stream was comprised of continuous sampling for about 47 minutes out of every
24 hour from June 9 – August 13, yielding more than 59,800 individual measurements, each with a 30 s duration. We identified
25 intervals when fire emissions had a dominant influence on trace gas variability at CRV tower, and used these intervals to derive
26 emission ratios. Analysis of these data indicate that smoldering processes may have a higher contribution to total wildfire emissions
27 from North American boreal forests than previous estimates derived from aircraft sampling. To quantify the spatial and temporal
28 variability of individual fires and their influence on CO, CH₄, and CO₂ at the CRV tower, we coupled a fire emissions inventory,
29 the Alaska Fire Emissions Database (AKFED) [Veraverbeke *et al.*, 2015] with an atmospheric transport model, the Polar Weather
30 Research and Forecasting Stochastic Time Integrated Lagrangian Transport (PWRF-STILT) model [Henderson *et al.*, 2015]. This
31 modeling analysis indicated that the number of 2015 wildfires sampled in our study is comparable to the total number of North
32 American boreal forest fires sampled in past work.

33 **2 Methods**

34 **2.1 CARVE (CRV) Tower Observations**

35 Atmospheric CO, CH₄, and CO₂ mole fractions were measured using a cavity ring-down spectrometer (CRDS, Picarro
36 models 2401 and 2401m) [Karion *et al.*, 2016] at the CRV tower in Fox, Alaska (64.986°N, 147.598°W, ground elevation 611m
37 above sea level). The tower is located about 20 km northeast of Fairbanks Alaska on top of a hill in hilly terrain (Figure 1), and
38 within the interior lowland and upland forested ecoregion in interior Alaska [Cooper *et al.*, 2006]. There are three separate inlets

1 on the CRV tower at different heights above ground level from which the spectrometer draws air for sampling. The spectrometer
2 samples air from the highest level for about 50 minutes out of every hour, and then draws air from the other two levels for 5 minutes
3 at each level [Karion *et al.*, 2016]. Standard reference gases are sampled every 8 hours for 5 minutes, and measurements are
4 removed for a time equivalent to three flushing volumes of the line, approximately 3 minutes, after a level change or switch to or
5 from a calibration tank. All raw 30 s average measurements were calibrated according to Karion *et al.* [2016].

6 We used observations from air drawn from the top intake height at a height of 32 m above ground level in our analysis
7 because this level had the highest measurement density and the smallest sensitivity to local ecosystem CO₂ fluxes near the tower
8 [Karion *et al.*, 2016]. We used gaps in this time series, created when the spectrometer cycled to the lower inlets and following
9 calibration, to separate the time series into discrete time intervals for the calculation of emission ratios. Each 30 s average
10 measurement within a 47-minute sampling interval served as an individual point in our calculation of an emission ratio described
11 below (Table 2).

12 **2.2 Emission Ratios, Emission Factors, and Modified Combustion Efficiency**

13 We isolated intervals when fire had a dominant influence on trace gas variability observed at CRV to calculate emission
14 ratios. An interval with dominant fire influence was defined as a continuous 47-minute measurement period that had: 1) a minimum
15 of at least 30 trace gas measurements (with each measurement representing a mean over 30 seconds), 2) a mean CO over the entire
16 interval exceeding 0.5 ppm, and 3) significant correlations between CO and CO₂, and between CH₄ and CO₂, with r^2 values for
17 both relationships exceeding 0.80.

18 For each interval, we required a sample size of at least 30 individual 30 s measurements. For each interval meeting this
19 criterion, we calculated the mean CO mole fraction and discarded intervals that had a mean CO less than 0.5 ppm. For each of the
20 intervals with mean CO that exceeded the 0.5 ppm threshold, we then extracted the 30 s measurement time series of CO, CH₄, and
21 CO₂ mole fractions and calculated correlation coefficients between the trace gas time series. Only intervals with high and
22 significant correlations between CO and CO₂ and between CH₄ and CO₂ ($r^2 > 0.80$; $p < 0.01$, $n > 30$) were retained, because
23 covariance among these co-emitted species is a typical signature of combustion [Urbanski, 2014]. Data from each of the intervals
24 that met the three criteria described above were used to compute emission ratios, emission factors, and MCE. These intervals are
25 reported in chronological order in Table 2.

26 We calculated background mole fractions of CO and CH₄ by taking an average of observations prior to any major fire
27 activity in interior Alaska during day of year (DOY) 160 – 162.5. This yielded a CO background of 0.110 ppm and a CH₄
28 background of 1.900 ppm. We modeled hourly CO₂ background mole fractions to account for the influence of net ecosystem
29 exchange (NEE) using a multi-variable linear regression model trained on CRV tower observations during 2012, a year with little
30 to no fire influence on trace gas variability. The variables used in the CO₂ model include DOY and hourly observations of
31 temperature, vapor pressure deficit, precipitation, latent heat flux, and hourly CO₂ observations from Barrow, AK (Figure 2).
32 Meteorological variables were acquired from the National Climatic Data Center Automated Weather Observing System for
33 Fairbanks International Airport (<http://www7.ncdc.noaa.gov/CDO/cdopoemain.cmd>). This location was chosen due to its
34 proximity to the CRV tower. We obtained 3-hourly latent heat flux estimates from the NOAA2.7.1 GLDAS/NOAH experiment
35 001 for version 2 of the Global Land Data Assimilation System (GLDAS-2) [Rodell *et al.*, 2015]. Hourly in situ CO₂ observations
36 from a clean air site at Barrow, AK were obtained from the Earth System Research Laboratory Global Monitoring Division
37 [Thoning *et al.*, 2007]. Our model assumed negligible influence from fossil fuel combustion on background mole fraction
38 variability. After training on data from the summer of 2012, the model was then run using 2015 input variables to calculate time

1 evolving CO₂ background mole fractions during our analysis period. In a final step, the hourly CO₂ model was linearly interpolated
 2 to have the same temporal resolution as the 30 s individual trace gas measurements.

3 We estimated an emission ratio (ER_X ; equation 1) by calculating the slope from a type II linear regression of CO or CH₄
 4 excess mole fractions (ΔX) relative to the CO₂ excess mole fraction (ΔCO_2) using all of the 30 s observations available within a
 5 single 47-minute sampling interval when fire had a dominant influence on tower trace gas variability (up to 95 pairs of
 6 measurements). Uncertainty estimates for each interval were estimated as the standard deviation of the slope of the regression. To
 7 estimate excess mole fractions (denoted with a Δ), we first removed background mole fractions (described above) before
 8 performing the regression analysis and obtaining the slope. The assumed background levels for CO and CH₄ did not influence this
 9 emission ratio estimate, because they were assumed to remain constant throughout the duration of each 47-minute interval (i.e.,
 10 they influenced the intercept but not the slope of the regression line). In a sensitivity analysis we found that the removal of the CO₂
 11 background, which did evolve within each 47-minute interval, had only a negligible effect, because the CO₂ background did not
 12 change rapidly over time. Since multiple fires were often burning simultaneously during the 2015 fire season, the emission ratios
 13 we report in Table 2 for each interval likely represent a composite of emissions from several fires.

$$14 \quad ER_X = \frac{\Delta X}{\Delta CO_2} = \frac{X_{Fire} - X_{Background}}{CO_2 Fire - CO_2 Background} \quad (1)$$

15 Emission factors (EF_X) were calculated using equation 2, where MM_X is the molar mass of CO or CH₄, MM_C is the molar
 16 mass of carbon, F_C is the mass fraction of carbon in dry biomass, 1000 is a factor to convert kg to g, ER_X is the emission ratio, and
 17 C_T is given by equation 3. The units for an emission factor are grams of compound emitted per kg dry biomass burned. In equation
 18 3, n is the number of carbon containing species measured, N_i is the number of carbon atoms in species i , and ΔX_i is the excess mole
 19 fraction of species i [Yokelson *et al.*, 1999; Akagi *et al.*, 2011]. Here we computed C_T by allowing i in equation 3 to cycle over
 20 CO₂, CO, and CH₄ ($n = 3$). We assumed the fraction of carbon in combusted fuels, F_C , was 0.45 [Santin *et al.*, 2015], but note that
 21 F_C can range from 0.45 – 0.55 [Akagi *et al.*, 2011].

$$22 \quad EF_X = \frac{MM_X}{MM_C} * F_C * 1000 * \frac{ER_X}{C_T} \quad (2)$$

$$23 \quad C_T = \sum_{i=1}^n N_i * \frac{\Delta X_i}{\Delta CO_{2i}} \quad (3)$$

24 We also calculated the MCE for each fire-affected interval. Modified combustion efficiency is defined as the excess mole
 25 fraction of CO₂ divided by the sum of the excess mole fractions of CO and CO₂ [Ward and Radke, 1993]. MCE was used to separate
 26 intervals into three categories: smoldering, mixed, or flaming. These categories reflect the dominant combustion process
 27 contributing to trace gas anomalies at the CRV tower during the summer of 2015. Periods with an MCE less than 0.85 were
 28 considered to consist of mostly smoldering combustion, periods with a MCE of greater than or equal to 0.85 and less than 0.92
 29 were classified as consisting of a mixture of smoldering and flaming combustion, and periods with an MCE greater than 0.92 were
 30 classified as flaming [Urbanski, 2014]. We performed this classification to allow for a visualization of how the sampled combustion
 31 processes varied from interval to interval (and day to day) during the 2015 fire season.

32 2.3 Transport Modeling

33 We coupled a fire emission model, the Alaskan Fire Emissions Database (AKFED) [Veraverbeke *et al.*, 2015] with an
 34 atmospheric transport model, the Polar Weather Research and Forecasting Stochastic Time Integrated Lagrangian Transport model

1 (PWRP-STILT) [Henderson *et al.*, 2015] to estimate fire contributions to trace gas variability at the CRV tower, following Wiggins
2 *et al.* [2016]. For this application, STILT [Lin *et al.*, 2007] was used to estimate the adjoint of PWRP [Skamarock *et al.*, 2005;
3 Chang *et al.*, 2014; Henderson *et al.*, 2015] during the summer of 2015 at the location of the CRV tower, to generate surface
4 influence functions that relate surface ecosystem fluxes from Alaska to trace mole fractions at CRV. These gridded influence
5 functions are known as footprints and have units of mole fraction per unit of surface flux ($\text{ppm}/(\mu\text{mol m}^{-2} \text{ s}^{-1})$). Here we emitted
6 fire emissions into the surface influenced volume of PWRP-STILT, which extends from the surface to the top of the planetary
7 boundary layer, with the assumption that fire emissions were equally distributed within the planetary boundary layer [Turqueti *et*
8 *al.*, 2007; Kahn *et al.*, 2008]. In a previous study using the same tower, a sensitivity study revealed plume injection height
9 contributed only minimally to variability in remote fire CO predictions at CRV with PWRP-STILT [Wiggins *et al.*, 2016].

10 Daily burned area in AKFED was mapped using thermal imagery from the Moderate Resolution Imaging
11 Spectroradiometer (MODIS) within fire perimeters from the Alaska Large Fire Database. Both above and belowground carbon
12 consumption were modeled as a function of elevation, day of burning, pre-fire tree cover, and difference normalized burn ratio
13 (dNBR) measurements derived from 500 m MODIS surface reflectance bands [Veraverbeke *et al.*, 2015]. AKFED predicted carbon
14 emissions from fires with a temporal resolution of 1 day and a spatial resolution of 450 m. We regridded AKFED to the same
15 spatial resolution as the atmospheric transport model (0.5°) for the model coupling. To account for diurnal variability in emissions,
16 here we imposed a diurnal cycle on daily emissions following Kaiser *et al.* [2009], where the diurnal cycle was the sum of a
17 constant and a Gaussian function that peaks in early afternoon with 90% of emissions occurring during the day (hours 0600 to
18 1800 local time) and 10% at night (hours 1800 to 0600 local time). Analysis of the sum of fire radiative power from all of the fire
19 detections in the MODIS MCD14ML C6 product showed that 83% of detected fire activity occurred during the daytime overpasses
20 (10:30am and 1:30pm) relative to the sum across both daytime and nighttime overpasses during the 2015 Alaskan wildfire season
21 (data not shown). The satellite observations, although temporally sparse (with only 4 over passes per day), were broadly consistent
22 with the diurnal cycle we prescribed for fire emissions in the model.

23 We convolved AKFED with the PWRP-STILT footprints to determine individual fire contributions to CO anomalies at
24 the CRV tower. This was achieved by calculating the total CO contribution from each individual 0.5° grid cell from the AKFED
25 \times PWRP-STILT combined model and utilizing the fire perimeters from the Alaska Large Fire Database (data provided by Bureau
26 of Land Management (BLM) Alaska Fire Service, on behalf of the Alaska Wildland Fire Coordinating Group (AWFCG) and
27 Alaska Interagency Coordination Center (AICC)) to identify the location of individual fires. AKFED uses the same fire perimeter
28 database for burned area and carbon emissions estimates [Veraverbeke *et al.*, 2015]. We determined an individual fire's
29 contribution to CO at the CRV tower by setting all emissions in AKFED for a particular grid cell to zero and rerunning the model
30 coupling with PWRP-STILT. The difference between the original model and the updated coupling that excluded emissions from
31 an individual fire was equal to the individual fire's contribution to CO at CRV tower, when integrated over the 2015 fire season.
32 Due to the 0.5° grid cell size used for model coupling, more than one fire perimeter existed in some of the individual grid cells. In
33 these cases, the contribution for each fire was determined by weighting the total signal contribution by fire size.

34 We also used the footprints from PWRP-STILT to quantify the contribution of day and night emissions and mean transit
35 times (Figure 3). The footprints are on a 0.5° latitude-longitude grid with a temporal resolution of 1 h during hours 0600 to 1800
36 (day) local time and 3 h during hours 1800 to 0600 local time (night). These functions provide an estimate of the impact of upwind
37 surface fluxes at different times in the past on CRV tower trace gas mole fraction measurements at a given time. We analyzed the
38 footprints for each interval in Table 2 to confirm CRV tower observations integrated emissions from multiple fires and captured
39 variability in emissions across the diurnal fire cycle. Overall, we found that 73% of the summer fire CO anomaly at CRV originated
40 from fire emissions that occurred during the day (0600 to 1800 local time) and 27% from emissions that occurred at night (1800 –

1 0600 local time). The footprints associated with each emission factor interval also were used to determine how much of the signal
2 was coming from burning on previous days. We found that more than 99% of the fire emissions that influenced CO at CRV
3 occurred within 3 days of an sampling interval used to derive an emission ratio, with 76% occurring within the first 24 hours, 21%
4 during the next 24 hours, and 3% occurring three days prior to the sampling interval.

5 **2.4 Comparison with Previous CO Emission Ratio Studies**

6 To investigate the possible influence of sampling strategy and differences associated with sampling in different ecosystem
7 types, we compiled available studies that report CO emission ratios for boreal forest fires and organized the studies into several
8 categories with common characteristics, including aircraft sampling of North American boreal forest wildfires, aircraft sampling
9 of North American boreal forest management or prescribed fires, combustion of North American boreal forest fuels measured in
10 the laboratory, and sampling of Siberian boreal fires from both aircraft and surface platforms (Table 1). In our analysis we included
11 original studies reported in Andreae (2019) and Akagi et al. (2011) and several others we found in a literature survey.

12 **3 Results**

13 **3.1 Emission Factors and Modified Combustion Efficiency**

14 During the 2015 Alaska fire season, we observed synchronized enhancements of CO, CH₄, and CO₂ well above
15 background concentrations at CRV from DOY 173 – 196 (Figure 4). We identified 55 individual fire-affected intervals in the
16 measurement time series (that each span about 47 minutes) and used these intervals to calculate emission ratios, emission factors,
17 and MCE (Figure 5; Table 2). CO/CO₂ emission ratios ranged from 0.025 to 0.272 and CH₄/CO₂ emission ratios ranged from 0.002
18 to 0.020. MCE varied between 0.786 and 0.976 (Table 2). CO emission factors ranged from 25 to 223 g kg⁻¹ dry biomass burned,
19 and CH₄ emission factors ranged from 1.2 to 10.7 g kg⁻¹ dry biomass burned.

20 The mean CO/CO₂ emission ratio was 0.141 ± 0.051 , the mean CO emission factor was 127 ± 40 g kg⁻¹ dry biomass
21 burned, and the mean MCE was 0.878 ± 0.039 . Concurrently, the mean CH₄/CO₂ emission ratio was 0.010 ± 0.004 and the mean
22 CH₄ emission factor was 5.32 ± 1.82 g kg⁻¹ dry biomass burned.

23 A strong linear relationship existed between the CH₄ emission factor and MCE across the different sampling intervals
24 (Figure 6). Linear relationships between CH₄ emission factors and MCE have also been observed in previous studies [*Yokelson et*
25 *al.*, 2007; *Burling et al.*, 2011; *Van Leeuwen and van der Werf*, 2011; *Yokelson et al.*, 2013; *Urbanski*, 2014; *Smith et al.*, 2014;
26 *Strand et al.*, 2016, *Guerette et al.*, 2018]. The relationship shown in Figure 6 implies MCE can be used to estimate CH₄ emissions
27 (and emissions of other closely related trace gases) from North American boreal forest wildfires when measurements of CH₄ are
28 not available.

29 We classified each fire-affected sampling interval as being associated with smoldering, mixed, or flaming combustion
30 processes using thresholds on MCE. This analysis revealed that intervals with different combustion phases were interspersed
31 throughout the fire season, with no clear progression over time, or clustering of flaming or smoldering processes during periods
32 with high or low levels of burning. We identified 12 smoldering intervals, 37 mixed intervals, and 6 flaming intervals throughout
33 the fire season (Figure 5, with examples shown in Figure 7). Smoldering intervals had a mean CO/CO₂ ratio of 0.214 ± 0.030 , a
34 mean CO emission factor of 183 ± 21 g kg⁻¹ dry biomass burned, a mean CH₄/CO₂ ratio of 0.014 ± 0.003 , a mean CH₄ emission
35 factor of 6.89 ± 1.18 g kg⁻¹ dry biomass burned, and a mean MCE of 0.824 ± 0.020 . Mixed intervals consisting of both smoldering
36 and flaming combustion had a mean CO/CO₂ emission ratio of 0.131 ± 0.024 , a mean CO emission factor of g kg⁻¹ dry biomass

1 burned, a mean CH₄/CO₂ emission ratio of 0.010 ± 0.003, a mean CH₄ emission factor of 5.28 ± 1.51 g kg⁻¹ dry biomass burned,
2 and a mean MCE of 0.884 ± 0.019. Flaming intervals had a mean CO/CO₂ emission ratio of 0.060 ± 0.020, a mean CO emission
3 factor of 59 ± 19 g kg⁻¹ dry biomass burned, a mean CH₄/CO₂ emission ratio of 0.004 ± 0.001, a mean CH₄ emission factor of 2.49
4 ± 0.78 g kg⁻¹ dry biomass burned, and a mean MCE of 0.944 ± 0.018 (Table 3).

5 In our primary analysis described above, each individual fire-influenced interval used to compute an emission ratio was
6 weighted equally in computing a season-wide mean. As a sensitivity analysis, we computed the mean emission ratios weighting
7 each interval according to its mean ΔCO mole fraction, and, alternately, according to its mean ΔCO₂ mole fraction. Weighting by
8 ΔCO caused the CO emission ratio to increase from 0.141 to 0.146 but did not change the CH₄ emission ratio. Weighting by ΔCO₂
9 caused the emission ratios to slightly increase, yielding a CO emission ratio of 0.144 and, again, no change in the CH₄ emission
10 ratio. Although the variation introduced from different weighting approaches was relatively small, the analysis highlights the
11 challenge of combining information from different individual fires, and the importance of moving toward flux-weighted estimates
12 in future work.

13 3.2 The Influence of Individual Fires on Trace Gas Variability at the CRV Tower

14 The forward model simulations combining AKFED fire emissions with PWRP-STILT confirmed that the elevated CO
15 signals at the CRV tower can be attributed primarily to boreal forest fire emissions (Figure 8) and not to fossil fuels or other CO
16 sources. The AKFED model had a Pearson's correlation coefficient of 0.61 with observed daily mean CO and had a low bias of
17 approximately 7%. Differences between the model simulations and observations were likely caused by errors in the magnitude and
18 timing of fire emissions within AKFED as well as the limited spatial resolution and incomplete representation of atmospheric
19 transport within PWRP-STILT. Nevertheless, the broad agreement between the model and the observations, including the timing
20 of the large burning interval between DOY 173 and 179, provides some confidence that our model can be used to explore the
21 influence and contribution of individual fires.

22 We identified 34 individual fires that contributed to at least 1% of the CO mole fraction time series at CRV tower over
23 the entire 2015 fire season (Figure 9; Figure 10; Table 3). The average distance of these fires from the CRV tower, weighted by
24 their fractional contribution, was 259 ± 134 km. Most of the fires were located to the west of Fairbanks, in the direction of the
25 prevailing summer surface winds. This analysis revealed that the CRV tower was sufficiently downwind to measure the integrated
26 impact of multiple fires on regional trace gas concentration anomalies, sampling air masses that were mixed through the full
27 planetary boundary layer and across several day-night cycles. The total CO emitted from these fires accounted for 75% of the
28 excess CO mole fraction signal during DOY 160 – 200. The remaining CO signal originated from many smaller fires that were
29 widely distributed across interior Alaska. The Tozitna fire was responsible for the greatest percentage of the total CO anomaly
30 integrated over the 2015 fire season at the CRV tower (accounting for more than 10% of the integrated CO anomaly at CRV). The
31 fires that contributed the most to the CO anomaly at CRV tower were not necessarily the closest fires to the tower or the largest
32 fires of the 2015 fire season in terms of burned area. Combined, however, this set of 34 fires accounted for 0.97 Mha, or
33 approximately 46% of the total burned area reported during the 2015 fire season [Veraverbeke *et al.*, 2017].

34 3.3 Comparison of emission ratios between sampling strategies

35 Previous studies sampled a total of 45 individual boreal forest fires for ΔCO/ΔCO₂ emission ratios or CO emission factors,
36 and additional measurements have been made by combusting fuels in a laboratory setting. Solely considering emission ratio
37 measurements from North American boreal forests (excluding boreal forests in Eurasia), the mean of aircraft sampling of wildfires

1 (0.102 ± 0.033, n=19) or management and prescribed fires (0.077 ± 0.022, n=14) were significantly lower than the mean derived
2 from tower measurements reported here along with earlier measurements from *Wiggins et al.* [2016] (0.141 ± 0.049, n=37) as
3 evaluated using a Student's t test. The mean emission ratio from Siberian boreal forest fires was 0.219 ± 0.048 (n=9), which was
4 significantly higher than the mean of emission ratios reported for boreal forest wildfires in North America (sampled either by
5 aircraft or tower).

6 **4 Discussion**

7 The most widely used emission factors for boreal forest fires are derived from syntheses that average together data from
8 individual field campaigns [*Andreae and Merlet*, 2001; *Akagi et al.*, 2011; *Andreae*, 2019]. Our mean emission factor for CO (127
9 ± 40 g kg⁻¹ dry biomass burned) is similar to the mean reported in past syntheses for boreal forests, including estimates by *Andreae*
10 [2019] (121 ± 47 g kg⁻¹ dry biomass burned) and *Akagi et al.* [2011] (127 ± 45 g kg⁻¹ dry biomass burned). Emission factors for
11 CH₄ were also similar to the estimates reported in these syntheses. Considering boreal forests as a whole, our measurements provide
12 a partial validation of the approach taken in previous compilations, which have attempted to combine information from different
13 sampling strategies and boreal forest ecoregions. The broad level of agreement provides confidence in the estimates of emission
14 factors for non-conserved species that cannot be measured using a remote tower sampling approach.

15 The observations summarized in Table 1 also show there are several important differences in boreal forest emission ratios
16 that exist as a function sampling strategy and ecoregion. Within North American boreal forests, the CRV observations we analyzed
17 here provide evidence that smoldering combustion contributes more to CO emissions than what has been estimated from previous
18 aircraft studies. Specifically, our mean CO emission ratio from the CRV tower is 39% higher (and significantly different at a p <
19 0.01 level using a Student's t test) than the mean derived from aircraft based measurements of 19 North American boreal wildfires
20 (Table 1). Although differences in reported emission ratios are expected between aircraft and ground based sampling approaches
21 [*Christian et al.*, 2007; *Burling et al.*, 2011; *Akagi et al.*, 2014; *Collier et al.*, 2016; *Benedict et al.*, 2017; *Selimovic et al.*, 2019],
22 several features of the CRV tower sampling are conducive to providing a regionally-representative mean estimate of emission
23 ratios during the 2015 Alaska fire season. First, we note that the CRV tower was located at a higher elevation (611 m above sea
24 level) than the core fire complex located in western Alaska and several hundreds of kilometers downwind. Multi-angle Imaging
25 SpectroRadiometer (MISR) satellite observations from Alaskan wildfires indicate most fire plumes reside within the planetary
26 boundary layer, which is typically between 1 and 3 km during midday in summer [*val Martin et al.*, 2010; *Wiggins et al.*, 2016].
27 Combining this vertical length scale with the mean horizontal distance of the 34 fires that most influenced CO at CRV (259 km),
28 we obtain a factor of about 100 for a back-of-the-envelope ratio of horizontal to vertical mixing processes. This ratio, together with
29 the simulated time delay of 1-2 days between emission and detection of CO anomalies at CRV (Figure 3), imply that mesoscale
30 atmospheric circulation played an important role in averaging together trace gas emissions from multiple fires before the air masses
31 were sampled (Figure 10). As a result, observations from the CRV tower represent a temporal integration of fire emissions over
32 day-night burning cycles as well as a spatial integration across flaming combustion at active fire fronts along with residual
33 smoldering combustion in soils that often persists for days after a fire front moves through an area. Collectively, the fires sampled
34 at CRV appeared to experience time-varying environmental conditions that were less ideal for flaming combustion than the fire
35 plumes sampled in past work by aircraft. This finding is consistent with remote tower observations of the black carbon to CO ratio
36 measured for wildfires from temperate North America [*Selimovic et al.*, 2019].

37 In contrast with remote tower sampling, aircraft-based studies often sample fires that have a strong contribution from
38 flaming combustion, which releases enough energy to generate well-defined plumes at an altitude accessible by the aircraft. This

1 methodology provides an opportunity to comprehensively measure the vertical and horizontal distribution of emissions from an
2 individual fire and their atmospheric evolution in a smoke plume. However, airborne sampling techniques are often limited to
3 daytime periods with good visibility, making it difficult to comprehensively measure emissions over a diurnal cycle or over the
4 full lifetime of a fire which may span several periods with inclement weather. Due to these sampling constraints, aircraft studies
5 are less likely to measure emissions from less energetic smoldering combustion, since these emissions are more likely to remain
6 near the surface [Ward and Radke, 1993; Selimovic et al., 2019]. Emissions from smoldering boreal forest fires can sometimes be
7 entrained in the convective columns of certain flaming fires and can be sampled by aircraft, but nighttime emissions or residual
8 smoldering emissions from fires that have weak convective columns usually cannot be measured in this way [Bertschi et al., 2003;
9 Burling et al., 2011]. While past studies have attempted to combine information from aircraft (more likely sampling flaming
10 combustion phases) with laboratory observations of emissions from smoldering combustion [Akagi et al., 2011], the balance of
11 these processes is well known to be sensitive to environmental conditions that can rapidly change over the lifetime of a wildfire;
12 this highlights the importance of designing sampling approaches that provide regionally-integrated estimates over the full duration
13 of a wildfire event or a regional fire complex.

14 During the latter half of June and early July of 2015, weather in Alaska was very hot and dry, allowing for a record number
15 of fires to rapidly expand in size, and yielding the second highest level of annual burned area in the observed record. The extreme
16 fire weather conditions would be expected to reduce fuel moisture content, thus promoting crown fires and flaming combustion
17 processes [e.g., Sedano and Randerson, 2014]. This raises the question of whether longer term monitoring of many normal and
18 low fire years (which tend to co-occur in cooler and wetter conditions) would provide evidence for an even larger role of smoldering
19 combustion compared to the estimates we report here for 2015. Another related question is whether even within a fire season, do
20 day-to-day or week-to-week variations in fire weather influence variability in emission ratios? We explored this latter question
21 with the datasets described here but were unable to uncover structural relationships between daily meteorological variables such
22 as vapor pressure deficit and CO emission ratios. Together, these questions represent important directions for future research and
23 emphasize the critical need of sustained long-term support for trace gas monitoring networks and field campaigns.

24 As a function of ecoregion, emission ratios from fires in boreal Eurasia tend to be higher than emission ratios from fires
25 in boreal North America, and are significantly different than tower or aircraft observations from North America when compared
26 using a Student's t test. Although more measurements are needed, higher CO emission ratios for Siberian fires appear consistent
27 with past work showing that boreal fire behavior is considerably different between North American and Eurasian continents as a
28 consequence of differences in tree species and their impacts on fire dynamics [Goldammer and Furyaev, 1996; Cofer et al., 1998].
29 Notably, as consequence of the presence of black spruce in many boreal forests of North America, fires tend to burn with a higher
30 fire radiative power and faster spread rate, traveling through the crowns of trees and inducing higher levels of tree mortality [Rogers
31 et al., 2015]. This occurs because black spruce is a well-known fire embracer, retaining dead branches that serve as ladder fuels
32 and carry fire into the overstory. Black spruce trees are absent from Siberia, where many pine and larch tree species lack ladder
33 fuels and are known to be fire resistors. In Siberian ecosystems ground fires are more common [Korovin, 1996; Rogers et al.,
34 2015], a finding that appears consistent with the higher CO emission ratios (and larger contribution of smoldering combustion)
35 shown in Table 1. Although emission factors from the Siberian boreal forest are often grouped together with emission factors from
36 North American boreal forest in biome-level syntheses [e.g., Andreae, 2019], both emission ratio and remote sensing observations
37 of fire severity suggest there may be enough evidence to separate these two ecoregions in future syntheses.

38 In Table 1 we also separated aircraft-based studies that measured emissions from wildfires from those that measured
39 emissions from prescribed slash and land management fires, where trees are bulldozed, dried and intentionally arranged to promote
40 maximum fuel consumption [Cofer et al., 1990; Cofer et al., 1998]. Land management fires consume dried aboveground fuels with

1 a different fuel structure and moisture content than fuels consumed in a wildfire, where combustion from soil organic material
2 layers is a dominant component of bulk emissions [Boby et al., 2010; Dieleman et al., 2020]. Although the number of land
3 management fires is relatively small, the mean from these studies suggest flaming processes are a more important contributor to
4 this fire type than for wildfires, and some consideration of this difference should be factored into regional and global syntheses.

5 Several additional studies report emission ratios from laboratory combustion of fuels collected from North American
6 boreal forests including biomass samples from black spruce, white spruce, and jack pine, as well as moss and surface organic
7 material (duff). The laboratory studies have considerable variability that can be attributed to the type of fuel combusted and fuel
8 moisture content. This work indicates duff consumption yields higher emission ratios for CO and CH₄ than combustion of black
9 spruce or jack pine needles and other fine fuels [Bertschi et al., 2003; McMeeking et al., 2009; Burling et al., 2011]. The fuels used
10 in laboratory studies are usually dried and burned individually, although some studies have attempted to mimic natural fires by
11 placing dried fine fuels on top of damp fuels that undergo residual smoldering combustion [Bertschi et al., 2003]. The structure,
12 composition, and moisture content of fuels are well known as key drivers of the composition and magnitude of emissions. Although
13 these laboratory studies provide valuable information on emissions from individual fuel components, they are not able to capture
14 the full complexity of a wildfire.

15 In the context of these comparisons among ecoregions and sampling strategies, it is important to recognize that tower-
16 based sampling strategies, including the methodology presented in this study, have important limits. Ground-based sites may
17 potentially miss some of the emissions injected above the planetary boundary layer. The fixed nature of this sampling technique
18 also restricts the range of sampling, because towers can only monitor upwind fires. Although the tower-based sampling strategy
19 allows for integration of emissions from fires across a range of environmental conditions and at different stages of fire life cycles,
20 it may not allow for emission ratio measurements of non-conserved species, including particulate matter and many fire-emitted
21 volatile organic compounds that have short lifetimes. The technique is also subject to higher uncertainty in the definition of
22 background mole fractions for fire-affected trace gases, because of the dilution and mixing of fire emissions that occurs during
23 transport. Thus, tower may not be a feasible or effective sampling methodology during years with low fire activity.

24 **5 Conclusions**

25 Using a remote tower downwind of a large regional fire complex in interior Alaska, we measured CO and CH₄ emission
26 factors from about 34 individual fires during the summer of 2015. This is comparable to the number of individual wildfires sampled
27 in North America in previous studies. Our results indicate smoldering combustion processes in North American boreal forest fires
28 contribute to more trace gas emissions than previous estimates derived from aircraft sampling. Together, the two-month near
29 continuous time series of CO₂, CO, and CH₄, along with the derived emission ratios reported here, may provide a means to test
30 models that couple together fire processes, emissions, and regional atmospheric transport.

31 Comparison of emission ratios reported here with observations derived other sampling strategies and ecoregions in
32 northern boreal forests provides directions for reducing future uncertainties. For boreal North America, our analysis of CRV tower
33 observations indicate CO emission ratios are likely higher what would be inferred from previous studies, although questions remain
34 regarding the representativeness of remote tower-based sampling. Given recent increases in data density for North America and
35 improvements in our understanding of differences in tree species composition and fire dynamics between North America and
36 Eurasia, it may be possible to reduce uncertainties in future syntheses by separately reporting emission factors for the two
37 continents. More data, particularly for Siberian fires, however, is needed to assess whether the continental differences in emission
38 ratios noted here are robust. Long-term monitoring from remote towers has the potential to provide new information about fire

1 complexes in other biomes, integrating across day-night variations in fire behavior, periods with different environmental
2 conditions, and across multiple fires in different stages of growth and extinction. In this context, more work is needed to find ways
3 to combine tower and aircraft sampling to attain accurate estimates of the total budget of fire-emitted trace gases and aerosols (i.e.,
4 estimating flux-weighted emission factors), given the large differences in data density and the different strengths and weaknesses
5 of the two approaches. To make progress on this issue, a closer integration is needed in future field campaigns between
6 measurements of pre-fire ecosystem state, fire behavior (temperature, fire radiative power, and spread rate), measurements of
7 emissions composition, and post-fire sampling of fuel consumption and combustion completeness during times when fire dynamics
8 are fundamentally different. This coordination across disciplines in both study design, data analysis, and modeling is rare and may
9 provide a path toward creating the observations needed to dynamically model the temporal evolution of the chemical composition
10 of wildland fire emissions over the lifetime of an individual fire and, within a region, during different phases of a fire season.

11 **Data availability**

12 The CRV tower observations and footprints used in our analysis are archived at the U.S. Oak Ridge National Laboratory Distributed
13 Active Archive Center for Biogeochemical Dynamics (<http://dx.doi.org/10.3334/ORNLDAAC/1316>).

14 **Author contribution**

15 Conceptualization: EBW and JTR. Data curation: CEM, SV, JMH, and AA. Investigation and Visualization: EBW. Writing –
16 original draft: EBW and JTR. Writing – review and editing: Everyone.

17 **Competing interest**

18 The authors declare that they have no conflict of interest.

19 **Acknowledgements**

20 E.B.W. thanks the U.S. National Science Foundation for a Graduate Research Fellowship (NSF 2013172241). The trace gas
21 observations, fire emissions time series, and WRF-STILT model were created through funding support to NASA's CARVE field
22 program led by C. Miller. JTR acknowledges additional NASA support from CMS (80NSSC18K0179) and IDS
23 (80NSSC17K0416) programs. We thank the staff of the NOAA Fairbanks Command and Data Acquisition Station that hosts CRV,
24 and especially Frank Holan and Marc Meindl, for their technical support. We also thank NOAA/ESRL/GMD staff, especially Phil
25 Handley, Jon Kofler, and Tim Newberger for ongoing remote maintenance of CRV.

26 **References**

27 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D. and Wennberg, P. O.: Emission
28 factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039-4072,
29 doi:10.5194/acp-11-4039-2011, 2011.

30

1 Akagi, S. K., Burling, I. R., Mendoza, A., Johnson, T. J., Cameron, M., Griffith, D. W. T., Paton-Walsh, C., Weise, D. R., Reardon,
2 J. and Yokelson, R. J.: Field measurements of trace gases emitted by prescribed fires in southeastern US pine forests using an
3 open-path FTIR system. *Atmos. Chem. and Phys.*, 14, 199-215, <http://dx.doi.org/10.5195/acp-14-199-2014>, 2014.
4
5 Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, 15, 955-
6 966, doi:10.1029/2000GB001382, 2001.
7
8 Andreae, M.O.: Emission of trace gases and aerosols from biomass burning—an updated assessment. *Atmos. Chem. Phys.*, 19,
9 8523-8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
10
11 Apps, M. J., Kurz, W. A., Luxmoore, R. J., Nilsson, L. O., Sedjo, R. A., Schmidt, R., Simpson, L. G. and Vinson, T. S.: Boreal
12 forests and tundra, *Water, Air, Soil Pollut.*, 70, 39-43, doi:10.1007/BF01104987, 1993.
13
14 Benedict, K. B., Prenni, A. J., Carrico, C. M., Sullivan, A. P., Schichtel, B. A., and Collett Jr., J. L.: Enhanced concentrations of
15 reactive nitrogen species in wildfire smoke, *Atmos. Environ.*, 148, 8–15, <https://doi.org/10.1016/j.atmosenv.2016.10.030>, 2017.
16
17 Bertschi, I., Yokelson, R. J., Ward, D. E., Babbitt, R. E., Susott, R. A., Goode, J. G. and Hao, W. M.: Trace gas and particle
18 emissions from fires in large diameter and belowground biomass fuels, *J. Geophys. Res. Atmos.*, 108, 8472,
19 doi:10.1029/2002JD002100, 2003.
20
21 Boby, L. A., Schuur, E. A. G., Mack, M. C., Verbyla, D. and Johnstone, J. F.: Quantifying fire severity, carbon, and nitrogen
22 emissions in Alaska’s boreal forest, *Ecol. Appl.*, 20, 1633-1647, doi:10.1890/08-2295.1, 2010.
23
24 Burling, I. R., Yokelson, R. J., Akagi, S. K., Urbanski, S. P., Wold, C. E., Griffith, D. W. T., Johnson, T. J., Reardon, J., Weise,
25 D. R.: Airborne and ground-based measurements of the trace gases and particles emitted by prescribed fires in the United States,
26 *Atmos. Chem. and Phys.*, 11, 12197-12216, 2011.
27
28 Chang, R.Y.W., Miller, C.E., Dinardo, S.J., Karion, A., Sweeney, C., Daube, B.C., Henderson, J.M., Mountain, M.E.,
29 Eluskiewicz, J., Miller, J.B. and Bruhwiler, L.M.: Methane emissions from Alaska in 2012 from CARVE airborne
30 observations, *Proc. Natl. Acad. Sci.*, 111, 47, 16694–16699, <https://doi.org/10.1073/pnas.1412953111>, 2014.
31
32 Christian, T. J., Yokelson, R. J., Carvalho Jr., J. A., Griffith, D. W. T., Alvarado, E. C., Santos, J. C., Neto, T. G. S., Veras, C. A.
33 G., and Hao, W. M.: The tropical forest and fire emissions experiment: Trace gases emitted by smoldering logs and dung on
34 deforestation and pasture fires in Brazil, *J. Geophys. Res.*, 112, D18308, doi:10.1029/2006JD008147, 2007.
35
36 Cofer, W. R., Levine, J. S., Sebacher, D. I., Winstead, E. L., Riggan, P. J., Stocks, B. J., Brass, J. A., Ambrosia, V. G., and Boston,
37 P. J.: Trace gas emissions from chaparral and boreal forest fires, *J. Geophys. Res.*, 94, 2255– 2259,
38 doi:[10.1029/JD094iD02p02255](https://doi.org/10.1029/JD094iD02p02255), 1989.
39

1 Cofer, W. R., Levine, J. S., Winstead, E. L. and Stocks, B. J.: Gaseous emissions from Canadian boreal forest fires, *Atmos. Environ.*
2 Part A, Gen. Top., 24, 1653-1659, doi:10.1016/0960-1686(90)90499-D, 1990.
3

4 Cofer, W. R., Winstead, E. L., Stocks, B. J., Goldammer, J. G. and Cahoon, D. R.: Crown fire emissions of CO₂, CO, H₂, CH₄, and
5 TNMHC from a dense jack pine boreal forest fire, *Geophys. Res. Lett.*, 25, 3919-3922, doi:10.1029/1998GL900042, 1998.
6

7 Collier, S., Zhou, S., Onasch, T., Jaffe, D., Kleinman, L., Sedlacek, A., Briggs, N., Hee, J., Fortner, E., Shilling, J., Worsnop, D.,
8 Yokelson, R., Parworth, C., Ge, X., Xu, J., Butterfield, Z., Chand, D., Dubey, M., Pekour, M., Springston, S., and Zhang, Q.:
9 Regional influence of aerosol emissions from wildfires driven by combustion efficiency: Insights from the BBOP campaign,
10 *Environ. Sci. Technol.*, 50, 8613-8622, doi:10.1021/acs.est.6b01617, 2016.
11

12 Cooper, D. J., Gallant, A. L., Binnian, E. F., Omernik, J. M. and Shasby, M. B.: Ecoregions of Alaska, *Arct. Alp. Res.*, 29, 494,
13 doi:10.2307/1551999, 2006.
14

15 de Groot, W. J., Flannigan, M. D. and Cantin, A. S.: Climate change impacts on future boreal fire regimes, *For. Ecol. Manage.*,
16 294, 35-44, doi:10.1016/j.foreco.2012.09.027, 2013.
17

18 Dieleman, C.M., Rogers, B.M., Potter, S., Veraverbeke, S., Johnstone, J.F., Laflamme, J., Solvik, K., Walker, X.J., Mack, M.C.
19 and Turetsky, M.R.: Wildfire combustion and carbon stocks in the southern Canadian boreal forest: Implications for a warming
20 world. *Glob. Change Bio.*, <https://doi.org/10.1111/gcb.15158>, 2020.
21

22 Duck, T. J., Firanski, B. J., Millet, D. B., Goldstein, A. H., Allan, J., Holzinger, R., Worsnop, D. R., White, A. B., Stohl, A.,
23 Dickinson, C. S. and van Donkelaar, A.: Transport of forest fire emissions from Alaska and the Yukon Territory to Nova Scotia
24 during summer 2004, *J. Geophys. Res. Atmos.*, 112, D10S44, doi:10.1029/2006JD007716, 2007.
25

26 Flannigan, M., Campbell, I., Wotton, M., Carcaillet, C., Richard, P. and Bergeron, Y.: Future fire in Canada's boreal forest:
27 paleoecology results and general circulation model - regional climate model simulations, *Can. J. For. Res.*, 31, 854-864,
28 doi:10.1139/x01-010, 2001.
29

30 Forster, C., Wandering, U., Wotawa, G., James, P., Mattis, I., Althausen, D., Simmonds, P., O'Doherty, S., Jennings, S. G.,
31 Kleefeld, C., Schneider, J., Trickl, T., Kreipl, S., Jäger, H. and Stohl, A.: Transport of boreal forest fire emissions from Canada to
32 Europe, *J. Geophys. Res. Atmos.*, 106, 22887-22906, doi:10.1029/2001JD900115, 2001.
33

34 French, N. H. F., Kasischke, E. S. and Williams, D. G.: Variability in the emission of carbon-based trace gases from wildfire in
35 the Alaskan boreal forest, *J. Geophys. Res. Atmos.*, 107, 8151, doi:10.1029/2001jd000480, 2002.
36

37 French, N. H. F., Goovaerts, P. and Kasischke, E. S.: Uncertainty in estimating carbon emissions from boreal forest fires, *J.*
38 *Geophys. Res. Atmos.*, 109, D14S08, doi:10.1029/2003JD003635, 2004.
39

1 Friedli, H. R., Radke, L. F., Prescott, R., Hobbs, P. V., and Sinha, P.: Mercury emissions from the August 2001 wildfires in
2 Washington State and an agricultural waste fire in Oregon and atmospheric mercury budget estimates, *Global Biogeochem.*
3 *Cycles*, 17, 1039, doi:[10.1029/2002GB001972](https://doi.org/10.1029/2002GB001972), 2003.

4

5 Fromm, M., Alfred, J., Hoppel, K., Hornstein, J., Bevilacqua, R., Shettle, E., R. Servranckx, R., Li, Z., and Stocks, B.:
6 Observations of boreal forest fire smoke in the stratosphere by POAM III, SAGE II, and lidar in 1998, *Geophys. Res.*
7 *Lett.*, 27, 1407–1410, <https://doi.org/10.1029/1999GL011200>, 2000.

8

9 Gillett, N. P., Weaver, A. J., Zwiers, F. W. and Flannigan, M. D.: Detecting the effect of climate change on Canadian forest fires,
10 *Geophys. Res. Lett.*, 31, L18211, doi:10.1029/2004GL020876, 2004.

11

12 Goldammer, J.G. and Furyaev, V.V. (Eds.): *Fires in Ecosystems of Boreal Eurasia*, Springer, Dordrecht 1996.

13

14 Goode, J. G., Yokelson, R. J., Ward, D. E., Susott, R. A., Babbitt, R. E., Davies, M. A. and Hao, W. M.: Measurements of excess
15 O₃, CO₂, CO, CH₄, C₂H₄, C₂H₂, HCN, NO, NH₃, HCOOH, CH₃COOH, HCHO, and CH₃OH in 1997 Alaskan biomass burning
16 plumes by airborne Fourier transform infrared spectroscopy (AFTIR), *J. Geophys. Res. Atmos.*, 105, 22147-22166,
17 doi:10.1029/2000jd900287, 2000.

18

19 Guerette, E., Paton-Walsh, C., Desservettaz, M., Smith, T. E. L., Volkova, L., Weston, C. J. and Meyer, C. P.: Emissions of trace
20 gases from Australian temperate forest fires: Emission factors and dependence on modified combustion efficiency, *Atmos. Chem.*
21 *and Phys.*, 18, 3717-3735, <http://dx.doi.org/10.5194/acp-18-3717-2018>, 2018.

22

23 Harden, J. W., Trumbore, S. E., Stocks, B. J., Hirsch, A., Gower, S. T., O’Neill, K. P. and Kasischke, E. S.: The role of fire in the
24 boreal carbon budget, *Glob. Chang. Biol.*, 6, 174-184, doi:10.1046/j.1365-2486.2000.06019.x, 2000.

25

26 Hayasaka, H., H. L. Tanaka, H. L., and Bieniek, P. A.: Synoptic-scale fire weather conditions in Alaska, *Polar Science*, 10, 217-
226, doi:[10.1016/j.polar.2016.05.001](https://doi.org/10.1016/j.polar.2016.05.001), 2016.

27

28 Henderson, J. M., Eluszkiewicz, J., Mountain, M. E., Nehr Korn, T., Chang, R. Y. W., Karion, A., Miller, J. B., Sweeney, C.,
29 Steiner, N., Wofsy, S. C. and Miller, C. E.: Atmospheric transport simulations in support of the Carbon in Arctic Reservoirs
30 Vulnerability Experiment (CARVE), *Atmos. Chem. Phys.*, 15, 4093-4116, doi:10.5194/acp-15-4093-2015, 2015.

31

32 Jaffe, D., Bertschi, I., Jaeglé, L., Novelli, P., Reid, J. S., Tanimoto, H., Vingarzan, R. and Westphal, D. L.: Long-range transport
of Siberian biomass burning emissions and impact on surface ozone in western North America, *Geophys. Res. Lett.*, 31, L16106,
doi:10.1029/2004GL020093, 2004.

33

34 Johnson, E. A.: *Fire and vegetation dynamics: studies from the North American boreal forest*, Cambridge University Press, New
York, USA, 1996.

35

36 Johnstone, J.F., Rupp, T.S., Olson, M. and Verbyla, D.: Modeling impacts of fire severity on successional trajectories and future
fire behavior in Alaskan boreal forests, *Landscape Ecology*, 26, 487-500, 2011.

1 Kahn, R. A., Chen, Y., Nelson, D. L., Leung, F. Y., Li, Q. B., Diner, D. J., and Logan, J. A.: Wildfire smoke injection heights:
2 Two perspectives from space, *Geophys. Res. Lett.*, 35, L04809, doi:10.1029/2007GL032165, 2008.

3 Kaiser, J., Suttie, M., Flemming, J., Morcrette, J. J., Boucher, O., and Schultz, M.: Global real-time fire emission estimates based
4 on space-borne fire radiative power observations, in: AIP conference proceedings, 1100, 645, 2009.

5 Karion, A., Sweeney, C., B Miller, J., E Andrews, A., Commane, R., Dinardo, S., M Henderson, J., Lindaas, J., C Lin, J., A Luus,
6 K., Newberger, T., Tans, P., C Wofsy, S., Wolter, S. and E Miller, C.: Investigating Alaskan methane and carbon dioxide fluxes
7 using measurements from the CARVE tower, *Atmos. Chem. Phys.*, 16, 5383-5398, doi:10.5194/acp-16-5383-2016, 2016.

8

9 Kasischke, E. S., Hyer, E. J., Novelli, P. C., Bruhwiler, L. P., French, N. H. F., Sukhinin, A. I., Hewson, J. H. and Stocks, B. J.:
10 Influences of boreal fire emissions on Northern Hemisphere atmospheric carbon and carbon monoxide, *Global Biogeochem.*
11 *Cycles*, 19, GB1012, doi:10.1029/2004GB002300, 2005.

12

13 Kasischke, E. S. and Turetsky, M. R.: Recent changes in the fire regime across the North American boreal region - Spatial and
14 temporal patterns of burning across Canada and Alaska, *Geophys. Res. Lett.*, 33, L09703, doi:10.1029/2006GL025677, 2006.

15

16 Konovalov, I. B., Beekmann, M., Kuznetsova, I. N., Yurova, A. and Zvyagintsev, A. M.: Atmospheric impacts of the 2010 Russian
17 wildfires: Integrating modelling and measurements of an extreme air pollution episode in the Moscow region, *Atmos. Chem. Phys.*,
18 11, 10031-10056, doi:10.5194/acp-11-10031-2011, 2011.

19

20 Korovin, G.N.: Analysis of distribution of forest fires in Russia, in: *Fires in Ecosystems of Boreal Eurasia*, edited by: Goldammer,
21 J.G. and Furyaev, V.V., Kluwer Academic, Netherlands, 112-128, 1996.

22

23 Laursen, K. K., Hobbs, P. V., Radke, L. F., and Rasmussen, R. A.: Some trace gas emissions from North American biomass fires
24 with an assessment of regional and global fluxes from biomass burning, *J. Geophys. Res.*, 97, 20687–20701,
25 doi:[10.1029/92JD02168](https://doi.org/10.1029/92JD02168), 1992.

26

27 Lin, J., Gerbig, C., Wofsy, S., Chow, V., Gottlieb, E., Daube, B., and Matross, D.: Designing Lagrangian experiments to measure
28 regional scale trace gas fluxes, *J. Geophys. Res.*, 112, D13312, doi:[10.1029/2006JD008077](https://doi.org/10.1029/2006JD008077), 2007.

29

30 McMeeking, G.R., Kreidenweis, S.M., Baker, S., Carrico, C.M., Chow, J.C., Collett Jr, J.L., Hao, W.M., Holden, A.S., Kirchstetter,
31 T.W., Malm, W.C. and Moosmüller, H.: Emissions of trace gases and aerosols during the open combustion of biomass in the
32 laboratory, *J. Geophys. Res.*, 114, D19210, doi:[10.1029/2009JD011836](https://doi.org/10.1029/2009JD011836), 2009.

33

34 McRae, D.J., Conard, S.G., Ivanova, G.A., Sukhinin, A.I., Baker, S.P., Samsonov, Y.N., Blake, T.W., Ivanov, V.A., Ivanov, A.V.,
35 Churkina, T.V. and Hao, W.: Variability of fire behavior, fire effects, and emissions in Scotch pine forests of central Siberia, *Mitig.*
36 *Adapt. Strateg. Global Change*, 11, 45-74, 2006.

37

38 Nance, J. D., Hobbs, P. V., Radke, L. F. and Ward, D. E.: Airborne measurements of gases and particles from an Alaskan wildfire,
39 *J. Geophys. Res. Atmos.*, 98, 14873-14882, doi:10.1029/93jd01196, 1993.

1
2 O'Shea, S. J., Allen, G., Gallagher, M. W., Bauguitte, S. J.-B., Illingworth, S. M., Le Breton, M., Muller, J. B. A., Percival, C. J.,
3 Archibald, A. T., Oram, D. E., Parrington, M., Palmer, P. I., and Lewis, A. C.: Airborne observations of trace gases over boreal
4 Canada during BORTAS: campaign climatology, air mass analysis and enhancement ratios, *Atmos. Chem. Phys.*, 13, 12451–
5 12467, <https://doi.org/10.5194/acp-13-12451-2013>, 2013.
6
7 Partain, J. L., Alden, S., Bhatt, U. S., Bieniek, P. A., Brettschneider, B. R., Lader, R. T., Olsson, P. Q., Rupp, T. S., Strader, H.,
8 Thoman, R. L., Walsh, J. E., York, A. D. and Ziel, R. H.: An assessment of the role of anthropogenic climate change in the Alaska
9 fire season of 2015, *Bull. Am. Meteorol. Soc.*, 97, S14-S18, doi:10.1175/BAMS-D-16-0149.1, 2016.
10
11 Peterson, D. A., Campbell, J. R., Hyer, E. J., Fromm, M. D., Kablick, G. P., Cossuth, J. H. and DeLand, M. T.: Wildfire-driven
12 thunderstorms cause a volcano-like stratospheric injection of smoke, *Clim. Atmos. Sci.*, 1, 30, doi:10.1038/s41612-018-0039-3,
13 2018.
14
15 Radke, L. F., Hegg, D. A., Hobbs, P. V., Nance, J. D., Lyons, J. H., Laursen, K. K., Weiss, R. E., Riggan, P. J. and Ward, D. E.:
16 Particulate and Trace Gas Emissions from Large Biomass Fires in North America, *Glob. Biomass Burn. Atmos. Clim. Biosph.*
17 *Implic.*, 209-224, 1991.
18
19 Rapalee, G., Trumbore, S.E., Davidson, E.A., Harden, J.W. and Veldhuis, H.: Soil carbon stocks and their rates of accumulation
20 and loss in a boreal forest landscape, *Global Biogeochem. Cycles*, 12, 4, <https://doi.org/10.1029/98GB02336>,
21 687-70, 1998.
22
23 Rodell, M., Beaudoin, H., and NASA/GSFC/HSL GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2. 0,
24 edited, Goddard Earth Sciences Data and Information Services Center (GES DISC) Greenbelt, MD, 2015.
25
26 Rogers, B. M., Soja, A. J., Goulden, M. L. and Randerson, J. T.: Influence of tree species on continental differences in boreal fires
27 and climate feedbacks, *Nat. Geosci.*, 8, 228-234, doi:10.1038/ngeo2352, 2015.
28
29 Ryan, K. C.: Dynamic interactions between forest structure and fire behavior in boreal ecosystems, in *Silva Fennica.*, 36, 13-39,
30 2002.
31
32 Santín, C., Doerr, S.H., Preston, C.M. and González-Rodríguez, G.: Pyrogenic organic matter production from wildfires: a missing
33 sink in the global carbon cycle. *Glob. Change Bio.*, 21, 1621-1633, <https://doi.org/10.1111/gcb.12800>, 2015.
34
35 Sedano, F. and Randerson, J. T.: Multi-scale influence of vapor pressure deficit on fire ignition and spread in boreal forest
36 ecosystems, *Biogeosciences*, 11, 3739–3755, <https://doi.org/10.5194/bg-11-3739-2014>, 2014.
37
38 Selimovic, V., Yokelson, R. J., McMeeking, G. R., and Coefield, S.: In situ measurements of trace gases, PM, and aerosol optical
39 properties during the 2017 NW US wildfire smoke event, *Atmos. Chem. Phys.*, 19, 3905–3926, [https://doi.org/10.5194/acp-19-](https://doi.org/10.5194/acp-19-3905-2019)
40 3905-2019, 2019.

1
2 Selimovic, V., Yokelson, R.J., McMeeking, G.R. and Coefield, S.: Aerosol mass and optical properties, smoke influence on O₃,
3 and high NO₃ production rates in a western US city impacted by wildfires, *Earth and Space Sci. Open Arc.*, 52,
4 <https://doi.org/10.1002/essoar.10501529.1>, 2020
5
6 Simpson, I. J., Akagi, S. K., Barletta, B., Blake, N. J., Choi, Y., Diskin, G. S., Fried, A., Fuelberg, H. E., Meinardi, S., Rowland,
7 F. S., Vay, S. A., Weinheimer, A. J., Wennberg, P. O., Wiebring, P., Wisthaler, A., Yang, M., Yokelson, R. J. and Blake, D. R.:
8 Boreal forest fire emissions in fresh Canadian smoke plumes: C1-C10 volatile organic compounds (VOCs), CO₂, CO, NO₂, NO,
9 HCN and CH₃CN, *Atmos. Chem. Phys.*, 11, 6445-6463, doi:10.5194/acp-11-6445-2011, 2011.
10
11 Skamarock, W.C., Klemp, J.B., Dudhia, J., Gill, D.O., Barker, D.M., Wang, W. and Powers, J.G.: A description of the Advanced
12 WRF Version 2, Tech. Note NCAR/TN-468+ STR, Natl. Cent. for Atmos. Res., Boulder, Colo., 2005.
13
14 Smith, T., Paton-Walsh, C., Meyer, C.P., Cook, G., Maier, S.W., Russell-Smith, J., Wooster, M. and Yates, C.P.: New emission
15 factors for Australian vegetation fires measured using open-path Fourier transform infrared spectroscopy-Part 2: Australian tropical
16 savanna fires., *Atmos. Chem. Phys.*, 14, 11335-11352, 2014.
17
18 Stocks, B., Wotton, B., Flannigan, M., Fosberg, M., Cahoon, D., and Goldammer, J.: Boreal forest fire regimes and climate change,
19 in: *Remote sensing and climate modeling: Synergies and limitations*, edited by: Beniston, M., and Verstraete, M. M., Springer,
20 Dordrecht, 233-246, <https://doi.org/10.1007/0-306-48149-9>, 2001.
21
22 Stockwell, C.E., Yokelson, R.J., Kreidenweis, S.M., Robinson, A.L., DeMott, P.J., Sullivan, R.C., Reardon, J., Ryan, K.C.,
23 Griffith, D.W. and Stevens, L.: Trace gas emissions from combustion of peat, crop residue, domestic biofuels, grasses, and other
24 fuels: configuration and Fourier transform infrared (FTIR) component of the fourth Fire Lab at Missoula Experiment (FLAME-
25 4). *Atmos. Chem. Phys.*, 14, 18, 9727-9754, <https://doi.org/10.5194/acp-14-9727-2014>, 2014.
26
27 Strand, T., Gullett, B., Urbanski, S., O'Neill, S., Potter, B., Aurell, J., Holder, A., Larkin, N., Moore, M. and Rorig, M.: Grassland
28 and forest understorey biomass emissions from prescribed fires in the south-eastern United States–RxCADRE 2012. *Int. J. Wildl.*
29 *Fire*, 25, 102-113, <https://doi.org/10.1071/WF14166>, 2016.
30
31 Susott, R. A., Ward, D. E., Babbitt, R. E., Latham, D. J.: The measurement of trace emissions and combustion characteristics for
32 a mass fire, *Global Biomass Burning: Atmospheric, Climatic, and Biosphere Implications*, MIT Press, Cambridge, MA, 245-257,
33 1991.
34
35 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern
36 circumpolar permafrost region, *Global Biogeochem. Cycles*, 23, GB2023, doi:[10.1029/2008GB003327](https://doi.org/10.1029/2008GB003327), 2009.
37
38 Thoning, K., Kitzis, D., and Crotwell, A.: Atmospheric carbon dioxide dry air mole fractions from quasi-continuous measurements
39 at Barrow, Alaska, Mauna Loa, Hawaii, American Samoa, and South Pole, 1973–2006, Version: 2007-10-01, National Oceanic
40 and Atmospheric Administration, Path: <ftp://ftp.cmdl.noaa.gov/ccg/co2/in-situ>, 2007.

1 Turquetly, S., Logan, J. A., Jacob, D. J., Hudman, R. C., Leung, F. Y., Heald, C. L., Yantosca, R. M., Wu, S., Emmons, L. K.,
2 Edwards, D. P. and Sachse, G. W.: Inventory of boreal fire emissions for North America in 2004: Importance of peat burning and
3 pyroconvective injection, *J. Geophys. Res. Atmos.*, 112, D12S03, doi:10.1029/2006JD007281, 2007.
4
5 Urbanski, S. P., Hao, W. M. and Baker, S.: Chapter 4 Chemical Composition of Wildland Fire Emissions, *Dev. Environ. Sci.*, 8,
6 79-107, doi:10.1016/S1474-8177(08)00004-1, 2008.
7
8 Urbanski, S.: Wildland fire emissions, carbon, and climate: Emission factors, *For. Ecol. Manage.*, 317, 51-60,
9 doi:10.1016/j.foreco.2013.05.045, 2014.
10
11 Val Martin, M., Logan, J. A., Kahn, R. A., Leung, F.-Y., Nelson, D. L., and Diner, D. J.: Smoke injection heights from fires in
12 North America: analysis of 5 years of satellite observations, *Atmos. Chem. Phys.*, 10, 1491–1510, [https://doi.org/10.5194/acp-10-](https://doi.org/10.5194/acp-10-1491-2010)
13 [1491-2010](https://doi.org/10.5194/acp-10-1491-2010), 2010.
14
15 Van Leeuwen, T. T. and van der Werf, G. R.: Spatial and temporal variability in the ratio of trace gases emitted from biomass
16 burning, *Atmos. Chem. Phys.*, 11, 3611-3629, doi:10.5194/acp-11-3611-2011, 2011.
17
18 Vasileva, A., Moiseenko, K., Skorokhod, A., Belikov, I., Kopeikin, V. and Lavrova, O.: Emission ratios of trace gases and particles
19 for Siberian forest fires on the basis of mobile ground observations. *Atmos. Chem. Phys.*, 17, 12303-12325,
20 <https://doi.org/10.5194/acp-17-12303-2017>, 2017.
21
22 Veraverbeke, S., Rogers, B. M. and Randerson, J. T.: Daily burned area and carbon emissions from boreal fires in Alaska,
23 *Biogeosciences*, 12, 3579-3601, doi:10.5194/bg-12-3579-2015, 2015.
24
25 Veraverbeke, S., Rogers, B. M., Goulden, M. L., Jandt, R. R., Miller, C. E., Wiggins, E. B. and Randerson, J. T.: Lightning as a
26 major driver of recent large fire years in North American boreal forests, *Nat. Clim. Chang.*, 7, 529-534, doi:10.1038/nclimate3329,
27 2017.
28
29 Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Johnstone, J. F., Rogers, B. M., Solvik, K., Turetsky, M. R. and Mack,
30 M. C.: Soil organic layer combustion in boreal black spruce and jack pine stands of the Northwest Territories, Canada, *Int. J. Wildl.*
31 *Fire*, 27, 125-134, doi:10.1071/wf17095, 2018.
32
33 Ward, D. E. and Radke, L. F.: Emissions Measurements from Vegetation Fires: A Comparative Evaluation of Methods and Results,
34 *Fire in the Environment: The Ecological Atmospheric and Climatic Importance of Vegetation Fires*, Dahlem Workshop Reports:
35 *Environmental Sciences Research Report 13*, John Wiley & Sons, Chischester, England. 53-76, 1993.
36
37 Wiggins, E. B., Veraverbeke, S., Henderson, J. M., Karion, A., Miller, J. B., Lindaas, J., Commane, R., Sweeney, C., Luus, K. A.,
38 Tosca, M. G., Dinardo, S. J., Wofsy, S., Miller, C. E. and Randerson, J. T.: The influence of daily meteorology on boreal fire
39 emissions and regional trace gas variability, *J. Geophys. Res. Biogeosciences*, 121, 2793-2810, doi:10.1002/2016JG003434, 2016.
40

1 Wotawa, G., Novelli, P. C., Trainer, M. and Granier, C.: Inter-annual variability of summertime CO concentrations in the Northern
2 Hemisphere explained by boreal forest fires in North America and Russia, *Geophys. Res. Lett.*, 28, 4575-4578,
3 doi:10.1029/2001GL013686, 2001.

4

5 Yokelson, R.J., Griffith, D.W. and Ward, D.E.: Open-path Fourier transform infrared studies of large-scale laboratory biomass
6 fires, *J. Geophys. Res. Atmos.*, 101, D15, <http://doi.org/10.1029/96JD01800>, 21067-21080, 1996.

7

8 Yokelson, R. J., Susott, R., Ward, D. E., Reardon, J., and Griffith, D. W. T.: Emissions from smoldering combustion of biomass
9 measured by open-path Fourier transform infrared spectroscopy, *J. Geophys. Res.*, 10, D15, 18865–18877,
10 doi:[10.1029/97JD00852](http://doi.org/10.1029/97JD00852), 1997.

11

12 Yokelson, R. J., Goode, J. G., Ward, D. E., Susott, R. A., Babbitt, R. E., Wade, D. D., Bertschi, I., Griffith, D. W. T., and Hao, W.
13 M.: Emissions of formaldehyde, acetic acid, methanol, and other trace gases from biomass fires in North Carolina measured by
14 airborne Fourier transform infrared spectroscopy, *J. Geophys. Res.*, 104, D23, 30109– 30125, doi:[10.1029/1999JD900817](http://doi.org/10.1029/1999JD900817), 1999.

15

16 Yokelson, R.J., Karl, T., Artaxo, P., Blake, D.R., Christian, T.J., Griffith, D.W., Guenther, A. and Hao, W.M.: The Tropical Forest
17 and fire emissions experiment: overview and airborne fire emission factor measurements. *Atmos. Chem. and Phys.*, 7, 19, 5175-
18 5196, <https://doi.org/10.5194/acp-7-5175-2007>, 2007.


19

20 Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., De Gouw, J., Akagi, S. K., Urbanski, S. P., Veres, P.,
21 Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, T. J., Hosseini, S., Miller, J. W., Cocker, D. R., Jung, H.
22 and Weise, D. R.: Coupling field and laboratory measurements to estimate the emission factors of identified and unidentified trace
23 gases for prescribed fires, *Atmos. Chem. Phys.*, 13, 89-116, doi:10.5194/acp-13-89-2013, 2013.

24


25 Young, A. M., Higuera, P. E., Duffy, P. A. and Hu, F. S.: Climatic thresholds shape northern high-latitude fire regimes and imply
26 vulnerability to future climate change, *Ecography*, 40, 606-617, doi:10.1111/ecog.02205, 2017.

1 **Figures**




2


3 **Figure 1.** The location of wildfires in Alaska during 2015, with color representing the day of burning estimated from the Alaska
4 Fire Emissions Database (AKFED). The black star denotes the location of CRV tower.




1
2 **Figure 2.** A) Observations of CO₂ mole fraction at the CRV tower in 2012 (black line) along with model estimates of the CO₂
3 background (green line) at CRV using the approach described in the main text. Very few fires occurred during 2012, and as a
4 consequence most of the CO₂ variability in the observations and in the model are associated with terrestrial net ecosystem exchange.
5 B) In 2015 wildfires in interior Alaska contributed significantly to CO₂ variability at the CRV tower, causing positive anomalies
6 in the observations shown in black, particularly between days 170 and 190. The modeled background for 2015 is shown in red.
7 The CO₂ mole fraction observations and model estimates have a 1 hour temporal resolution.




1
 2 **Figure 3.** Distribution of transit times representing the difference between the time when CO was emitted by a fire and the time the
 3 CO anomaly reached the CRV tower, as estimated by multiplying footprints from PWRP-STILT with fire emissions from AKFED.
 4 Only times when fire emission ratios were calculated were used in the analysis.




1
 2 **Figure 4.** Trace gas observations at the CRV tower during the summer of 2015 for A) CO, B) CH₄, and C) CO₂ mole fractions. The
 3 trace gas observations are shown at a 30 s temporal resolution. Daily active fire detections derived from the MODIS sensors on
 4 Terra and Aqua satellites (MCD14ML C6) are shown in panel D.



1
 2 **Figure 5.** CRV tower observations of A) CO, B) CH₄, and C) CO₂ are shown along with intervals used to calculate emission ratios
 3 (shown in color). The primary combustion process is noted with blue for smoldering, purple for mixed, and red for flaming. The
 4 trace gas observations are shown at a 30 s temporal resolution.




1
 2 **Figure 6.** Relationship between CH₄ emission factor and modified combustion efficiency (MCE). The strong linear relationship
 3 indicates that periods with more smoldering combustion (and a lower MCE) produce significantly higher levels of CH₄ emissions.
 4 The relationship was defined by a slope of -46.77 ± 4.70 , a Y intercept of 46.37 ± 4.13 g kg⁻¹ dry biomass burned, an r^2 of 0.54,
 5 and a significance value of $p < 0.01$.




1
 2 **Figure 7.** Examples of intervals used to calculate emission ratios. The flaming combustion example is from DOY 177, the mixed
 3 example is from DOY 177, and smoldering example is from DOY 175. These intervals correspond to events 27, 25, and 19 in
 4 Table 2. The trace gas measurements are shown at a 30 s temporal resolution.

1




2

3 **Figure 8.** CRV observations of CO (black) compared with the modeled CO anomaly from fires (red) derived from the PWRP-
4 STILT atmospheric model driven by AKFED fire emissions. The trace gas observations and model predictions are shown at a 1
5 hour temporal resolution.



1
 2 **Figure 9.** Individual fire contributions to the total fire season integral of CO anomalies measured at the CRV tower, as determined
 3 by convolving footprints from PWRP-STILT with fire emissions from AKFED. The location of CRV is shown as a black circle.
 4 Fire perimeters are shown in black.



1
 2 **Figure 10.** A) Top 5 individual fire contributions to the CO anomalies simulated at the CRV tower. The black line shows original
 3 PWRP-STILT \times AKFED model, pink denotes contributions from the Tozitna fire, green from the Kobe fire, blue from the Blair
 4 fire, gold from the Aggie Creek fire, and purple from the Spicer Creek fire. B) The total CO anomaly from the 34 fires that
 5 contributed to at least 1% of the modeled CO anomaly at CRV tower (red) compared to sum of all fire shown in black derived
 6 from original PWRP-STILT \times AKFED simulation (black).

1 **Tables**

2 **Table 1.** Comparison of CO emission ratio and modified combustion efficiency (MCE) from previous studies that sampled
 3 emissions from boreal forest fires. The studies are organized according to wildfire domain (North America or Siberia), management
 4 practice (wildfire or management fire), and sampling approach (aircraft, laboratory, or surface tower). Siberian studies are indicated
 5 as aircraft studies (A), surface based studies (S), or a combination of the two (A & S). The CO emission ratio column has units of
 6 ppmv ppmv⁻¹ and uses CO₂ as the reference gas. MCE was calculated as 1/(1 + ΔCO/ΔCO₂) when not directly reported in the
 7 study. The weighted mean of emission ratios and MCE for all previous studies is shown in the row labeled fire-weighted mean,
 8 with each study weighted according to the number of fires sampled.

Study	ΔCO/ΔCO ₂ Emission Ratio	Modified Combustion Efficiency	Number of fires sampled
<u>North American wildfires sampled by aircraft</u>			
Cofer et al., 1989	0.069 ± 0.004	0.935 ± 0.004	1
Cofer et al., 1998	0.140 ± 0.012	0.878 ± 0.009	1
Friedli et al., 2003	0.100 ± 0.020	0.909 ± 0.017	1
Goode et al., 2000	0.085 ± 0.008	0.922 ± 0.007	4
Laursen et al., 1992	0.050 ± 0.007	0.953 ± 0.006	1
Nance et al., 1993	0.078 ± 0.012	0.928 ± 0.011	1
O'Shea et al., 2013	0.150 ± 0.024	0.871 ± 0.012	4
Radke et al., 1991	0.116 ± 0.087	0.896 ± 0.075	1
Simpson et al., 2011	0.110 ± 0.070	0.901 ± 0.061	5
Fire-weighted mean	0.102 ± 0.033	0.908 ± 0.027	19
<u>North American management fires sampled by aircraft</u>			
Cofer et al., 1990	0.086 ± 0.008	0.921 ± 0.007	2
Cofer et al., 1998	0.095 ± 0.016	0.913 ± 0.013	7
Radke et al., 1991	0.047 ± 0.032	0.956 ± 0.030	4
Susott et al., 1991	0.060 ± 0.061	0.943 ± 0.058	1
Fire-weighted mean	0.077 ± 0.022	0.929 ± 0.020	14
<u>North American fuels sampled in the laboratory</u>			
Yokelson et al., 1997 ^a	0.208 ± 0.039	0.827 ± 0.083	-
Yokelson et al., 1997 ^b	0.231 ± 0.068	0.813 ± 0.167	-
Yokelson et al., 1997 ^c	0.162	0.860	-
Bertschi et al., 2003 ^d	0.151 ± 0.040	0.870 ± 0.030	-
Burling et al., 2010 ^e	0.209	0.827	-
McMeeking et al., 2009 ^e	0.153 ± 0.032	0.867 ± 0.074	-
McMeeking et al., 2009 ^f	0.045 ± 0.005	0.957 ± 0.012	-
McMeeking et al., 2009 ^c	0.030	0.971	-
Stockwell et al., 2014^f	0.043 ± 0.004	0.959 ± 0.008	-
Stockwell et al., 2014^g	0.245 ± 0.005	0.803 ± 0.009	-
Mean	0.143 ± 0.028	0.875 ± 0.053	
<u>Siberian wildfires – sampled by aircraft or surface tower</u>			
Cofer et al., 1998 (A)	0.224 ± 0.036	0.817 ± 0.025	1
McRae et al., 2006 (A & S)	0.249 ± 0.064	0.800 ± 0.043	6
Vasileva et al., 2017 (S)	0.126 ± 0.007	0.888 ± 0.005	2
Fire-weighted mean	0.219 ± 0.048	0.822 ± 0.033	9
<u>North American wildfires sampled by surface tower</u>			
Wiggins et al., 2016	0.128 ± 0.023	0.887 ± 0.018	3
This study	0.142 ± 0.051	0.878 ± 0.039	34
Fire-weighted mean	0.141 ± 0.049	0.879 ± 0.027	37

-
- 1 ^a Moss (Alaska), ^b Peat (Alaska), ^c White Spruce (Alaska), ^d Duff Jack Pine/Black Spruce (Canada), ^e Duff Black Spruce (Alaska),
 - 2 ^f Black Spruce (Alaska), and ^g Peat (Canada).
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24

1 **Table 2.** Intervals with elevated trace gas mole fractions at CRV associated with fire emissions. Columns show the number of 30
2 s measurements used to calculate emission factors for each interval (N), the time of the interval (units of day of year (DOY)),
3 emission ratios (ppmv ppmv⁻¹), emission factor (g kg⁻¹ dry biomass burned), and modified combustion efficiency (MCE). The
4 primary combustion process is denoted as flaming, mixed, or smoldering using thresholds on MCE defined in the text.

Interval number	N	Time of Event (DOY)	CO Emission Ratio	CO Emission Factor	CH ₄ Emission Ratio	CH ₄ Emission Factor	MCE	Combustion Phase
1	82	173.27 - 173.30	0.161 ± 0.004	144 ± 4	0.012 ± 0.0003	6.1 ± 0.2	0.861 ± 0.004	Mixed
2	95	173.32 - 173.35	0.151 ± 0.004	136 ± 4	0.011 ± 0.0002	5.8 ± 0.2	0.869 ± 0.004	Mixed
3	95	173.36 - 173.39	0.141 ± 0.003	128 ± 3	0.010 ± 0.0002	5.5 ± 0.1	0.877 ± 0.003	Mixed
4	83	173.40 - 173.43	0.149 ± 0.008	135 ± 8	0.011 ± 0.0005	5.5 ± 0.3	0.870 ± 0.008	Mixed
5	95	173.45 - 173.48	0.130 ± 0.006	120 ± 6	0.009 ± 0.0004	5.0 ± 0.3	0.885 ± 0.006	Mixed
6	95	173.84 - 173.87	0.136 ± 0.008	124 ± 8	0.014 ± 0.0009	7.3 ± 0.5	0.880 ± 0.008	Mixed
7	85	174.27 - 174.30	0.170 ± 0.008	152 ± 8	0.008 ± 0.0003	4.3 ± 0.2	0.855 ± 0.008	Mixed
8	95	175.15 - 175.18	0.08 ± <0.001	78 ± 0.3	0.004 ± <1e4	2.3 ± <0.1	0.926 ± 1e4	Flaming
9	95	175.19 - 175.22	0.143 ± 0.007	131 ± 7	0.008 ± 0.0004	4.2 ± 0.3	0.875 ± 0.007	Mixed
10	58	175.23 - 175.25	0.091 ± 0.002	87 ± 2	0.005 ± 0.0002	2.5 ± 0.1	0.916 ± 0.002	Mixed
11	88	175.27 - 175.30	0.091 ± 0.001	87 ± 1	0.005 ± 0.0001	2.9 ± <0.1	0.917 ± 0.001	Mixed
12	95	175.32 - 175.35	0.153 ± 0.003	138 ± 4	0.009 ± 0.0002	4.5 ± 0.1	0.867 ± 0.003	Mixed
13	89	175.40 - 175.44	0.187 ± 0.012	164 ± 12	0.013 ± 0.0008	6.4 ± 0.5	0.842 ± 0.012	Smoldering
14	95	175.66 - 175.70	0.060 ± 0.003	59 ± 3	0.005 ± 0.0002	2.6 ± 0.1	0.943 ± 0.003	Flaming
15	55	175.75 - 175.77	0.129 ± 0.001	119 ± 1	0.009 ± 0.0001	4.5 ± 0.1	0.886 ± 0.001	Mixed
16	35	175.77 - 175.79	0.237 ± 0.015	198 ± 15	0.017 ± 0.0010	8.1 ± 0.6	0.809 ± 0.014	Smoldering
17	95	175.80 - 175.83	0.147 ± 0.002	133 ± 2	0.011 ± 0.0001	5.5 ± 0.1	0.872 ± 0.002	Mixed
18	95	175.88 - 175.91	0.155 ± 0.003	139 ± 3	0.009 ± 0.0002	4.9 ± 0.2	0.866 ± 0.003	Mixed
19	95	175.92 - 175.96	0.198 ± 0.004	172 ± 4	0.012 ± 0.0001	6.1 ± 0.1	0.835 ± 0.004	Smoldering
20	80	175.98 - 176.00	0.193 ± 0.003	169 ± 3	0.011 ± 0.0001	5.4 ± 0.1	0.838 ± 0.003	Smoldering
21	95	176.06 - 176.09	0.119 ± 0.007	111 ± 7	0.008 ± 0.0004	4.4 ± 0.3	0.893 ± 0.007	Mixed
22	85	177.06 - 177.09	0.108 ± 0.001	102 ± 1	0.010 ± 0.0001	5.3 ± <0.1	0.902 ± 0.001	Mixed
23	75	177.11 - 177.14	0.122 ± 0.002	113 ± 2	0.011 ± 0.0001	5.6 ± 0.1	0.892 ± 0.002	Mixed
24	95	177.15 - 177.18	0.129 ± 0.001	119 ± 1	0.010 ± 0.0001	5.5 ± 0.1	0.886 ± 0.001	Mixed
25	95	177.19 - 177.22	0.102 ± 0.002	96 ± 2	0.008 ± 0.0002	4.4 ± 0.1	0.908 ± 0.002	Mixed
26	58	177.23 - 177.25	0.148 ± 0.011	134 ± 12	0.012 ± 0.0009	6.0 ± 0.5	0.871 ± 0.011	Mixed
27	94	177.27 - 177.31	0.060 ± 0.002	59 ± 2	0.004 ± 0.0001	2.3 ± 0.1	0.944 ± 0.002	Flaming
28	95	177.80 - 177.83	0.094 ± 0.002	89 ± 2	0.008 ± 0.0001	4.1 ± 0.1	0.914 ± 0.002	Mixed
29	95	177.88 - 177.91	0.120 ± 0.006	111 ± 6	0.020 ± 0.0012	10.7 ± 0.7	0.893 ± 0.006	Mixed
30	93	177.92 - 177.96	0.164 ± 0.006	146 ± 7	0.018 ± 0.0007	8.9 ± 0.4	0.859 ± 0.006	Mixed
31	95	184.23 - 184.26	0.232 ± 0.014	196 ± 15	0.013 ± 0.0007	6.5 ± 0.4	0.811 ± 0.014	Smoldering
32	80	186.49 - 186.52	0.025 ± 0.002	25 ± 2	0.002 ± 0.0001	1.2 ± 0.1	0.976 ± 0.002	Flaming
33	64	188.07 - 188.09	0.188 ± 0.012	165 ± 13	0.013 ± 0.0008	6.6 ± 0.5	0.842 ± 0.012	Smoldering
34	95	188.10 - 188.13	0.106 ± 0.002	100 ± 2	0.008 ± 0.0002	4.5 ± 0.1	0.904 ± 0.002	Mixed
35	54	188.14 - 188.16	0.109 ± 0.001	102 ± 1	0.008 ± 0.0001	4.3 ± <0.1	0.902 ± 0.001	Mixed
36	64	188.20 - 188.22	0.104 ± 0.004	99 ± 4	0.008 ± 0.0003	4.2 ± 0.2	0.906 ± 0.004	Mixed
37	52	188.23 - 188.25	0.080 ± 0.007	77 ± 7	0.006 ± 0.0004	3.2 ± 0.2	0.926 ± 0.007	Flaming
38	95	188.40 - 188.44	0.194 ± 0.003	169 ± 3	0.012 ± 0.0002	6.1 ± 0.1	0.837 ± 0.003	Smoldering
39	95	188.45 - 188.48	0.131 ± 0.004	120 ± 4	0.013 ± 0.0006	6.9 ± 0.3	0.884 ± 0.004	Mixed
40	36	188.53 - 188.55	0.146 ± 0.002	132 ± 2	0.012 ± 0.0001	6.0 ± 0.1	0.873 ± 0.002	Mixed
41	54	188.59 - 188.61	0.163 ± 0.002	145 ± 2	0.012 ± 0.0001	6.3 ± 0.1	0.860 ± 0.002	Mixed
42	95	188.62 - 188.65	0.179 ± 0.002	158 ± 2	0.014 ± 0.0002	6.9 ± 0.1	0.848 ± 0.002	Smoldering
43	74	188.66 - 188.69	0.214 ± 0.011	183 ± 12	0.015 ± 0.0008	7.4 ± 0.5	0.824 ± 0.011	Smoldering
44	95	188.71 - 188.74	0.138 ± 0.005	126 ± 5	0.010 ± 0.0004	5.1 ± 0.2	0.879 ± 0.005	Mixed
45	95	188.75 - 188.78	0.055 ± 0.003	54 ± 3	0.006 ± 0.0002	3.3 ± 0.1	0.948 ± 0.003	Flaming
46	95	188.79 - 188.83	0.272 ± 0.009	223 ± 10	0.012 ± 0.0005	5.7 ± 0.3	0.786 ± 0.009	Smoldering
47	52	188.84 - 188.85	0.120 ± 0.002	112 ± 2	0.009 ± 0.0001	4.9 ± 0.1	0.893 ± 0.002	Mixed
48	39	188.86 - 188.87	0.091 ± 0.002	87 ± 2	0.007 ± 0.0001	4.0 ± 0.1	0.916 ± 0.002	Mixed
49	59	189.03 - 189.05	0.154 ± 0.012	139 ± 13	0.010 ± 0.0008	5.3 ± 0.5	0.867 ± 0.012	Mixed
50	95	189.27 - 189.31	0.149 ± 0.008	135 ± 9	0.011 ± 0.0005	5.6 ± 0.3	0.871 ± 0.008	Mixed
51	30	189.34 - 189.35	0.090 ± 0.009	86 ± 9	0.006 ± 0.0005	3.2 ± 0.3	0.917 ± 0.009	Mixed
52	89	189.49 - 189.52	0.165 ± 0.009	147 ± 9	0.012 ± 0.0007	6.1 ± 0.4	0.858 ± 0.009	Mixed
53	48	195.10 - 195.12	0.212 ± 0.019	181 ± 20	0.016 ± 0.0014	8.0 ± 0.9	0.825 ± 0.018	Smoldering
54	37	195.12 - 195.13	0.262 ± 0.027	215 ± 28	0.020 ± 0.0020	9.5 ± 1.2	0.792 ± 0.026	Smoldering
55	95	195.14 - 195.17	0.140 ± 0.007	128 ± 8	0.010 ± 0.0006	5.5 ± 0.3	0.877 ± 0.007	Mixed
Mean			0.142 ± 0.051	127 ± 40	0.010 ± 0.0038	5.3 ± 1.8	0.878 ± 0.039	

5

1 **Table 3.** All fires that contributed to at least 1% of the total CO anomaly observed at CRV, in order from largest CO contribution
2 to smallest CO contribution. The distance column represents the distance of the center of the fire perimeter to CRV tower.
3 Contribution is the percent contribution to the total integral of fire CO at CRV for the entire 2015 fire season. Some fires were
4 grouped together if they were inside the same 0.5° grid cell during model coupling. For those cases, individual fire contribution
5 to the CO anomaly observed at CRV tower was weighted based on fire size.

	Fire Name	Distance (km)	Contribution (%)	Total Hectares	Fuel Type	Ignition Source
1	Tozitna	229	10.74	31652	Black Spruce	Lightning
2	Kobe	119	7.20	3444	Black Spruce	Lightning
3	Blair	82	6.31	15217	Black Spruce	Lightning
4	Aggie Creek	41	5.63	12829	Black Spruce	Lightning
5	Spicer Creek	195	5.30	39761	Black Spruce	Lightning
6	Blind River	252	3.87	24608	Black Spruce	Lightning
7	Holtnakatna	404	3.44	90308	Mixed	Lightning
8	Blazo	514	3.39	49106	Black Spruce	Lightning
9	Big Creek 2	351	3.23	126637	Black Spruce	Lightning
10	Chitanana River	241	3.12	17483	Black Spruce	Lightning
11	Sea	309	3.06	172	Black Spruce	Human
12	Sushgitit Hills	276	2.92	111712	Black Spruce	Lightning
13	Big Mud River 1	254	2.72	42076	Black Spruce	Lightning
14	Lost River	347	2.58	21088	Black Spruce	Lightning
15	Munsatli 2	302	2.36	40682	Black Spruce	Lightning
16	FWA Small Arms Complex	19	2.31	740	Black Spruce	Prescribed
17	Tobatokh	280	2.24	21868	Black Spruce	Lightning
18	Trail Creek	363	2.24	11939	Black Spruce	Lightning
19	Lloyd	201	2.22	26818	Black Spruce	Lightning
20	Isahultila	342	2.17	60445	Black Spruce	Lightning
21	Nulato	499	2.17	449	Black Spruce	Lightning
22	Three Day	472	2.17	39378	Black Spruce	Lightning
23	Hay Slough	188	1.90	37007	Black Spruce	Lightning
24	Rock	316	1.83	3714	Other	Lightning
25	Sulukna	329	1.77	6760	Black Spruce	Lightning
26	Titna	273	1.77	12415	Black Spruce	Lightning
27	Quinn Creek	657	1.49	2002	Other	Lightning
28	Harper Bend	188	1.45	17555	Black Spruce	Lightning
29	Hard Luck	328	1.43	5230	Black Spruce	Lightning
30	Fox Creek	369	1.42	2346	Black Spruce	Lightning
31	Bering Creek	280	1.36	45654	Black Spruce	Lightning
32	Eden Creek	324	1.16	18614	Black Spruce	Lightning
33	Falco	390	1.10	1817	Mixed	Lightning
34	Jackson	202	1.00	2969	Black Spruce	Lightning

6