

Interactive comment on “Evaluating Trends and Seasonality in Modeled PM_{2.5} Concentrations Using Empirical Mode Decomposition” by Huiying Luo et al.

Huiying Luo et al.

marina.astitha@uconn.edu

Received and published: 31 July 2020

Reply to interactive comments on “Evaluating Trends and Seasonality in Modeled PM_{2.5} Concentrations Using Empirical Mode Decomposition”

Anonymous Referee #3

This manuscript presented an evaluation of the WRF-CAMQ model simulated temporal trends through a detailed comparison with observation using improved CEEMDAN method. The comparison was based on measurements of PM_{2.5} and its key components, i.e., sulphate, nitrate, ammonium, chloride, organic carbon, and elemental

Printer-friendly version

Discussion paper



carbon, made at three ground monitoring stations in US from t 2002 to 2008. It is clearly demonstrated that the improved CEEMDAN approach can decompose the observed and simulated temporal trends, which allows to extract more information from the comparisons of individual temporal modes. For example, the authors concluded that the model can better simulate the rate of change of the multi-year trend than the absolute magnitude. At the same time, model can generally reproduce the amplitudes of the sub-seasonal and annual variations for PM2.5, sulphate, and OC. This study revealed that it appears there is a temporal phased shift between the observed and model simulated PM2.5, OC, and EC as large as a half year. It is further suggested that this phase shift indicted “a need for proper temporal allocation of emissions”. In general, the manuscript is well organized.

We thank the reviewer for the positive assessment of our manuscript and for providing constructive feedback to help improve the quality of the manuscript. We have addressed all questions and suggestions in our response as well as in the text or figures, as necessary. Please see detailed responses below and the marked-up version of the revised manuscript.

This reviewer believes that this is an important work which can potentially help identifying model deficiencies. However, there several concerns needed to be addressed:

1) The authors correctly stated that EMD is a widely used methodology in various field. At the same time, this reviewer would like to suggest that the authors should consider adding some brief high-level descriptions of the method. This will improve the manuscript's readability, especially for those who are not familiar with EMD methods. It is also important to clearly state the criteria how the modes are determined and separated. The statement in line 134-135, “to achieve best mode separation”, leaves much room for interpretation. The discussion on determination of t_p and t_m is interesting and thorough. It does, however, leave an impression that the evaluation of t_p and t_m is somewhat uncertain and is not completely deterministic. This reviewer would like to suggest adding additional text to discuss if the determination of t_p and t_m is suffi-

[Printer-friendly version](#)[Discussion paper](#)

ciently accurate or useful for model assessment to identify issues in the processes at the similar time scale as decomposed t_p and/or t_m . This will strengthen the manuscript to demonstrate the usefulness of the improved CEEMDAN approach in model assessments.

The decomposition process and parameters controlling the decomposition have been added in Section 3.1 as suggested. The “best mode separation” is also further explained following the reviewer’s suggestion.

CEEMDAN is a technique that is particularly suitable to analyze non-linear and non-stationary time series data. The decomposed time series of speciated and total PM_{2.5} reveal the agreement/disagreement between observations and model simulations at various intrinsic temporal scales without any predetermined assumptions on the data. Both t_p and t_m represent approximate estimates of the characteristic scale of an IMF, where non-linear and non-stationary processes with close temporal scales could exist. For t_p (from the revised text): “The peak estimates can be biased if more than one high-power frequency is located closely within one IMF. Thus, the power spectrum and t_p is only used as a fast screening tool to determine if a desired decomposition is accomplished.” For t_m : “As the frequency decreases, the mean period estimates become less accurate because of the limited time span compared with the length of the cycle and should be carefully interpreted.” We have added the following test in Section 4.1: “Since each IMF represents a non-stationary process, the mean period t_m is only an estimate of its characteristic scale. Evaluation of t_m might not necessarily be able to identify issues with corresponding model simulations, and it does not indicate any information on the magnitude or the phase of the time series, which is more important and will be further discussed in Sections 4.3 to 4.4.”

2) Section 2 (starting from line 74) provided a good discussion on how the observation data sets are selected. It is equally important to discuss the temporal resolution of model in terms of the driving factors, e.g., emissions. This will give readers a sense if one should expect if the model should reproduce observations at certain temporal

[Printer-friendly version](#)[Discussion paper](#)

scale. For example, if the emissions are given in yearly average, one would consider the impact of the lack emission temporal variability on the comparison of seasonal and/or sub-seasonal trends.

We added the following text in Section 2: “Annual emissions for the CMAQ simulations were estimated using the methodology described in Xing et al. (2013). Briefly, the National Emissions Inventory (NEI) for 1990, 1995, 1996, 1999, 2001, 2002 and 2005 and a number of sector-specific long-term databases containing information about trends in activity data and emission controls were used to create county-level annual emissions for a total of 49 emission sectors. Prior to being used as input to the CMAQ simulations, these annual emissions were then temporally and spatially allocated to provide hourly emissions based on monthly, weekly, and diurnal temporal cross-reference and profile data from the 2005 NEI modeling platform. These profile data vary by emissions source and sometimes by state and county and are generally based on surveys and extrapolation of activity data which can be subject to uncertainty. Exceptions to the use of 2005 NEI platform temporal profile data for temporal allocation were emissions from electric generating units (EGU) which directly used measured hourly emissions after 1995 and wildfire emissions that used climatological monthly, weekly, and diurnal profiles for temporal allocation.”

The large discrepancy in the magnitude of some long-term trend component seen in Fig. 6 is likely pointing to the systematic bias in the annual emission estimations as discussed in Xing et al. (2013): “. . .since this study mainly focused on trends rather than the absolute value in each individual year, some sectors (e.g., industrial processes) and sub-sectors (types of combustion and stoves) may not have been well considered or examined.” The intra-annual emission allocation could possibly impact the model performance at the seasonal and sub-seasonal scales. This discussion of the impact of emissions on the long-term trend has been added in Section 4.2.

3) This reviewer believes that the concluding remark of “indicating the need for proper allocation of emissions” is an important conclusion. However, it was not adequately

[Printer-friendly version](#)[Discussion paper](#)

justified. There are many controlling factors and processes. The authors should have provided more discussions to illustrate how they narrowed to emissions as the likely factor. It should also be pointed out that SOA is typically a large component of OC. Changes in emissions to affect OC will likely have implications on O₃.

We would like to clarify that our illustrative application of the new methodology to PM_{2.5} time series at three specific sites does not allow us to conclude that errors in the temporal allocation of PM emissions are the controlling factors for disagreements between observed and modeled annual cycle. While we believe that they do play a role as discussed below, we also know that the CMAQ version used for these simulations has underestimated the formation of SOA, which would also affect the modeled annual cycle of OC (e.g. Appel et al., 2017; Murphy et al., 2017; Xu et al., 2018). Because of the underestimation of SOA, OC in the simulations analyzed here has an overestimated relative contribution of primary OC which, in turn, makes its temporal variations analyzed by CEEMDAN sensitive to the temporal allocation of primary PM and specifically primary OC emissions. The full statement partially quoted by the reviewer points to both factors “indicating the need for proper allocation of emissions and an updated treatment of organic aerosols compared to the earlier model version used in this set of model simulations”. Without running a new set of decadal simulations with a newer version of the model and/or modified temporal allocation of emissions, we are unable to determine the relative importance of these factors at the sites examined. However, if such simulations were to be performed in the future, the CEEMDAN methodology can help demonstrate the benefits of updated emissions allocations and/or the SOA process representation.

4) The authors presented detailed trend analysis on PM_{2.5} and its components. It is also scientifically interesting to understand the relative contribution of each component and their contribution to the identified temporal variability, which are useful to gain insights into controlling factors. This reviewer would like to suggest the authors to consider addition of the trend analysis on the relative contribution of sulphate, nitrate,

[Printer-friendly version](#)[Discussion paper](#)

ammonium, organic carbon, and elemental carbon to PM_{2.5}. More specific to the manuscript, it would be much easier to interpret the results shown in Table 1, 2, and 3 if the relative contribution of each component is known.

Yes, it would be useful to explicitly show the importance of each component in driving the trend of total PM_{2.5} in both observations and model simulations. The time series of the concentration share of each component (e.g. OC/Total PM_{2.5} %) is added in Fig. S6 in the supplement. However, the decomposition of the concentration share is not included since there is not much change in the percentage share in its trend component (few percentages at most in very limited cases) and the ratio does not necessarily have strong seasonality because of the phase difference in specific component and total PM_{2.5}. Thus, including the trend component of time variant share of the ratio would only complicate the interpretation of the results. Instead, we have added a new Table 1 (see below) to show the overall concentration share of each component for both observations and model simulations to reflect the relative importance of different species.

5) In general, model evaluation is designed to improve model. It is difficult to relate the comparison results presented in this manuscript to specific model deficiencies in description of the chemical/physical processes and/or issues in model data sets, meteorological field and/or emission data. As sulphate, OC, nitrate are controlled by very different chemical processes, this reviewer would like to encourage the authors to further explore the difference in the comparison results for these species, which may reveal additional insights into the process-level model deficiencies.

We thank the reviewer for recognizing the potential of the proposed methodology in helping identify problems in the specific processes and/or model input. However, without running a new set of decadal simulations with a newer version of the model and/or modified temporal allocation of emissions, we cannot determine specific model deficiencies and/or issues in the model input data sets.

[Printer-friendly version](#)[Discussion paper](#)

Specific Comments: 1) Figure 3 is hard to read because of log-log scale. It may be better to change the x-axis to the IMF number and y-axis to the ratio between model and observation characteristic scales. A second y-axis can be added to show the absolute characteristic scales for each IMF.

We thank the reviewer for the suggestion. However, because of the large discrepancies in the scales of IMFs (few days to thousands of days), log scale has to be employed to show the scales for all IMFs. Given that the characteristic periods are not easy to read from the plot, we provided the average characteristic periods for sub-seasonal and seasonal IMFs in the text. Moreover, since “not all IMFs from observation are being simulated and vice versa”, a figure is needed for each site to show the characteristic scales (at least for the last few IMFs) separately for observations and model simulations. Thus, we have moved the inlet figures to Figure 3d-f for clarity and added the explanation in the caption. Adding a second y-axis and showing only observed characteristic scale would result in a very busy plot and we will not be able to achieve the second point above. Please find our revision to the figure in the manuscript and below.

2) Section 4.2. Figure 6 shows some variation in time-derivatives. At the same, this reviewer would like to argue that about half of cases shown in the figure can be well approximated by linear assumption. The authors should comment on this aspect.

Linear assumption is useful in many cases, and linear trends do provide a general idea of magnitude of the change as well as whether the linear trend is significant or not. EMD is particularly useful for analyzing meteorological and pollutant time series, which are non-linear and non-stationary. The decomposed trend components can provide the exact time span and magnitude of a decreasing/increasing change throughout time. If we take the trend component of observed OC at ATL as an example, the OC level is stable at around $4.5 \mu\text{g}/\text{m}^3$ in 2002 and 2003 and decreases at varying rates during 2004-2007.

Please also note the supplement to this comment:

<https://www.atmos-chem-phys-discuss.net/acp-2019-1079/acp-2019-1079-AC1-supplement.pdf>

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-1079>, 2020.

ACPD

Interactive
comment

Printer-friendly version

Discussion paper



Table 1. Concentration share (%) of different components in total $PM_{2.5}$. It is estimated by dividing the mean trend components of each species by that of total $PM_{2.5}$ for both OBS and CMAQ, multiplied by 100. The concentration share of the remainder species (*Rem*) is estimated by subtracting all the available species share from 100 to compensate for the small discrepancies caused by the rounding up process and uncertainty in the mode decomposition. "-" indicates the data is not available (same applies for all other tables).

		SO _x	NO _x	NH ₃	OC	EC	Cl	Rem
QURE	OBS	38	7	-	19	5	1	30
	CMAQ	19	15	-	14	5	1	47
RENO	OBS	7	13	5	46	11	-	20
	CMAQ	11	4	2	30	7	-	45
ATL	OBS	28	6	10	24	8	-	24
	CMAQ	22	10	8	17	9	-	33

Fig. 1.

Printer-friendly version

Discussion paper






Fig. 3. The characteristic scales (τ_{m}) resolved in the IMFs of observed and simulated total and speciated $\text{PM}_{2.5}$ for (a, d) QURE, (b, e) RENO and (c, f) ATL. In (a-c), IMF1 to the last pair of IMFs with increasing characteristic periods are shown from bottom left to top right. Mean periods of IMFs with scales longer than a year are being displayed in (d-f) with the same shapes as in the legend above to show the characteristic scales of all decomposed IMFs given that not all IMFs from observations are being simulated and vice versa. In the (d-f), species decomposed from observations are shown with smaller filled shapes, while species decomposed from simulations are represented by larger open shapes in slightly darker shades.

Fig. 2.

C10

Printer-friendly version

Discussion paper

