

1 **LIMS observations of lower stratospheric ozone in the southern polar springtime of 1978**

2
3 Ellis Remsberg¹, Virginia L. Harvey², Arlin Krueger³, Larry Gordley⁴,
4 John C. Gille⁵, and James M. Russell III⁶

5
6 ¹Science Directorate, NASA Langley Research Center, 21 Langley Blvd, Mail Stop 401B,
7 Hampton, VA 23681, USA

8 ²Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, 3665
9 Discovery Drive, Boulder, CO 80303, USA

10 ³Emeritus Senior Scientist, Code 614 Atmospheric Chemistry and Dynamics Laboratory, NASA
11 Goddard Space Flight Center, Greenbelt, MD 20771, USA

12 ⁴GATS, Inc., 11864 Canon Blvd., Suite 101, Newport News, VA 23606, USA

13 ⁵Senior Scientist Emeritus, National Center for Atmospheric Research, P.O. Box 3000, Boulder,
14 CO 80307-3000, USA

15 ⁶Endowed Professor and Co-Director, Center for Atmospheric Sciences, Hampton University,
16 Hampton, VA 23668, USA

17
18
19
20 Correspondence to: Ellis Remsberg (ellis.e.remsberg@nasa.gov)

21
22 For submission to Atmospheric Chemistry and Physics Journal

23 February~~December~~, 2020~~19~~

24 Abstract

25 The Nimbus 7 limb infrared monitor of the stratosphere (LIMS) instrument operated from
26 October 25, 1978, through May 28, 1979. This ~~note-paper~~ focuses on its Version (V6) data and
27 indications of ozone loss infor the lower stratosphere of the southern hemisphere, subpolar
28 region during the last week of October 1978. We provide profiles and maps that show V6 ozone
29 values of only 2 to 3 ppmv at 46 hPa within the edge of the polar vortex ~~at 46 hPa~~ near 60°S
30 from late October through mid-November 1978. There are also low values of V6 nitric acid (~3
31 to 6 ppbv) and nitrogen dioxide (<1 ppbv) at the same locations, indicating that conditions were
32 suitable for a chemical loss of Antarctic ozone some weeks earlier. These “first light” LIMS
33 observations provide the earliest, space-based view of conditions within the lower stratospheric
34 ozone layer of the southern polar region in springtime.

35

36 1 Introduction and historical context

37 The Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) provided the first daily image of
38 total ozone for the Southern Hemisphere (SH) on November 1, 1978. That image in Figure 1
39 shows an equatorward extension of the region of low ~~column~~ polar, total column ozone (TCO)
40 between 90°E and 135°E. Minimum TCO ~~polar-ozone~~ is of the order of 2750 Dobson units (DU)
41 at (75°S, 9270°E) on this day. As a comparison, Farman et al. (1985) reported ground-based
42 measurements of total ozone of about 225 DU on November 1 for 1980-1984 at Halley Bay
43 (76°S, 333°E) and of about 270 DU at Argentine Islands (65°S, 296°E) (see also TOMS total
44 ozone values of Table 2 in Stolarski et al. (1986)). We note, however, that those values are
45 higher than 220 DU, which is a threshold definition for “ozone hole” conditions (WMO, 2018).

46

47 There are very few observations of lower stratospheric ozone above Antarctica prior to
48 November 1978, especially for the months of September and October when the seasonal loss of
49 ozone is most significant (WMO, 2018). The historic Nimbus 7 Limb Infrared Monitor of the
50 Stratosphere (LIMS) experiment (Gille and Russell, 1984) provided data ~~for~~ middle
51 atmosphere temperature, geopotential height (GPH), ozone, water vapor (H₂O), nitric acid vapor
52 (HNO₃), and nitrogen dioxide (NO₂) from October 25, 1978, through May 28, 1979, for

53 scientific analysis and for comparisons with atmospheric models (e.g., Langematz et al., 2016).
54 Remsberg et al. (2007) provide a description of its Version 6 (V6) ozone profiles. The mapping
55 of the V6 profiles to the LIMS Level 3 product employs a sequential estimation algorithm with a
56 relaxation time of about 2.5 days for analyses of its zonal, 6-wavenumber Fourier coefficients at
57 each of 28 pressure levels of the middle atmosphere (Remsberg and Lingenfelter, 2010). We
58 then generated daily, polar stereographic plots of V6 ozone and HNO₃ on pressure surfaces
59 based on a gridding (2° latitude and 5.625° longitude) from those coefficients.

60

61 This note focuses on the character of the polar vortex and of the V6 ozone, HNO₃, and NO₂ in
62 that region of the lower stratosphere during the last week of October 1978. ~~T~~Although the LIMS
63 measurements extend to only 64°S, due to the orbital inclination of Nimbus 7 and to the viewing
64 geometry of the LIMS instrument (Gille and Russell, 1984). ~~W~~e will show that the profiles
65 and pressure surface maps indicate that there was a loss of SH polar ozone during some weeks
66 earlier in the springtime. Section 2 contains plots that show a loss of ozone inside the vortex in
67 late October. Section 3 reports on evidence for a denitrification of the air in the same region,
68 indicating that there was ~~likely~~ a chemical loss of ozone some weeks earlier. Section 3 also
69 presents time versus longitude or Hovmöller diagrams that reveal ~~the~~ good correspondence
70 forecontinuity of the low ozone and HNO₃ values within the vortex region well into November.
71 Section 4 summarizes the findings.

72

73 **2 Antarctic ozone from late October to early November 1978**

74 Figure 2 shows SH polar plots of V6 ozone mixing ratios at 46.4 hPa for October 26 and for
75 November 1, where the orbital measurements of LIMS extend only to 64°S. The plot at right
76 shows that there are minimum ozone values of about 2.6 ppmv near 120°E and 315°E at 60°S ~~on~~
77 November 1, which agrees reasonably with the locations of low total ozone from the TOMS
78 image of Fig. 1. Ozone is of order 3.5 to 4 ppmv at most other longitudes. Low ozone occurs
79 within the edges of the polar vortex, based on the concurrent GPH field from the operational
80 ECMWF Re-Analysis or ERA-40 products (Uppala et al., 2005). The bold contour in Fig. 2
81 denotes the edge of the vortex, in the manner of ~~as defined by~~ Harvey et al. (2002). We define

82 the vortex edge as the streamfunction contour coincident with maximum wind speed that also
83 encloses a region of rotation. Meek et al. (2017) showed that this definition of the vortex edge is
84 in good agreement with the PV-gradient based definition of Nash et al. (1996). We note that
85 daily plots of GPH are also available from LIMS V6, ~~but~~ However, they exhibit a discontinuous
86 anomaly at the 46-hPa level for the vortex region between October ~~29~~30 and 31, due to an
87 interpolation of National Meteorological Center (NMC) GPH~~likely from the early NOAA~~
88 Climate Prediction Center (CPC) analyses supplied to the Nimbus 7 Project and at 50 hPa-used
89 for the baseline pressure level of 50 hPa for the V6 GPH product (Remsberg et al., 2004). V6
90 geometric height and GPH profiles~~lots~~ above and below that level are the result of a hydrostatic
91 integration of the LIMS-retrieved temperature versus pressure profiles of T(p). ~~farther away~~
92 ~~from~~ Maps of V6 GPS farther away from the 50-hPa~~that~~ level are very similar to those from
93 ERA-40.

94

95 LIMS began its daily observations one week earlier than TOMS or on October 25, and the left
96 plot of Fig. 2 shows that the ozone for October 26 at 31°E is about half of that at 119°E on
97 November 1. The vortex on October 26 extends toward lower latitudes from about 60°S, 40°E.
98 Both the vortex and region of low ozone deform and undergo a clockwise rotation from October
99 26 onward, such that their low values extend equatorward at 120°E and at 315°E on November
100 1. Bodeker et al. (2002) reported that the edge of the vortex often extends to near 60°S during
101 October, and Stolarski et al (1986, their Fig. 1) and Hassler et al. (2011) reported on an
102 analogous clockwise rotation of the vortex during ~~mid-October of 1984~~.

103

104 **3 Findings of denitrification of the vortex air in late October**

105 The location of the vortex edge is helpful in deciding which V6 species profiles one ought to
106 examine with regard to any constraints from HNO₃ and NO₂ on the ozone chemistry. As an
107 example, Fig. 3 shows V6 Level 2 ozone profile segments from 11.4 to 88 hPa for two locations
108 on October 26, where ozone is now presented in units~~in terms~~ of partial pressure (in mPa) for a
109 better delineation of its relative changes in the subpolar lower stratosphere. Estimates of
110 accuracy for single V6 ozone profiles are 14%, 26%, and 34% for 10 hPa, 50 hPa, and 100 hPa,

111 respectively (see row (g) of Table 1 in Remsberg et al., 2007). The V6 ozone profile (black
112 solid) at 54.9°S, 119°E is just outside the October 26 vortex, as shown by the black dot in Fig. 2,
113 and its ozone values are nominal for subpolar latitudes. The largest contribution to total ozone
114 from that profile in Fig. 3 occurs at the 68-hPa level. A second V6 ozone profile (solid red) is
115 from 59.5°S, 31°E, and it is in a region of lower GPH as shown by the red dot in Fig. 2. Its
116 ozone decreases rapidly from ~8.0 mPa at the 53-hPa level to 2.6 mPa at the 88-hPa level,
117 indicating a significant loss of ozone in the lower stratosphere sometime prior to October 26.
118 Komhyr et al. (1988, their Fig. 10) and Gernandt (1987) show from ozonesonde measurements
119 that most of the observed losses of ozone for the mid-1980s occurred in the vortex in September
120 and early October. Therefore, we also include in Fig. 3 an ozonesonde profile (solid green) from
121 Syowa station (69°S, 40°E—the green dot in Fig. 2) for September 3, 1978, perhaps before there
122 were any pronounced losses of ozone. Its ozone profile values are intermediate of those for the
123 two V6 profiles of October 26.

124

125 Loss of ozone due to reactive chlorine chemistry proceeds effectively in the presence of air that
126 has undergone denitrification (Müller et al., 2008). Lambert et al. (2016) somewhat loosely set
127 an HNO₃ threshold ~~for it~~ of < 5 ppbv for indicating denitrification at 46 hPa, based on
128 Microwave Limb Sounder (MLS) data of 2008. Nitrous oxide is the source molecule for odd
129 nitrogen (mainly HNO₃) in the lower stratosphere, and its tropospheric values have grown by
130 only a small amount from 1975 (~296 ppbv) through 2008 (~322 ppbv) (WMO, 2018); the
131 HNO₃ threshold of 5 ppbv should also be representative for 1978. Thus, in Figure 3 we also
132 show the accompanying V6 profiles of HNO₃ and nighttime NO₂ for the same two locations on
133 October 26. HNO₃ and NO₂ at 31°E are a half (or 3 ppbv) and a third (or < 1 ppbv),
134 respectively, of those at 119°E below about the 31-hPa level. Thus, both species indicate that
135 there was a denitrification of the air in the vortex region and a likely loss of ozone due to reactive
136 chlorine chemistry in the presence of polar stratospheric clouds (PSCs) several weeks earlier
137 (Solomon, 1999; WMO, 2018). Although the V6 temperatures at 31°E on October 26 ~~were no~~
138 ~~colder than~~ 206 K (at 53 hPa), it is normal to find temperatures in the Antarctic vortex that are
139 below the chlorine activation threshold value of 1953 K and in the presence of PSCs during
140 September and early October (~~Drdla and Müller, 2012;~~ WMO, 2018).

141

142 Figure 4 shows the corresponding V6 plots of HNO₃ at 46 hPa in terms of its mixing ratios,
143 which have an estimated accuracy of ~9% (Remsberg et al., 2010, Table 10). There are very low
144 values of HNO₃ on October 26 poleward of 60°S and from 31°E to at least 90°E, indicating an
145 earlier conversion of HNO₃ from vapor to condensed phase and the sedimentation of larger
146 HNO₃ containing particles rather than an advection of low HNO₃ from lower latitudes. Low
147 HNO₃ mixing ratios-values are also present within the vortex region on November 1. Analogous
148 polar plots of the nighttime NO₂ fields are quite noisy (not shown) due to the large uncertainties
149 for tangent layer NO₂ in the lower stratosphere. Nevertheless, most of the odd nitrogen reservoir
150 at 46 hPa comes from HNO₃, not NO₂. Together, they indicate the extent of denitrification of the
151 air in the vortex region during late October 1978.

152

153 We show in Figs. 5 and 6 the details of the changing ozone and nitric acid from late October
154 through November. To investigate whether some of the low ozone and HNO₃ values might be
155 due to advection from lower latitudes, we show in Figure. 5 displays the time/longitude or
156 Hovmöller diagrams for both species at 60°S; thick black contours indicate the vortex edge and
157 dotted horizontal lines“+” the vortex interior. The occurrence of lowest species mixing ratios
158 shows clearly-values in the vortex region ~~shows clearly~~ in late October. Figure. 6 extends the
159 findings of Fig. 5 through the end of November, and there is an eastward progression of the
160 region of low values from late October to early November. Reduced mixing ratios of those
161 species occur inside~~The occurrence of low species values within~~ the vortex-region remains good
162 until about November 25, as expected for chemicals that are tracers of air motions in the lower
163 stratosphere. The vortex distorts and then exhibits a stationary wave-1 pattern from November 5
164 onward, where height is lowesthaving lowest heights near 0°E. Mixing of air across the vortex
165 edge appears slow for both ozone and HNO₃ during that time.

166

167 **4 Summary and concluding remarks**

168 We find low V6 ozone mixing ratios-values of order 2 to 3 ppmv at 60°S within the edge of the
169 polar vortex at 46 hPa during the last week of October and well into November 1978. There is

170 good agreement between the V6 ozone map at 46 hPa and the TOMS image of total ozone in the
171 region of the vortex on November 1. Low V6 HNO₃ ~~mixing ratios~~ ~~values~~ of order 3 to 6 ppbv at
172 the same locations indicate ~~denitrification~~ ~~that and~~ conditions ~~that~~ were suitable for a chemical
173 loss of Antarctic ozone some weeks earlier. ~~We note that~~ ~~The~~ equivalent effective stratospheric
174 chlorine (EESC) values used to predict conditions for the depletion of ozone in 1980 are about
175 twice those of 1950, while the 1980 values are only half those of 2000 (Newman et al., 2007). In
176 hindsight and based on the LIMS V6 dataset, we conclude that there was very likely some
177 halogen-catalyzed loss of ozone in the southern polar vortex in winter/spring of 1978. a
178 chemical process was likely responsible for springtime losses of ozone above Antarctica even in
179 the late 1970s. Yet, those ozone losses in the SH spring ~~of 1978~~ were not to the low level of a
180 true “ozone hole” (<220 DU total ozone). We also conclude that the LIMS V6 Level 2 profiles
181 and the daily-analyzed maps from their Level 3 zonal coefficients represent useful comparison
182 data for model simulations of the changes in Antarctic ozone in spring 1978.

183

184 **Data Availability**

185 The LIMS V6 data archive is at the NASA EARTHDATA site of EOSDIS and its website:

186 <https://search.earthdata.nasa.gov/search?q=LIMS>). Nimbus 7 TOMS ozone is at

187 https://disc.gsfc.nasa.gov/datacollection/TOMSN7L2_008.html. ECC ozone [sonde ozone](#)

188 profiles are available from the World Ozone and Ultraviolet Radiation Data Centre or WOUDC

189 at <https://woudc.org/data/explore.php>. ECMWF Re-Analysis (ERA-40) data are accessible

190 through <https://climatedataguide.ucar.edu/climate-data/era40>.

191

192 *Author Contributions.* ER and VLH wrote the manuscript and prepared the figures with input

193 from all the other co-authors. AK provided information about the TOMS ozone images. LG led

194 the development of the LIMS version 6 algorithms. JCG and JMR are the [Co-Principal](#)

195 Investigators of the LIMS experiment. They also commented on the new insight from the

196 findings about ozone and nitric acid of October 1978.

197

198 *Acknowledgements.* VLH acknowledges support from NASA LWS grant NNX14AH54G,

199 NASA HGI grant NNX17AB80G, and NASA HSR grant 80NSSC18K1046. EER carried out

200 his work while serving as a Distinguished Research Associate within the Science Directorate at

201 NASA Langley.

202

203 **References**

204 Bodeker, G. E., Struthers, H., and Connor, B. J.: Dynamical containment of Antarctic ozone
205 depletion, *Geophys. Res. Lett.*, 29, 2-1 to 2-4, <https://doi.org/10.1029/2001GL014206>, 2002.

206
207 ~~Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the~~
208 ~~polar stratosphere, *Ann. Geophys.*, 30, 1055-1073, <https://doi.org/10.5194/angeo-30-1-2012>,~~
209 ~~2012.~~

210
211 Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica
212 reveal seasonal ClOx/NOx interaction, *Nature*, 315, 207-210,
213 <https://www.nature.com/articles/315207a0.risNature>, 1985.

214
215 Gernandt, H.: The vertical ozone distribution above the GDR research base, Antarctica in 1985,
216 *Geophys. Res. Lett.*, 14, 84-66, 1987.

217
218 Gille, J. C. and Russell III, J. M.: The limb infrared monitor of the stratosphere: experiment
219 description, performance, and results, *J. Geophys. Res.*, 84, 5125-5140,
220 <https://doi.org/10.1029/JD089iD04p05125>, 1984.

221
222 Harvey, V.L., Pierce, R. B., Fairlie, T. D., and Hitchman, M. H.: A climatology of stratospheric
223 polar vortices and anticyclones, *J. Geophys. Res.*, 107(D20), 4442,
224 <https://doi.org/10.1029/2001JD001471>, 2002.

225
226 Hassler, B., Bodeker, G. E., Solomon, S., and Young, P. J.: Changes in the polar vortex: effects
227 on Antarctic total ozone observations at various stations, *Geophys. Res. Lett.*, 38, L01805,
228 [doi:10.1029/2010GL045542](https://doi.org/10.1029/2010GL045542), 2001.

229

230 Komhyr, W. D., Oltmans, S. J., and Grass, R. D.: Atmospheric ozone at South Pole, Antarctica,
231 in 1986, *J. Geophys. Res.*, 93, 5167-5184, <https://doi.org/10.1029/JD093iD05p05167>, 1988.

232

233 Lambert, A., Santee, M. L., and Livesey, N. J.: Interannual variations of early winter Antarctic
234 polar stratospheric cloud formation and nitric acid observed by CALIOP and MLS, *Atmos.*
235 *Chem. Phys.*, 16, 15219-15246, <https://doi.org/10.5194/acp-16-15219-2016>, 2016.

236

237 Langematz, U., Schmidt, F., Kunze, M., Bodeker, G. E., and Braesicke, P.: Antarctic ozone
238 depletion between 1960 and 1980 in observations and chemistry-climate model simulations,
239 *Atmos. Chem. Phys.*, 16, 15619-15627, <https://doi:10.5194/acp-16-15619-2016>, 2016.

240

241 Meek, C. E., Manson, A. H., and Drummond, J. R.: Comparison of Aura MLS stratospheric
242 chemical gradients with north polar vortex edges calculated by two methods, *Adv. Space Res.*,
243 60, 1898-1904, <http://dx.doi.org/10.1016/j.asr.2017-06.009>, 2017.

244

245 Müller, R., Grooß, J.-U., Lemmon, C., Heinze, D., Dameris, M., and Bodeker, G.: Simple
246 measures of ozone depletion in the polar stratosphere, *Atmos. Chem. Phys.*, 8, 251-264, 2008.

247

248 Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective determination
249 of the polar vortex using Ertel's potential vorticity, *J. Geophys. Res.*, 101, 9471-9478, 1996.

250

251 Newman, P. A., Daniel, J. S., Waugh, D. W., and Nash, E. R.: A new formulation of equivalent
252 effective stratospheric chlorine (EESC), *Atmos. Chem. Phys.*, 7, 4537-4552, 2007.

253

254 Remsberg, E., and Lingenfelter, G.: LIMS Version 6 Level 3 dataset, NASA-TM-2010-216690,
255 available at <http://www.sti.nasa.gov> (last access: 17 September 2019), 13 pp., 2010.

256

257 [Remsberg, E. E., Gordley, L. L., Marshall, B. T., Thompson, R. E., Burton, J., Bhatt, P., Harvey,](#)
258 [V. L., Lingenfelter, G., and Natarajan, M.: The Nimbus 7 LIMS version 6 radiance conditioning](#)
259 [and temperature retrieval methods and results, J. Quant. Spectros. Rad. Transf., 86, 395-424,](#)
260 [doi:10.1016/j.jqsrt.2003.12.007, 2004.](#)

261

262 Remsberg, E., Lingenfelter, G., Natarajan, M., Gordley, L., Marshall, B. T., and Thompson, E.:
263 On the quality of the Nimbus 7 LIMS version 6 ozone for studies of the middle atmosphere, J.
264 Quant. Spectros. Rad. Transf., 105, 492-518, doi:10.1016/j.jqsrt.2006.12.005, 2007.

265

266 Remsberg, E., Natarajan, M., Marshall, B. T., Gordley, L. L., Thompson, R. E., and
267 Lingenfelter, G. L.: Improvements in the profiles and distributions of nitric acid and nitrogen
268 dioxide with the LIMS version 6 dataset, Atmos. Chem. Phys., 10, 4741-4756,
269 <https://doi.org/10.5194/acp-10-4741-2010>, 2010.

270

271 Solomon, S.: Stratospheric ozone depletion: a review of concepts and history, Rev. Geophys., 37,
272 275-316, <https://doi.org/10.1029/1999RG900008>, 1999.

273

274 Stolarski, R. S., Krueger, A. J., Schoeberl, M. R., McPeters, R. D., Newman, P. A., and Alpert, J.
275 C.: Nimbus 7 satellite measurements of the springtime Antarctic ozone decrease, Nature, 322,
276 808-811, <https://doi.org/10.1038/322808a0>, 1986.

277


278 Uppala, S. M., KÅllberg, P. W., Simmons, A. J., Andrae, U., Da Costa Bechtold, V., Fiorino,
279 M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S.,
280 Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Van De

281 Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher,
282 M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M.,
283 Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W.,
284 Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The
285 ERA-40 reanalysis, *Q. J. Roy. Meteorol. Soc.*, 131, 2961–3012,
286 <https://doi.org/10.1256/qj.04.176>, 2005.

287


288 WMO (World Meteorological Organization), *Scientific Assessment of Ozone Depletion: 2018*,
289 Global Ozone Research and Monitoring Project — Report No. 58, 588 pp., Geneva, Switzerland,
290 2018.

291



200 250 300 350 400 450 500
 NIMBUS-7/TOMS COLUMN OZONE (matm-cm)

293



200 250 300 350 400 450 500
 NIMBUS-7/TOMS COLUMN OZONE (matm-cm)


294

295

296 Figure 1—Southern Hemisphere image of total column ozone (TCO) from TOMS for November
297 1, 1978. Longitude orientation is 0°E to the right and 90°E at the bottom; latitude circles
298 (dotted) have a spacing of 10 degrees. White areas indicate where there ~~are discrete~~ data
299 ~~voids~~ or no measurements. Ozone units of matm-cm are equivalent to Dobson units
300 (DU), where 1 DU is 2.687×10^{20} molecules-m⁻². Black contours are TCO at intervals of 20
301 matm-cm.

302

303




304

305 Figure 2—V6 ozone mixing ratios at 46.4 hPa for October 26 and November 1, 1978. Polar
306 plots extend from 30°S to the Pole and longitude is in °E with 0° at right. Bold contours denote
307 the vortex edge from ERA-40. The superposed, three colored dots correspond to the locations of
308 profiles on October 26 (black and red) and on September 3 (green) in Fig. 3.

309

310

311




312

313 Figure 3—V6 Level 2 species profiles for 59.5°S, 31°E (red) and 54.9°S, 119.4°E (black) on
314 October 26, 1978, and from an ozonesonde at Syowa (69°S, 40°E—green) on September 3,
315 1978. Ozone (solid) has units of millipascals (mPa), while HNO₃ (dashed) and NO₂ (dot-dashed)
316 have units of ppbv.

317


318



319

320 Figure 4—As in Fig. 2, but for V6 HNO₃.


321



323

324 Figure 5—Time/longitude or Hovmöller plots of LIMS ozone (left) and HNO₃ (right) for 60°S
 325 and 46 hPa. The ERA-40 vortex edge shows as thick black contours, and the vortex interior has
 326 horizontal dotted lines“+” symbols.

327



329

330 Figure 6—As in Fig. 5, but extended in time from October 25 to November 30, 1978.

331

Reply (in small CAPS) to comments of Referee #1

Anonymous Referee #1

GENERAL COMMENTS

First, it is great to see that 'old' data such as those obtained from LIMS are still be re-worked and used in analyses. Previous analyses, based on chemistry-climate models (e.g. Langematz et al...), have shown that ozone destruction through heterogeneous halogen-catalysed chemical reactions was occurring over Antarctica well before 1980 which is often (erroneously) considered as when Antarctic ozone depletion started and which is why many studies consider a 'return to 1980 values' as indicative of a recovery of the ozone layer over Antarctica from the effects of ozone depleting substances. This is one of the few papers to provide observational evidence of halogen-catalysed ozone depletion occurring over Antarctica prior to 1980. I think that this point should be made more strongly in the paper. It is made, almost in passing, around line 162 but I believe it should be highlighted in the abstract.

WE WRITE IN THE REVISED ABSTRACT AND THE SUMMARY THAT THERE WAS VERY LIKELY SOME HALOGEN-CATALYZED LOSS OF OZONE IN THE SOUTHERN POLAR VORTEX IN WINTER/SPRING OF 1978.

SPECIFIC COMMENTS

Figure 1: I wonder whether this is an older version of the TCO distribution from TOMS on 1 November 1978? It looks different to the file I have on my computer. Sure enough, when I go and download the raw data file from GSFC, here is the header:

Day: 305 Nov 1, 1978 Production V70 NIMBUS-7/TOMS OZONE Asc LECT: 11:49 AM Longitudes: 288 bins centered on 179.375 W to 179.375 E (1.25 degree steps) Latitudes : 180 bins centered on 89.5 S to 89.5 N (1.00 degree steps)

and here is the data line for 45.5°S:

```
3563573563493543543543513553613623603573543513503453453453463433453523523
3573553533543573643723803863943973953963973973983983883783813813833843833
383378387383377376 0360364364363351343336338341342344350350351 0 0 0 0
0 0 0 0353353354355358358359363358356361370372371366367365364359360359
357358364362361363365364364365366366 0 0 0 0 0 0 0368368368365363
3593583573533573513563533503473453483453463453443413473473533563543533473
3433363343333413443473533563673723793914114224184094054144043943763653543
3393363293233233173183213233213223183143102962973063103183233263353433493
3543583693733743773743713623553423393343243113063073103093163173253343373
3373303253223203153153183213243233213173143143153173163183213223223233233
328330333332332339340340 0 0 0359366367376377382383388392400407384360353
346346348347332329335338343343343351347 lat = -45.5
```

It shows some missing data but not as much as is apparent in your Figure 1 and certainly no data missing just west of the international date line. So why the discrepancy between the TCO field shown in your Figure 1 and the TOMS data stored on the GSFC server?

Line 40: You state that 'Minimum polar ozone is of the order of 250 Dobson units (DU) at (75°S, 270°E) on this day' When I go and look at the actual TOMS data file for that day at 74.5°S and 75.5°S at 270°E (which is 90°W; cell number 72 in TOMS-world) I see values in excess of 400 DU. These are quite different to the value of 250 DU that you are reporting here. What is the source of this difference? Perhaps you mean 90°E, but even

then the lowest ozone value is 270 DU.

YOUR DATA SOURCE APPEARS TO BE TOMS VERSION 7 (NOTE PRODUCTION V70 IN HEADER LINE), WHILE WE USE VERSION 8 DATA FOR FIGURE 1--

https://acdisc.gesdisc.eosdis.nasa.gov/data/nimbus7_toms_level3/tomsn7l3dtoz_008/1978/L3_OZONE_N7T_19781101.TXT

THE HEADER LINES ON ITS .TXT FILE FOR NOVEMBER 1 ARE

DAY: 305 Nov 1, 1978 NIMBUS-7/TOMS NRT OZONE GEN:04.119 V8 ALECT: 11:49
AM LONGITUDES: 288 BINS CENTERED ON 179.375 W TO 179.375 E (1.25 DEGREE STEPS)
LATITUDES : 180 BINS CENTERED ON 89.5 S TO 89.5 N (1.00 DEGREE STEPS).

THE DATA FOR -45.5 ARE BELOW AND ARE VERY SIMILAR TO, BUT NOT EXACTLY AS FROM YOUR SOURCE.

```
354354350344347352351349354362364361358357353352347346343340337339346347349
353353353356356363374376385395397398400400398397395387375379379378379380380
381384389383371369 0364369368365351338332336339340341346346347 0 0 0 0
0 0 0 0348348348348353354356359356353356363365362360358357359353352353
352352358357356359359359359358359360 0 0 0 0 0 0 0363363363360357
353354352351355355351351344341342342340340342341341346347352355352349346342
339335334332340342341351358368373382393411421415406402410400391374361351341
333331322316317314317322323321319317315305293295305311322326328335344353354
353354364374377376372368361354342339332324309303302307305310311321330330332
333330323318318312310313317321322317315314314316317317318322324322320322324
327328329330330337338339 0 0 0356359359373375376378381384395403379357345
340346334337314317327332339336344344342 LAT = -45.5
```

HOWEVER, WE DID NOTICE THAT THERE SHOULD BE ONLY THREE DISCRETE DATA VOID REGIONS AT THIS LATITUDE. WE CHECKED ON OUR CODE AND FOUND A BUG IN THE WAY WE HANDLED THE ZEROS. OUR REVISED FIGURE 1 (SEE BELOW) IS MORE IN LINE WITH THE TCO VALUES FROM YOUR V7 DATA SOURCE. WITH REGARD TO YOUR COMMENT ABOUT OUR SENTENCE AT LINE 40, WE AGREE THAT WE SHOULD HAVE SAID THAT THE MINIMUM OZONE IS 270 DU AND IS LOCATED NEAR 75°S, 90°E. THANK YOU FOR CHECKING ABOUT FIGURE 1 AND FOR ASKING WHETHER WE MADE AN ERROR.

Lines 82-83: When you say that 'We note that daily plots of GPH are also available from LIMS V6' do you mean that LIMS also retrieves temperature and pressure profiles from which GPH fields are calculated? or do you mean that GPH fields are provided (from some other source) along with the LIMS data? If the latter, can you please describe the source of those GPH fields. Thank you.

THE LIMS V6 GPH FIELDS AT 50 HPA ARE FROM NMC ANALYSES PROVIDED TO THE NIMBUS 7 PROJECT. THOSE NMC 50 HPA FIELDS REPRESENTED A BASE LEVEL FOR THE PREVIOUS LIMS V5 PRODUCT, AND THOSE SAME FIELDS PROVIDE A SCALING FOR BOTH THE GEOMETRIC AND GEOPOTENTIAL HEIGHTS FOR THE V6 PROFILES (SEE REMSBERG ET AL., JQSRT, 2004). HEIGHTS AND GPH PROFILES ABOVE AND BELOW THE 50-HPA LEVEL ARE OBTAINED HYDROSTATICALLY USING THE LIMS RETRIEVED T(P) PROFILES. WE ARE ADDING COMMENTS TO MAKE THIS CLEARER IN THE REVISED MANUSCRIPT.

Lines 93-95: You may also find the following paper relevant and of interest: Hassler, B., G. E. Bodeker, S. Solomon, and P. J. Young (2011), Changes in the polar vortex: Effects on Antarctic total ozone observations at various stations, Geophysical

THIS PAPER IS DEFINITELY PERTINENT AND WE REFER TO ITS FINDINGS IN SECTION 3 OF THE REVISED MANUSCRIPT. THANK YOU FOR BRINGING IT TO OUR ATTENTION.

Line 101: I would suggest replacing 'is now in terms of partial pressure' with 'is now presented in units of partial pressure'.

WE MADE THIS CHANGE.

Line 107: Are formally derived uncertainties on the LIMS measurements available? If so they should be quoted here.

REMSBERG ET AL. (2007, TABLE 1, ROW G) CONTAINS ESTIMATES OF ACCURACY FOR THE V6 OZONE PROFILES. THEY ARE OF ORDER 14% AT 10 hPa, 26% AT 50 hPa, AND 34% AT 100 hPa. WE MAKE NOTE OF THAT IN THE REVISED MANUSCRIPT.

Line 117: For clarity I suggest replacing 'for it' with 'for indication of denitrification'.

WE MADE THIS CHANGE.

Line 128: I always thought that the chlorine activation threshold on PSCs was 195K not 193K?

WE DECIDED TO CITE A TEMPERATURE THRESHOLD OF 195 K, BASED ON FIG. 4-1 OF WMO (2018), ALTHOUGH DRDLA AND MÜLLER (2012) INDICATE THAT IT IS EVEN LOWER THAN THAT IN THE PRESENCE OF STS PARTICLES.

Line 147: It was not clear to me what was meant by 'remains good'? Can you please describe that more specifically.

WE NOW WRITE "REDUCED CONCENTRATIONS OF THOSE SPECIES ARE PRESENT INSIDE THE VORTEX UNTIL ABOUT NOVEMBER 25"...

Line 149: By 'lowest heights' to you mean 'lowest amplitudes' of the wave-1?

WE CORRECT THE SENTENCE TO READ "AND WHERE THE HEIGHT IS LOWEST NEAR 0°E".

Line 272: Would it not be better here to state that $1\text{DU} = 2.687 \times 10^{20}$ molecules/m².

THE NUMERICAL VALUE AND UNITS FOR DU ARE NOW IN THE FIGURE CAPTION.

GRAMMAR AND TYPOGRAPHICAL ERRORS

Line 105: 59.5°S not just 59.5°

Line 109: Should this be 'ozonesonde' rather than just 'sonde'?


Line 127: Replace 'no colder than' with 'no lower than'. I was always taught 'the air is cold - temperatures are low'. There can be no more a cold temperature than a heavy temperature.

Line 144: Replace 'lowest species values' with 'lowest species concentrations' and again on line 147.

Line 170: Replace 'ECC ozone' with 'ECC ozonesonde ozone'.

WE MADE ALL THE ABOVE CHANGES.

Day: 305 Nov 1, 1978



Revised Figure 1

Interactive comment on “LIMS observations of lower stratospheric ozone in the southern polar springtime of 1978” by Ellis Remsberg et al.

Anonymous Referee #2

Received and published: 21 January 2020

I enjoyed reading this article as it contains an interesting analysis of (now-) historic measurements. I am happy to recommend it for publication once the following comments have been addressed.

53-59 add text about the quality of the ozone data 63 say that the latitude limit is due to its orbit (assuming it is!) 144 I prefer 'is' to 'shows clearly' 153 'low LIMS V6' Figure 5 I find the + symbols hard to see. Perhaps just say dotted lines. Also the text around line 144 says 'late October'. Given that the lowest values are in the first part of the period, I wonder if this should be rephrased - 'in this period'? I also wonder if Fig 5 is really needed.

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-1086>, 2019.

Author responses to comments from Referee #2

Thank you for expressing your interest in the findings of our manuscript.

Lines 53-59—We are adding estimates of profile accuracy.

Line 63—The southern latitude limit is due to the orbital inclination of Nimbus 7 and to the viewing geometry of the LIMS instrument.

Lines 144-153 and Figure 5—We prefer to keep both Figs. 5 and 6. Figure 5 shows details of the changing ozone and nitric acid in late October, while Figure 6 shows that their wave-1 patterns at 60S become nearly stationary in November. We rewrite “dotted horizontal lines” in the figure caption. We also delete the opening clause of line 141 and now draw our conclusions regarding advection from lower latitudes based on distributions of the species in Figs. 2 and 4.