

Response to RC2

Review of "Analysis of 24 years of mesopause region OH rotational temperature 1 observations at Davis, Antarctica. Part 2: Evidence of a quasi-2 quadrennial oscillation (QOO) in the polar mesosphere" by French et al.

This manuscript reports on the detection of a quasi-quadrennial oscillation detected in the SH polar winter OH nightglow, MLS and SABER temperatures. The feature is characterized after the removal of the solar signal and a linear trend as described in Part 1 of the study. The signal has a counterpart in the NH according to MLS data. The authors suggest a relationship between the temperature QOO and the changes in the mesospheric meridional winds and the corresponding changes in downwelling through adiabatic cooling and heating. The authors also show a significant correlation with the SST and provide an explanation of the connection with the mesospheric QOO through a changing GW filtering by modulated PWs.

I found the paper interesting in general and well written, the methodology used is convincing, and the results are new. Therefore, I think it is in principle suitable for publication in ACP. However, I must admit that I sometimes got lost when reading the discussion in Sect. 4. There are several places where it is not clear to me what the authors want to state. There are cases where mechanisms that are clearly not connected to the feature (for example, due to their time-scale) are even described in detail. The discussion should be more concise in order to make the reading more fluent and the reasoning more understandable. It should provide briefer but still comprehensive information only on mechanisms that may potentially be connected to the mesospheric QOO. Also, the authors should try to use a more cautious wording because they only suggest the origin of the feature through reasonable connections but do not provide a solid proof. In summary, in order to improve the paper, I urge the authors to re-write and significantly shorten section 4. Besides taking into account the suggestions listed below, I think the paper would that way be ready for publication in ACP.

We thank the Reviewer RC2 for your considered and helpful review. We have revised and re-organised the discussion in section 4 to hopefully improve the flow and comprehension of our study and the discussion. Please find specific responses to your review below. References to figures and sections are as per the reviewed manuscript, but be aware figure order and sections have been rearranged in the revised manuscript.

Comments:

L19 "peak-to-peak" amplitude

Added as suggested

L27 suggesting -> suggests

Corrected

L27 That the authors have found a correlation between the QOO and the CO vertical transport does not necessarily mean that a "substantial part of the QOO is the result of adiabatic heating and cooling driven by the meridional flow" but just that the QOO is plausibly linked, at least, to the

change in meridional flow. Please, provide some quantification proving such substantial contribution or re-write the sentence.

'Plausibly linked' is a better term to describe the relationship. Thank you. We will use that. We have added the correlation coefficients for the OH temperature and radar derived meridional wind at 86 km (from Fig 7) and from the CO correlations in Fig 8. The text in this section has been re-written and now reads

"We find a significant anti-correlation between the QO temperature and the meridional wind at 86 km altitude measured by a medium frequency spaced antenna radar at Davis ($R^2 \sim 0.516$; poleward flow associated with warmer temperatures at $\sim 0.83 \pm 0.21$ K/ms⁻¹). The QO signal is also marginally correlated with vertical transport as determined from evaluation of carbon monoxide (CO) concentrations in the mesosphere ($R^2 \sim 0.18$ at 0.73 ± 0.45 K/ppbvCO). Together this relationship suggests that the QO is plausibly linked to adiabatic heating and cooling driven by the meridional flow.

L30 In case it is not confirmed by modelling, please, replace 'suggest a tidal or planetary wave influence in' by 'is consistent with tidal or planetary waves influencing'

Modified sentence as suggested.

L32: 'potentially' -> 'plausibly'

Modified as suggested.

L40, insert comma after Antarctica

Inserted comma.

L78. Close parenthesis

Parenthesis closed.

L147 'Retrieved' might be ambiguous in this context. Do you mean 'selected'?

Changed wording to 'selected'.

L148 Did you select both daytime and nighttime MLS profiles? If so, did you check mean biases with respect to nighttime only profiles?

Yes, both daytime and nighttime profiles were selected as there are only ~2 profiles per day, however for our high latitude winter comparison the vast majority of the samples are night-time profiles. The latest version of the Aura/MLS data quality document (https://mls.jpl.nasa.gov/data/v4-2_data_quality_document.pdf) does not raise temperature biases between the ascending and descending passes as an issue over the recommended pressure level range.

There is a small diurnal temperature variation in the upper atmosphere (largely ozone heating). We have measured the diurnal variation to be <2K in the Davis OH temperatures.


Xu et al., 2010 (Jiyao Xu, A. K. Smith, Guoying Jiang and Wei Yuan. "Seasonal variation of the Hough modes of the diurnal component of ozone heating evaluated from Aura Microwave Limb Sounder observations" JGR, VOL. 115, D10110, doi:10.1029/2009JD013179, 2010) report on the diurnal variation in MLS temperatures due to ozone heating throughout the atmosphere which shows a minimum in the mesopause region (their Fig 2.) with the comment that "at 90 km, not all of the absorbed solar energy immediately appears as heat in the atmosphere. Substantial portions of the


incident solar energy are stored as chemical potential energy (i.e., as photodissociation products or other reactant species) and released as heat during exothermic reactions, which can take place during day or night.”

L153 The selection of this altitude is somehow artificial. The OH-layer altitude changes with time. Explain using SABER data the year-to-year change of this altitude and its effect on MLS data. Also explain the relationship between the QOQ in temperature and the QOQ in altitude.

The selection of the 0.0046hPa native MLS pressure level as an appropriate representation of an OH equivalent temperature for comparison with the Davis measurements is detailed in Part 1 of this study. This was also a reviewer concern in Part 1 and the following is paraphrased from the response provided.

In selecting the 0.0046hPa level we compared the Davis OH winter average anomaly with MLS [AMJJAS average] anomaly over a range of altitude and pressure levels. (see plots below, the first are altitude ranges, the second pressure levels, both temperatures and the anomaly are shown).





Altitude ranges 78-83km, 83-88km and 85-90km and pressure ranges 0.00215hPa and 0.00464hPa are all in reasonable agreement (<5K in absolute terms) with the OH temperatures, but we know there are biases with MLS (see French and Mulligan, 2010), and we know that the Davis OH temperatures are ~2 K high using LWR transition probabilities compared to those computed with the experimentally measured transition probability ratios determined in French et al., 2000.

We have previously reported, and routinely compare our measured OH temperatures with both Aura/MLS and SABER profiles. In particular, French and Mulligan, 2010 examined biases between Davis OH temperatures and both Aura/MLS and SABER. A significant limitation of SABER for comparisons with Davis observations is the yaw cycle sampling of the satellite. Comparable observations over Davis are confined to two intervals (day-of-year 75-140 and 196-262) and days prior to 106 and after 259 are outside the OH winter averaging interval. Therefore only days 106-140


and 195-259 are comparable and a large part of the winter months is not sampled. As a consequence SABER winter averages do not fit the OH observations at Davis as well as MLS.

In any case for this study, these biases are removed by comparing anomalies. We calculate the Chi-Square goodness of fit parameter between the OH winter average anomaly with the Aura/MLS anomalies. The 0.0046hPa pressure level yields the smallest chi-sqr (14.8) compared to a layer centred on the traditional 87km altitude level (85-90km chisqr=18.8).


This difference is small, but we prefer the pressure level comparison because we agree that the geometrical altitude of the layer varies. Since the hydroxyl layer position is primarily controlled by collisional quenching with O₂ and N₂ on the bottom-side of the layer, and reaction with atomic oxygen on the top-side of the layer it is the concentration (density) of the reacting species that governs the layer position. Therefore it is reasonable to compare with MLS pressure (proportional to density) levels than on geometrical altitude levels.

With regard to the change of altitude determined by SABER. We have examined the variation in OH layer height over this period in Part 1 of this study (acp-2019-1001; Figure 7 and related text). The overall trend is not significant, the mean winter VER peak altitude is 85.6 km and the standard deviation is 0.32 km, bearing in mind that the OH temperatures are integrated over the 8km FWHM of the layer shape, and for Aura/MLS the FWHM of the averaging kernel at this height is approximately 15 km.

The Figure below shows a comparison of SABER VER (altitude of peak) and corresponding pressure value (mb) for the years 2002-2018 (day 106 – 259 of each year) at Davis Station. The OH peak occurs at pressures in the range 0.00255 hPa to 0.003 hPa (2.55×10^{-3} - 3.0×10^{-3} Pa). This value lies between two of the Aura MLS levels (0.00464 hPa and 0.00215 hPa) on which the averaging kernels are centered and is in excellent agreement with the Davis MLS 4.2 Temperature PRES plot (page 4 above). An inverse relationship between altitude and pressure at the OH peak is clearly evident, and justifies the selection of a pressure level comparison for OH temperatures over an altitude level.




In a global context, we can also examine the altitude (or pressure) of OH peak as a function of latitude, the figure below shows the variation of the altitude of the OH peak as a function of latitude and day-of-year for the year 2005 from SABER data. The overall pattern shown here is repeated year after year with only minor changes in detail.



Based on the two figures above, the MLS averaging kernel centered on 0.00464 hPa would appear to be a good representative for the temperature of the OH layer.

The relationship between pressure and geopotential height (GPH) is examined below using the MLS data set. Global decreases in GPH anomaly (between ~110 to 240 metres/decade) at the 0.0046 hPa pressure level are consistent with a contraction of the underlying atmosphere and also consistent with the SABER trend in OH mean winter layer altitude for Davis shown in Fig 7. (200metres/decade) and discussed in the text in part 1 of this study.

MLS slope of fit to residual geopotential height at 0.0046 hPa



L184 What is the error in temperature due to a 1km-vertical shift?

The precise answer to this question depends on the vertical temperature profile and the OH volume emission rate vertical profile at the time of the measurement. (The OH-equivalent temperature (T_{VER}) is determined from a SABER temperature profile weighted with the volume emission rate profile (VER) measured simultaneously.) Using a typical temperature profile at Davis in the middle of the observing season, a 1 km -vertical shift in the OH-layer at 87 km would result in a temperature error of less than 1 K (added to the manuscript). This study uses long-term averages however and we note from above that the standard deviation in emission peak altitude for mean winter profiles is ~320 metres (over 16 years).

L190 Your figure 2b shows that the seasonal variation of the QO is significant, further, its year-to-year change is quite important. Indeed, according to the colour lines in Fig.1b, the large 2011 residual temperature seems to be influenced by the lack of measurements during days 141-195, with lower temperatures. This suggests that, due to a potential sampling bias, comparing SABER data with the ground-based OH dataset shown in Fig.1a, that includes days 141-195, might be misleading. Also there should be a bias in the derived trend and solar term due to that fact. I suggest to compare just with results from ground-based OH using only the coincident days, or at least FMA+ASO data in Fig. 1b, even if this needs from another figure. This may lead to a better SABER and Davis OH agreement. Indeed, I was expecting a better agreement than with MLS, given that SABER takes into account potential variations of the layer altitude.

Figure 2b) does not show any seasonal variation in the QO. This is the MLS polar cap average for the months AMJJAS at 0.00464hPa. We assume you are actually referring to figure 1b).

Figure 1b) shows the variation in detrended Davis OH temperatures for the intervals FMA, MJJ and ASO compared to the winter mean [AMJJAS]. This is / n{ SABER data which has the missing days 141-195 due to the yaw cycle ~~ZfZxt~~ year (not just 2011) .

The comparison with SABER data in Fig 1a) is provided only to demonstrate that the QO pattern exists in the SABER OH-equivalent temperatures in the vicinity of Davis. We have stated that the SABER data exclude days 141-195 from the winter averaging window (~~ZfZxt~~ ~~tZM~~ L197) and so do not believe the comparison is misleading.

The derived solar response and long-term trend from SABER OH equivalent temperatures in the vicinity of Davis is reported and discussed in part 1 of this study (acp-2019-1001). We have also previously compared Davis-OH and SABER temperatures in French and Mulligan (2010). The focus of this manuscript is to explore the QO feature and the SABER temperatures provided in Fig 1a serve only to support the existence of a QO pattern in winter average temperatures. We are aware that the yaw cycle data gap in SABER observations may yield a bias in the direct comparison with OH temperatures, but this is not a study of satellite biases.

L190. I wouldn't say that not observing from 21 May to 14 July means that 'SABER samples the same days'. Please, correct.

Modified sentence to read "Thus only days 106-140 and 196-259 are comparable between SABER and Davis-OH over the winter interval and days 141 - 195 (21 May to 14 July) are excluded."

L193 What trend and solar terms do OH data provide when not using days 141-195?

The terms are :

Davis OH (1995-2018)

$L = -1.68 \pm 0.68$ K/decade ($-0.26 > L > -3.09$ K/decade 95% confidence)

$S = 4.11 \pm 1.35$ K/100sfu ($6.9 > S > 1.31$ K/100sfu 95% confidence)

if days 141-195 are omitted from each year (1995-2018) and

Davis OH (2002-2018)

$L = -0.912 \pm 1.16$ K/decade ($1.58 > L > -3.40$ K/decade 95% confidence)

$S = 5.40 \pm 3.03$ K/100sfu ($9.55 > S > 1.25$ K/100sfu 95% confidence)

if days 141-195 are omitted from each year (2002-2018) to compare with the SABER trends in the vicinity of Davis.

SABER OH-equivalent T_VER (2002-2018)

$L = -0.76 \pm 1.06$ K/decade ($1.52 > L > -3.03$ K/decade 95% confidence)

$S = 3.38 \pm 1.87$ K/100sfu ($7.39 > S > -0.63$ K/100sfu 95% confidence)

for SABER T_VER within 500km of Davis (given yaw cycle excludes days 141-195)

This comparison is of interest for the trend assessment in Part 1 of this study but is not particularly relevant to the discussion of the QO in this part.

L197. According to a sentence written four lines above this one, the SABER T_VER trend is -0.77K/decade. This sounds contradictory with the -0.13 K/year mentioned in L197.

Line 194 is the result of the solar cycle and linear trend model fit, whereas line 197 is the trend in the raw temperatures (i.e., no solar component removed) in order to compare directly with the trend in the VER peak height. Edited the sentence to make this clear.

L198 This anti-correlation has a strong seasonal dependency, being significant in late autumn and very small in early spring for their case (e.g., Garcia-Comas et al., 2017) this. It is therefore not surprising that you get only slight anti-correlation when mixing temperatures for the two intervals day 75-140 and 196-

See response to L262 below and Figure of the anti-correlation for days 106-140 plus days 196-259 (top plot), days 106-140 only (middle plot) and days 196-259 only (lower plot). Days prior to day 106, and after day 259 are not considered because they are outside the observing season at Davis station.

L208 End sentence with a dot.

corrected

L250 Is this MLS temperature also?


Yes, this correlates the Aura/MLS polar cap mean temperature (indicated by the green ring in Fig 2b) with each MLS grid box, on several pressure levels. The sentence has been modified slightly to explicitly state this.

L262 What happens to the temperature-altitude anti-correlation if you separate these two intervals? This should also help to answer my previous comment on the effect of OH altitude variation on temperature.

The temperature-altitude anti-correlation for SABER DoY 106-140 is -4.6 ± 1.4 K/km and is significant at the 95% level ($R^2 = 0.43$), whereas the corresponding value for SABER DoY 196-259 is -3.2 ± 1.3 K/km and is also significant at the 95% level ($R^2 = 0.29$). In view of the difference between the two intervals, we calculated the solar and linear terms as in the case of the Davis OH data.

Summary of trends from SABER T _{OH} -equivalent (2002-2018) at Davis						
	DOY 106-140, 196-259	Significant at 95%	DOY 106-140 only	Significant at 95%	DOY 196-259 only	Significant at 95%
Solar term, <i>S</i>	3.4 ± 1.8 K/100 sfu	No	6.2 ± 2.3 K/100 sfu	Yes	1.2 ± 2.3 K/100 sfu	No
Linear trend, <i>L</i>	-0.77 ± 1.05 K/decade	No	-0.9 ± 1.3 K/decade	No	-0.1 ± 1.3 K/decade	No

Again, this comparison is of interest for the trend assessment in Part 1 of this study but is not particularly relevant to the discussion of the QGO in this part.




L260 Why the detrended OH temperature is compared with the GPH anomaly instead of the detrended GPH?

These are GPH composites, binned by the detrended Davis OH temperature into warm, mid and cold years, not time correlations. ie the detrended temperatures are only used to define the composite bins. The linear trend in the GPH does not affect the wave pattern observed in the results.


L294 Perhaps, this is more like a WN1 + WN2 structure. The WN2 is more clearly seen at 50hPa.

Agreed, it certainly appears as a WN2 pattern in the warm years meridional wind but more WN1 like in the cold and intermediate years. Modified this sentence to make this point.



L308 I do not think the feature is a wave-3 structure but more a wave-2. Note that the apparent wave-3 is due to the equirectangular projection.

Agreed, zonal wind at 10 hPa is not a wave 3. Corrected to wave-2.



L347 Do you mean 10m meridional and zonal winds?

Yes 10 m or 'near-surface' - as described in that sentence.



L350 I do not see a clear correlation between sea ice and the OH QQQ, except for mesoscale areas. The QQQ is however more clearly correlated with the SST in Fig.6b or the winds in Figs 6c-6d. That the sea ice concentration should be related to the SST and the winds seems not really big news. Then, what is the additional information from OH-QQQ and the sea ice correlation here? If it is difficult to provide a satisfactory answer, I would suggest changing the title of section 4.1 to "SST and winds", suppressing Fig.6a and the long discussion below on the sea ice QQQ (L376-L386).

We have added time series for the correlation between the regions of maximum anti-correlation between Davis-OH and the sea-ice cover to the supplementary material (see figures below). Region A has a correlation coefficient -0.49 with Davis and -0.46 with MLS. Region B has a correlation coefficient -0.56 with Davis and -0.65 with MLS.

We do not see sea ice cover as a potential source of the mesospheric QQQ. "Correlation does not imply causation". Rather, we note that there are regions that are significantly correlated with the

Davis QOQ variation (regions A and B below) that appear to have a wave-2 structure and suggest that possibly the meridional winds could drive both observations ie “a persistent northward (southward) flow on one side of a circulation anomaly could increase (decrease) sea ice due to the associated flow of relatively cold (warm) air from higher (lower) latitudes and expansion compaction) of the ice edge.”

This section on correlations with sea ice cover has been revised, suppressed, moved from section 4.1 to 4.4 and renamed “Relationship with SST and Antarctic sea ice”.



L359 Do you mean cyclonic circulation anomalies? Note that you show correlations in Figure 6 and not winds.

Inserted ‘anomalies’

L366 I think the reasoning is in general correct but what the authors show here is a strong connection between the QJO and the SST and also links to the surface circulation. The phenomenon where the origin resides cannot be determined from this type of analysis.

Agreed, we only suggest “this could hint as to the origin” and provide a plausible possibility to explore with other analyses.

L369 Again, unless I missed something, I do not understand the interest of mentioning the sea ice in this work unless the authors see it as a potential source of their OH QJO. Please, provide a more convincing argument or suppress.

See response to L350 above.

L385 QJO temperature signal at what level?

Generally speaking the QJO in the upper mesosphere. We have shown the QJO detected at Davis is correlated with the SH polar cap between 0.0046 hPa, down to 0.1hPa. We have added “in the upper mesosphere” to the sentence to clarify.

L405 colder 'mesospheric' temperatures


Inserted 'mesospheric'.

L414-426 I do not really understand the need for this discussion. Even if the tides are affected by the QJO, there is no impact of QJO-ENSO on OH QJO. Could you please extend on the point here?

We find distinctly different wave patterns in the ERA5 geopotential and meridional wind anomalies during warm and cold years of the QJO so some factor is modifying the behaviour of these parameters on a QJO timescale. Is this possibly tides or planetary waves ?. This discussion makes the point that interaction between tides with the QJO and ENSO can have an impact at inter-annual timescales as reported by Baldwin et al (2019) and Liu (2016). Is a similar interaction occurring to produce these different wave patterns on a QJO timescale?. The discussion merely poses this as a possibility by corollary with the QJO and ENSO interaction.

L415 Not particularly but only in the cold years.

We see a SH wave-1 structure in the composite cold years, but also possible in the mid years. However, have removed the word 'particularly'



L423 'longitudinal' wave patterns

Inserted 'longitudinal'.

Figure 7a. Is that really detrended OH T, as the x-axis title states? Please, write also in the caption "zonal mean meridional wind anomaly". Also, this figure is somehow redundant with Fig. 1. Winds could be overplotted there.

No, they are residual temperatures as stated in the caption. Thank you for picking that up, the y-axis title has been corrected. The meridional wind anomaly is that measured at Davis it is $1/n$ a zonal mean. Thanks for the suggestion, we agree the figure is somewhat replicated from figure 1 but with the addition of the mean meridional winds. The winds could be added as a separate panel in fig 1 but we prefer to keep this separated as the radar winds are not introduced until the discussion in section 4.2.




Figure 7b. I think Fig. 7b is not needed. The correlation is clearly seen in Fig. 7a. Remove Fig 7b or, at least, move it to the supplement.

Figure 7b provides the regression coefficients and R^2 values for the meridional wind to OH, Aura/MLS and SABER temperatures referred to in the text. As the relationship of the OH temperature QO to the meridional wind has the most significant correlation (R^2 0.51) of the factors compared this is a key finding in the investigation and we prefer to keep this figure (time-series and slope of the relationship) in the manuscript.

L439 Do you mean the background 'meridional' wind?

Inserted 'meridional'.

L438-442 I think it is well known that the mesospheric poleward circulation is connected to downwelling and adiabatic heating below 100km. I do not think the references credited by the authors are the first ones showing that.

We agree with the comment. These references are selected because they are among the first to report observing the effect in OH temperatures, which are the data in which the QGO was identified initially in this study. We have added a reference to the earlier work of Garcia and Solomon (1985).

L444 'the adiabatic action of the residual meridional circulation' sounds confusing and may be misinterpreted


The sentence has been modified to avoid possible confusion. "and the long-term linear trend is due to the adiabatic action of the residual meridional circulation."

L452-L454 Please, remove this sentence. It is not necessary to provide the sources and the sinks of CO. Just knowing that it is a dynamical tracer due to its long lifetime, particularly, during polar winter, is enough.

The sentence has been removed.

L464 I find the anti-correlation between temperature and the CO trends interesting. This might be out of the scope of this paper and not worth to mention in the paper but: What are the errors of the CO measurements? Is the instrumental drift characterized? Can the authors provide a link for the trend anti-correlation? Perhaps CO2 increase?

Uncertainties in the CO measurements are of the order of 0.3-0.4 ppmv at 0.00464hPa and 0.05-0.07 ppmv at 0.1 hPa. Error bars have been added to figure 8 as shown below.



Instrumental drift and calibration is addressed by the Aur/MLS science team (https://mls.jpl.nasa.gov/products/co_product.php) but is beyond the scope of this work. There are both dynamical and chemical factors that could contribute to the trend in CO. Perhaps there are trends in the large scale upwelling and downwelling via meridional flow that occur over the polar caps and, as you suggest perhaps the trend is associated with CO₂ increases. This deserves further study, but is outside of this manuscript aims.

L473 Would the correlation be better if the CO detrended data were used? Given the nature of the link between CO and T QGOs that the authors are providing, does it make sense to provide 8 maps with projections for selected altitudes instead of just one plot with a lat-z cross section? On the

other hand, if the QO is clearly exhibited in the SH polar cap average (see Fig. 2 and Fig 3), why the correlation with T and CO QO is not ubiquitous south of 65S?

The correlation coefficient with detrended CO and Temperatures is described on L463. We have considered a Lat vs Z plot but this necessarily requires zonal averages whereas Fig 9 shows that the patterns are zonally asymmetric. These levels and the plot format are provided so that they can be compared directly to the plots in Fig 2,3 and 4 (mesosphere levels). With regard to the polar cap we describe in the text “temperature and CO are significantly positively correlated over most of Antarctica, which is consistent with Fig. 8 and our hypothesis that the QO temperature variation is an adiabatic response (i.e. increased (decreased) temperature is associated with increased (decreased) CO concentration due to descent (ascent)). This general positive correlation over Antarctica is also seen for CO at 0.1 hPa (Fig. 9b) and is apparent though less clear for CO at 1 hPa (Fig. 9c).”

L495 Temperatures at what pressure level?

inserted “(the 0.0046hPa pressure level)”

L498-500 I would not call that "somewhat smaller". There is a very weak QO in the NH winter.

We have modified this section substantially and the sentence now reads “we see little evidence of a QO variation in the NH winter (blue line with linear fit)” referring to Fig 10 (below)

Section 4.5 Why is this section called "inter-hemispheric coupling"? Certainly, Figure 10 shows NH and SH residual temperatures but their connection is not discussed at all.

We have modified the section heading to Hemispheric comparison as your point is correct.

L530 If that is the case, why the QO is largest during the SH summer?

We do not have measurements over the summer to confirm the Aura/MLS observation of a larger QO in the summer months [ONDJFM] in Fig 10. We do state “We have attempted to examine the seasonal variability of the QO signal by dividing averages into intervals FMA, MJJ, ASO (also plotted in Fig 1b). While these shorter term averages obviously suffer from greater uncertainty, there is a suggestion that the QO is strongest over the winter months MJJ, mid-range in ASO and less apparent in the FMA interval.”

Davis is not near the GW hot spot of the Antarctic Peninsula and we also comment from L543 “Gravity wave energy is generally weak in summer but in winter, gravity waves have large

amplitudes and are distributed around the polar vortex in the upper stratosphere and mesosphere. The wave energy is not zonally uniformly distributed but is concentrated on the leeward side of the Southern Andes and Antarctic Peninsula. Energy propagation extends several thousand kilometres eastwards which explains the gravity wave distribution around the polar vortex in winter.”

L544 Please, state this is true for SH.

That this refers to the SH is stated in the preceding sentence on L543.

L550-554 I do not think that results from Sato et al. (2012) (showing a close to annular structure around the Arctic edge, with maximum values over the Antarctic Peninsula) are well reproduced by Fig. 2a and 2b (showing more or less homogeneous correlations over the Arctic and a lack of correlation over the Peninsula).

The polar projection plots of the SH at 0.01 hPa (not 0.1 hPa as stated in the original manuscript) and 0.0046 hPa shown in Figure 2(c) in this work show a striking resemblance to the pattern in the SH winter months shown in Sato et al. (2012).

Compare the polar projection plots of June, July, August and September from Figure 2 of Sato et al. (2012) shown below with Figure 2(b) at 0.01 hPa and above from this work (also shown below). Please note that the Sato et al. (2012) plots have longitude 0° at the top of each projection plot, while the SH in Figure 2 of this work have 180° at the top of each projection plot. (This is corrected in the revised manuscript)





FIG. 2. Polar stereographic projection maps of gravity wave potential energy (shading) in the SH for each month at 10 hPa. Monthly mean zonal winds are shown by thick contours at an interval of 40 m s^{-1} .



P554-558 Could the authors provide some conclusion after these sentences and make clear the relationship with their work?

A concluding sentence has been added to the text after these sentences. “The results of Sato et al. (2012), together with the focussing of GWs into the polar night wind reported by Wright et al. (2016, 2017) suggests GWs as a plausible explanation for the asymmetry in the polar projection plots of the QQO correlation coefficient at 0.01 hPa and above. “

L562-570 The vertical and latitudinal structure of Zhang et al.'s TO is similar to the QBO. However, the authors showed no correlation of their QQO with the QBO. Then, why this discussion of that TO if the four-year oscillation is not even discussed in Zhang et al.? Please, clarify the interest of this discussion or remove. Please, also note that, even if Zhang et al. showed no evidence of the four-year oscillation at 45 km, they did at 85km.

As the referee has noted, Zhang et al. (2017) found evidence of a four year oscillation at 85 km in their analysis. This is stated explicitly in the second sentence of the paragraph, which is the reason that we include it in our discussion. Since we have found evidence of the QQO at different altitudes, we note that Zhang et al. (2017) also found a four-year oscillation at 25 km. Zhang et al. (2017) do not discuss the 4-year oscillation (as we state), presumably because it has a much lower amplitude than the periods on which they focus. We make mention of the three-year oscillation simply because Zhang et al. (2017) suggest that it may arise from a modulation of the QBO by the semiannual oscillation. This struck us as interesting, and we included it here for that reason. We examined our data for evidence of a similar explanation for the QQO, but we did not find any evidence of it (and so we do not include it here).

L570-572 From the discussion in the previous section, I thought that the authors were suggesting that mesospheric QQO was related to orographic GWs.

The previous section entitled “Gravity wave interaction” noted the similarity of the pattern of GW potential energy at 10 hPa in Fig 2. of Sato et al. (2012) with the polar cap correlation plots of

Aura/MLS temperature (at 0.01 hPa (and above)) with the QGO. This was interpreted as a possible clue to the origin of the QGO without being specific about the period. The section entitled "Mechanisms for a 4-year cycle" discusses reports which include references to periods in the vicinity of a 4-year cycle.

L573-575 Did Liu et al. find any QGO in their GW potential energy at the equator. Figure 2b shows a QGO there at e.g. 81km

Liu et al. (2017) do not refer to a QGO component in their results.

L579-585 That a changing eddy diffusion is not needed anymore to explain SABER CO2 trends has nothing to do with its potential link with a QGO.

The three sentences dealing with this point have been removed.

L594 Replace the first two dots by a comma

replaced as advised

L596 Offermann et al. note in their introduction periods ranging from around 2 yrs to 11 yrs. What periods do the authors refer to in this sentence? All of them?

This sentence has been modified to focus on periods at 3.4 and 5.5 years that are closest to the QGO.

L623-624 If the polar cap also shows a QGO of similar amplitude and in phase with Davis (as you write in the next sentence, it is obvious that the Davis QGO is positively correlated with the polar cap QGO. Please, re-write.

The first point (L620-621) notes that the OH-layer equivalent temperatures of both Aura/MLS and SABER in the vicinity of the station (within 500km) agree with the Davis OH temperature QGO in amplitude, period and phase.

The following point (L622-624) notes that the Davis QGO signal is positively correlated with the Aura/MLS 0.0046 hPa level temperature field large over a large part of the polar cap and southern Indian ocean (and where it is anti-correlated in the Southern Ocean).

both points have been re-written to make this distinction clearer and the negative correlation with the NH identified as a separate point.

L631 extends vertically "at least" from the mesopause

Added text as suggested

L646 Would a significant variability in atmospheric tides be actually expected at this high polar latitudes?

No, good point, tidal variability would not be expected to be significant (tides less than 2K) but potentially their interactions with longer term oscillatory modes is responsible for the different wave patterns seen during warm and cold years of the QGO. We suggest only a potential role. See also response to L414-426.

L653-655 I do not think the authors prove this connection to be "most likely".

This point is modified to the word 'possibly' and merged with the previous point.

L669-672 Is the Davis QGO anti-correlated with the NLC boundary latitude reported by Russell et al.?

NLC data from Russell et al., 2014 are for the NH and span 2005-2011. Their fig 4 shows the NLC occurrence frequency anomaly for these years in 5° latitude bands and fig 5 the corresponding MLS and SABER [MJJA] temperatures. From our Fig 10 (see below), MLS NH summer [AMJJAS] averages show warm years in 2006 and 2009 and cold years in 2007-08 and 2010-11 when we would expect increased NLC occurrence. There is an indication of an NLC increase in 2007-08 and 2011 in their fig 4 panels h and i for the high latitude bands but the variation is small. The difficulty with the temperature/NLC correlations at high latitude are nicely explained in their summary “As latitude increases, the cloud frequency increase weakens significantly, and in high polar latitudes, the cloud frequency remains roughly constant over the 10 years examined here. This is expected because under highly saturated conditions, i.e., when temperature is far below the frost point, the cloud frequency is close to 100%, so there is no room for further increases, and the temperature control of cloud frequency no longer holds.”

The low latitude boundary anomaly is provided in their fig 7 (reproduced right) and in summary “The results show a statistically significant increase in the number of PMCs each season in the latitude range 40°N–55°N for the 10 year period examined. Increases in cloud frequency appear to be driven by the corresponding temperature decreases over the same time period. During this time, solar activity decreased from an active to a quiet period, which might have been partially responsible for the temperature decrease over this time period.” There is possible evidence of a QGO in the SABER data, but generally not enough data to be definitive.




Figure 7. Interannual variability of the PMC low-latitude boundary anomaly (in degrees) from its 10 year average. The anomaly is first calculated on a daily basis, and then, a seasonal average is obtained for each year. The seasonal length used here is from –10 DFS to 50 DFS when PMCs are widespread enough to cover all longitudes. The error bars are the standard error of the mean. The results show a slightly negative but statistically insignificant change in the low-latitude PMC boundary over the 10 year period used in this study.

Fig 2 caption. I think the caption for b) is not correct. If "as for a)", shouldn't it be just "correlation of the SH polar cap average", without "0.0046hPa"?

- a) correlates the **0.0046hPa grid box temperature over Davis** with the MLS temperature field
- b) correlates the **0.0046hPa polar cap average temperature** with the MLS temperature field both show the same plot layout of time-series and projection maps (which was the As for a) reference). We have removed the “As for a)” and specified the correlation in b) explicitly in the figure caption.

L921 What green circle?

The green circle on the SH projection map at 0.0046hPa which indicates the polar cap averaging region. It has been made a bit thicker for visibility. Figure 2 has been redone.

Figure 6. Please, indicate the maximum and minimum values in the legend of the color scale for the lower panels. Are the two color scales the same? If the answer is yes, just keep one of them. If the answer is no, use the same color scale for the four panels.

Figure 6 has been redone with the same color scale for all panels.

