



Trends in N₂O and SF₆ mole fraction in archived air samples from Cape Meares, Oregon (USA) 1978–1996

Terry C. Rolfe¹ and Andrew L. Rice¹

¹Department of Physics, Portland State University, Portland, 97201, United States

5 *Correspondence to:* Terry C. Rolfe (trolfe@pdx.edu)

Abstract. Quantifying historical trends in atmospheric greenhouse gases (GHG) is important to understanding changes in their budgets and for climate modeling which simulates historic and projects future climate. Archived samples analyzed using updated measurement techniques and calibration scales can reduce uncertainties in historic records of GHG mole fractions and their trends in time. Here, we present historical measurements of two important GHG, nitrous oxide (N₂O) and sulfur hexafluoride (SF₆), collected at the midlatitude northern hemisphere station Cape Meares, Oregon (USA, 45.5° N, 124° W) between 1978 and 1996 in archived air samples from the Oregon Health and Science University – Portland State University (OHSU–PSU) Air Archive. N₂O is the third most important anthropogenically forced GHG behind carbon dioxide (CO₂) and methane (CH₄). SF₆ has a low abundance in the atmosphere, but is one of the most powerful GHG known. Measurements of atmospheric N₂O made during this period are available for select locations but prior to mid-1990 have larger uncertainties than more recent periods due to advancements made in gas chromatography (GC) methods. Few atmospheric SF₆ measurements pre-1990 exist, particularly in the northern hemisphere. The GC system used to measure N₂O and SF₆ concentrations in this work is designed to be fully automated, capable of running up to 15 samples per batch. Measurement precision (1σ) of N₂O and SF₆ is 0.16% and 1.1% respectively. Samples were corrected for detector response non-linearity when measured against our reference standard, determined to be 0.14 ppb ppb⁻¹ in N₂O and 0.03 ppt ppt⁻¹ in SF₆. The concentration of N₂O in archived samples is found to be 301.5 ± 0.3 ppb in 1980 and rises to 313.5 ± 0.3 ppb in 1996. The average growth rate over this period is 0.78 ± 0.03 ppb yr⁻¹ (95% CI). Seasonal amplitude is statistically robust, with a maximum anomaly of 0.3 ppb near April and a minimum near November of -0.4 ppb. Measurements of N₂O match well with previously reported values for Cape Meares and other comparable locations. The concentration of SF₆ in analyzed samples is found to be 0.85 ± 0.03 ppt in 1980 and rises to 3.83 ± 0.03 ppt in 1996. The average growth rate over this period is 0.17 ± 0.01 ppt yr⁻¹ (95% CI). Seasonality is statistically robust and has an annual peak anomaly of 0.04 ppb near January and a minimum anomaly of -0.03 ppt near July. These are unique SF₆ results from this site and represent a significant increase in SF₆ data available during the 1980s and early 1990s at any location. The concentration and growth rate of SF₆ measured compares well to other northern hemisphere measurements over this period. From these N₂O and SF₆ measurements, overall we conclude that sample integrity is robust in the OHSU-PSU Air Archive.



1 Introduction

Anthropogenic sources of greenhouse gases (GHG) have altered the atmospheric composition resulting in a significant climate forcing near 3 W m^{-2} since 1750 (Myhre et al. 2013). Measurements of GHG concentrations since the industrial revolution constrain global budget uncertainties and are central to interpret recent changes to source and sink processes (Prinn et al. 2000; Khalil et al. 2002; Saikawa et al. 2014). When projecting future GHG concentrations, many additional factors must be included in models such as climate feedback effects and possible changes in transport processes. Uncertainties in model predictions can be minimized if GHG measurements are precise and span many different latitudes (Meinshausen et al. 2017).

When historical timeseries records are not available, past atmospheric GHG abundance can be evaluated using archived air samples and ice core and firn air. One significant advantage of ice core and firn air for measuring past atmospheric concentrations of GHG is that samples may be collected today that represent past conditions. However, ice core and firn samples are difficult to obtain due to the remoteness of the locations where the samples are collected (Greenland and Antarctica) and provide limited spatial information. Temporal uncertainties also must be evaluated when measuring ice core and firn samples due to diffusion and gravitational separation (Ishijima et al. 2007); samples are best represented by a mean age. By contrast, archived air samples are discrete in time and space, making them very valuable for evaluating past atmospheric abundance at specific periods in time. However, few air archives are available today. Archive samples may also contain storage artifacts that can contaminate historical records.

Nitrous oxide (N_2O) is the third most important GHG with anthropogenic sources after carbon dioxide (CO_2) and methane (CH_4). Today, the concentration of N_2O is close to 330 ppb with a trend averaging 0.75 ppb yr^{-1} over the last 30 years (Ciais et al., 2013). N_2O has a large global warming potential (GWP), 298 times that of CO_2 over a 100-year period and a global radiative forcing estimated at 0.19 W m^{-2} since 1750 (Myhre et al., 2013). The long lifetime (~ 120 years) results in most emitted N_2O reaching the stratosphere, where photooxidation is the major source of stratospheric NO_x ("active nitrogen"). NO_x is the main natural catalyst of ozone (O_3) destruction (Crutzen 1970).

Anthropogenic sources of N_2O account for roughly 40% of all N_2O emissions, with natural sources accounting for the other 60% (Ciais et al., 2013). Bottom-up calculations estimate anthropogenic production of $6.9 (2.7-11.1) \text{ TgN yr}^{-1}$ and natural production of $11 (5.4-19.6) \text{ TgN yr}^{-1}$. The uncertainty in these estimations is large, with 1σ error nearly $\pm 50\%$. Together with atmospheric measurements, top-down modelling better constrains the N_2O budget and reduces uncertainty in the global source. Sources of N_2O calculated this way estimate anthropogenic and natural source production of $6.5 (5.2-7.8) \text{ TgN yr}^{-1}$ and $9.1 (8.1-10.1) \text{ TgN yr}^{-1}$, respectively (Prather et al., 2012).

There are three major natural sources and six major anthropogenic sources of N_2O . Natural sources of N_2O are natural soils ($3.3-9.0 \text{ TgN yr}^{-1}$), oceans ($1.8-9.4 \text{ TgN yr}^{-1}$), and atmospheric chemistry ($0.3-1.2 \text{ TgN yr}^{-1}$) (Note: sources include the minimum and maximum estimates provided from bottom-up calculations in Ciais et al., 2013). By far, the largest anthropogenic source is agriculture, producing $1.7-4.8 \text{ TgN yr}^{-1}$, followed by industrial and fossil fuel sources ($0.2-$



1.8 TgN yr⁻¹), biomass burning (0.2-1 TgN yr⁻¹), rivers and estuaries (0.1-2.9 TgN yr⁻¹), atmospheric deposition (0.4-1.3 TgN yr⁻¹), and human excreta (0.1-0.3 TgN yr⁻¹) (Ciais et al. 2013). More constraints on source production provided through atmospheric measurements are needed to improve estimates of individual source magnitudes.

The main loss mechanism for N₂O is destruction in the stratosphere through photolysis and the reaction with O(¹D) (Prather et al. 2015). Soils and the oceans can act as sinks for N₂O through microbial processes, however because the production of N₂O is greater than what is consumed, the global net flux is positive. Estimates of the stratospheric sink account for 11.9 (11.0-12.8) TgN yr⁻¹ (Ciais et al., 2013).

Rising global concentrations of N₂O are due to the imbalance between the sources and the sinks. Based on a top-down constraint, the imbalance between sources and sinks is 3.6 (3.5-3.8) TgN yr⁻¹ (Ciais et al., 2013).

Models have shown that future climate conditions will likely amplify N₂O production, meaning a linear increase in time may under-predict future concentrations based on the current rate of change (Khalil and Rasmussen 1983; Stocker et al. 2013). To minimize uncertainty in the N₂O budget and in model projections, precise measurements of current and past atmospheric conditions from multiple global locations are needed. Measurements of atmospheric N₂O made prior to mid-1990 have larger uncertainties than more recent periods due to advancements made in gas chromatography (GC) methods (Prinn et al. 2000; Jiang et al. 2007). To reduce uncertainty during this period, archived samples may be analyzed using updated measurement techniques. Additionally, measurements of the isotopic composition of N₂O in archived samples can constrain the N₂O budget and changes in time due to characteristic isotopic effects in sources and sinks (Park et al. 2012; Snider et al. 2015).

Sulfur hexafluoride (SF₆) is an extremely potent GHG with a GWP of 22800 (over 100 years compared to CO₂) and a lifetime of ~3200 years (Ravishankara et al. 1993). While SF₆ is one of the strongest GHG controlled under emission regulations, it has a low global concentration (7.29 ppt in 2011), so it does not add significantly to climate forcing by itself (Myhre et al. 2013).

Sources of SF₆ are anthropogenic, with main uses being high voltage insulation, magnesium production and semiconductor manufacture (Maiss and Brenninkmeijer 1998; Olivier et al. 2005). Global production in 2008 was estimated to be 7.16 Gg yr⁻¹ (Levin et al. 2010). With a very low solubility and no reactivity in the lower atmosphere, the only known sink for SF₆ is loss in the mesosphere.

With almost all of the SF₆ that has been emitted since the industrial revolution to the atmosphere still present, global emissions can be accurately determined from observations of atmospheric concentration. Due to its long lifetime and anthropogenic origins, SF₆ is used as a validity check for atmospheric transport models (Levin and Hesshaimer, 1996; Patra et al., 2009). It has been estimated that 94% of all SF₆ emissions originate in the northern hemisphere (Maiss et al. 1996), explaining a north-south hemisphere gradient in SF₆ concentration of about 0.4 ppt (Levin et al. 2010).

Observations of the SF₆ growth rate have been reported by several studies (Levin et al. 2010; Rigby et al. 2010; Hall et al. 2011). The trend in SF₆ has varied over the last 30+ years and while the magnitude of the growth rate differs slightly between sample locations, several features are prominent. From the early 1970s to the mid-1990s, the trend steadily



increased from 0.1 ppt yr⁻¹ and peaked near 0.26 ppt yr⁻¹. The trend then slowly declined to ~0.20 ppt yr⁻¹ until the early 2000s, when the trend increased again. The inferred global emission of SF₆ from the trend increases nearly linearly from 2 Gg yr⁻¹ in the late 1970s to over 6 Gg yr⁻¹ in 1994-1995 (Levin et al. 2010; Rigby et al. 2010).

Reported atmospheric measurements of SF₆ before the year 1987 are few. In the southern hemisphere, Cape Grim, Tasmania (41° S, 145° E) archive measurements date back to 1978 (Levin et al. 2010). Northern hemisphere measurements are reported dating from 1973 from Trinidad Head, CA (41° N, 121° W), but few are prior to 1990 (Rigby et al. 2010). A more complete record of past SF₆ atmospheric concentrations is desirable.

The Oregon Health & Science University–Portland State University (OHSU-PSU) air archive includes archived air samples collected from Cape Meares, Oregon (45.5° N, 124.0° W) in the late 1970s, 1980s, and 1990s by the Department of Environmental and Biomolecular Systems, Oregon Graduate Institute of Science and Engineering (currently OHSU). The samples were collected by air liquefaction, where ~1000 L of air (STP) was compressed to 3000 kPa into 33 L electropolished stainless steel canisters. Today, archive samples are stored at Portland State University and contain pressures ranging from 60-2000 kPa (Rice et al. 2016). Here, we present details of the analytical technique employed and results from the analysis of 159 Cape Meares air samples from the OHSU-PSU air archive.

2 Methods

2.1 Gas chromatography analytical system

The gas chromatography (GC) analytical system (Fig. 1) employed at Portland State University for measuring N₂O and SF₆ in archived air samples is based on the configuration used by Hall et al. (2007) and references therein. We use an Agilent model 6890N gas chromatograph fitted with a micro-electron capture detector (μECD, Agilent Technologies, Santa Clara, CA). Peak separation is achieved by two Poropak Q 80/100 mesh columns (1.8 m × 2 mm i.d. pre-column, 3.7 m × 2 mm i.d. analytical column). The carrier gas is P5 (99.999%, Airgas, Portland, OR) equipped with O₂ and hydrocarbon traps (Restek, Bellefonte, PA) to further reduce impurities and found to significantly improve baseline signal stability. Two six-port switching valves (V₁ and V₂), a four-port switching valve (V₃), and a 16-port multi-position valve (Valvo Instrument Company Inc., Houston, TX) are controlled through Chemstation (V1.A, Agilent Technologies Inc., Santa Clara, CA).

A sample run begins in “back-flush” mode, with carrier gas flushing the pre-column in the reverse analytical direction to remove the build-up of water on the analytical column that would otherwise eventually elute to the μECD and affect signal baseline. A 16-port multi-position valve is used to introduce pressurized samples into the system; a 2-way electric valve (Clippard, Cincinnati, OH) is used to stop sample flow to the sample loop and prevent sample loss. Samples initially pass through a desiccant trap (Perma Pure, Toms River, NJ), before flushing a 10 ml sample loop at 60 ml min⁻¹ for 1.5 minutes. At this time, V₃ rotates, which places the system in “front-cut” mode and allows the sample loop to equilibrate. V₁ rotates at 1.75 minutes and allows the carrier gas to carry the sample N₂O and SF₆ to the pre-column where separation from O₂ and H₂O occurs. After O₂ elutes through the pre-column to vent, at 3 minutes V₂ rotates and places the pre-column



in line with the analytical column, transferring N₂O and SF₆ to the analytical column. At 4.25 minutes, the sample has reached the analytical column and V₁, V₂, and V₃ rotate. This begins the back-flush of the pre-column while the analytes are carried to the μ ECD on the analytical column.

Oven and detector temperatures are maintained at 56° C and 310° C respectively. Carrier gas flow rates are 40 ml min⁻¹ maintained by electronic pressure control of the 6890N. N₂O peak retention time is 6.1 minutes and SF₆ peak retention time is 7.0 minutes (Fig. 2). Peak integration is accomplished via Chemstation based on peak height.

All measurements of N₂O and SF₆ are made relative to a calibrated whole air sample on the NOAA-06 N₂O scale and NOAA-14 SF₆ scale (NOAA Tank CB11406-A, 328.71 ± 0.5 ppb N₂O, 8.76 ± 0.06 ppt SF₆), here-on referred to as the NOAA reference gas. Each sample is analysed 6 times and bracketed by 6 reference gas runs used to measure instrument response and track signal drift. The GC- μ ECD analytical system was evaluated for precision, reproducibility, and linearity prior to its application to measure mole fraction in historic archive samples.

2.2 Precision and reproducibility of analytical system

Precision of measurement was determined by comparing residuals from sets of 6 gas analyses. Histogram distributions in figure 3 show 180 residuals (expressed as a percent relative standard deviation) collected from 30 sets of 6 measurements of N₂O (Fig. 3a) and SF₆ (Fig. 3b) of the NOAA reference gas. Both N₂O and SF₆ compare well to a normal distribution (black dashed lines), with chi-square goodness of fit p-values of 0.16 and 0.35, respectively. For N₂O, 1 σ = 0.16% while for SF₆, 1 σ = 1.1%. This corresponds to an uncertainty of ± 0.52 ppb for N₂O and ± 0.10 ppt for SF₆.

Measurement reproducibility was evaluated by repeatedly measuring a dry air sample (Breathing Air, Airgas, Portland, OR) against the NOAA reference gas and evaluating consistency from the standard deviation of the results. The sample was measured 18 times over a two-week period with mean measured concentrations of N₂O and SF₆ of 390.9 ppb and 13.2 ppt, respectively. The standard deviations in N₂O and SF₆ measurements are 0.46 ppb and 0.11 ppt respectively, which are indistinguishable from 1 σ precision for a set of 6 NOAA reference gas measurements.

2.3 Linearity of the GC- μ ECD system

To ensure accurate results for this work, the detector response was evaluated over the mole fraction range expected for N₂O and SF₆ in the OHSU-PSU Air Archive. The range in northern hemisphere N₂O mole fraction between 1978 and 1996 is between 295 and 314 ppb (Prinn et al. 2000; Ciais et al. 2013). Archived air sample measurements of northern hemisphere SF₆ mole fraction from Trinidad Head, CA measure below 1 ppt in the 1970s and rise to nearly 4 ppt in 1997; southern hemisphere measurements from Cape Grim, Tasmania and the South Pole show a similar range (Levin et al. 2010; Rigby et al. 2010).

A series of manometric dilutions were prepared from the NOAA reference gas at Portland State University to evaluate the μ ECD response over historical N₂O and SF₆ mole fraction sample range. To characterize the N₂O response, the



N₂O reference gas was diluted with ultra-pure air (zero grade, Airgas, Portland, OR; N₂O and SF₆ at concentrations below detection limits) using capacitance manometers (MKS Instruments, Andover, MA; range 0-10 torr and 0-1000 torr) into 3L electropolished stainless steel canisters (precision ± 0.01%). The range of N₂O concentrations produced in 3 L canisters was 32.2 - 321.4 ppb. Error introduced from the manometric process is small when compared to measurement uncertainty
5 (maximum 1σ error of ± 0.07 ppb for N₂O).

To characterize of the SF₆ response at low ppt concentrations requires consideration of the effect of the falling N₂O tail on the chromatogram baseline. To properly account for this interference, SF₆ dilutions at low concentrations (0.6 - 6.0 ppt) must have a N₂O concentration that reflects expected concentrations in archived samples (302–314 ppb). Prepared dilutions of SF₆ included the addition of an aliquot of 1 ppm N₂O (±5%, Scott Specialty Gases, St. Louis, MO) into the
10 canister prior to dilution with ultra-pure air. The maximum error (1σ) in SF₆ introduced from the manometric process is small (0.001 ppt) compared to measurement uncertainty. All dilution samples were measured at PSU on the GC-μECD system over several weeks to account for instrument drift. Tables 1 and 2 provides dilution sample pressures, calculated and observed μECD response, and measured N₂O and SF₆ mole fractions with the error in measurement used to characterize the GC-μECD linearity.

15 Results of linearity experiments are shown in figure 4. For N₂O, a slope of 0.870 ± 0.028 (95% CI) is found over the data range 289.7 - 328.7 ppb, most relevant for this work. A linear fit is a good model for the deviation from expected over this range ($R^2 = 0.964$); additional polynomial terms are not statistically robust. This results in sample measurements deviating from expected by ~ 0.14 ppb ppt⁻¹ N₂O difference from the NOAA reference. For the range of the N₂O in the OHSU-PSU air archive, all N₂O samples are adjusted for a linear correction of the form:

$$20 [N_2O]_x = a_1[N_2O]_y + a_2 \quad (1)$$

$$a_1 = 1.146 \pm 0.037 \text{ (95\% CI)} \quad (2)$$

$$a_2 = -47.95 \pm 11.49 \text{ (95\% CI)} \quad (3)$$

Where $[N_2O]_y$ is the measured N₂O mole fraction and $[N_2O]_x$ is the corrected value. The slope and y-intercept, as well as their 95% confidence intervals, are represented by a_1 and a_2 respectively

25 The entire NOAA reference gas dilution range for N₂O (32 - 321 ppb) results in a deviation that can be adequately modeled using a 3rd degree polynomial. The linear fit discussed above is indistinguishable from the full 3rd degree polynomial over the N₂O concentration range of the OHSU-PSU Air Archive. However, if measuring N₂O samples with a difference of more than 80 ppb compared to the NOAA reference gas, the full 3rd degree polynomial is necessary to correct for the non-linear response in the μECD.

30 For SF₆, the prepared sample range over which the linear correction is applied is 0.59 – 8.76 ppt, most relevant for this work. The slope of the SF₆ linear fit is 0.971 ± 0.017 (95% CI) and is a good model for the deviation from expected over this range ($R^2 = 0.9995$). This results in a deviation from expected of ~ 0.03 ppt ppt⁻¹ SF₆ difference from the NOAA reference when measuring samples. All SF₆ measurements are adjusted for a linear correction of the form:

$$[SF_6]_x = b_1[SF_6]_y + b_2 \quad (4)$$



$$b_1 = 1.03 \pm 0.018 \text{ (95\% CI)} \quad (5)$$

$$b_2 = -0.297 \pm 0.099 \text{ (95\% CI)} \quad (6)$$

Where $[SF_6]_Y$ is the measured SF_6 mole fraction and $[SF_6]_X$ is the corrected value. The slope and y-intercept, as well as their 95% confidence intervals, are represented by b_1 and b_2 , respectively.

5 Detector response non-linearity has been evaluated in previous work by other groups on GC-ECD systems. For N_2O , deviations from expected of ~ 0.2 ppb ppb^{-1} difference from the reference gas are reported (Schmidt et al. 2001; Hall et al. 2007). These are similar to the value reported here for the μ ECD. Over larger ranges, a similar non-linear response curve is also reported. SF_6 non-linearity reported in Levin et al. (2010) has a similar curvature to the full N_2O non-linear response found here and previously discussed. Yet, this curvature is not observed to be significant over the range of SF_6 dilutions
10 conducted here.

3 Results and Discussion

3.1 Air archive mole fractions of N_2O and SF_6

Measurements of N_2O and SF_6 mole fraction from 159 samples of the OHSU-PSU Air Archive were initially filtered for analysis using a 7 median absolute deviation (7MAD) noise filter to remove far outliers. Polynomial fits (1st
15 degree for N_2O and 2nd degree for SF_6) were then applied to the data. Residual values outside of 2σ for N_2O and 3σ for SF_6 were removed for further data analysis. The entire process removed 12 data points for N_2O and 4 data points for SF_6 used in analysis.

Deseasonalized measurements of N_2O and SF_6 from Cape Meares are shown in figure 5a and 5b, respectively. A locally weighted linear regression (LOWESS) is used to smooth the data using a 3-year smoothing window (Cleveland and
20 Devlin 1988). The confidence intervals around regressions are calculated by bootstrapping residual variability 1000 times. The regression results in a concentration of 301.5 ± 0.3 ppb (1σ) in 1980 and increasing roughly linearly to the mid 1990s, where the concentration is 313.5 ± 0.3 ppb (1σ) in 1996.

Observations of N_2O mole fraction match well with previously published measurements of N_2O from Cape Meares between 1978 and 1998 of 301.2 ppb in 1980 and 313 - 314.5 ppb in 1996 on the SIO-1998 N_2O scale (Prinn et al. 1990;
25 Prinn et al. 2000; Khalil et al. 2002). The N_2O scale difference between SIO-1998 and NOAA-06 is minimal (Hall et al. 2007). Additional measurements by the Advanced Global Atmospheric Gases Experiment (AGAGE) and NOAA/ESRL (on the SIO-1998 N_2O and NOAA-06 N_2O scales, respectively) are reported from comparable sample locations. Trinidad Head, CA (41° N, 121° W), Mace Head, Ireland (53° N, 10° W), and Niwot Ridge, CO (40° N, 106° W) all measure ~ 313 ppb in 1996 (Prinn et al 2000; Hall et al. 2007). Together, these comparisons indicate the N_2O in the archived samples has stored
30 well.

Measured SF_6 concentration in archived Cape Meares samples is determined to be 0.85 ± 0.03 ppt (1σ) in 1980 and increases to a concentration of 3.83 ± 0.03 ppt (1σ) in 1996. Cape Meares does not have previously reported measurements



of SF₆ to compare with directly. Measurements of SF₆ from Trinidad Head, CA are reported to be ~0.85 ppt in 1980 and ~3.73 ppt in 1996 on the SIO-2005 SF₆ scale (Rigby et al. 2010). To convert to the NOAA-06 SF₆ scale, values measured on the SIO-2005 SF₆ scale are divided by a conversion factor of 0.9991 (Hall et al. 2014). In 1996, values of 3.87 ppt, 3.87 ppt, and 3.78 ppt are reported for Alert, Canada (82° N, 62° W), Barrow, AK (71° N, 157° W), and Niwot Ridge, CO 5 respectively on the NOAA-06 SF₆ scale (Hall et al. 2011). At these SF₆ concentrations, the difference between the NOAA-06 scale and the NOAA-14 scale is minimal. Cape Meares SF₆ measured values compare well with these northern hemisphere locations.

In the northern hemisphere, maximum background concentration measurements of SF₆ are reported from mid-to-high latitudes. For the year 1994, measurements from Fraserdale, Canada (50° N, 82° W) are reported to be 0.14 ppt higher 10 than samples measured from Izaña, Tenerife (28° N, 16° W) (Maiss et al. 1996). This difference is explained by the vast majority of SF₆ emissions coming from the mid-latitudes in the northern hemisphere (Maiss and Brenninkmeijer 1998; Levin et al. 2010; Rigby et al. 2010). The measured SF₆ concentrations from Cape Meares, also a midlatitude NH site, appear to fit in well with the expected meridional gradient when comparing to previously mentioned reported values.

Southern hemisphere measurements of SF₆ from archived atmospheric samples from Cape Grim, Tasmania (41° S, 15 145° E) and Neumayer, Antarctica (70° S, 8° W) are ~0.6 – 0.7 ppt in 1980 and ~3.4 – 3.5 ppt in 1996 on SIO-2005 and University of Heidelberg SF₆ scales (Levin et al. 2010, Rigby et al. 2010). As with the SIO-2005 SF₆ scale, the NOAA-06 and University of Heidelberg scale differences are small. To convert to the NOAA-06 SF₆ scale, values measured on the University of Heidelberg SF₆ scale are divided by a conversion factor of 0.9954 (Hall et al. 2014). Including a scale 20 correction, Cape Meares SF₆ measurements are higher than Cape Grim and Neumayer during this period by 0.2 - 0.4 ppt. Much or all of this difference can be explained by an interhemispheric north-south difference of 0.3 – 0.4 ppt (Levin et al. 2010).

3.2 Growth rate in N₂O and SF₆

The mean secular trend between 1978 and 1996 for N₂O and SF₆ is 0.78 ± 0.03 ppb yr⁻¹ (95% CI) and 0.17 ± 0.01 ppt yr⁻¹ (95% CI) respectively, determined by applying a linear fit to deseasonalized data. These trends translate to annual 25 increases of ~ 0.25% yr⁻¹ and ~ 10% yr⁻¹ for N₂O and SF₆, respectively. Annual trends for N₂O and SF₆ at Cape Meares, Oregon are determined from the derivative of the deseasonalized localized regression (Fig. 5 c&d). Uncertainty bands are generated from regressions of bootstrapped variability. Data points represent the mean annual trend with error bars equal to $\pm 1\sigma$ of the trend over the year.

The mean annual trend in N₂O (Fig. 5c) ranges between 0.6 ppb yr⁻¹ and 1.2 ppb yr⁻¹. All years between 1980 and 30 1996 show a positive rate of change significant at the 95% confidence level. The uncertainty in the annual trend is smallest in the early 1980s, at ± 0.15 ppb yr⁻¹ (95% CI), where there are largest numbers of data (~50% of samples are between 1980



and 1985). After 1985, uncertainty in the annual trend becomes ± 0.5 ppb yr⁻¹ (95% CI). This relatively large uncertainty results in an annual growth rate that is statistically indistinguishable between years.

A previously reported secular trend of N₂O reported between 1978 and 1998 for Cape Meares is 0.74 ± 0.02 ppb yr⁻¹, indistinguishable from our result (Prinn et al. 2000). The global secular trend of N₂O for the period 1985 to 1996 reported by Khalil et al. (2002) is 0.69 ± 0.03 ppb yr⁻¹, also compatible with our trend at Cape Meares.

The SF₆ annual trend (Fig. 5d) from the Cape Meares analysis increases from 0.07 ± 0.03 ppt yr⁻¹ (95% CI) in 1980 to 0.26 ± 0.05 ppt yr⁻¹ (95% CI) in 1994. The average rate of change in the growth rate (second derivative of mole fraction vs. time) over this period is 0.014 ppt yr⁻². The increase in growth rate over this period is statistically significant at high levels of confidence (95%). After 1994, we measure a decrease in the growth rate, though this decline is not statistically significant at high levels of confidence over this short time interval.

Comparable trends in SF₆ measured at other locations are available for the mid-1990s. The average global growth rate of SF₆ in 1994 was reported at 0.23 ppt yr⁻¹ in the northern hemisphere (Maiss et al. 1996). Alert, Canada and Izaña, Tenerife are observed to have maximum trends of 0.26 ppt yr⁻¹ in mid-1994 and at the beginning of 1995, respectively, compatible with results presented here (Levin et al. 2010). The localized maximum in growth rate in ~1994 observed here is present in some southern hemisphere observations of SF₆ at a similar time as well; Neumayer, Antarctica shows a maximum trend in 1995-1996 of 0.25 ppt yr⁻¹ (Levin et al. 2010). This finding is consistent with a peak in SF₆ emissions as reported by the European Database for Global Atmospheric Research (EDGAR, v4.2).

Another feature observed in the SF₆ trend from Cape Meares is a local maximum in the growth rate near 1987 (Fig. 5d). Notably however, not all data sets agree. The growth rate reported from Neumayer, Antarctica has this feature during a similar period (Levin et al. 2010), but the trend reported at Cape Grim, Tasmania does not show this local maximum (Rigby et al. 2010). Due to the large uncertainty from the few archived samples available during that time period, this local maximum is not statistically distinguishable from surrounding years at high levels of confidence in the Cape Meares analysis and this result is merely suggestive. Additional evidence is needed to corroborate this finding.

3.3 Seasonality in N₂O and SF₆ mole fraction

Seasonal behaviour for N₂O and SF₆ are shown in figure 6 determined from residuals to the secular trend. The N₂O seasonal cycle at Cape Meares shows a maximum near April and May of 0.3 ppb and an extended minimum from September through December of -0.4 ppb. Although there is considerable uncertainty surrounding monthly means, the difference between the spring maximum and fall minimum is statistically robust at high levels of confidence (2-sample KS test p-value = 0.003).

The seasonal amplitude matches well with previously reported northern hemisphere magnitudes of ± 0.4 ppb (Liao et al. 2004). Seasonal phase is also similar to Cape Meares at other mid-latitude northern hemisphere sites. N₂O seasonality reported at Mace Head, Ireland has a maximum near April and a minimum near August and September (Nevison et al. 2004;



Jiang et al. 2007) and Trinidad Head, CA seasonality has a maximum near late May and a broad minimum from September to January (Nevison et al. 2007).

In general, N₂O seasonal amplitude is known to vary strongly with latitude, e.g., 0.29 ppb at the South Pole (90° S, 102° W) and 1.15 ppb at Alert, Canada (Jiang et al. 2007). This is attributed in part to the stronger branch of the Brewer Dobson circulation in the northern hemisphere which also explains the high latitude minimums in late-summer months related to the influx of N₂O depleted air from the stratosphere during the spring (Liao et al. 2004; Nevison et al. 2004). Aside from atmospheric circulation, N₂O seasonality may also be influenced by regional sources. Leuker et al. (2003) suggested local maximums at Trinidad Head may reflect the influence of strong coastal upwelling. Similarly located in the Eastern Pacific, Cape Meares may also be subject to coastal upwelling influences. Isotopic analysis or modeling of transport effects and source influence would be useful to help interpret seasonal behaviour of N₂O at Cape Meares.

Seasonality for SF₆ shows a maximum between December and February of 0.04 ppt and a minimum near July of 0.03 ppt. The difference between the winter maximum and summer minimum is statistically significant (2-sample KS test p-value = 0.004). SF₆ seasonality has not previously been reported for Cape Meares.

Some seasonality in northern hemisphere observations of SF₆ is reported in the literature at select locations. Barrow, AK has a minimum in September and October with a broad maximum from December to June (Patra et al. 2009). Alert, Canada shows a strong minimum in October, though a maximum is not clearly defined (Wilson et al. 2014). Continental sites such as Niwot Ridge show large interannual variability (IAV) but have little distinguishable seasonality (Patra et al. 2009).

SF₆ seasonality at Cape Grim has been reported to have amplitude of ± 0.01 ppt with a maximum in September and October and a minimum in near February (Nevison et al. 2007; Wilson et al. 2014). The seasonal phase of Cape Grim is nearly anti-phase of Cape Meares reported here, though the amplitude is a factor of 4 smaller at Cape Grim. Similar to N₂O, seasonal amplitude is expected to be larger in the northern hemisphere than in the southern hemisphere (Nevison et al. 2007). Because sources of SF₆ are a-seasonal and sinks are essentially zero in the troposphere, the driving force behind the observed seasonality in SF₆ is considered to be atmospheric transport. Processes such as convection, vertical diffusion, boundary layer mixing, and shifts in the ITCZ can potentially influence the observed seasonality at a location. Modeling atmospheric transport effects on SF₆ at Cape Meares could help confirm amplitude and phase reported here.

4 Conclusions

We have measured 159 samples from the OHSU-PSU Air Archive collected at Cape Meares, Oregon (45.5° N, 124.0° W) for N₂O and SF₆ mole fraction using GC-μECD spanning April 1978 to December 1996. The GC-μECD system is designed to be fully automated, capable of running multiple pressurized samples per run. Measurement precision of N₂O and SF₆ is 0.16% and 1.1% respectively. Sample concentrations were corrected for detector response non-linearity when



measured against our reference gas. The linearity correction was found to be $0.14 \text{ ppb ppb}^{-1}$ and $0.03 \text{ ppt ppt}^{-1}$ for N_2O and SF_6 , respectively.

Analysis of archived air samples finds the mole fraction of N_2O in 1980 to be $301.5 \pm 0.3 \text{ ppb}$ (1σ) and rises to $313.5 \pm 0.3 \text{ ppb}$ (1σ) in 1996. The average growth rate over this period is $0.78 \pm 0.03 \text{ ppb yr}^{-1}$ (95% CI). Seasonality shows peak amplitude of 0.3 ppb near April and minimum amplitude of -0.4 ppb near November and is statistically robust. Our measurements of N_2O were found to match well with previously reported values for Cape Meares and other comparable northern hemisphere mid-latitude locations.

For SF_6 , the concentration in 1980 is found to be $0.85 \pm 0.03 \text{ ppt}$ (1σ), increasing to $3.83 \pm 0.03 \text{ ppt}$ (1σ) in 1996. The average growth rate over this period is $0.17 \pm 0.01 \text{ ppt yr}^{-1}$ (95% CI). Seasonality shows peak amplitude of 0.04 ppt near January and minimum amplitude of -0.03 ppt near July. There are no previous reported measurements of SF_6 from Cape Meares to compare against directly. SF_6 measurements compare well to other northern hemisphere measurements from Levin et al. (2010), Rigby et al. (2010), and Hall et al. (2011) over similar time periods when including spatial variability. From these N_2O and SF_6 measurements, we can conclude the sample integrity is robust within the OHSU-PSU Air Archive from Cape Meares, Oregon. Resulting dataset of SF_6 , in particular, contributes to a better characterization of historic SF_6 growth rate and its atmospheric variability over this period of dramatic growth.

Competing interests

The authors declare that they have no conflict of interest.

Author Contribution

Both authors (Terry Rolfe and Andrew Rice) worked closely together in the development and implementation of the GC technique used to make the mole fraction measurements of N_2O and SF_6 . Data curation and analysis was also completed by both authors. The original draft of this manuscript was prepared by Terry Rolfe with Andrew Rice responsible for the review and editing of the manuscript.

Acknowledgements

This study was supported by the US National Science Foundation (Atmospheric and Geospace Sciences grant 0952307). The authors would like to recognize Johnathan Radda for his assistance developing the GC- μECD technique used to measure the samples, R. A. Rasmussen for help establishing the OHSU-PSU air archive, and Christopher Butenhoff and M. A. K. Khalil for their assistance with data analysis and interpretation.



Data availability: A supplementary dataset of N₂O and SF₆ mole fractions at Cape Meares, Oregon measured for this work from the OSHU-PSU Air Archive are available to the scientific community (upon publication) and may be obtained by contacting the corresponding author.

References

- 5 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M. (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 10 Crutzen, P.: The influence of nitrogen oxides on the atmospheric ozone content, Quarterly Journal of the Royal Meteorological Society, 96(408), 320-325, 1970.
- Cleveland, W. S., and Devlin, S. J.: Locally weighted regression: An approach to regression analysis by local fitting, Journal of the American Statistical Association, 83(403), 596-610, 1988.
- 15 EDGAR, Emission Database for Global Atmospheric Research (EDGAR): European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL), release version 4.2., <http://edgar.jrc.ec.europa.eu>, 2013.
- Hall, B. D., Dutton, G. S., and Elkins, J. W.: The NOAA nitrous oxide standard scale for atmospheric observations. J. Geophys. Res.-Atmos., 112, D09305, doi:10.1029/2006JD007954, 2007.
- 20 Hall, B. D., Dutton, G. S., Mondeel, D. J., Nance, J. D., Rigby, M., Butler, J. H., Moore, F. L., Hurst, D. F., and Elkins, J. W.: Improving measurements of SF₆ for the study of atmospheric transport and emissions, Atmos. Meas. Tech., 4(11), 2441-2451, 2011.
- Hall, B. D., Engel, A., Mühle, J., Elkins, J. W., Artuso, F., Atlas, E., Aydin, M., Blake, D., Brunke, E.-G., Chiavarini, S., Fraser, P. J., Happell, J., Krummel, P. B., Levin, I., Loewenstein, M., Maione, M., Montzka, S. A., O'Doherty, S., Reimann, S., Rhoderick, G., Saltzman, E. S., Scheel, H. E., Steele, L. P., Vollmer, M. K., Weiss, R. F., Worthy, D., and Yokouchi, Y.: Results from the International Halocarbons in Air Comparison Experiment (IHALACE), Atmos. Meas. Tech., 7(2), 469-490, doi:10.5194/amt-7-469-2014, 2014.
- 25 Ishijima, K., Sugawara, S., Kawamura, K., Hashida, G., Morimoto, S., Murayama, S., Aoki, S., and Nakazawa, T.: Temporal variations of the atmospheric nitrous oxide concentration and its $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ for the latter half of the 20th century reconstructed from firn air analyses. J. Geophys. Res.-Atmos., 112(D3), 2007.
- 30 Jiang, X., Ku, W. L., Shia, R.-L., Li, Q., Elkins, J. W., Prinn, R. G., and Yung, Y. L.: Seasonal cycle of N₂O: Analysis of data, Global Biogeochem. Cy., 21, GB1006, doi:10.1029/2006GB002691, 2007.



- Khalil, M. A. K., and Rasmussen, R. A.: Increase and seasonal cycles of nitrous oxide in the earth's atmosphere. *Tellus*, 35B, 161-169, 1983.
- Khalil, M. A. K., Rasmussen, R. A., and Shearer, M. J.: Atmospheric nitrous oxide: patterns of global change during recent decades and centuries, *Chemosphere*, 47, 807–821, 2002.
- 5 Levin, I. and Hesshaimer, V.: Refining of atmospheric transport model entries by the globally observed passive tracer distributions of 85krypton and sulfur hexafluoride (SF₆), *J. Geophys. Res.-Atmos.*, 101(D11), 16745-16755, doi:10.1029/96JD01058, 1996.
- Levin, I., Naegler, T., Heinz, R., Osusko, D., Cuevas, E., Engel A., Ilmberger, J., R. Langenfelds, R. L., Neining, B., Rohden, C. v., Steele, L. P., Weller, R., Worthy, D. E., and Zimo, S. A.: The global SF₆ source inferred from long-
10 term high precision atmospheric measurements and its comparison with emission inventories, *Atmos. Chem. and Phys.*, 10, 2655–2662, 2010.
- Liao, T., Camp, C., and Yung, Y.: The seasonal cycle of N₂O, *Geophys. Res. Lett.*, 31, L17108, doi:10.1029/2004GL020345, 2004.
- 15 Lueker, T., Walker, S., Vollmer, M., Keeling, R., Nevison, C., Weiss, R., and Garcia, H: Coastal upwelling air-sea fluxes revealed in atmospheric observations of O₂/N₂, CO₂, and N₂O, *Geophys. Res. Lett.*, 30(6), 1292, doi:10.1029/2002GL016615, 2003.
- Maiss, M. and Brenninkmeijer, C.: Atmospheric SF₆: Trends, sources, and prospects, *Environ. Sci. Technol.*, 32(20), 3077–3086, 1998.
- Maiss, M., Steele, L. P., Francey, R., Fraser, P., Langenfelds, R., Trivett, N., and Levin, I.: Sulfur hexafluoride – a powerful
20 new atmospheric tracer, *Atmos. Environ.*, 30(10), 1621–1629, doi:10.1016/1352-2310(95)00425-4, 1996.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O'Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse gas concentrations for climate modelling (CMIP6). *Geosci. Model
25 Dev.*, 10(5), 2057-2116, 2017.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., Qin, D., Plattner, G.-K.,
30 Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M. (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Nevison, C., Kinnison, D., and Weiss, R.: Stratospheric influences on the tropospheric seasonal cycles of nitrous oxide and chlorofluorocarbons, *Geophys. Res. Lett.*, 31(20), doi:10.1029/2004GL020398, 2004.



- Nevison, C., Mahowald, N., Weiss, R., and Prinn, R.: Interannual and seasonal variability in atmospheric N₂O, *Global Biogeochem. Cy.*, 21(3), doi:10.1029/2006GB002755, 2007.
- Olivier, J., Van Aardenne, J., Dentener, F., Pagliari, V., Ganzeveld, L., and Peters, J.: Recent trends in global greenhouse gas emissions: regional trends 1970–2000 and spatial distribution of key sources in 2000, *Environ. Sci.*, 2(2-3), 81-99, doi:10.1080/15693430500400345, 2005.
- 5 Park, S., Croteau, P., Boering, K. A., Etheridge, D. M., Ferretti, D., Fraser, P. J., Kim, K.-R., Krummel, P. B., Langenfelds, R. L., van Ommen, T. D., Steele, L. P., Trudinger, C. M.: Trends and seasonal cycles in the isotopic composition of nitrous oxide since 1940, *Nat. Geosci.*, 5(4), 261-265, doi:10.1038/NGEO1421, 2012.
- Patra, P. K., Takigawa, M., Dutton, G. S., Uhse, K., Ishijima, K., Lintner, B. R., Miyazaki, K., and Elkins, J. W.: Transport mechanisms for synoptic, seasonal and interannual SF₆ variations and “age” of air in troposphere, *Atmos. Chem. and Phys.*, 9, 1209–1225, 2009.
- 10 Prather, M. J., Holmes, C. D., and J. Hsu, J.: Reactive green house gas scenarios: Systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39, L09803, doi:10.1029/2012GL051440, 2012.
- Prather, M. J., Hsu, J., DeLuca, N. M., Jackman, C. H., Oman, L. D., Douglass, A. R., Fleming, E. L., Strahan, S. E., Steenrod, S. D., Søvdé, O. A., Isaksen, I. S. A., Froidevaux, L., and Funke, B.: Measuring and modeling the lifetime of nitrous oxide including its variability, *J. Geophys. Res.-Atmos.*, 120(11), 5693-5705, doi:10.1002/2015JD023267, 2015.
- 15 Prinn, R., Cunnold, D., Rasmussen, R., Simmonds, P., Alyea, F., Crawford, A., Fraser, P., and Rosen, R.: Atmospheric emissions and trends of nitrous oxide deduced from 10 years of ALE-GAGE data, *J. Geophys. Res.*, 95(D11), 18369-18385, 1990.
- 20 Prinn, R., Weiss, R., Fraser, P., Simmonds, P., Cunnold, D., Alyea, F., O’Doherty, S., Salameh, P., Miller, B. R., Huang, J., Wang, R. H. J., Hartley, D. E., Harth, C., Steele, L. P., Sturrock, G., Midgley, P. M., and McCulloch, A.: A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *J. Geophys. Res.*, 105(D14), 17751-17792, doi:10.1029/2000JD900141, 2000.
- 25 Ravishankara, A., Solomon, S., Turnipseed, A., and Warren, R.: The atmospheric lifetimes of long-lived halogenated species, *Science*, 259, 194–199, 1993.
- Rice, A. L., Butenhoff, C. L., Teama, D. G., Röger, F. H., M. Khalil, M. A. K., and Rasmussen, R. A.: Atmospheric methane isotopic record favors fossil sources flat in 1980s and 1990s with recent increase, *P. Natl. Acad. Sci. USA*, 113(39), 10791-10796, 2016.
- 30 Rigby, M., Muhle, J., Miller, B. R., Prinn, R. G., Krummel, P. B., Steele, L. P., Fraser, P. J., Salameh, P. K., Harth, C. M., Weiss, R. F., Grealley, B. R., O’Doherty, S., Simmonds, P. G., Vollmer, M. K., Reimann, S., Kim, J., Kim, K.-R., Wang, H. J., Olivier, J. G. J., Dlugokencky, E. J., Dutton, G. S., Hall, B. D., and Elkins, J. W.: History of atmospheric SF₆ from 1973 to 2008, *Atmos. Chem. and Phys.*, 10, 10305–10320, 2010.



- Saikawa, E., Prinn, R. G., Dlugokencky, E., Ishijima, K., Dutton, G. S., Hall, B. D., Langenfelds, R., Tohjima, Y., Machida, T., Manizza, M., Rigby, M., O'Doherty, S., Patra, P. K., Harth, C. M., Weiss, R. F., Krummel, P. B., van der Schoot, M., Fraser, P. J., Steele, L. P., Aoki, S., Nakazawa, T., and Elkins, J. W.: Global and regional emissions estimates for N₂O, *Atmos., Chem., and Phys.*, 14, 4617–4641, 2014.
- 5 Schmidt, M., Glatzel-Mattheier, H., Sartorius, H., Worthy, D. E., and Levin, I.: Western European N₂O emissions: A top-down approach based on atmospheric observations, *J. Geophys. Res.*, 106(D6), 5507-5515, 2001.
- Snider, D. M., Venkiteswaran, J. J., Schiff, S. L., Spoelstra, J.: From the ground up: Global nitrous oxide sources are constrained by stable isotope values, *PLOS ONE*, 10(3), doi:10.1371/journal.pone.011895, 2015.
- Stocker, B. D., Roth, R., Joos, F., Spahni, R., Steinacher, M., Zeahle, S., Bouwman, L., and Prentice, I. C.: Multiple
10 greenhouse-gas feedbacks from the land biosphere under future climate change scenarios, *Nat. Clim. Change*, 3(7), 666-672, doi:10.1038/nclimate1864, 2013.
- Sulfur Hexafluoride (SF₆) WMO Scale: https://www.esrl.noaa.gov/gmd/ccl/sf6_scale.html, last access: 17 June 2018.
- Wilson, C., Chipperfield, M. P., Gloor, M., and Chevallier, F.: Development of a variational flux inversion system
15 (INVICAT v1.0) using the TOMCAT chemical transport model, *Geosci. Model Dev.*, 7(5), 2485-2500, 2014.




Figure 1. Schematic view of the analytical system for sample evaluation. The system is shown in “back-flush” mode. V_1 = Valve 1, V_2 = Valve 2, V_3 = Valve 3.




Figure 2. Sample chromatogram showing N_2O peak at a retention time of 6.1 minutes and SF_6 peak at a retention time of 7.0 minutes. Upper-right corner inlay shows an enlarged plot of the SF_6 peak.




Figure 3. Precision in measurement for N_2O (a) and SF_6 (b) expressed as percent relative standard deviation from 30 sets of 6 measurements of the NOAA reference gas. The black dotted line represents a normal distribution curve with the same mean and standard deviation. The standard deviation for N_2O and SF_6 is 0.16% and 1.1%, respectively.




Figure 4. Measurement linearity from plots of measured mole fraction vs. expected mole fraction of N₂O (a) and SF₆ (b). Expected mole fraction is calculated from the NOAA reference mole fraction (328.71 ppb N₂O and 8.76 ppt SF₆) after dilution with ultra-pure air. Error bars represent 1 σ total uncertainty.




Figure 5. Deseasonalized measurements of mole fraction versus date of collection, N_2O (a) and SF_6 (b), and annual trends in time from Cape Meares, Oregon, N_2O (c) SF_6 (d). Error bars are 1σ uncertainty. The solid black lines are LOWESS fit to the data using a smoothing window of 3 years and shaded areas are 95% confidence intervals in the LOWESS fit calculated from bootstrapping residual variability 1000 times.




Figure 6. Seasonality for N₂O (a) and SF₆ (b) calculated from the residuals of observed data points to the secular trend. The black line is a LOWESS fit to residuals with a smoothing window of 1 month. Data points show observed monthly mean residual after binning by month with error bars representing standard error within the month. Shaded areas are 95% CI
5 calculated from 1000 bootstrapped LOWESS fits while including the measurement uncertainty to each data point.



Table 1. Characteristics of 12 manometric N₂O dilution samples prepared at Portland State University.

Canister ID	^a . P _{Ref} (kPa)	^b . P _{Total} (kPa)	^c . Expected N ₂ O Response	^d . Measured N ₂ O Response	^e . Measured N ₂ O (ppb)	^f . N ₂ O 1σ (ppb)
1.7	97.0	132.4	0.7327	0.7673	252.21	0.78
1.14	36.6	132.2	0.2767	0.3262	107.22	0.45
1.5	61.5	132.9	0.4627	0.5157	169.52	0.51
2.14	21.0	132.6	0.1585	0.1941	63.79	0.27
2.7	13.0	132.5	0.0978	0.1239	40.73	0.44
2.5	80.5	132.2	0.6092	0.6535	214.80	0.52
3.7	127.0	132.8	0.9559	0.9618	316.15	0.83
3.5	117.0	132.8	0.8813	0.8981	295.20	0.87
3.14	123.8	132.7	0.9326	0.9423	309.75	0.85
4.5	129.7	132.7	0.9778	0.9813	322.56	0.96
4.14	119.1	132.9	0.8959	0.9085	298.62	0.71
4.7	120.9	132.5	0.9129	0.9226	303.26	0.80

- a.** P_{Ref} is the NOAA reference gas pressure (in kPa) introduced to the canister.
b. P_{Total} is the final pressure (in kPa) of the canister after balancing with ultra-pure air.
5 c. Expected response is calculated from the P_{Ref}/P_{Final} fraction.
d. Measured N₂O response of the μECD.
e. Measured N₂O in ppb.
f. N₂O 1σ (ppb) is from combined uncertainty of sample and surrounding NOAA reference.

10 Table 2. Characteristics of 9 manometric SF₆ dilution samples prepared at Portland State University.

Canister ID	^a . P _{Ref} (kPa)	^b . P _{Scotty} (kPa)	^c . P _{Total} (kPa)	^d . Expected SF ₆ Response	^e . Measured SF ₆ Response	^f . Measured SF ₆ (ppt)	^g . SF ₆ 1σ (ppt)
1.14	97.0	-	132.4	0.7327	0.7476	6.55	0.12
3.5	117.0	-	132.8	0.8812	0.8943	7.83	0.24
3.14	123.8	-	132.7	0.9326	0.9414	8.25	0.10
1.1	29.6	31.0	132.6	0.2230	0.2443	2.14	0.06
1.18	11.9	36.9	133.0	0.0896	0.1199	1.05	0.07
1.28	8.9	37.6	132.3	0.0674	0.1153	1.01	0.06
2.1	18.7	34.8	131.9	0.1418	0.1644	1.44	0.08
2.18	75.9	16.0	132.2	0.5740	0.5879	5.15	0.15
2.28	52.9	23.5	132.1	0.4002	0.4110	3.60	0.12

- a.** P_{Ref} is the NOAA reference gas pressure (in kPa) introduced to the canister.
b. P_{Scotty} is the 1 ppm N₂O balanced with He (in kPa) introduced to the canister.
c. P_{Total} is the final pressure (in kPa) of the canister after balancing with ultra-pure air.
15 d. Expected SF₆ response is calculated from the P_{Ref}/P_{Final} fraction.
e. Measured SF₆ response of the μECD.



f. Measured SF₆ in ppt.

g. SF₆ 1σ (ppt) is from combined uncertainty of sample and surrounding NOAA reference.