

Editor comments on ACP-2019-1169, 2nd revision of Kuilman et al.

The abstract and the manuscript reads now much better.

We thank the editor for her positive feedback and for her further comments.

On the abstract I have the following two comments (P1, L46):

- (1) Abbreviation SST has not been introduced.
- (2) Further, it has not been mentioned anywhere yet that changed SSTs have been considered. Either you add a sentence here or you add that the SSTs have been changed as well in line 32 where you write what is done in this study.

This has been implemented:

"In this study, we investigate the response of the middle atmosphere to a doubling of the CO₂-concentration and the associated changes in sea surface temperatures (SSTs) using the Whole Atmosphere Community Climate Model (WACCM)."

On the main text I have the following comments:

P5, L218: What do you mean with the latter is not considered in this study? No chemistry is considered!? Please rephrase the sentence to be clearer.

This sentence has been rephrased:

"The SSTs might be slightly different when they would be generated using a model that also includes atmospheric chemistry, however, this aspect is not considered in this study."

P5, L229-230: This sentence is also very weird. Why do they do not play a role in your experiments? The question from the referee was quite simple and could have been answered with yes or no. Either CFCs are considered or not. Since you use pre-industrial conditions I would assume that these are either zero or low since to my knowledge anthropogenic production (and thus the increase) of CFCs started later than 1850. This is something which can easily be checked. So please check and rephrase the sentence accordingly.

It has been checked the CFC are zero (there are some values 10^{-22} mol/mol, but mostly zeros), this also doesn't change between the control run and the perturbation runs. So, the short answer to the reviewer's question would be no, these are not considered.

The text has now been rewritten:

"The compset used in this experiment and all the following ones is still F_1850, which means that other radiatively and chemically active gases, such as ozone, will change only because of the changes in the CO₂-concentration, due to WACCM's interactive chemistry. This also means that the effects of chlorofluorocarbons (CFCs) are not considered in our experiments, as anthropogenic production of CFCs started later than 1850."

P6, Table 1: The naming in the table is quite misleading and I needed a while to understand how your four experiments differentiate from each other. I would suggest to do the following change (and simplification) to the table to be clearer:

Experiment CO2 SSTs

C1 PI PI

C2 double PI

S1 PI double (or high)

S2 double double (or high)

The simplification has been implemented.

Further, it would be more logical if the naming would be C1, S1, S2 and S3 since you have one control run and three scenarios.

This has been implemented.

P7, L298-312: This text part is also quite difficult to read. I would suggest that you repeat here the sentences from the appendix what R, S and the deltas are.

This has been implemented.

"In which \vec{R} represents the vertical profile of the net long-wave radiation emitted by each layer in the atmosphere and by the surface. \vec{S} is the vertical profile of the solar radiation absorbed by each layer. The matrix $\left(\frac{\partial \vec{R}}{\partial \vec{T}}\right)$ is the Planck feedback matrix, in which the vertical profiles of the changes in the divergence of radiative energy fluxes due to a temperature change are represented. ΔT represents the temperature change."

Further, I would suggest to move the second paragraph to the appendix or to make a list with bullets for CO2, H2O, O3, cloud and Albedo where then is written what has been done for each species or you just simplify the text. In general the difference between control and experiment is considered, so just mention then what is the difference for the species. In your text especially the phrase "the other variables" is not clear and rather confusing.

The authors agree that the former formulation was confusing. The following has been added to the appendix:

The factors $\Delta(\vec{S} - \vec{R})_{CO_2}$, $\Delta(\vec{S} - \vec{R})_{O_3}$, $\Delta(\vec{S} - \vec{R})_{H_2O}$, $\Delta(\vec{S} - \vec{R})_{albedo}$ and $\Delta(\vec{S} - \vec{R})_{cloud}$ in eqs (1-5) are calculated by inserting the output variables from WACCM in the radiation code of CFRAM. Here, one takes the output variables from the control run, apart from the variable that is related to the direct forcing or the feedback. The table below shows which variables have been taken from the perturbation runs for each feedback.

Direct forcing/feedback	Changed variables in the radiation code
CO ₂	CO ₂
Ozone	O ₃
Water vapour	Specific humidity Surface pressure Surface temperature Dew point temperature
Albedo	Downwelling solar flux at surface

	<i>Net solar flux at surface</i>
<i>Cloud</i>	<i>Cloud fraction Cloud ice Cloud liquid amount</i>

Table A1: The variables from the perturbation runs inserted in the radiation code of CFRAM to calculate the temperature change in response to the changes in CO₂, O₃, water vapour, cloud and albedo.

The main text now refers to the appendix in the following way:

Here, one takes the output variables from the control run, apart from the variable that is related to the direct forcing or the feedback. Table A1 in the Appendix shows which variables from the perturbation runs have been inserted in the radiation code of CFRAM in order to calculate $\Delta(\vec{S} - \vec{R})_{CO_2}$, $\Delta(\vec{S} - \vec{R})_{O_3}$, $\Delta(\vec{S} - \vec{R})_{H_2O}$, $\Delta(\vec{S} - \vec{R})_{albedo}$ and $\Delta(\vec{S} - \vec{R})_{cloud}$ and eventually the associated temperature changes.

P9, 392: add “(C1)” after control simulation.

This has been added (also for the other simulations):

“Figure 1: The total change in temperature in July (top) and January (bottom) for (a,d) the simulation with high CO₂ and SSTs (S3), (b,e) the simulation with high CO₂ (S1), (c,f) the simulation with high SSTs (S2), all as compared to the pre-industrial control simulation (C1). The dotted regions indicate the regions where the data reaches a confidence level of 95%. The black line indicates the tropopause height for the experiments S3 (a,d), S1 (b,e) and S2 (c,f).”

P14, Figure 5 caption: Shouldn't it read “S1 - C1” and “C2 - C1”, respectively? The same holds for Fig7, Fig 9, Fig. 10, Fig 11 and Fig 13.

Thanks for this comment, this was indeed incorrect. It has now been corrected.

P18, L697: Be more precise than “higher up”. Above which level?

This has been clarified:

“Fig. 8 shows that above 1 hPa, there are also large percentage changes in water vapour. However, the absolute concentration of water vapour is small there, which explains why there is no temperature response to these changes.”

P24, L870: Sentence not clear. Please rephrase. Which exact mechanism are you talking about? What exactly is out of the scope of the paper? Do you mean to investigate the exact mechanism why the non-LTE effects are small is outside the scope of the paper?

This has been rewritten:

“We find that there are also some small temperature changes due to non-LTE effect above 0.1 hPa. How the non-LTE effects exactly cause the small temperature changes in this region is outside the scope of this paper and needs further investigation.”

P26, L897: Which errors? Do you mean the bars in the bar chart of Figure 13?

Yes, that is correct. I have made this clearer in the text.

"As in Fig. 11, the 'Error'-column in Fig. 13 shows the difference between temperature change in WACCM and the sum of the calculated temperature responses in CFRAM."

We have also clarified this bit in the description in Fig. 11:

"This term shows the difference between temperature change in WACCM and the sum of the calculated temperature responses in CFRAM (see eq. 9 in section 2.3)."

P30, L1122: What is the PRP method? Please add more details.

This method and abbreviation had introduced in the introduction. Now more details have been given at this point in the paper as well:

"This is also commonly done in the partial radiative perturbation (PRP) method, in which partial derivatives of the model top of the atmosphere radiation are evaluated with respect to changes in model parameters by diagnostic rerunning the model's radiation code (Bony et al., 2006)."

Other technical issues:

P2, L64: the -> this
This has been corrected.

P3, L122: these -> "the" or "the respective"
This has been corrected.

P3, L126: use plural, thus "forcing and feedbacks"
This has been corrected.

P3, L143: such idealized -> such "an" idealized
This has been corrected.

P6, L264: effect -> affect
This has been corrected.

P9, L391: add "to" before "the"
This has been corrected.

P10, L433: Past should be used here: "In this section, we have discussed....."
It was meant to be in the current section (4), we will discuss. This has now been changed in the text to make it clearer:

"In the following subsections 4.1-4.5, we will discuss the meridional-vertical profiles of the temperature responses to the direct forcing and the various feedbacks during

July and January. In section 5, we will discuss regional and global means of partial temperature changes due to feedbacks."

P10, L444: add "is found" after CO₂-concentration.
This has been added.

P11, L452: singular should be used here -> run
This has been corrected.

P13, L536: add "by" so that it reads "could be caused by an increase...."
This has been corrected.

P14, L545: add "the" so that it reads "...due to the.....".
This has been corrected.

P15, L589: add "scale" so that it reads ".....synoptic scale waves"
This has been added.

P16, L645: add "concentration" after CO₂.
This has been added.

P17, L652: add "to" before "the" so that it reads "...with respect to the....."
This has been added.

P18, L698: add "vapour" so that it reads "water vapour".
This has been added.

P20, L759: stratosphere -> stratospheric
This has been corrected.

P20, L760: until -> up to
This has been changed.

P21, L771: add "average" so that it reads "by taking the average in this way".
This has been changed:

"By calculating the average in this way, we can directly compare the vertical values in different regions of the atmosphere."

P22, L814: add "is" so that it reads "is a bit smaller".
This has been added.

P22, L822-824: "before" appears in the sentence twice. I would suggest to delete the latter one.
The latter 'before' has been removed.

P22, L832: move "mostly" before "contributes" so that it reads "which mostly contributes...".
'Mostly' has been moved.

P24, L874: “for due” is not correct, this “for” is obsolete here. Please remove.
'For' has been removed.

P26, L890: add “the” so that it reads “shows the”
'The' has been added.

P26, L931-932: “before” twice. One is therefore obsolete. I would suggest to remove the latter one.
'Before' has been removed.

Using the climate feedback response method to quantify climate feedbacks in the middle atmosphere in WACCM

Maartje Sanne Kuilman¹, Qiong Zhang², Ming Cai³, Qin Wen^{1,4}

1. Department of Meteorology and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden
2. Department of Physical Geography and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden
3. Department of Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee, Florida, USA
4. Laboratory for Climate and Ocean-Atmosphere Studies (LaCOAS), Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, China

Corresponding author: Maartje Sanne Kuilman (maartje.kuilman@misu.su.se)

Abstract

Over recent decades it has become clear that the middle atmosphere has a significant impact on surface and tropospheric climate. A better understanding of the middle atmosphere and how it reacts to the current increase of the concentration of carbon dioxide (CO₂) is therefore necessary. In this study, we investigate the response of the middle atmosphere to a doubling of the CO₂-concentration and the associated changes in sea surface temperatures (SSTs) using the Whole Atmosphere Community Climate Model (WACCM). We use the climate feedback response analysis method (CFRAM) to calculate the partial temperature changes due to an external forcing and climate feedbacks in the atmosphere. As this method has the unique feature of additivity, these partial temperature changes are linearly addable. In this study, we discuss the direct forcing of CO₂ and the effects of the ozone, water vapour, cloud, albedo and dynamical feedbacks. As expected, our results show that the direct forcing of CO₂ cools the middle atmosphere. This cooling becomes stronger with increasing height: the cooling in the upper stratosphere is about three times as strong as the cooling in the lower stratosphere. The ozone feedback yields a radiative feedback that mitigates this cooling in most regions of the middle atmosphere. However, in the tropical lower stratosphere and in some regions of the mesosphere, the ozone feedback has a cooling effect. The increase in CO₂-concentration causes the dynamics to change. The temperature response due to this dynamical feedback is small in the global average, although there are large temperature changes due to this feedback locally. The temperature change in the lower stratosphere is influenced by the water vapour feedback and to a lesser degree by the cloud and albedo feedback. These feedbacks play no role in the upper stratosphere and the mesosphere. We find that the effects of the changed SSTs on the middle atmosphere are relatively small as compared to the effects of changing the CO₂. However, the changes in SSTs

Deleted: to a doubling CO₂ in

are responsible for dynamical feedbacks that cause large temperature changes. Moreover, the temperature response to the water vapour feedback in the lower stratosphere is almost solely due to changes in the SSTs. As CFRAM has not been applied to the middle atmosphere in this way before, this study also serves to investigate the applicability as well as the limitations of this method. This work shows that CFRAM is a very powerful tool to study climate feedbacks in the middle atmosphere. However, it should be noted that there is a relatively large error term associated with the current method in the middle atmosphere, which can be for a large part be explained by the linearization in the method.

1. Introduction

The increase of concentration of carbon dioxide in the atmosphere forms a major perturbation to the climate system. It is commonly associated with lower-atmospheric warming. However, in the middle atmosphere, the increase of CO₂ leads to a cooling of **this** region instead. This cooling has been well documented and is found by both model studies and observations (e.g. Manabe and Wetherald, 1975; Ramaswamy *et al.*, 2001; Beig *et al.*, 2003).

Deleted: the

The middle atmosphere is not only affected by the increase in CO₂-concentration, but also by the decrease in ozone-concentration. The depletion of ozone (O₃) also effects the temperature in the stratosphere and leads to a cooling (Shine *et al.*, 2003). A better understanding of the effect of the increased CO₂-concentration on the middle atmosphere, will help to distinguish the effects of the changes CO₂- and O₃-concentration.

Another major motivation for this study is the emerging evidence that the middle atmosphere has an important influence on surface and tropospheric climate (Shaw and Shepherd, 2008). It has, for example, been shown that cold winters in Siberia are linked to changes in the stratospheric circulation (Zhang *et al.*, 2018).

Nowack *et al.* (2015) has found that there is an increase in global mean surface warming of about 1°C when the ozone is prescribed at pre-industrial levels, as compared with when it is evolving in response to an abrupt 4xCO₂ forcing. It should be noted that the exact importance of changes in ozone seems to be dependent on both the model and the scenario (Nowack *et al.*, 2015) and is not found by all studies (Marsh *et al.*, 2016).

As the effect is found to be rather large in some studies, and absent in other, there is a need for a better understanding of the behaviour of the middle atmosphere in response to changing CO₂ conditions, as the ozone concentration is influenced by this. Ozone is an example of a climate feedback, a process that changes in response to a change in CO₂-concentration and in turn dampens or amplifies the climate response to the CO₂ perturbation.

These climate feedbacks are a challenging subject of study, as observed climate variations might not be in equilibrium, multiple processes are

operating at the same time and moreover the geographical structures and timescales of different forcings differ. However, feedbacks form a crucial part of understanding the response of the atmosphere to changes in the CO₂-concentration.

Various methods have been developed to study these feedbacks, such as the partial radiative perturbation (PRP) method, the online feedback suppression approach and the radiative kernel method (Bony *et al.*, 2006 and the references therein). These methods study the origin of the global climate sensitivity (Soden and Held, 2006; Caldwell *et al.*, 2016; Rieger *et al.*, 2017). The focus of these methods is on changes in the global mean surface temperature, global mean surface heat and global mean sensible heat fluxes (Ramaswamy *et al.*, 2019).

These methods are powerful for this purpose; however, they are not suitable to explain temperature changes on spatially limited domains. They neglect non-radiative interactions between feedback processes and they only account for feedbacks that directly affect the radiation at the top of the atmosphere (TOA).

The climate feedback-response analysis method (CFRAM) is an alternative method which takes into account that the climate change is not only determined by the energy balance at the top of the atmosphere, but is also influenced by the energy flow within the Earth's system itself (Cai and Lu, 2009, Lu and Cai, 2009). The method is based on the energy balance in an atmosphere-surface column. It solves the linearized infrared radiation transfer model for the individual energy flux perturbations. This makes it possible to calculate the partial temperature changes due to an external forcing and the internal feedbacks in the atmosphere. It has the unique feature of additivity, such that these partial temperature changes are linearly addable.

As a practical diagnostic tool to analyse the role of various forcings and feedbacks, CFRAM has been used widely in climate change research on studying surface climate change (Taylor *et al.*, 2013; Song and Zhang, 2014; Hu *et al.*, 2017; Zheng *et al.*, 2019). CFRAM has been applied to study the middle atmosphere climate sensitivity as well (Zhu *et al.*, 2016). In their study, Zhu *et al.* (2016) have adapted CFRAM and applied it to both model output, as well as observations. The atmospheric responses during solar maximum and minimum were studied and it was found that the variation in solar flux forms the largest radiative component of the middle atmosphere temperature response.

In the present work, we apply CFRAM to climate sensitivity experiments performed with the Whole Atmosphere Community Climate Model (WACCM), which is a high-top global climate system model, including the full middle atmosphere chemistry.

We investigate the middle atmosphere response to CO₂-doubling. We acknowledge that such an idealized equilibrium simulation cannot reproduce the complexity of the atmosphere, in which the CO₂-concentration is changing

Deleted: these

Deleted: forcing

Deleted: feedback

gradually. However, simulating a double CO₂-scenario still allows us to identify robust feedback processes in the middle atmosphere.

There are two aspects of the middle atmosphere response to CO₂-doubling: there is the effect of the changes in CO₂-concentration directly, as well as the changes in sea surface temperature (SST) which are in itself caused by the changes in CO₂-concentration. It is useful to investigate these aspects separately, as former should be robust, while the effect of the changed SST depends on the changes in tropospheric climate, which can be expected to depend more on the model.

In this study, we investigate the effects of doubling the CO₂-concentration and the accompanying sea surface temperature change on the temperature in the middle atmosphere as compared to the pre-industrial state. We use CFRAM to calculate the radiative contribution to the temperature change due to changes in carbon dioxide directly as well as due to changes in ozone, water vapour, albedo, and clouds. We refer to the changes in ozone, water vapour, albedo and clouds in response to changes in the CO₂-concentration as the ozone, water vapour, albedo and cloud feedbacks.

Deleted: ,

The circulation in the middle atmosphere is driven by waves. Wave forcing drives the temperatures in the middle atmosphere far away from radiative equilibrium. In the mesosphere, there is a zonal forcing, which yields a summer to winter transport. In the polar winter stratosphere, there is a strong forcing that consists of rising motion in the tropics, poleward flow in the stratosphere and sinking motion in the middle and high latitudes. This circulation is referred to as the 'Brewer-Dobson circulation' (Brewer, 1949; Dobson, 1956).

Dynamical effects make important contributions to the middle-atmosphere energy budget, both through eddy heat flux divergence and through adiabatic heating due to vertical motions. It is therefore important that we also consider changes to the middle-atmosphere climate due to dynamics. We refer to this as the 'dynamical feedback' (Zhu *et al.*, 2016).

The main goal of this paper is to calculate the contribution to the temperature change due to changes in carbon dioxide directly as well as due to changes in ozone, water vapour, albedo, clouds and dynamics in the middle atmosphere under a double CO₂-scenario using CFRAM. Our intention is not to give a complete account of the exact mechanisms behind the changes in ozone, water vapour, albedo, clouds and dynamics.

2. The model and methods

2.1 Model description

The Whole Atmosphere Community Model (WACCM) is a chemistry-climate model, which spans the range of altitudes from the Earth's surface to about 140 km (Marsh *et al.*, 2013). The model consists of 66 vertical levels with irregular vertical resolution, which ranges from ~1.1 km in the troposphere,

198 1.1–1.4 km in the lower stratosphere, 1.75 km at the stratosphere and 3.5 km
199 above 65 km. The horizontal resolution is 1.9° latitude by 2.5° longitude.

200 WACCM is a superset of the Community Atmospheric Model version 4
201 (CAM4) developed at the National Center for Atmospheric Research (NCAR).
202 Therefore, WACCM includes all the physical parameterizations of CAM4
203 (Neale *et al.*, 2013), and a well-resolved high-top middle atmosphere. The
204 orographic gravity wave (GW) parameterization is based on *McFarlane*
205 (1987). WACCM also includes parameterized non-orographic GWs, which are
206 generated by frontal systems and convection (Richter *et al.*, 2010). The
207 parameterization of non-orographic GW propagation is based on the
208 formulation by Lindzen (1981).

209 The chemistry in WACCM is based on version 3 of the Model for Ozone and
210 Related Chemical Tracers (MOZART3). This model represents chemical and
211 physical processes from the troposphere until the lower thermosphere.
212 (Kinnison *et al.*, 2007). In addition, WACCM simulates chemical heating,
213 molecular diffusion and ionization and gravity wave drag.

214 2.2 Experimental set-up

215 In this study, the F_1850 compset (component set) of the model is used, i.e.
216 the model assumes pre-industrial (PI) conditions. This compset simulates an
217 equilibrium state, which means that it runs a perpetual year 1850. Four
218 experiments have been performed for this study (see Table 1).

219 Experiment C1 is the control run, with the pre-industrial CO₂ concentration
220 (280 ppm) and forced with pre-industrial ocean surface conditions such as
221 sea surface temperatures and sea ice. These SSTs are generated from the
222 CMIP5 pre-industrial control simulation by the fully coupled Earth system
223 model CESM. The atmospheric component of CESM is the same as WACCM,
224 but does not include stratospheric chemistry (Hurrell *et al.*, 2013). The SSTs
225 might be slightly different when they would be generated using a model that
226 also includes atmospheric chemistry, however, this aspect is not considered
227 in this study.

Deleted: temperature

Deleted: (referred to SSTs from now on).

Deleted: This latter

228 Experiment S1 represents the experiment with the CO₂ concentration doubled
229 as compared to the pre-industrial state (560 ppm) and forced with the same
230 pre-industrial SSTs as in experiment C1. In WACCM, the CO₂-concentration
231 does not double everywhere in the atmosphere. Only the surface level CO₂
232 mixing ratio is doubled, and elsewhere in the atmosphere is calculated
233 according to WACCM's chemical model.

Deleted: C2

234 The compset used in this experiment and all the following ones is still F_1850,
235 which means that other radiatively and chemically active gases, such as
236 ozone, will change only because of the changes in the CO₂-concentration,
237 due to WACCM's interactive chemistry. This also means that the effects of
238 chlorofluorocarbons (CFCs) are not considered in our experiments, as
239 anthropogenic production of CFCs started later than 1850.

Deleted: Chlorofluorocarbons

Deleted:), which have a major impact on the ozone concentration in the real atmosphere, don't play a role

247 In experiment S2, we simulate the scenario, in which there is the SSTs forcing
 248 from the coupled CESM for double CO₂ condition. This means that the sea
 249 surface temperatures are higher than in the PI run, and there is less sea ice.
 250 However, in this experiment the CO₂-concentration is kept at the pre-industrial
 251 value of 280 ppm. S3 represents the experiment with the CO₂-concentration
 252 in the atmosphere doubled to 560 ppm and the SSTs prescribed for the
 253 double CO₂-climate. Experiment C1, S1, S2 and S3 will be also referred to
 254 hereafter by PI, the simulation with high CO₂, the simulation with high SSTs
 255 and the simulation with high CO₂ and SSTs, respectively.

256 The experimental setup of this study is similar to the setup performed with the
 257 Canadian Middle Atmosphere Model (CMAM) by *Fomichev et al. (2007)* and
 258 with the Hamburg Model of the Neutral and Ionized Atmosphere
 259 (HAMMONIA) by *Schmidt et al. (2006)*. The HAMMONIA model is coupled to
 260 the same chemical model as WACCM: MOZART3. The setup in their study is
 261 similar, however, in their study, they double the CO₂-concentration from 360
 262 ppm to 720 ppm, while in our study, we double from the pre-industrial level of
 263 CO₂ (280 ppm).

264 Note that experiment S2 and S1 are not representing scenarios that could
 265 happen in the real atmosphere. These experiments have been used to study
 266 the effect of the SSTs separately. Experiment S3 doesn't take into account
 267 other (anthropogenic) changes in the atmosphere not caused by changes in
 268 the CO₂-concentration and the SSTs.

269 All the simulations are run for 50 years, of which the last 40 years are used for
 270 analysis. In the all results shown, we have used the 40 year mean of our
 271 model data.

272 Table 1. Set-up of the model experiments.

Experiment	CO ₂	SSTs from CESM equilibrium run
C1	<u>PI</u>	<u>PI</u>
<u>S1</u>	<u>Double</u>	<u>PI</u>
<u>S2</u>	<u>PI</u>	<u>High</u>
<u>S3</u>	<u>Double</u>	<u>High</u>

273 2.3 Climate feedback-response analysis method (CFRAM)

274 In this study, we aim to quantify the different climate feedbacks that may play
 275 a role in the middle atmosphere in a double CO₂-climate. For this purpose, we
 276 apply the climate feedback-response analysis method (CFRAM) (*Lu and Cai,*
 277 *2009*).

278 As briefly discussed in the introduction, traditional methods to study climate
 279 feedbacks are based on the energy balance at the top of the atmosphere
 280 (TOA). This means that the only climate feedbacks that are taken into
 281 consideration are those that affect the radiative balance at the TOA. However,
 282 there are other thermodynamic and dynamical processes that do not directly
 283 affect the TOA energy balance, while they do yield a temperature response in
 284

Deleted: S1

Deleted: , but

Deleted: CO₂-concentration

Deleted: S2

Deleted: C2,

Deleted: S2

Deleted: S1

Deleted: C2

Deleted: S2

Deleted: 280 ppm

Deleted: control

Deleted: C2

Moved (insertion) [1]

Deleted: 560 ppm

Deleted: control

Deleted: S1

Deleted: 280 ppm

Moved up [1]: Double

Deleted: CO₂ run

Deleted: S2

Moved (insertion) [2]

Deleted: 560 ppm

Moved up [2]: Double

Deleted: CO₂ run

Deleted: effect

the atmosphere.

Contrary to TOA-based methods, CFRAM considers all the radiative and non-radiative feedbacks that result from the climate system due to response to an external forcing. This means that CFRAM starts from a slightly different definition of a feedback process. Note also that as the changes in temperature are calculated simultaneously, the vertical mean temperature or lapse rate feedback per definition do not exist in CFRAM.

Another advantage of CFRAM is that it allows for measuring the magnitude of a certain feedback in units of temperature. We can actually calculate how much of the temperature change is due to which process. The '*climate response*' in the name of this method refers to the changes in temperature in response to the climate forcings and climate feedbacks.

We refer to the Appendix for the complete formulation of CFRAM diagnostics using outputs of WACCM. Based on the linear decomposition principle, we can solve Eq. (A12) for each of the terms on its right-hand side. This yields the partial temperature changes due to each specific process namely:

$$\Delta T_{CO_2} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta(S - R)_{CO_2} \quad (1)$$

$$\Delta T_{O_3} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta(S - R)_{O_3} \quad (2)$$

$$\Delta T_{H_2O} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta(S - R)_{H_2O} \quad (3)$$

$$\Delta T_{albedo} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta(S - R)_{albedo} \quad (4)$$

$$\Delta T_{cloud} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta(S - R)_{cloud} \quad (5)$$

In which R represents the vertical profile of the net long-wave radiation emitted by each layer in the atmosphere and by the surface. S is the vertical profile of the solar radiation absorbed by each layer. The matrix $\left(\frac{\partial R}{\partial T} \right)$ is the Planck feedback matrix, in which the vertical profiles of the changes in the divergence of radiative energy fluxes due to a temperature change are represented. ΔT represents the temperature change.

The factors $\Delta(S - R)_{CO_2}$, $\Delta(S - R)_{O_3}$, $\Delta(S - R)_{H_2O}$, $\Delta(S - R)_{albedo}$ and $\Delta(S - R)_{cloud}$ are calculated by inserting the output variables from WACCM in the radiation code of CFRAM. Here, one takes the output variables from the control run, apart from the variable that is related to the direct forcing or the feedback. Table A1 in the Appendix shows which variables from the perturbation runs have been inserted in the radiation code of CFRAM in order

Formatted: Font colour: Text 1

to calculate $\Delta(S - R)_{CO_2}$, $\Delta(S - R)_{O_3}$, $\Delta(S - R)_{H_2O}$, $\Delta(S - R)_{albedo}$ and $\Delta(S - R)_{cloud}$ and eventually the associated temperature changes.

Similarly, to equations (1)-(5), we also calculate the temperature change due to non-local thermal equilibrium (non-LTE) processes and the dynamical feedback. We calculate the terms $\Delta(S - R)_{non-LTE}$ and Δdyn in Eq. (A4) and (A7).

$$\Delta T_{non-LTE} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta(S - R)_{non-LTE} \quad (6)$$

$$\Delta T_{dyn} = \left(\frac{\partial R}{\partial T} \right)^{-1} \Delta dyn \quad (7)$$

The calculated partial temperature changes can be added, their sum being equal to the total temperature change. It is important to note that this does not mean that the individual processes are physically independent of each other.

$$\Delta T_{CFRAM} = \Delta T_{O_3} + \Delta T_{H_2O} + \Delta T_{albedo} + \Delta T_{cloud} + \Delta T_{non-LTE} + \Delta T_{dyn} \quad (8)$$

The linearization done for equations (A9) and (A10) introduces an error between the temperature difference as calculated by CFRAM and as seen in the model output. Another source of error is that the radiation code of the CFRAM calculations is not exactly equal to the radiation code of WACCM.

$$\Delta T_{CFRAM} = \Delta T_{WACCM} - \Delta T_{error} \quad (9)$$

For more details on the CFRAM method, please refer to *Lu and Cai (2009)*.

Note that the method used in this study differs from the Middle Atmosphere Climate Feedback Response Analysis Method (MCFRAM) used by *Zhu et al. (2016)*. The major difference is that in this study, we perform the calculations using the units of energy fluxes (Wm^{-2}) instead of converting to heating rates (Ks^{-1}). In other words, we use Wm^{-2} as the units of heating rates for the layer between two adjacent vertical levels. Because the radiative heating rates are the net radiative energy fluxes entering the layer, it is rather natural and straightforward (i.e., without dividing the mass in the layer to convert it to units of Ks^{-1}) to have the same units of heating rates (convergence) as the radiative energy fluxes. Another difference is that our method is not applicable above 0.01 hPa (~80 km), while *Zhu et al. (2016)* added molecular thermal conduction to the energy equation, to perform the calculations beyond the mesopause.

3. Temperature responses in a double CO₂ scenario

As described in section 2.2, four experiments were performed with WACCM: a simulation with pre-industrial conditions (experiment C1), a simulation with changed SSTs only (experiment S2), a simulation with only a changed CO₂-

Formatted: Font: Italic, Font colour: Accent 1

Deleted: This means that for the direct forcing of CO₂, one takes the CO₂ from the perturbation run, while one takes the other variables from the control run. For the ozone feedback, one takes the ozone from the perturbation run. For the water vapour feedback, one takes the specific humidity, surface pressure, surface temperature and dew point temperature. While for the albedo feedback, one takes the downwelling solar flux at surface and net solar flux at surface from the perturbation run and the other variables from the control run. ¶
For the cloud feedback, one takes the cloud fraction, cloud ice and cloud liquid amount from the perturbation run. For all these feedbacks, one takes the other variables from the control run. ¶

Deleted: LTE

Deleted: 1

concentration (experiment [S1](#)) and a final simulation with both changed SSTs and CO₂-concentration (experiment [S3](#)).

Figure 1 shows the zonal mean temperature changes for the different experiments with respect to the pre-industrial state, as modelled by WACCM. The results reach a statistical significance of 95% for the whole middle atmosphere domain in the experiments [S3](#)-C1 and [S1](#)-C1, and most of the middle atmosphere for experiment [S2](#)-C1. For this figure, as well as for all the results shown in this paper, we have used the 40 year mean of our data.

In line with what was shown in earlier studies (e.g. *Akmaev, 2006; Fomichev et al., 2007*), we observe that an increase in CO₂ causes a cooling in the middle atmosphere with the exception of the cold summer upper mesosphere region. We also observe that changing the SSTs alone, while leaving the CO₂-concentration at the pre-industrial levels (Fig 1c and 1f) also yields significant temperature changes over a large part of the middle atmosphere and contributes to the observed warming in the cold summer mesopause region.

As found previously by *Fomichev et al. (2007)* and *Schmidt et al. (2006)*, we find that the sum of the two separate temperature changes in the experiment with changed CO₂ only and changed SSTs only (experiment [S1](#) and [S2](#)) is approximately equal to the changes observed in the combined simulation (experiment [S3](#)). *Shepherd (2008)* has explained this phenomenon as follows: climate change affects the middle atmosphere in two ways: either radiatively through in situ changes associated with changes in CO₂ or dynamically through changes in stratospheric wave forcing, which are primarily a result of changing the SSTs (*Shepherd, 2008*). Even though the radiative and dynamic processes are not independent, these processes are seen to be approximately additive (*Sigmond et al., 2004; Schmidt et al., 2006; Fomichev et al., 2007*).




Figure 1: The total change in temperature in July (top) and January (bottom) for (a,d) the simulation with high CO₂ and SSTs ([S3](#)), (b,e) the simulation with high CO₂ ([S1](#)), (c,f) the simulation with high SSTs ([S2](#)), all as compared to the

Deleted: C2

Deleted: S2

Deleted: S2

Deleted: C2

Deleted: S1

Deleted: C2

Deleted: S1

Deleted: S2

Deleted: ,

Deleted: ,

Deleted: ,

pre-industrial control simulation, (C1). The dotted regions indicate the regions where the data reaches a confidence level of 95%. The black line indicates the tropopause height for the experiments S3 (a,d), S1 (b,e) and S2 (c,f).

4. Meridional-vertical profiles of partial temperature changes

The CFRAM makes it possible to separate and estimate the temperature responses due to an external forcing and various climate feedbacks, such as ozone, water vapour, cloud, albedo and dynamical feedbacks. Note that for the ozone, water vapour, cloud and albedo feedback, we can only calculate the radiative part of the feedback. The response to dynamical changes is calculated in a separate term.

This can be understood as we use the Fu-Liou radiative transfer model (*Fu and Liou*, 1992, 1993) to do offline calculations of the total local thermal equilibrium (LTE) radiative heating rate perturbation fields between the control experiment C1 and one of the other three experiments (i.e., S1, or S2, or S3). We use the standard outputs of atmospheric compositions (e.g., CO₂ and O₃) and thermodynamic fields (e.g., pressure, temperature, water vapour, clouds, surface albedo) as well as partial LTE radiative heating rate perturbation fields due to perturbations in individual atmospheric composition or thermodynamic fields (e.g., the terms on the right hand side of (A.9) except the first term).

We use the difference between the offline calculation of the total LTE radiative heating rate perturbations and the original total LTE radiative heating rate perturbations derived directly from the standard WACCM outputs as the error term of our offline LTE radiative heating perturbations. We note that the standard WACCM output fields also include non-LTE radiative heating fields, but do not include non-radiative heating rates. Therefore, we use the sum of the total LTE radiative heating rate perturbations and non-LTE radiative heating fields derived from the standard WACCM output fields to infer non-radiative heating rate perturbations under the equilibrium condition, namely Eq. (A.8).

We should also note that, because we are using an atmosphere-only model, in our experiment, the external forcing is either the change in CO₂ concentration or the change in SSTs or both. In an atmosphere-ocean model (such as CESM) and, of course, in reality, the changes in sea surface temperature and sea ice distributions are responses to the changed CO₂ concentration.

In the following subsections 4.1-4.5, we will discuss the meridional-vertical profiles of the temperature responses to the direct forcing and the various feedbacks during July and January. In section 5, we will discuss regional and global means of partial temperature changes due to feedbacks.

4.1 Direct temperature response to CO₂

Figure 2 shows the zonal mean temperature change due to the increase in CO₂. We see that increasing CO₂ leads to a cooling almost everywhere in the

Deleted: .

Deleted: S2

Deleted: C2

Deleted: S1

Deleted: C2, or

Deleted: this section

Deleted: the next

middle atmosphere, except at the high latitudes in the cold summer upper mesosphere, where we see a warming instead. The higher the temperature, the more cooling due to the increasing CO₂-concentration *is found* (Shepherd, 2008). The reason for this is that the outgoing longwave radiation strongly depends on the Planck blackbody emission (Zhu *et al.*, 2016).




Figure 2: Partial temperature change due to the direct forcing of CO₂ for July (top) and January (bottom) due to the doubling of the atmospheric CO₂-concentration, as calculated by CFRAM, using experiment S1 and C1. The black line indicates the tropopause height for the S1 run (with double CO₂-concentration).

Changing the SSTs does not lead to a change in CO₂-concentration, therefore the temperature response to changes in CO₂ is not present for the run with only changed SST (Figures not shown).

4.2 Ozone feedback

Ozone plays a crucial role in the chemical and radiative budget of the middle atmosphere. The distribution of ozone in the middle atmosphere is determined by both chemical and dynamical processes. Most of the ozone production takes place in the tropical stratosphere, as a result of photochemical processes, which involve oxygen. Meridional circulation then transports ozone to other parts of the middle atmosphere (Langematz, 2019). The production of ozone is largely balanced by catalytic destruction cycles involving NO_x, HO_x and Cl_x radicals. HO_x dominates ozone destruction in the mesosphere and lower stratosphere, while NO_x and Cl_x dominate this process in the middle and upper stratosphere (e.g. Cariolle, 1983).

Since the 1970s ozone in the middle atmosphere began to decline globally, due to increased production of ozone depleting substances (ODSs) (Brühl and Crutzen, 1988). The Montreal Protocol, adopted in 1987 to stop this threat, eventually led to a slow recovery of the stratospheric ozone over the recent two decades (WMO, 2018; Langematz, 2019). In our study, we don't consider the effect of anthropogenic ODSs since pre-industrial times (Langematz, 2019).

In this study, we are interested in the temperature response to changes in ozone concentration induced by the increased CO₂ concentration and/or the

Deleted: C2

Deleted: C2 runs

changes in SST in WACCM. Under enhanced CO₂ concentrations, the ratio between O₃ and O mixing ratios is generally shifted toward a higher concentration of ozone, which is caused by the strong temperature dependency of the ozone production reaction ($O + O_2 + M \rightarrow O_3 + M$).

Fig. 3 shows the percentage changes in O₃-concentration when the CO₂-concentration and/or the SSTs change. The results reach a statistical significance of 95% for the whole middle atmosphere domain in the experiments S3-C1 and S1-C1, and most of the middle atmosphere for experiment S2-C1.

We find, as expected, that an increase in CO₂ leads to an increase of ozone in most of the middle atmosphere. The increase of O₃ is about 20% around 2 hPa in the tropical region for experiment S3 with respect to C1. This corresponds with what is seen by *Fomichev et al., (2007)*, however they find that the increase in ozone in January is a bit lower in this region (around 15%, see their Figure 7).

There are some regions where the O₃-concentration is decreasing. In the tropical lower stratosphere, a decrease of about 20% is seen, in the summer polar mesosphere (around 0.01 hPa) ozone decreases by 3%, while in the mesosphere (around 0.02 hPa), ozone decreases by over 30%. Fig. 3c and f show that changing the SSTs also has a significant impact on the ozone concentration. A complete account of the ozone changes is out of the scope of this paper.




Figure 3: The percentage change in the zonal and monthly mean ozone concentration for July (top) and January (bottom) due to (a,d) combined effect of the CO₂ increase and SSTs changes (experiments S3 -C1), (b,e) the doubling of the atmospheric CO₂-concentration (experiments S1 - C1) and the (c,f) SSTs (experiments S2 - C1), as simulated by WACCM. The dotted regions indicate the regions where the data reaches a confidence level of 95%. The tropopause height is indicated as in Fig. 1.

Deleted: S2

Deleted: C2

Deleted: S1

Deleted: S2

Deleted: S2

Deleted: C2

Deleted: S1

597 As we will discuss in the next section, an enhanced concentration of CO₂ also
598 leads to changes in the dynamics in the middle atmosphere. The stratospheric
599 Brewer-Dobson circulation is projected to strengthen, which would lead to an
600 increase in the poleward transport of ozone. We will also see that an increase
601 in CO₂-concentration leads to stronger summer pole-to-winter pole flow in the
602 mesosphere.

603 Figure 4 shows the percentage change in the zonal and monthly mean
604 concentration of Cl, NO, O, OH, CH₃, NO_x and N₂O in July due to **the**
605 combined effect of the CO₂ increase and SSTs changes (experiment **S3** vs
606 C1), as simulated by WACCM. The patterns in January look similar (not
607 shown). These results reach a statistical significance of 95% for the whole
608 middle atmosphere domain.

Deleted: S2

609 We would like to point out that the changes in these constituents are only
610 brought about by the CO₂-concentration and/or the SSTs. We still use the
611 F_1850 compset and the only difference between the runs is the forcing in
612 CO₂ and SSTs. The changes in chemical constituents look very similar to
613 those found by *Schmidt et al.* (2006) who performed a similar experiment as
614 discussed in section 2.2, see their Figure 20. Note that Fig. 4 shows the
615 changes due to both the CO₂ increase and SSTs changes, while their Figure
616 20 shows the percentage changes due **to** the changes in CO₂-concentration
617 only and also only above 1 hPa.

618 As in *Schmidt et al.* (2006), we see an decrease in atomic oxygen (O) mixing
619 ratio at high summer latitudes around 0.01 hPa (see Fig. 4c), which results
620 from increased upwelling. This increase in O leads to a decrease in ozone in
621 this region. We also see decrease of ozone concentration in the winter polar
622 region around 0.1 hPa (approximately 65 km). This could be caused **by** an
623 increase of NO and for a small part by Cl mixing ratios, which result from a
624 stronger subsidence of NO and Cl rich air, as suggested in Schmidt et al.
625 2006. As stated before, complete discussion of the changes in ozone
626 concentration is out of the scope of this paper and the changes in other
627 constituents shown in Figure 4 are shown for reference only.




Figure 4: The percentage change in the zonal and monthly mean concentration of Cl (a), NO (b), O (c), OH (d), CH₄ (e) and NO_x (f) and N₂O (g) in July due to the combined effect of the CO₂ increase and SSTs changes (experiment (S3 vs C1), as simulated by WACCM. The dotted regions indicate the regions where the data reaches a confidence level of 95%. The tropopause height is indicated as in Fig. 1.

Deleted: S2

What is new in this study, is that we can calculate the temperature responses due to the changes in ozone concentration. These temperature responses are shown in Figure 5. It can be seen that there is a warming in the regions where there is an increase of the O₃-concentration, while there is a cooling for the regions with a decrease of the O₃-concentration. However, this is not the case for the winter polar region, where there is no sunlight. Note that the temperature responses to the changes in CO₂- and O₃- concentration behave differently in this respect: the temperature responses due to the direct forcing of CO₂ follow the temperature distribution quite closely, while the temperature responses due to O₃ follow the ozone concentration, as also seen by Zhu et al., (2016).




Figure 5: Partial temperature responses to changes in O_3 -concentration, as calculated by CFRAM, in July (top) and January (bottom) due to the (a,d) combined effect of the CO_2 increase and SSTs changes (experiment S3), (b,e) the doubling of the atmospheric CO_2 -concentration (experiment S1) and the (c,f) SSTs (experiment S2). The tropopause height is indicated as in Fig. 1.

Deleted: S1

Deleted: experiments C2

Deleted: experiments S1

4.2 Dynamical feedback

The zonal mean residual circulation forms an important component of the mass transport by the Brewer-Dobson circulation (BDC). It consists of a meridional (v^*) and a vertical (w^*) component as defined in the Transformed Eulerian Mean (TEM) framework. The residual circulation consists of a shallow branch, which controls the transport of air in the tropical lower stratosphere, as well as a deep branch in the mid-latitude upper stratosphere and mesosphere.

Both of these branches are driven by atmospheric waves. In the winter hemisphere, planetary Rossby waves propagate upwards into the stratosphere, where they break and deposit their momentum on the zonal mean flow, which in turn induces a meridional circulation. The two-cell structure in the lower stratosphere, which is present all-year round, is driven by synoptic scale waves. The circulation is also affected by orographic gravity wave drag in the stratosphere and by non-orographic gravity wave drag in the upper mesosphere (Oberländer *et al.*, 2013).

Most climate models show that the BDC and the upwelling in the equatorial region will speed up due to an increase in CO_2 -concentration (Butchart *et al.*, 2010). It has been shown that the strengthening of the Brewer-Dobson circulation in the lower stratosphere is caused by changes in transient planetary and synoptic scale waves, while the upper stratospheric changes are due to changes in the propagation properties for gravity waves (Oberländer *et al.*, 2013).

684 It has been explained that the increased stratospheric resolved wave drag is
 685 caused by an increase of the meridional temperature gradient in the
 686 stratosphere, which leads to a strengthening of the upper flank of the
 687 subtropical jets. This in turn shifts the critical layers for Rossby wave breaking
 688 upward, which allows for more Rossby waves to reach the lower stratosphere,
 689 where they break and deposit their momentum, enhancing the BDC
 690 (Shepherd and McLandress, 2011)

691 The differences in the meridional component of the residual circulation (\bar{v}^*)
 692 between the different simulations are shown in Fig. 6. These data are
 693 averaged over the 40 years of data. The results reach a statistical
 694 significance of 95% almost the whole area above 1 hPa for the experiments
 695 S1-C1, for the experiment S2-C1 the results reach a statistical significance of
 696 95% in most of the area below this level. The experiments S3-C1 show the
 697 largest region of statistical significance, apart from some regions below 1 hPa.

Deleted: C2


Deleted: S1

Deleted: S2

698 Figure 6b and 6e show that only doubling the CO₂-concentration leads to a
 699 stronger pole-to-pole flow in the mesosphere. Changing the SSTs also leads
 700 to changes in the residual circulation as can be seen in Fig. 6c and 6f.
 701 Oberländer et al. (2013) have shown that the rising CO₂-concentration affects
 702 the upper stratospheric layers, while the signals in the lower stratosphere are
 703 almost completely due to changes in sea surface temperature.

704 The warmer sea surface temperatures affect the dynamics in the middle
 705 atmosphere. It has for example been shown that higher SSTs in the tropics
 706 leads to an amplification in deep convection, which enhances the generation
 707 of quasi-stationary waves (Deckert and Dameris, 2008). Enhanced SSTs lead
 708 to an enhanced dissipation of planetary waves, as well as an enhanced
 709 dissipation of orographic and non-orographic waves in the upper stratosphere
 710 (Oberländer et al., 2013).

711



712

713 Figure 6: Changes in the zonal and monthly mean transformed Eulerian-mean
 714 residual circulation horizontal velocity \bar{v}^* for July (top) and January (bottom)
 715 due to (a,d) combined effect of the CO₂ increase and SSTs changes

(experiments [S3](#) - C1), (b,e) the doubling of the atmospheric CO₂-concentration (experiments [S1](#) - C1) and the (c,f) SSTs (experiments [S2](#) - C1), as simulated by WACCM. The dotted regions indicate the regions where the data reaches a confidence level of 95%. The tropopause height is as indicated in Fig. 1.

Deleted: S2

Deleted: C2

Deleted: S1

We are interested in the temperature responses due to the dynamical feedbacks in the different experiments. These temperature responses are shown in Figure 7. Figure 7b and 7e show that there is cooling in the summer mesosphere, while there is warming in the winter mesosphere, which is consistent with a stronger summer-to-winter pole flow.

Figure 7c and 7f show the temperature responses due to changes in the SSTs. It is seen that there is mostly a warming in the summer mesosphere and mostly a cooling in the winter hemisphere, which would weaken the effect of the changed CO₂-concentration. Most of the temperature responses in the lower stratosphere are caused by the changes in SSTs, as expected.




Figure 7: Partial temperature responses to changes in dynamics, as calculated by CFRAM, in July (top) and January (bottom) due to the (a,d) combined effect of the CO₂ increase and SSTs changes ([experiment S3](#)), (b,e) the doubling of the atmospheric CO₂-concentration ([experiment S1](#)) and the (c,f) SSTs ([experiment S2](#)). The tropopause height is indicated as in Fig. 1.

Deleted: S1

Deleted: experiments C2

Deleted: experiments S1

In summary, doubling the CO₂-concentration leads to a stronger pole-to-pole flow in the mesosphere, which leads to cooling of the summer mesosphere and a warming of the winter mesosphere. Changing the SSTs weakens this effect, but leads to temperature changes in the stratosphere and lower mesosphere.

4.4 Water vapour feedback

Figure 8 shows how the water vapour is changing in the middle atmosphere if the CO₂-concentration is increased and/or the SSTs are changed with respect to the pre-industrial control run. In WACCM, increasing the CO₂-concentration alone leads to a decrease of water vapour in most of the middle atmosphere (Fig. 8b and f). The results reach a statistical significance of 95% for the whole middle atmosphere domain in the experiments S3-C1 and S2-C1, and most of the middle atmosphere for experiment S1-C1, apart from the winter hemisphere region around 0.1 hPa.

The amount of water vapour in the stratosphere is determined by transport through the tropopause as well as by the oxidation of methane in the stratosphere itself. The transport of the water vapour in the stratosphere is mainly a function of the tropopause temperature (Solomon *et al.*, 2010). In WACCM, we see a decrease in temperature in the tropical tropopause for the double CO₂ experiment of about -0.25 K. The cold temperatures in the tropical tropopause lead to a reduction of water vapour of between 2 and 8% due to freeze-drying in this region.

It can be seen that using the SSTs from the doubled CO₂-climate leads to an increase in water vapour almost everywhere in the middle atmosphere as compared to PI (Fig. 8c and f). In WACCM, forcing with SSTs from a double CO₂-climate is observed to lead to a higher and warmer tropopause, which can explain this increase of water vapour. However, it should be noted that models currently have a limited representation of the processes determining the distribution and variability of lower stratospheric water vapour. Minimum tropopause temperatures are not consistently reproduced by climate models (Solomon *et al.*, 2010; Riese *et al.*, 2012). At the same time, observations are not completely clear about whether there is a persistent positive correlation between the SST and the stratospheric water vapour (Solomon *et al.*, 2010).




Figure 8: The percentage changes in the zonal and monthly mean water vapour mixing ratio for July (top) and January (bottom) due to (a,d) combined effect of the CO₂ increase and SSTs changes (experiments S3 - C1), (b,e) the doubling of the atmospheric CO₂-concentration (experiments S1 - C1) and the (c,f) SSTs (experiments S2 - C1), as simulated by WACCM. The dotted

Deleted: S2

Deleted: S1

Deleted: C2

Deleted: S2 -

Deleted: C2

Deleted: S1

regions indicate the regions where the data reaches a confidence level of 95%. The tropopause height is as indicated in Fig. 1.

Figure 9 shows the temperature responses due to the changes in water vapour as calculated by CFRAM. It can be seen that in the regions where there is an increase in the water vapour, there is a cooling, and vice versa. This can be understood as increasing the water vapour in the middle atmosphere leads to an increase in longwave emissions in the mid and far-infrared by water vapour. This in turns leads to a cooling of the region. Similarly, a decrease in water vapour leads to a warming of the region (Brasseur and Solomon, 2005). Fig. 8 shows that above 1 hPa, there are also large percentage changes in water vapour. However, the absolute concentration of water vapour is small there, which explains why there is no temperature response to these changes.

Deleted: Higher up in the atmosphere




Figure 9: Partial temperature responses to changes in water vapour, as calculated by CFRAM, in July (top) and January (bottom) due to the (a,d) combined effect of the CO₂ increase and SSTs changes (experiment S3), (b,e) the doubling of the atmospheric CO₂-concentration (experiment S1) and the (c,f) SSTs (experiment S2). The tropopause height is indicated as in Fig. 1.

Deleted: S1

Deleted: experiments C2

Deleted: experiments S1

Water vapour plays a secondary but not negligible role in determining the middle atmosphere climate sensitivity. In the lower stratosphere, H₂O contributes considerably to the cooling in this region. Above 30 hPa, the water vapour contribution to the energy budget is negligible, as also seen by Fomichev et al. (2007).

4.5 Cloud and albedo feedback

Forcing the model with SSTs from the double CO₂-climate (as in experiment S2 and S3) yields an overall increase in the cloud cover in the upper troposphere, while this is not the case if one only increases the CO₂ concentration (as in experiment S1). Figure 10 shows the temperature

Deleted: S1

Deleted: S2

Deleted: C2

responses to changes in cloud (left) and albedo (right) in July (top) and January (bottom) for experiment [S2](#), as calculated by CFRAM.

Deleted: S1

Fig. 10 shows in the tropical region, there is a warming due to changes in clouds, while there is a cooling at higher latitudes in July (see Figure 10a). In January, the pattern looks slightly different (see Figure 10c). These temperature changes are due to changes in the balance between the increased reflected shortwave radiation and the decrease of outgoing longwave radiation.

We also see an effect of the changes in surface albedo in the stratosphere (see Figure 10b and d). The cooling in the summer polar stratosphere shown in Figure 10b and d is due to radiative changes. We suggest that this cooling is due to a decrease in surface albedo, which would lead to less shortwave radiation being reflected. However, more research is needed.




Figure 10: Partial temperature responses to changes in clouds (left) and albedo (right), as calculated by CFRAM, in July (top) and January (bottom) due to the SSTs (experiment [S2](#)). The tropopause height is indicated as in Fig. 1.

Deleted: S1

Cloud and albedo feedbacks due to changes in clouds and surface albedo play a crucial role in determining the tropospheric and surface climate (*Boucher et al., 2013, Royer et al., 1990*). However, it is clear that these feedbacks play only a very small role in the middle atmosphere temperature response to the doubling of CO_2 and SSTs.

5. Regional and global means of partial temperature changes due to feedbacks

To study the relative importance of the different feedback processes globally we show the average change in global mean temperature for the lower stratosphere, the upper stratosphere and the mesosphere for the S3 experiment with the changed CO₂-concentration and changed SSTs in Figure 11. We also show the average change in temperature in the polar regions (90°S-70°S and 70°N-90°N), the tropics (20°S-20°N) for the lower and upper stratosphere and the mesosphere.

In order to calculate the lower stratospheric temperature changes, we take the average value of the temperature change from the tropopause up to 24 hPa. The pressure level of the tropopause is simulated in WACCM for each latitude and longitude, we use this pressure level to demarcate between the troposphere and stratosphere. We consider 24 hPa as a crude estimate for the boundary between the lower and upper stratosphere.

The tropopause is not exactly at the same pressure level in the perturbation experiments as compared to the pre-industrial control run (C1). We always take the tropopause of the perturbation experiment which is a bit higher at some latitudes, to make sure that we do not use values from the troposphere. We add the values for each latitude up and take the average. This average is not mass weighted. By calculating the average in this way, we can directly compare the vertical values in different regions of the atmosphere. The temperature changes in the upper stratosphere and in the mesosphere are calculated in the same way, but then for the altitudes 24 hPa-1 hPa and 1 hPa-0.01 hPa respectively.

Figure 11 shows the radiative feedbacks due to ozone, water vapour, clouds, albedo and the dynamical feedback, as well as the small contribution due to the Non-LTE processes in column 'NLTE', as calculated by CFRAM. The 'total'-column shows the temperature changes in WACCM, while the column 'error' shows the difference between temperature change in WACCM and the sum of the calculated temperature responses in CFRAM. Note that the range of values on the y-axis is not the same for the different subplots.

Figure 11 shows that the temperature change in the lower stratosphere due to the direct forcing of CO₂ is around 3 K in the global mean. There is a stronger cooling in the tropical region of about 4 K in July and 3.5 K in January. We also observe that there is a cooling of about 1 K due to ozone feedback in the tropical region while there is a slight warming taking place in the summer hemispheres in both January and July. We also see that the temperature change in the lower stratosphere is influenced by the water vapour feedback. There is a cooling of about 0.5 K in the lower stratosphere, apart from in the southern polar area. There is some small influence from the cloud and albedo feedback, which can be negative or positive (see also Fig. 9).

In the upper stratosphere, the cooling due to the direct forcing of CO₂ is with about 9 K in the global mean considerably stronger than in the lower

Deleted: S2

Deleted: stratosphere

Deleted: until

Deleted: taking

922 stratosphere. The cooling is stronger in the summer polar regions, where the
923 cooling due to the direct forcing of CO₂ reaches 11K. In the winter polar
924 region, this cooling is only about 8K.

925
926 The water vapour, cloud and albedo feedback play no role in the upper
927 stratosphere nor in the mesosphere. The ozone feedback results in the
928 positive partial temperature changes in the upper stratosphere, of about 2 K in
929 the global mean. The changes in ozone don't result in temperature changes in
930 the winter hemisphere, as discussed in section 4.2.

931
932 The picture in the mesosphere is similar as in the upper stratosphere. The
933 main difference is that the temperature changes are larger. The global
934 temperature change due to direct forcing of CO₂ is about 15 K. The O₃-
935 feedback results in a partial temperature changes of about 3 K in the
936 mesosphere in the global mean. The temperature change due to ozone in the
937 equatorial mesosphere is about 4 K, while the warming due to ozone in the
938 summer polar region is a bit smaller: around 3K. Just like in the upper
939 stratosphere water vapour, cloud and albedo feedback play no role.

940
941 We see, that the ozone feedback generally yields a radiative feedback that
942 mitigates the cooling, which is due to the direct forcing of CO₂. This has been
943 suggested in earlier studies, such as *Jonsson et al., 2004, Dietmüller et al.,*
944 *2014*. With CFRAM, it is possible to quantify this effect and to compare it with
945 the effects of other feedbacks in the middle atmosphere. Note that no other
946 method before has been able to quantify how much of the temperature
947 change in the middle atmosphere is due to the different feedback processes.

Deleted: before

948
949 The temperature response due to dynamical feedbacks is small in global
950 average: less than 1 K. This can be understood as waves generally do not
951 generate momentum and heat, but redistribute these instead (*Zhu et al.,*
952 *2016*). However, the local responses to dynamical changes in the high
953 latitudes are large, as we have seen in section 4.2. There are some very small
954 temperature responses due to non-LTE effects as well, which mostly
955 contribute to the temperature change in the mesosphere.

Deleted: mostly

956
957 The error term is relatively large, as can be seen from the rightmost column in
958 Fig.11. This term shows the difference between temperature change in
959 WACCM and the sum of the calculated temperature responses in CFRAM
960 (see eq. 9 in section 2.3). In CFRAM, we assumed that the radiative
961 perturbations can be linearized by neglecting the higher order terms of each
962 thermodynamic feedback and the interactions between these feedbacks, this
963 yields an error.

Deleted: The error term is relatively large.

Formatted: Font: 12 pt

964
965 Cai and Lu (2009) show that this error is larger in the middle atmosphere than
966 for similar calculations in the troposphere. In the middle atmosphere, the
967 density of the atmosphere is smaller, which leads to smaller numerical values
968 of the diagonal elements of the Planck feedback matrix. As a result, the linear
969 solution is very sensitive to forcing in the middle atmosphere. Another part of
970 the error is due to the fact that the radiative transfer model used in the offline

Formatted: Font: 12 pt

Formatted: x_body, Line spacing: At least 12,65 pt, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

974 CFRAM calculations is different than the radiative transfer model used in the
975 climate simulations with WACCM.
976





Figure 11: The mean temperature responses to the changes in CO₂ and various feedback processes in the lower stratosphere from the tropopause height up to 24 hPa (a,b), upper stratosphere from 24-1 hPa (c, d) and in the mesosphere from 1-0.01 hPa (e,f) in July (a, c, e) and January (b, d, f) in the polar regions (90°S-70°S and 70°N-90°N), the tropics (20°S-20°N) and the global mean, for S3 experiment (double CO₂ and changed SSTs). Note that the range of values on the y-axis is not the same for the different subplots.

In addition, the vertical profiles of the temperature responses to the direct forcing of CO₂ and the feedbacks are shown in Figure 12. Here, one can see that the increase in CO₂ leads to a cooling over almost the whole middle atmosphere; an effect that increases with height. We also observe that in the summer upper mesosphere regions, the increased CO₂-concentration leads to a warming. The changes in ozone concentration in response to the doubling of CO₂ lead to a warming almost everywhere in the atmosphere. In some places, this warming exceeds 5 K. In the polar winter the effect of ozone is small due to lack of sunlight.

There is also a relatively large temperature response to the changes in dynamics. In Fig. 12, it can be seen that there is a cooling in the summer mesosphere, while there is warming in the winter mesosphere. The water vapour, cloud and albedo feedback play only a very small role in the middle atmosphere, as we observed in Figure 11. We find that there are also some small temperature changes due to non-LTE effect above 0.1 hPa. How the non-LTE effects exactly cause the small temperature changes in this region is outside the scope of this paper and needs further investigation.

Temperature responses to different feedback processes



Deleted: until

Deleted: S1

Deleted: The Non-LTE effects

Deleted: , but start

Deleted: play a small role

Deleted: ,

Deleted: exact mechanism of which

Figure 12: Vertical profiles of the temperature responses to the changes in CO₂ and various feedback processes in July (top) and January (bottom) due to double CO₂ and changed SSTs in the atmosphere between 200 and 0.01 hPa, for regions from 50° N/S poleward and the tropics (20°S-20°N), as calculated by CFRAM.




Figure 13: The mean temperature responses to the changes in CO₂ and various feedback processes in July (a,c) and January (b,d) in the upper stratosphere between 24 and 1 hPa, for polar regions (90°S-70°S and 70°N-90°N), the tropics (20°S-20°N) and the global mean for the experiment with double CO₂ (S1) (a,b) and changed SSTs (S2) (c,d) separately.

Deleted: for

Deleted: C2

Deleted: S1

Figure 13 shows the temperature responses in the upper stratosphere for the experiment with double CO₂ (a,b) and changed SSTs (c,d) separately. This has been done to give insight in the temperature response of the CO₂ and the SST separately. These temperature changes were calculated in the same way as for Fig. 11. Again also, the 'total'-column shows the temperature changes as simulated by WACCM, the columns CO₂, O₃, H₂O, cloud, albedo, dynamics, the column 'N₂O' shows the temperature responses due to non-LTE processes as calculated by CFRAM. As in Fig. 11, the 'Error'-column in Fig. 13 shows the difference between temperature change in WACCM and the sum of the calculated temperature responses in CFRAM.

We learn from this figure that the effects of the changed SSTs on the upper stratosphere are relatively small as compared to the effects of changing the CO₂. We show the temperature changes for the upper stratosphere as an example. In the lower stratosphere and the mesosphere, we see the same pattern: the effect of the CO₂ on the temperature is generally much larger than the effect of the SSTs on the temperature. This finding is consistent with the study of Fomichev *et al.* (2007), where it is concluded that the impact of changes in SSTs on the middle atmosphere is relatively small and localized as compared to the combined response of changing the CO₂-concentration and the SSTs.

The changes in SSTs are, however, responsible for large temperature changes as a result of the dynamical feedbacks, especially in the winter hemispheres, where there is a temperature response of 4K. A similar figure for the lower stratosphere (not shown) shows that the temperature response to the water vapour feedback is almost solely due to changes in the SSTs and not the direct forcing of CO₂.

Earlier, we discussed that the sum of the two separate temperature changes in the experiment with double CO₂ and changed SSTs is approximately equal to the changes observed in the combined simulation. We find that the same is true for the temperature responses to the different feedback processes.

5. Discussion and conclusions

In this study, we have applied the climate feedback response analysis method to climate sensitivity experiments performed by WACCM. We have examined the middle atmosphere response to CO₂ doubling with respect to the pre-industrial state. We investigated the combined effect of doubling CO₂ and subsequent warming SSTs, as well as the effects of separately changing the CO₂ and the SSTs. It is important to note that no other method before has been able to quantify how much of the temperature change in the middle atmosphere is due to the different feedback processes.

It was found before that the sum of the two separate temperature changes in the experiment with only changed CO₂ and only changed SSTs is, at first approximation, equal to the changes observed in the combined simulation

Deleted: ¶
¶

Deleted: on-

Deleted: LTE

Deleted: Error

Deleted: before

1084 (see e.g. *Fomichev et al. (2007) and Schmidt et al. (2006)*). This is also the
1085 case for WACCM.

1086
1087 We have found that, even though changing the SSTs yields significant
1088 temperature changes over a large part of the middle atmosphere, the effects
1089 of the changed SSTs on the middle atmosphere are relatively small as
1090 compared to the effects of changing the CO₂ without changes in the SSTs.

1091 We have given an overview of the mean temperature responses to the
1092 changes in CO₂ and various feedback processes in the lower stratosphere,
1093 upper stratosphere and in the mesosphere in January and July. We find that
1094 the temperature change due to the direct forcing of CO₂ increases with
1095 increasing height in the middle atmosphere. The temperature change in the
1096 lower stratosphere due to the direct forcing of CO₂ is around 3 K. There is a
1097 stronger cooling in the tropical lower stratosphere of about 4 K in July and 3.5
1098 K in January.

1099 In the upper stratosphere, the cooling due to the direct forcing of CO₂ is about
1100 9 K, which is considerably stronger than in the lower stratosphere. The
1101 cooling is stronger in the summer polar regions, where the cooling reaches a
1102 value of 11K, than in the winter polar region, where the cooling is only about
1103 8K. In the mesosphere, the cooling due to the direct forcing of CO₂ is even
1104 stronger: 15 K.

1105 The ozone concentration changes due to changes in the CO₂-concentration
1106 as well as by changes in the SSTs. The temperature changes caused by this
1107 change in ozone concentration generally mitigate the cooling caused by the
1108 direct forcing of CO₂. However, in the tropical lower stratosphere and in some
1109 regions of the mesosphere, the ozone feedback cools these regions further. In
1110 the tropical lower stratosphere, for example, there is a cooling of 1K due to
1111 the ozone feedback.

1112
1113 We also have seen that the global mean temperature response due to
1114 dynamical feedbacks is small in the global average in all regions: less than 1
1115 K. However, local responses to the changes in dynamics can be large.
1116 Doubling the CO₂-concentration leads to a stronger summer-to-winter-pole
1117 flow, which leads to a cooling of the summer mesosphere and a warming of
1118 the winter mesosphere. Changing the SSTs weakens this effect in the
1119 mesosphere, but affects the temperature response in the stratosphere and
1120 lower mesosphere.

1121
1122 Using CFRAM on WACCM data shows that the change in water vapour leads
1123 to a cooling of up to 2 K in the lower stratosphere. It should be noted that
1124 climate models currently have a limited representation of the processes
1125 determining the distribution and variability of lower stratospheric water vapour.
1126 This means that the temperature response to the water vapour feedback
1127 might be different using a different model. We have also seen a small effect of
1128 the cloud and albedo feedback on the temperature response in the lower
1129 stratosphere, while these feedbacks play no role in the upper stratosphere
1130 and the mesosphere.

Deleted: ¶

The results seen in this study are consistent with earlier findings. As in *Shepherd et al.*, (2008), we find that the higher the temperature at a region in the atmosphere, the more cooling there is seen due to the direct feedback of CO₂. We find, as in *Zhu et al.*, (2016) that the temperature responses due to the direct forcing of CO₂ follow the temperature distribution quite closely, while the temperature responses due to O₃ follow the changes in ozone concentration instead.

We have also seen that the ozone feedback generally yields a radiative feedback that mitigates the cooling, which is due to the direct forcing of CO₂, which is consistent with earlier studies such as *Jonsson et al.*, (2004), *Dietmüller et al.*, (2014). CFRAM is the first study that allows for calculating how much of the temperature response is due to which feedback process.

The next step would be to investigate the exact mechanisms behind the feedback processes in more detail. Some processes can influence the different feedback processes, such as ozone depleting chemicals influencing the ozone concentration and thereby the temperature response of this feedback. A better understanding of the effect of the increased CO₂-concentration on the middle atmosphere, will help to distinguish the effects of the changes CO₂- and O₃-concentration.

There is also a need for a better understanding of how different feedbacks in the middle atmosphere affect the surface climate. As discussed in the introduction, the exact importance of ozone feedback on the global mean temperature is currently not clear (*Nowack et al.*, 2015, *Marsh et al.*, 2016). A similar analysis as in this paper can be performed to quantify the effects of feedbacks on the surface climate.

In conclusion, we have seen that CFRAM is an efficient method to quantify climate feedbacks in the middle atmosphere, although there is a relatively large error due to the linearization in the model. The CFRAM allows for separating and estimating the temperature responses due to an external forcing and various climate feedbacks, such as ozone, water vapour, cloud, albedo and dynamical feedbacks. More research into the exact mechanisms of these feedbacks could help us to understand the temperature response of the middle atmosphere and their effects on the surface and tropospheric climate better.

Appendix: Formulation of CFRAM diagnostics using outputs of WACCM

The mathematical formulation of CFRAM is based on the conservation of total energy (*Lu and Cai*, 2009). At a given location in the atmosphere, the energy balance in an atmosphere-surface column can be written as:

$$R = S + Q^{conv} + Q^{turb} - D^v - D^h + W^{fric} \quad (A1)$$

R represents the vertical profile of the net long-wave radiation emitted by each layer in the atmosphere and by the surface. S is the vertical profile of the solar

radiation absorbed by each layer. Q^{turb} is the convergence of total energy fluxes in each layer due to turbulent motions, Q^{conv} is convergence of total energy fluxes into the layers due to convective motion. D^v is the large-scale vertical transport of energy from different layers to others. D^h is the large-scale horizontal transport within the layers and W^{fric} is the work done by atmospheric friction. All terms in (A1) have units of Wm^{-2} .

Due to an external forcing (in this study, the change in CO_2 -concentration and/or change in SSTs), the difference in the energy flux terms then becomes:

$$\Delta R = \Delta F^{ext} + \Delta S + \Delta Q^{conv} + \Delta Q^{turb} - \Delta D^v - \Delta D^h + \Delta W^{fric} \quad (A2)$$

In which the delta (Δ) stands for the difference between the perturbation run and the control run.

CFRAM takes advantage of the fact that the infrared radiation is directly related to the temperatures in the entire column. The temperature changes in the equilibrium response to perturbations in the energy flux terms can be calculated. This is done by requiring that the temperature-induced changes in infrared radiation balance the non-temperature induced energy flux perturbations.

Equation (A2) can also be written as:

$$\Delta(S - R)_{total} + \Delta dyn = 0 \quad (A3)$$

The term $\Delta(S - R)$ can be calculated as the longwave heating rate and the solar heating rate are output variables of the model simulations. We take the time mean of the WACCM data and perform the calculations for each grid point of the WACCM data. This means that in the end, we will have the temperature changes at each latitude, longitude and height.

We then calculate the difference in these heating rates for the perturbation simulation and the control simulation.

We use the term $\Delta(S - R)_{total}$ to calculate the dynamics term Δdyn .

$$\Delta dyn = -\Delta(S - R)_{total} \quad (A4)$$

WACCM provides us with a heating rate in Ks^{-1} . For the CFRAM calculations, we need the energy flux in Wm^{-2} . We can calculate the energy flux by multiplying with the mass of different layers in the atmosphere and the specific heat capacity.

$$\Delta(S - R) = \Delta(S - R)_{(WACCM)} * mass_k * c_p \quad (A5)$$

1229 In which $\Delta(S - R)$ is the difference in the shortwave radiation (\overline{S}) and
 1230 longwave radiation (\overline{R}) between the perturbation run and the control run as a
 1231 flux in Wm^{-2} , while $\Delta(S - R)_{(WACCM)}$ is this difference as heating rate in Ks^{-1} in
 1232 WACCM, with $mass_k = \frac{p_{k+1} - p_k}{g}$ with p in Pa, $c_p = 1004 \text{ J kg}^{-1} \text{ K}^{-1}$ the
 1233 specific heat capacity at constant pressure and g the gravitational
 1234 acceleration 9.81 ms^{-2} .

1235
 1236 WACCM includes a non-local thermal equilibrium (non-LTE) radiation scheme
 1237 above 50 km. It consists of a long-wave radiation (LW) part and a short-wave
 1238 radiation (SW) part which includes the extreme ultraviolet (EUV) heating rate,
 1239 chemical potential heating rate, CO_2 near-infrared (NIR) heating rate, total
 1240 auroral heating rate and non-EUV photolysis heating rate.

1241 Therefore, we split the term $\Delta(S - R)_{total}$ in an LTE and a non-LTE term:

$$1243 \Delta(S - R)_{total} = \Delta(S - R)_{LTE} + \Delta(S - R)_{non-LTE} \quad (A6)$$

1244
 1245 WACCM provides us with the total longwave heating rate as well as the total
 1246 solar heating rate and the non-LTE longwave and shortwave heating rates for
 1247 the different runs. This means that we can calculate the term $\Delta(S - R)_{non-LTE}$
 1248 as well, where we again need to convert our result from Ks^{-1} to Wm^{-2} :

$$1250 \Delta(S - R)_{non-LTE} = \Delta(S - R)_{non-LTE(WACCM)} mass_k * c_p \quad (A7)$$

1251 This term can be inserted in equation (3):

$$1254 \Delta(S - R)_{LTE} + \Delta(S - R)_{non-LTE} + \Delta dyn = 0 \quad (A8)$$

1255
 1256 The central step in CFRAM is to decompose the radiative flux vector, using a
 1257 linear approximation.

1258
 1259 We start by decomposing the LTE infrared radiative flux vector ΔR

$$1261 \Delta R_{LTE} = \frac{\partial R}{\partial T} \Delta T + \Delta R_{CO_2} + \Delta R_{O_3} + \Delta R_{H_2O} + \Delta R_{albedo} + \Delta R_{cloud} \quad (A9)$$

1262
 1263 where ΔR_{CO_2} , ΔR_{O_3} , ΔR_{H_2O} , ΔR_{albedo} , ΔR_{cloud} are the changes in infrared
 1264 radiative fluxes due to the changes in CO_2 , ozone, water vapour, albedo and
 1265 clouds, respectively.

1266
 1267 For equation (A9), we assumed that radiative perturbations can be linearized
 1268 by neglecting the higher order terms of each thermodynamic feedback and
 1269 the interactions between these feedbacks. This is also commonly done in the
 1270 partial radiative perturbation (PRP) method, in which partial derivatives of the
 1271 model top of the atmosphere radiation are evaluated with respect to changes
 1272 in model parameters by diagnostic rerunning the model's radiation code (Bony
 1273 *et al.*, 2006).

1274

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

The term $\frac{\partial R}{\partial T} \Delta T$ represents the changes in the IR radiative fluxes related to the temperature changes in the entire atmosphere-surface column. The matrix $\frac{\partial R}{\partial T}$ is the Planck feedback matrix, in which the vertical profiles of the changes in the divergence of radiative energy fluxes due to a temperature change are represented.

We calculate this feedback matrix using the output variables of the perturbation and the control run of WACCM and inserting these in the CFRAM radiation code: atmospheric temperature, surface temperature, reference height temperature, ozone, surface pressure, solar insolation, downwelling solar flux at the surface, net solar flux at the surface, dew point temperature, cloud fraction, cloud ice amount, cloud liquid amount, ozone and specific humidity.

Similarly, the changes in the LTE shortwave radiation flux can be written as the sum of the change in shortwave radiation flux due to the direct forcing of CO₂ and the different feedbacks:

$$\Delta S_{LTE} = \Delta S_{CO_2} + \Delta S_{O_3} + \Delta S_{H_2O} + \Delta S_{albedo} + \Delta S_{cloud} \quad (A10)$$

Similarly, to equation (A9), we perform a linearization.

Substituting (A9) and (A10) in equation (A8) yields:

$$\Delta(S - R)_{CO_2} + \Delta(S - R)_{O_3} + \Delta(S - R)_{H_2O} + \Delta(S - R)_{albedo} + \Delta(S - R)_{cloud} - \frac{\partial R}{\partial T} \Delta T + \Delta(S - R)_{non-LTE} + \Delta dyn = 0 \quad (A11)$$

This can be written as:

$$\Delta T = \left(\frac{\partial R}{\partial T} \right)^{-1} \left\{ \Delta(S - R)_{CO_2} + \Delta(S - R)_{O_3} + \Delta(S - R)_{H_2O} + \Delta(S - R)_{albedo} + \Delta(S - R)_{cloud} + \Delta(S - R)_{non-LTE} + \Delta dyn \right\} \quad (A12)$$

As described in the main text of this paper, we can solve Eq. (A12) for each of the terms on its right-hand side, based on the linear decomposition principle. This yields the partial temperature changes due to each specific process. The factors $\Delta(S - R)_{CO_2}$, $\Delta(S - R)_{O_3}$, $\Delta(S - R)_{H_2O}$, $\Delta(S - R)_{albedo}$ and $\Delta(S - R)_{cloud}$ in eqs (1-5) are calculated by inserting the output variables from WACCM in the radiation code of CFRAM. Here, one takes the output variables from the control run, apart from the variable that is related to the direct forcing or the feedback. The table below shows which variables have been taken from the perturbation runs for each feedback.

Direct forcing/feedback	Changed variables in the radiation code
CO ₂	CO ₂

<u>Ozone</u>	<u>O₃</u>
<u>Water vapour</u>	<u>Specific humidity</u> <u>Surface pressure</u> <u>Surface temperature</u> <u>Dew point temperature</u>
<u>Albedo</u>	<u>Downwelling solar flux at surface</u> <u>Net solar flux at surface</u>
<u>Cloud</u>	<u>Cloud fraction</u> <u>Cloud ice</u> <u>Cloud liquid amount</u>

Table A1: The variables from the perturbation runs inserted in the radiation code of CFRAM to calculate the temperature change in response to the changes in CO₂, O₃, water vapour, cloud and albedo.

Acknowledgements

The computations and simulations were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at National Supercomputer Center (NSC) in Linköping.

Hamish Struthers NSC is acknowledged for assistance concerning technical aspects in making the WACCM code run on NSC supercomputer Tetralith. We thank Qiang Zhang for helping to make the radiation model code applicable to WACCM model data.

Competing interests

The authors have no competing interests to declare.

References

- Akmaev, R.A., Fomichev, V.I. and Zhu, X.: Impact of middle-atmospheric composition changes on greenhouse cooling in the upper atmosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, 68, 1879-1889, doi:10.1016/j.jastp.2006.03.008, 2006.
- Beig, G., et al.: Review of mesospheric temperature trends, *Reviews of Geophysics*, 41(4), doi:10.1029/2002RG000121, 2003.
- Bony, S., and co-authors: How well do we understand and evaluate climate change feedback processes?, *Journal of Climate*, 19(15), 3445–3482, doi:10.1175/JCLI3819.1, 2006.
- Boucher, O., Randall, D., and co-authors: Clouds and Aerosols, in: *Climate Change: The Physical Science Basis. Contribution of Working Group I to IPCC AR5*, edited by: Stocker T.F. and coauthors., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

1356 Brasseur, G. P., and Solomon, S.: *Aeronomy of the middle atmosphere,*
1357 *Chemistry and physics of the stratosphere*, Springer, New York, 2005.

1358 Brewer, A. W.: Evidence for a world circulation provided by the measurements
1359 of helium and water vapour distribution in the stratosphere. *Quarterly Journal*
1360 *of the Royal Meteorological Society*, 75(326), 351-363,
1361 doi:10.1002/qj.49707532603, 1949.

1362 Brühl, C., & Crutzen, P. J.: Scenarios of possible changes in atmospheric
1363 temperatures and ozone concentrations due to man's activities, estimated
1364 with a one-dimensional coupled photochemical climate model, *Climate*
1365 *Dynamics*, 2(3), 173-203, doi: /10.1007/BF01053474, 1988.

1367 Butchart, N., and co-authors: Chemistry–climate model simulations of twenty-
1368 first century stratospheric climate and circulation changes, *Journal of Climate*,
1369 23(20), 5349–5374, doi:10.1175/2010JCLI3404.1, 2010.

1370 Caldwell, P.M., Zelinka, M.D., Taylor, K.E., Marvel, K.: Quantifying the
1371 sources of intermodal spread in equilibrium climate sensitivity, *Journal of*
1372 *Climate*, 29, 513-524, doi:10.1175/JCLI-D-15-0352.1, 2016.

1373 Cai, M., and Lu, J.: A new framework for isolating individual feedback
1374 processes in coupled general circulation climate models. Part II: Method
1375 demonstrations and comparisons, *Climate dynamics*, 32(6), 887-900,
1376 doi:10.1007/s00382-008-0424-4, 2009.

1377
1378 Cariolle, D.: The ozone budget in the stratosphere: Results of a one-
1379 dimensional photochemical model, *Planetary and Space Science*, 31(9),
1380 1033-1052, doi:10.1016/0032-0633(83)90093-4, 1983.

1381
1382 Deckert, R. and Dameris, M.: Higher tropical SSTs strengthen the tropical
1383 upwelling via deep convection, *Geophysical Research Letters*, 35(10), doi:
1384 10.1029/2008GL033719, 2008.

1385
1386 Dobson, G. M. B.: Origin and distribution of the polyatomic molecules in the
1387 atmosphere. *Proceedings of the Royal Society of London. Series A.*
1388 *Mathematical and Physical Sciences*, 236(1205), 187-193,
1389 doi:10.1098/rspa.1956.0127, 1956.

1390
1391 Fomichev, V.I., Jonsson, A.I., De Grandpre, J., Beagley, S.R., McLandress,
1392 C., Semeniuk, K., Shepherd, T.G.: Response of the middle atmosphere to
1393 CO₂ doubling: Results from the Canadian Middle Atmosphere Model, *Journal*
1394 *of Climate*, 20(7), 1121-1141, doi:10.1175/JCLI4030.1, 2007.

1395
1396 Fu, Q., and Liou, K. N.: On the correlated k-distribution method for radiative
1397 transfer in nonhomogeneous atmospheres. *Journal of the Atmospheric*
1398 *Sciences*, 49(22), 2139-2156, doi:10.1175/1520-
1399 0469(1992)049<2139:OTCDMF>2.0.CO;2, 1992.

1400
1401

1402 Fu, Q., and Liou, K. N.: Parameterization of the radiative properties of cirrus
 1403 clouds. *Journal of the Atmospheric Sciences*, 50(13), 2008-2025, doi:
 1404 10.1175/1520-0469(1993)050<2008:POTRPO>2.0.CO;2 1993.
 1405
 1406 Hu, X., Y. Li, S. Yang, Y. Deng and Cai. M.: Process-based decomposition of
 1407 the decadal climate difference between 2002-13 and 1984-95, *J. Climate*, 30,
 1408 4373–4393, doi: 10.1175/JCLI-D-15-0742.1, 2017.
 1409
 1410 Hurrell, J.W., et al.: The Community Earth System Model: A framework for
 1411 collaborative research. *Bulletin of the American Meteorological Society*, 94(9),
 1412 1339-1360, doi: 10.1175/BAMS-D-12-00121.1, 2013.

 1413 Jaiser, R., K. Dethloff and Handorf, D.: Stratospheric response to Arctic sea
 1414 ice retreat and associated planetary wave propagation changes, *Tellus A:
 1415 Dynamic Meteorology and Oceanography*, 65(1), 19375, doi:
 1416 10.3402/tellusa.v65i0.19375, 2013.

 1417 Jonsson, A.I., de Grandpré, J., Fomichev, V.I., McConnell, J.C., Beagley,
 1418 S.C.: Doubled CO₂-induced cooling in the middle atmosphere: Photochemical
 1419 analysis of the ozone radiative feedback, *Journal of Geophysical Research*,
 1420 109, D24103, doi:10.1029/2004JD005093, 2004

 1421 Kinnison, D.E., Brasseur, G.P., Walters, S., Garcia, R.R. Marsh, D.R, Sassi,
 1422 F., Harvey, V.L., Randall, C.E., Emmons, L., Lamarque, J.F., Hess, P.,
 1423 Orlando, J.J., Tie, X.X., Randall, W., Pan, L.L., Gettelman, A., Granier, C.,
 1424 Diehl, T., Niemeijer, Y., Simmons, A.J.: Sensitivity of chemical tracers to
 1425 meteorological parameters in the MOZART-3 chemical transport model,
 1426 *Journal of Geophysical Research: Atmospheres*, 112, D20302,
 1427 doi:10.1029/2006JD007879, 2007.
 1428
 1429 Langematz, U.: Stratospheric ozone: down and up through the
 1430 anthropocene, *ChemTexts* 5, 8, doi/10.1007/s40828-019-0082-7, 2019.

 1431 Lindzen, R.S.: Turbulence stress owing to gravity wave and tidal breakdown,
 1432 *Journal of Geophysical Research: Oceans*, 86.C10, 9707–9714,
 1433 doi:10.1029/JC086iC10p09707, 1981.

 1434
 1435 Lu, J., and Cai, M.: A new framework for isolating individual feedback
 1436 processes in coupled general circulation climate model. Part I: Formulation.
 1437 *Climate dynamics*, 32 (6), 873–885, doi:10.1007/s00382-008-0425-3, 2009.
 1438
 1439 Manabe, S., & Wetherald, R. T.: The effects of doubling the CO₂
 1440 concentration on the climate of a general circulation model. *Journal of the
 1441 Atmospheric Sciences*, 32(1), 3-15, 1975.

 1442 Marsh, D.R., Mills, M.J. Kinnison, D.E., Lamarque, J.F., Calvo, N., Polvani,
 1443 L.M.: Climate change from 1850 to 2005 simulated in CESM1(WACCM),
 1444 *Journal of Climate*, 26, 7372–7391, doi:10.1175/JCLI-D-12-00558.1, 2013.

1445 Marsh, D.R., Lamarque, J.-F., Conley, A.J. and Polvani, L.M., Stratospheric
 1446 ozone chemistry feedbacks are not critical for the determination of climate
 1447 sensitivity in CESM1(WACCM), *Geophysical Research Letters*, 43, 3928–
 1448 3934, doi:10.1002/2016GL068344, 2016.

1449 McFarlane, N.A., The effect of orographically excited wave drag on the
 1450 general circulation of the lower stratosphere and troposphere, *Journal of*
 1451 *Atmospheric Sciences*, 44(14), 1775–1800,
 1452 doi:10.1175/15200469(1987)044<1775:TEOOEG>2.0.CO;2, 1987.

1453 Neale, R., Richter, J., Park, S., Lauritzen, P., Vavrus, S., Rasch, P. and
 1454 Zhang, M.: The mean climate of the Community Atmosphere Model (CAM4) in
 1455 forced SST and fully coupled experiments, *Journal of Climate*, 26, 5150–
 1456 5168, doi:10.1175/JCLI-D-12-00236.1, 2013.

1457 Nowack, P.J., Abraham, N.L., Maycock, A.C., Braesicke, P., Gregory, J.M.,
 1458 Joshi, M.M., Osprey, A., Pyle, J.A.: A large ozone-circulation feedback and its
 1459 implications for global warming assessments, *Nature Climate Change*, 5 (1),
 1460 41-45, 2015, doi:10.1038/NCLIMATE2451, 2015.

1461 Oberländer, S., Langematz, U. and Meul, S.: Unraveling impact factors for
 1462 future changes in the Brewer-Dobson circulation, *Journal of Geophysical*
 1463 *Research: Atmospheres*, 118, 10296-10312, doi:10.1002/jgrd.50775, 2013.

1464
 1465 Ramaswamy, V., Collins, W., Haywood, J., Lean, J., Mahowald, N., Myhre,
 1466 G., Naik, V., Shine, K.P., Soden, B., Stenchikov, G., Storelvmo, T., 2019:
 1467 Radiative forcing of climate: The historical evolution of the radiative forcing
 1468 concept, the forcing agents and their quantification, and application,
 1469 *Meteorological Monographs* 59, 14.1 14.99,
 1470 doi:10.1175/AMSMONOGRAPHS-D-19-0001.1
 1471

1472 Ramaswamy, V., et al.: Stratospheric temperature trends: Observations and
 1473 model simulations, *Reviews of Geophysics* 39.1, 71-122, doi:
 1474 10.1029/1999RG000065, 2001.

1475
 1476 Richter, J.H., Sassi, F., Garcia, R.R.: Toward a physically based gravity wave
 1477 source parameterization in a general circulation model, *Journal of the*
 1478 *Atmospheric Sciences*, 67, 136–156, doi:10.1175/2009JAS3112.1, 2010.

1479 Rieger, V.S., Dietmüller, S., Ponater, M.: Can feedback analysis be used to
 1480 uncover the physical origin of climate sensitivity and efficacy differences?,
 1481 *Climate Dynamics* 49, 2831-2844, doi: 10.1007/s00382-016-3476-x 2017.

1482 Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., &
 1483 Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS
 1484 composition and related radiative effects, *Journal of Geophysical Research:*
 1485 *Atmospheres*, 117(D16), doi:10.1029/2012JD017751, 2012.

1486

1487 Royer, J.F., Planton, S., Déqué, M.: A sensitivity experiment for the removal
 1488 of Arctic sea ice with the French spectral general circulation model, *Climate*
 1489 *Dynamics*, 5(1), 1-17, doi:10.1007/BF00195850, 1990.

1490
 1491 Schmidt, H., Brasseur, G.P, Charron, M., Manzini, E., Giorgetta, M.A., Diehl,
 1492 T., Fomichev, V., Kinnison, D., Marsh, D., Walters, S., The HAMMONIA
 1493 Chemistry Climate Model: Sensitivity of the mesopause region to the 11-year
 1494 solar cycle and CO₂ doubling, *Journal of Climate*, 19(16), 3903-3931,
 1495 doi:10.1175/JCLI3829.1, 2006.

1496
 1497 Shaw, T.A., and Shepherd T.G.: Atmospheric science: Raising the
 1498 roof, *Nature geoscience*, 1(1), 12, doi:10.1038/ngeo.2007.53, 2008.

1499
 1500 Shepherd, T.G.: Dynamics, stratospheric ozone and climate change,
Atmosphere-Ocean, 46,1, 117-138, doi:10.3137/ao.460106, 2008.

1501
 1502 Shepherd, T.G., and McLandress, C.: A robust mechanism for strengthening
 1503 of the Brewer–Dobson circulation in response to climate change: critical layer
 1504 control of subtropical wave breaking, *Journal of the Atmospheric Sciences*,
 68, 4, 784-797, doi:10.1175/2010JAS3608, 2011.

1505
 1506 Shine, K. P., et al.: A comparison of model-simulated trends in stratospheric
 1507 temperatures, *Quarterly Journal of the Royal Meteorological Society: A journal*
 1508 *of the atmospheric sciences, applied meteorology and physical*
 1509 *oceanography* 129(590),1565-1588, doi:10.1256/qj.02.186, 2003.

1510
 1511 Sigmond, M., Siegmund, P.C., Manzini, E. and Kelder, H.: A simulation of the
 1512 separate climate effects of middle-atmospheric and tropospheric CO₂
 1513 doubling, *Journal of Climate*, 17(12), 2352-2367, doi:10.1175/1520-
 0442(2004)017<2352:ASOTSC>2.0.CO;2, 2004.

1514
 1515 Soden, B., Held, I.M.: An assessment of climate feedbacks in coupled ocean-
 1516 atmosphere models, *Journal of Climate*, 19, 3354-3360,
 doi:10.1175/JCLI3799.1, 2006.

1517
 1518 Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M.,
 1519 Sanford, T. J., & Plattner, G. K.: Contributions of stratospheric water vapor to
 1520 decadal changes in the rate of global warming, *Science*, 327(5970), 1219-
 1521 1223, doi: 10.1126/science.1182488, 2010.

1522
 1523 Song, X., and Zhang, G.J.: Role of climate feedback in El Niño-like SST
 1524 response to global warming, *Journal of Climate*, 27, 7301–7318,
 1525 doi:10.1175/JCLI-D-14-00072.1, 2014

1526
 1527 Taylor, P.C., Cai, M., Hu, A., Meehl, J., Washington, W. and Zhang, G.J.: A
 1528 decomposition of feedback contributions to polar warming amplification,
Journal of Climate, 26, 7023–7043, doi:10.1175/JCLI-D-12-00696.1, 2013.

1529

1530 WMO (World Meteorological Organization), *Scientific Assessment of Ozone*
 1531 *Depletion: 2018*, Global Ozone Research and Monitoring Project – Report No.
 1532 58, 588 pp., Geneva, Switzerland, 2018.

1533
 1534 Zhang, P., Wu, Y., Simpson, I.R., Smith, K.L., Zhang, X., De, B., and
 1535 Callaghan, P.: A stratospheric pathway linking a colder Siberia to Barents-
 1536 Kara Sea sea ice loss, *Science Advances*, 4(7), eaat6025, doi:
 1537 10.1126/sciadv.aat6025, 2018.

1538
 1539 Zheng, J., Zhang, Q., Li, Q., Zhang, Q. and Cai, M., Contribution of sea ice
 1540 albedo and insulation effects to Arctic amplification in the EC-Earth Pliocene
 1541 simulation, *Climate of the Past*, 15, 291-305, doi:10.5194/cp-15-291-2019,
 1542 2019.

1543
 1544 Zhu, X., Yee, J.-H., Cai, M., Swartz, W.H., Coy, L., Aquila, V., Garcia, R.,
 1545 Talaat, E.R.: Diagnosis of middle-atmosphere climate sensitivity by the
 1546 climate feedback-response analysis method, *Journal of Atmospheric*
 1547 *Sciences*, 73(1), 3-23, doi:10.1175/JAS-D-15-0013.1, 2016.