Response to Anonymous Referee #1

General Comment. The paper evaluates the impact of trans-regional transport of aerosol from the Po Plain to the Aosta Valley (north-western Italian Alps), by means of both the analysis of experimental data and numerical simulations. The paper is complete, well-written and can be of interest for the community. Therefore in my opinion it is worth publishing in Atmospheric Chemistry and Physics, after a few minor comments are addressed.

We thank the reviewer for taking the time to revise our manuscript and for his/her pertinent comments. Our point-to-point reply is given hereafter (the text in italics represents a citation of the revised manuscript and the figure references follow the updated numbering).

Referee's comment 1. Meteorological model: the Authors say (pag. 8, line 7) that "we used a nudged, high-resolution variant, called COSMO-I2, covering Italy". However, in my opinion 2.8 km is not high-resolution in complex terrain. The floor of the Aosta Valley is 2-3 km wide, so meteorological phenomena triggered by the orography may be not well resolved by the model at this resolution, also considering the fact that the actual model resolution is 6-8 times the grid cell (Skamarock 2004). In fact authors say that the model surface altitude of Aosta urban area is 900 m a.s.l., whereas the actual altitude is 600 m a.s.l. For example Schmidli et al. (2018) showed that at 2.2-km resolution the COSMO model poorly simulates valley winds, while at 1.1-km resolution the diurnal cycle of the valley winds is well represented. Similarly, Giovannini et al. (2014) showed that 2-km resolution can be considered as the limit for a good representation of valley winds in narrow Alpine valleys.

Author's response 1. We understand the point raised by the reviewer. In the submitted version, we used the terminology "high resolution", as this appears in the COSMO web page (http://www.cosmo-model.org/content/tasks/operational/default.htm), where COSMO is indeed defined as a "high-resolution" model.

However, we accept the reviewer's remark and thus removed the term "high-resolution" in the revised text. Also, in Sect. 3.1 we now discuss the advantages of COSMO-I2 version over COSMO-ME, with reference to the high*er* resolution of the former (but not "high" in absolute terms). The revised text reads as follows:

The forecasts, inclusive of the complete set of parameters (such as the 3-D wind velocity used here) for eight time steps (from 00 to 21 UTC), are disseminated daily in two different configurations by the meteo-rological operative centre – air force meteorological service (COMET): a lower-resolution version (COSMO-ME, 7 km horizontal grid and 45 levels vertical grid, 72 hours integration), covering central and southern

Europe, and a nudged, higher-resolution version (COSMO-I2 or COSMO-IT, 2.8 km, 65 vertical levels, 2 runs/day), covering Italy (orange rectangle in Fig. 1a). Owing to the complex topography of the Aosta Valley, and the consequent need to resolve as much as possible the atmospheric circulation at small spatial scales, we used the latter version in the present work (cf. Sect. 4.5 for a discussion about possible effects of the finite model resolution in complex terrain).

Furthermore, following the reviewer's comment, we now preferably use the term "grid spacing/step" instead of "resolution" throughout the revised text.

We also added the suggested references in Sect. 4.5:

Although the COSMO-I2 grid spacing is certainly in line with that of other state-of-the-art operational limited-area models, in our complex terrain it could be insufficient to appropriately resolve local meteorological phenomena triggered by the valley orography (Wagner et al., 2014), also considering that the actual model resolution is 6–8 times the grid cell (Skamarock, 2004). For example, Schmidli et al. (2018) show that at 2.2-km grid step the COSMO model poorly simulates valley winds, while at 1.1-km grid step the diurnal cycle of the valley winds is well represented. Similarly, Giovannini et al. (2014) show that 2km step can be considered as the limit for a good representation of valley winds in narrow Alpine valleys. Smoothed digital elevation model (DEM) used within COSMO and FARM could also play a direct role in the detected underestimation. In fact, as mentioned in Sect.3.2, the model surface altitude of the Aosta urban area is 900 m a.s.l., whereas the actual altitude is about 580 m a.s.l. (Table 1). The adjacent cells are given an even higher altitude, owing to the fact that the valley floor and the neighbouring mountain slopes are not properly resolved at the current resolution. This results in an apparent lower elevation of the sites located in flatter and wider areas, such as Aosta, while the real topography profile at the bottom of the valley presents a monotonic increase from the Po plain to Aosta. Therefore, it is expected that, just by better reproducing the orography, a higher resolution would allow to better resolve the horizontal advection of aerosol-laden air from much lower altitudes on the plain.

Most of these issues were extensively addressed in the companion paper (Diémoz et al., 2019). In that study, COSMO-I2 was shown to be capable of reproducing the mountain-plain wind patterns observed at the surface both on average (cf. Fig. S1 in the companion paper) and in specific case studies (Fig. S13 therein). Nevertheless, it was also found to slightly anticipate in time and overestimate the easterly diurnal winds in the first hours of the afternoon and to overestimate the nighttime drainage winds (katabatic winds, ventilating the urban atmosphere and reducing pollutant loads). These limitations were mostly attributed to the finite resolution of the model.

RC2. Building on the previous consideration, it is not easy to evaluate the performance of the chemical model without a validation of the meteorological model. The Authors attribute most of the discrepancies with respect to measurements to deficiencies in the boundary conditions of the chemical model. However, this statement is difficult to be demonstrated without a complete validation of the modelling chain. Moreover, increasing the boundary conditions by a factor 4 reduces the mean bias, but does not reduce the RMSE (Table 3). So it seems that there is still a compensation of underestimations and overestimations.

AR2. One of the most relevant updates in the revised manuscript is the study of the chemical transport model performances in a flatter terrain compared to the complex orography of the mountain valley. This is accomplished by including the PM_{10} dataset measured in a station at the boundary of the domain, Ivrea. This addition clearly showed that the same PM_{10} underestimation of the model is also obtained over flat terrain, demonstrating this is not driven by inaccuracies of the simulated wind fields over the more complex terrain of Aosta. Therefore, Sect. 4.5 has been modified to better explain this point and investigate the relative importance of both emissions (inventories/unaccounted physical processes) and transport (performances of the meteorological model) in generating the model-measurements differences:

The observed model-measurement discrepancies might originate from (1) an incomplete representation of the inventory sources (emission component), (2) inaccurate NWP modelling of the meteorological fields, and notably the wind (transport component), or a combination of (1) and (2). Some details on these aspects are provided in the following.

1. Emissions. Inaccuracies in the (local and national) emission inventories could degrade the comparison between simulations and observations. Based on results shown in Fig. 17(a,b), we decided to investigate the sensitivity of our simulations in Aosta-Downtown to the magnitude of the external contributions (boundary conditions). In particular, we used a simplified approach to speed up the calculations: assuming that the contribution from outside the regional domain and the local emissions add up without interacting, we simulated the surface PM_{10} concentrations turning on/off the boundary conditions (dark- and light-blue lines in Fig. 16) to roughly estimate the only contribution from sources outside the regional domain. Interestingly, the difference between the two FARM runs correlates well with the advection classes observed by the ALC (Fig. 18). This clear correlation between the simulations and the experimentally-determined atmospheric conditions is a first good indication that the NWP model used as input to the CTM yields reasonable meteorological inputs to FARM. Then, to further explore the sensitivity to the boundary conditions, we gradually increased, by a weighting factor W, the PM_{10} concentration from outside the boundaries of the domain trying to match the observed values. In Figs. 17(c, d) we show the results obtained with W=4. This exercise show that the overall mean bias error is much reduced compared to the original simulations (W=1, Figs. 17(a, b)), especially for the winter and autumn seasons, while slight overestimations are now visible for summer and spring. Also the annually-averaged MBE, NMBE and NMSD improve (Table 3), whilst the other statistical indicators remain stable or even slightly worsen.

Clearly, our simplified test has major limitations. (...)

2. Although the COSMO-12 grid spacing is certainly in line with that of other state-of-the-art operational limited-area models, in our complex terrain it could be insufficient to appropriately resolve local meteorological phenomena triggered by the valley orography (Wagner et al., 2014), also considering that the actual model resolution is 6–8 times the grid cell (Skamarock, 2004). (...)

Also, to disentangle the role of factors (1) and (2) discussed above, we evaluated the model performances in a flatter area of the domain, where simulations are expected to be less affected by the complex orography of the mountains. We therefore compared PM_{10} simulations and measurements in Ivrea (Fig. 16b). This test is also useful to operate the CTM model at the boundaries of the domain, with special focus on the emissions from the "boundary conditions". Model underestimation in both the warm and cold seasons is evident. Incidentally, it can be noticed that concentrations simulated without "boundary conditions" are nearly zero, since the considered cell is far from the strongest local sources and most part of the aerosol comes from outside the regional domain. As done for Aosta (...), Fig. 19(a,b) presents the absolute and relative differences between the model and the observations in Ivrea. The overall NMBE is -0.73, which rather well corresponds to the underestimation factor W=4 (or NMBE=-0.75) found for Aosta–Downtown and other sites in the valley. Since it is expected that in this flat area the NWP model is able to better resolve the circulation compared to the mountain valley, this result suggests poor CTM performances to be mostly related to underestimated emissions at the boundaries rather than to incorrect transport. In addition to this general underestimation, a large scatter between observations and simulations can be noticed in Fig. 16, the linear correlation index between simulations and observation in Ivrea being rather low (ρ =0.54). If such a large scatter, due to the erroneous boundary conditions, is already detectable at the border of the domain, it is likely that at least part of the RMSE reported in Table 3 for Aosta–Downtown can be attributed to inaccuracies in the national inventory.

We report the new figures 16 and 19, previously mentioned in the text, here below. Fig. 17 is shown in AR3.



Figure 16: Long-term (1 year) comparison between PM₁₀ surface measurements and simulations (FARM) in Aosta–Downtown (a) and Ivrea (b).



Figure 19: Absolute (a) and relative (b) differences between simulated and observed PM_{10} concentrations at the surface in Ivrea.

Note that, in accordance with the inclusion of the additional measurement site of Ivrea, Sect. 2 has been integrated as follows:

We also consider in our analysis the PM_{10} records collected in the city of Ivrea, a site in the Po Plain (31 µg m^{-3} average PM_{10} in 2017) located just outside the Aosta Valley, in the Italian Piedmont region (...). This was done to check if and how much inaccuracies of the model in reproducing the aerosol loads over the Aosta Valley are due to difficulties in simulating the wind field in such a complex terrain. In fact, the city centre of Ivrea (24000 inhabitants) is approximately in the middle between the measurement site (south of the city) and the nearest cell of our model domain (north of the city), the distance between these two points being only 2 km. The altitude of the cell (from the digital elevation model used in our simulations) is 241 m a.s.l., i.e. approximately the real one (i.e., 243 m a.s.l). (...) The station of Ivrea features, among other instruments, a TCR Tecora Charlie/Sentinel PM_{10} sampler. PM_{10} concentrations are then determined by a gravimetric technique.

Finally, following the reviewer's comment, we also mention in the revised manuscript that *an additional contribution to the observed RMSE could still be given by errors in modelling transport due to the coarse resolution of the NWP model, although we are not able to quantify the relative role of the two factors. To unravel this issue, high-resolution models (grid step 0.5 km, e.g. Golzio and Pelfini, 2018; Golzio et al., 2019) are being tested on the investigated area and their results will be addressed in future studies.*

RC3. In Figure 17 it would be interesting to see also a normalized mean bias (i.e. a mean bias normalized by the average concentration measured for each class).

AR3. Figure 17 has been modified taking the reviewer's advice into account (see panels b and d). References to the Donnas station have been removed following comment RC20 by Referee#2. The new figure is reported below.



Figure 17: Absolute (a,c) and relative (b,d) differences between simulated and observed PM_{10} concentrations at the surface in Aosta–Downtown. The mean bias error (MBE) and the normalised mean bias error (NMBE) for each case are reported in the plot titles. First row: FARM simulations as currently performed by ARPA. Second row: the PM_{10} concentrations from outside the boundaries of the domain were multiplied by a factor W=4.

RC4. Page 1, line 4: ... 3-year period ... Section 4.3: The two final sentences begin with "Finally". Page 36, line 18: ... this regular air mass transport ...

AR4. Thank you for these technical remarks. All suggested changes have been accepted in the revised manuscript.

RC5. References [...]

AR5. All recommended references have been included in the revised manuscript.

References

- Diémoz, H., Barnaba, F., Magri, T., Pession, G., Dionisi, D., Pittavino, S., Tombolato, I. K. F., Campanelli, M., Della Ceca, L. S., Hervo, M., Di Liberto, L., Ferrero, L., and Gobbi, G. P.: Transport of Po Valley aerosol pollution to the northwestern Alps – Part 1: Phenomenology, Atmos. Chem. Phys., 19, 3065– 3095, doi:10.5194/acp-19-3065-2019, 2019.
- Giovannini, L., Antonacci, G., Zardi, D., Laiti, L., and Panziera, L.: Sensitivity of Simulated Wind Speed to Spatial Resolution over Complex Terrain, Energy Proced., 59, 323–329, doi:10.1016/j.egypro.2014.10. 384, 2014.
- Golzio, A. and Pelfini, M.: High resolution WRF over mountainous complex terrain: testing over the Ortles Cevedale area (Central Italian Alps), in: Conference: First National Congress, Italian Association of Atmospheric Sciences and Meteorology (AISAM), AISAM, 2018.
- Golzio, A., Ferrarese, S., Cassardo, C., Diolaiuti, G. A., and Pelfini, M.: Resolution improvement in WRF over complex mountainous terrain, in preparation for Boundary-Layer Meteorol., 2019.
- Schmidli, J., Böing, S., and Fuhrer, O.: Accuracy of Simulated Diurnal Valley Winds in the Swiss Alps: Influence of Grid Resolution, Topography Filtering, and Land Surface Datasets, Atmosphere, doi:10. 3390/atmos9050196, 2018.
- Skamarock, W. C.: Evaluating Mesoscale NWP Models Using Kinetic Energy Spectra, Mon. Weather Rev., 132, 3019–3032, doi:10.1175/MWR2830.1, 2004.
- Wagner, J. S., Gohm, A., and Rotach, M. W.: The Impact of Horizontal Model Grid Resolution on the Boundary Layer Structure over an Idealized Valley, Mon. Weather Rev., 142, 3446–3465, doi:10.1175/ MWR-D-14-00002.1, 2014.

Response to Anonymous Referee #2

General Comment. The manuscript presents a comprehensive analysis of meteorological (standard weather parameters, ceilometer) and air quality (PM10 characterization) measurements over a threeyear period in the Aosta Valley in north-western Italy. The study convincingly demonstrates that a non-negligible fraction of the particulate matter concentration observed at receptor sites in the valley is advected from the neighboring Po plain, occasionally providing a decisive contribution to the exceedance of concentration limit values. The study also shows that advected aerosols differ from the locally emitted ones in their physical-chemical properties. Results from a chemical transport modelling chain are evaluated, demonstrating that the effects of advection from the Po plain can be represented only qualitatively from state-of-the art operational model chains.

Methodology and results are scientifically sound. The manuscript is written in good English and figures are generally of adequate quality. Suggestions for a few minor modifications should be taken into account before publication can be recommended. Details are provided below. Please note that the assessment is based mostly on the meteorological aspects of the study, reflecting the expertise of the reviewer.

We thank the reviewer for taking the time to revise our manuscript and for his/her pertinent comments. Our point-to-point reply is given hereafter (the text in italics represents a citation of the revised manuscript and the figure references follow the updated numbering).

Referee's comment 1. The superficial reader might conclude that pollution in the Aosta valley mostly depends on advection from the Po plain. While this is true to some extent, one should also consider that easterly advection is most common in summer (Figure 3), that is, when PM10 concentrations in Aosta are at their yearly minimum (Figure 8). Based on these contrasting statements, another superficial reader (with an opposite bias) would conclude that easterly advection actually cleans the air. The latter is obviously a flawed argument, still it shows that it is important to carefully delimit the message of the study. A balanced perspective on the whole issue could be the following: easterly advection from the Po valley, which is most frequent in summer but possible in all seasons, may combine with adverse weather conditions (low-level inversions) locally worsened by the valley topography, greatly enhancing pollution levels. This is especially true in the coldest part of the year, when emissions are highest. I feel that, in the introductory and concluding remarks, the authors could do a better job in explaining these subtleties.

Author's response 1. We thank the reviewer for raising this possible ambiguity. In the revised manuscript (Sect. 4.3.2), we now split the impact of transport on PM_{10} levels by season. In this way, we better show

that the overall impact actually depends on the coupling between frequency and severity of the episodes, and also on the background local pollution levels. The revised text reads as follows:

On a seasonal basis, the absolute impact of air masses transport depends on the coupling between emissions (stronger in winter and weaker in summer) and weather regimes (e.g., thermal winds occurring more frequently in summer/autumn with respect to winter/spring, Sect. 4.1). This results in a PM_{10} contribution of nearly 6 µg m⁻³ in winter and autumn, 4 µg m⁻³ in summer, and 3 µg m⁻³ in spring. In terms of relative contribution, this also depends on the "background" PM_{10} levels in Aosta. It is therefore highest in summer (32%), when thermally-driven fluxes are more frequent and local emissions lower, and lowest in winter (16%), when thermal winds are less frequent and local emissions higher. Intermediate relative values are found in spring and summer (27% and 28%, respectively). In specific episodes, advections from the Po basin may still produce an increase of (...)

Following the reviewer's comment, we also updated the Conclusions accordingly:

Using their relevant scores as a proxy, we estimate an average 25% contribution of these transported nitrate and sulfate-rich components to the Aosta PM_{10} . This impact varies on a seasonal basis. The relative contribution of non-local PM_{10} is highest in summer (32%), when advection is most frequent and local PM_{10} is lowest, while it is lowest in winter (16%), when advection is least frequent and local PM_{10} is highest. In absolute terms the reverse occurs and the impact of transport is found to be highest in winter/autumn, reaching levels of 50 µg m⁻³ in some episodes (thus exceeding alone the EU legislative limits). This occurs due to the superposition of advected particles with higher background concentrations in the coldest part of the year, when emissions are highest and pollution is further enhanced by adverse weather conditions, i.e. low-level inversions, locally worsened by the valley topography. Note, though, that in this study the impact of Po Valley advection on air quality was mostly quantified for the Aosta–Downtown station. In rural and remote sites of the region this non-local contribution is expected to be even larger, in relative terms (...)

Finally, where we now specify already in the Introduction that the impact on air quality depends *on the interplay between frequency and severity of the episodes*.

RC2. The abstract is very long. ACP doesn't seem to set length limits, but it is in the interest of the authors to provide a more concise formulation. Please refrain from using references to published literature in the abstract.

AR2. Thank you for this suggestion. We agree with it. The abstract has been shortened by more than 700 characters and bibliographic references have been removed. It now reads:

This work evaluates the impact of trans-regional aerosol transport from the polluted Po basin on particulate matter levels (PM_{10}) and physico-chemical characteristics in the northwestern Alps. To this purpose, we exploited a multi-sensor, multi-platform database over a 3-year period (2015–2017) accompanied by a series of numerical simulations. The experimental setup included operational (24/7) vertically resolved aerosol profiles by an Automated LiDAR-Ceilometer (ALC), vertically integrated aerosol properties by a sun/sky photometer, and surface measurements of aerosol mass concentration, size distribution and chemical composition. This experimental set of observations was then complemented by modelling tools, including Numerical Weather Prediction (NWP), Trajectory Statistical (TSM) and Chemical Transport (CTM) models, plus Positive Matrix Factorisation (PMF) on both the PM_{10} chemical speciation analyses and particle size distributions. In a first companion study, we showed and discussed through detailed case studies the 4-D phenomenology of recurrent episodes of aerosol transport from the polluted Po basin to the northwestern Italian Alps. Here we draw more general and statistically significant conclusions on the frequency of occurrence of this phenomenon, and on the quantitative impact of this regular, winddriven, aerosol-rich "atmospheric tide" on PM_{10} air quality levels in this alpine environment. Based on an original ALC-derived classification, we found that an advected aerosol layer is observed at the receptor site

(Aosta) in 93% of days characterized by easterly winds (i.e., from the Po basin) and that the longer the time spent by air masses over the Po plain the higher this probability. Frequency of these advected aerosol layers was found to be rather stable over the seasons with about 50% of the days affected. Duration of these advection events ranges from few hours up to several days, while aerosol layer thickness ranges from 500 up to 4000 m. Our results confirm this phenomenon to be related to non-local emissions, to act at the regional scale and to largely impact both surface levels and column-integrated aerosol properties. In Aosta, PM_{10} and AOD values increase respectively up to a factor of 3.5 and 4 in dates under the Po Valley influence. Pollution transport events were also shown to modify the mean chemical composition and typical size of particles in the target region. In fact, increase in secondary species, and mainly nitrate- and sulfate-rich components, were found to be effective proxies of the advections, with the transported aerosol responsible for at least 25% of the PM₁₀ measured in the urban site of Aosta, and adding up to over 50 μ g m⁻³ during specific episodes (thus exceeding alone the EU established daily limit). From a modelling point of view, our CTM simulations performed over a full year showed that the model is able to reproduce the phenomenon, but markedly underestimates its impact on PM_{10} levels. As a sensitivity test, we employed the ALC-derived identification of aerosol advections to re-weight the emissions from outside the boundaries of the regional domain in order to match the observed PM_{10} field. This simplified exercise indicated that an increase of such "external" emissions by a factor of 4 in the model is needed to halve the model PM_{10} maximum deviations and to significantly reduce the PM_{10} normalised mean bias forecasts error (from -35% to 5%).

RC3. Page 1, lines 17-18. "maximises" → "is highest", "minimizes" → "is lowest".

AR3. The wording has been changed in the new abstract (see AR2).

RC4. Page 2, line 27. There is a double blank space before "WMO".

AR4. Corrected.

RC5. Page 3, Figure 1. Please align image borders (merely for aesthetic reasons).

AR5. Done. The updated Fig. 1 is shown here below. This also includes other modifications as required by the revision process.



Figure 1: True colour corrected reflectance from MODIS Terra satellite (http://worldview.earthdata.nasa. gov) on 17 March 2017. (a) Italy, with indication of the Alpine and Po Valley regions, of the Aosta Valley FARM regional domain (light blue rectangle), and the COSMO-I2 domain (orange rectangle, approximately corresponding to the boundaries of the national inventory). (b) Zoom over the Aosta Valley. The circle markers represent the sites of 1) Aosta–Downtown; 2) Aosta–Saint-Christophe; 3) Donnas; 4) Ivrea. A thin aerosol layer over the Po basin starting to spread out into the Alpine valleys is visible in both figures.

RC6. Page 3, line 15. Remove "the main of which are".

AR6. Removed.

RC7. Page 4, line 31. Thermally-driven winds include a night-time component as well (drainage flows along slopes and downvalley winds), which can ventilate the urban atmosphere and reduce pollutant loads. Are nocturnal breezes not observed in Aosta?

AR7. Thanks for pointing this out. Indeed, we extensively studied nocturnal breezes in the companion paper (Diémoz et al., 2019), since they can be relevant for the daily cycle of pollutant concentrations. Some of those results are now mentioned in the revised manuscript (Sect. 4.5):

(...) Most of these issues were extensively addressed in the companion paper (Diémoz et al., 2019). In that study, COSMO-I2 was shown to be capable of reproducing the mountain-plain wind patterns observed at the surface both on average (cf. Fig. S1 in the companion paper) and in specific case studies (Fig. S13 therein). Nevertheless, it was also found to slightly anticipate in time and overestimate the easterly diurnal winds in the first hours of the afternoon and to overestimate the nighttime drainage winds (katabatic winds, ventilating the urban atmosphere and reducing pollutant loads). These limitations were mostly attributed to the finite resolution of the model.

We have also updated the Introduction, which now reads:

The topography of the area triggers some of the most common weather regimes in the mountains, such as thermally-driven, *up-valley (daytime) and down-valley (nighttime) winds, up-slope (daytime) and*

down-slope (nighttime) winds, "Foehn" winds (Seibert, 2012) from the west, and frequent temperature inversions during wintertime anticyclonic days (...)

RC8. Page 4, line 32. Strictly speaking, Foehn is not necessarily a rain-shadow wind. Even if the (quite inaccurate) textbook picture of Foehn emphasizes the impact of upstream latent heat release due to condensation, Foehn is in most cases connected only to low-level blocked flow upstream of the orographic obstacle. Foehn may develop even if there is no upstream precipitation. Warming mostly occurs through the adiabatic descent of unblocked air from levels above the mountain tops (see for instance this DOI: 10.1127/0941-2948/2012/0398).

AR8. Thank you for this remark. We have removed the term "rain-shadow" from the text and we have included the suggested reference (the revised paragraph is already reported in AR7).

RC9. Page 7, table 1. Footnotes b and c. Unclear: is this the average data availability, or are data regularly available with this periodicity?

AR9. A short sentence has been added to Sect. 2 to better describe the *laboratory schedule (i.e., metal analyses on two consecutive days, EC/OC on the following two days, followed by two sequences metal-metal-EC/OC)*. Therefore, *over a 10 days period EC/OC concentrations are routinely provided on 4 days on average and metal concentrations on 6 days*.

The two footnotes of Table 1 now read: *The analysis is performed on 4* (or 6) *out of 10 days according to the laboratory schedule.*

RC10. Page 8, line 7. I am not sure that COSMO-I2 is a "high-resolution" model. Its grid spacing is certainly in line with that of other state-of-the-art operational limited-area-models. However, it is barely sufficient to appropriately resolve thermally-driven circulations. See for instance this DOI: 10.1175/MWR-D-14-00002.1. Please consider also comment 21 below.

AR10. We understand the point raised by the reviewer. In the submitted version, we used the terminology "high resolution", as this appears in the COSMO web page (http://www.cosmo-model.org/ content/tasks/operational/default.htm), where COSMO is indeed defined as a "high-resolution" model.

However, we accept the reviewer's remark and thus removed the term "high-resolution" in the revised text. Also, in Sect. 3.1 we now discuss the advantages of COSMO-I2 version over COSMO-ME, with reference to the high*er* resolution of the former (but not "high" in absolute terms). The revised text reads as follows:

The forecasts, inclusive of the complete set of parameters (such as the 3-D wind velocity used here) for eight time steps (from 00 to 21 UTC), are disseminated daily in two different configurations by the meteorological operative centre – air force meteorological service (COMET): a lower-resolution version (COSMO-ME, 7 km horizontal grid and 45 levels vertical grid, 72 hours integration), covering central and southern Europe, and a nudged, higher-resolution version (COSMO-I2 or COSMO-IT, 2.8 km, 65 vertical levels, 2

runs/day), covering Italy (orange rectangle in Fig. 1a). Owing to the complex topography of the Aosta Valley, and the consequent need to resolve as much as possible the atmospheric circulation at small spatial scales, we used the latter version in the present work (cf. Sect. 4.5 for a discussion about possible effects of the finite model resolution in complex terrain).

Furthermore, following the reviewer's comment, we now preferably use the term "grid spacing/step" instead of "resolution" throughout the revised text.

We have also included some of the reviewer's comments in Sect. 4.5:

Although the COSMO-I2 grid spacing is certainly in line with that of other state-of-the-art operational limited-area models, in our complex terrain it could be insufficient to appropriately resolve local meteorological phenomena triggered by the valley orography (Wagner et al., 2014), also considering that the actual model resolution is 6–8 times the grid cell (Skamarock, 2004). For example, Schmidli et al. (2018) show that at 2.2-km grid step the COSMO model poorly simulates valley winds, while at 1.1-km grid step the diurnal cycle of the valley winds is well represented. Similarly, Giovannini et al. (2014) show that 2km step can be considered as the limit for a good representation of valley winds in narrow Alpine valleys. Smoothed digital elevation model (DEM) used within COSMO and FARM could also play a direct role in the detected underestimation. In fact, as mentioned in Sect.3.2, the model surface altitude of the Aosta urban area is 900 m a.s.l., whereas the actual altitude is about 580 m a.s.l. (Table 1). The adjacent cells are given an even higher altitude, owing to the fact that the valley floor and the neighbouring mountain slopes are not properly resolved at the current resolution. This results in an apparent lower elevation of the sites located in flatter and wider areas, such as Aosta, while the real topography profile at the bottom of the valley presents a monotonic increase from the Po plain to Aosta. Therefore, it is expected that, just by better reproducing the orography, a higher resolution would allow to better resolve the horizontal advection of aerosol-laden air from much lower altitudes on the plain.

RC11. Page 9, lines 26-30. How sensitive are the TSM model results to the ad-hoc method of weighting the concentration fields on the number of trajectories through each cell? How is the weighting performed, exactly? Does it only reduce noise in the spatial fields, as stated, or does it also affect their magnitude?

AR11. It only reduces noise. To show this we added a figure in the Supplement (Fig. S7):



Figure S7: Same as Fig. 13 in the main paper, but without any weighting applied to the backtrajectory cells. (a) Output of the Concentration Field TSM, using the sum of the contributions from PMF nitrate- and sulfate-rich modes as concentration variable at the receptor (Aosta–Downtown). (b) CF based on the traffic and heating mode from PMF. Trajectories are cut at the borders of the COSMO-I2 domain.

The new figure (Fig. S7) was obtained without any weighting. No substantial differences can be noticed compared to Fig. 13.



Figure 13: (a) Output of the Concentration Field Trajectory Statistical Model, using the sum of the contributions from PMF nitrate- and sulfate-rich modes as concentration variable at the receptor (Aosta–Downtown). (b) Concentration Field based on the traffic and heating mode from PMF. Trajectories are cut at the borders of the COSMO-I2 domain.

We also tried to better describe the weighting procedure in the revised Sect. 3.2:

To reduce statistical noise, every value P_{ij} of the resulting map was then multiplied by a weighting factor, w_{ij} , linearly varying from 0 (for $N_{ij} \le 20$ end points in a cell) to 1 (for $N_{ij} \ge 200$), i.e. $w_{ij} = min\{1, \frac{max[0, N_{ij} - 20]}{180}\}$. To provide an idea of the effect of this weighting procedure, TSM maps without any weighting (i.e., $w_{ij} = 1$) have been included in the Supplement (Fig. S7). RC12. Page 10, line 20. "...was chosen based on the Q/Qexp ratio". Please clarify.

AR12. Section 3.3 now includes a short explanation of the Q/Q_{exp} ratio. The latter is the ratio between the objective function obtained with the selected number of factors (Q, introduced before) and its expected value (Q_{exp}). Elevated (i.e., > 2) Q/Q_{exp} ratios could indicate that some samples and/or species are not well modelled and could be better explained by adding another source (Norris and Duvall, 2014).

RC13. Page 13, table 2. It seems to me that the criteria outlined here are not mutually exclusive. For instance, would a 13-hour period with wind speed exceeding 4 m/s be classified as "downwind" or as "foehn"? I also have a few concerns about the terminology.

(A) "Upwind" and "downwind" typically identify position relative to the wind and to a specific location (e.g. "when the flow is southerly, the city is downwind of the mountains"). In this context, it might be more appropriate to speak of "Channeled synoptic flow from the west/east". In addition, it would be appropriate to specify wind direction ranges for the two classes.

(B) "Breezes" have by definition wind speed > 1.5 m/s (Beaufort scale). This would exclude wind calms at night. It might be better to speak of "Diurnal wind system".

(C) "Stability" refers to the thermal structure of the atmosphere. Because winds are emphasized here, it might be appropriate to speak of "Wind calm".

(D) The directional range for the Foehn class probably deserves being explained.

(E) Please use wind "direction" instead of "provenance". "Direction" refers conventionally to the direction the wind comes from.

AR13. Thanks for the remark, we acknowledge that the description of the criteria was not clear enough in the original manuscript. Actually, in our classification code the two classes (formerly called "Downwind" and "Foehn") were and are mutually exclusive. For example, in the case presented by the reviewer, the episode would be classified as "Foehn", since this category takes precedence over the synoptic winds. We now better specify this point in the updated Table 2 (below).

(A), (B) (C) and (E): the terminology suggested by the reviewer is now used throughout the revised manuscript (both the text and the figures have been updated).

(B): it must be noticed that nighttime and daytime conditions must be met for the same day. This is done in order to discriminate the "Diurnal wind systems" (inactive at night) from synoptic winds. A brace and a footnote in the revised Table 2 should make it clear, now.

(D): as explained in a new footnote, the directional range for the Foehn cases was determined experimentally by comparison with manual classification from a trained weather observer.

The changes to Table 2 are shown hereafter:

Primary classification	Secondary classification	Definition		
Easterly winds	Channelled synoptic flow from the east	$ v > 1 \text{ m s}^{-1}, \Delta t > 12 \text{ h}, \text{ wind direction } 0-180^{\circ}$ $\begin{cases} \text{Night (19-8 UTC)}^{a}: \text{ calm wind } (v < 1 \text{ m s}^{-1}) \end{cases}$		
	Diurnal wind systems (east)	$\begin{cases} & \text{or low westerly wind } (\boldsymbol{v} < 1.5 \text{ m s}^{-1}) \\ & \text{Day } (8-19 \text{ UTC})^{\text{a}}: & \text{easterly wind } (\boldsymbol{v} > 1.5 \text{ m s}^{-1}), \Delta t > 4 \text{ h} \end{cases}$		
Westerly winds	Foehn (west)	$ v > 4 \text{ m s}^{-1}$, $\Delta t > 4 \text{ h}$, wind direction $< 25^{\circ}$ or $> 225^{\circ}$ ^b , RH< 40%		
	Channelled synoptic flow from the west	no Foehn, $ \boldsymbol{v} >1~{\rm ms^{-1}},$ Δt >12 h, wind direction 180–360 $^\circ$		
Wind calm	_	$ v < 1 \text{ m s}^{-1}, \Delta t > 20 \text{ h}$		
Other	Precipitation	Accumulated daily precipitation $> 1 \text{ mm}$, $\Delta t > 4 \text{ h}$		
	Unclassified	Every day that does not fall in the previous categories		

Table 2. Classification of weather regimes used in the study, based on wind speed (|v|), wind direction, daily duration of the condition (Δt) and other measured meteorological variables.

^a Both conditions must be met in order to discriminate such diurnal wind systems (inactive at night) from synoptic winds.

^b The directional range for the Foehn cases was determined experimentally by comparison with manual classification from a trained weather observer.

RC14. Page 14, figure 4. In summer, about 70% of the days feature breeze systems, but aerosol-layer days are only about 50% (Figure 3). What about the rest? Are easterly breezes occasionally "clean"?

AR14. A comment about this (only apparent) discrepancy has been added to Sect. 4.1:

For cases of easterly winds and no aerosol layer found (7%), possible reasons are:

- the diurnal wind, although clearly detected by the surface network, was of too short duration or too weak to transport the polluted air masses from the Po basin to Aosta–Saint-Christophe (at least ~40 km must be travelled by the air masses along the main valley). In our record, this was for example the case of 14 and 18 September 2015, and 17 June 2016;
- back-trajectories were indeed channelled along the central valley, however they did not come from the Po basin, but rather from the other side of the Alps. This was the case, for instance, of 20 August 2015, in which air masses came from the Divedro Valley (north-east of the Aosta Valley) after crossing the Simplon pass (between Switzerland and Italy).

These exceptions also help understand why, in summer, about 70% of the days feature breeze systems (Fig. 4) while aerosol layers are detected only about 50% of the days (Fig. 3). Indeed, some part (7%, Fig. 5a) of this 20% discrepancy can be attributed to the above events, the remaining 13% being likely associated with days with a complex aerosol structure (not classified in Fig. 2) or to days affected by low clouds and/or desert dust events. These days were therefore labelled as "Other" in Fig. 3.

RC15. Page 15, line 14. It would be useful to explain where the Divedro Valley is. The great majority of readers wouldn't have a clue otherwise.

AR15. Done, the updated text is cited in the previous reply (AR14).

RC16. Page 16, line 15. Please remove the comma between "simple" and "forecasting".

AR16. Done.

RC17. Page 19, line 4. Reference to Teixeira et al 2016. The interested reader wouldn't find much in this editorial. More appropriate references could be those available at these DOIs: 10.3389/feart.2015.00077 and 10.3390/atmos9030102.

AR17. The bibliographic reference has been changed according to the reviewer's comment.

RC18. Page 25, line 5. The dashed continuous line in Figure 12a is very hard to see. Consider replacing with a colored line, or with a filled area in the background.

AR18. Thank you for this suggestion. The dashed line has been replaced with a filled area. The new figure is presented below:



Figure 12: (a) Temporal chart of the contribution of secondary (nitrate-rich and sulfate-rich) modes to the total PM_{10} . The shaded area represents the estimated local production of secondary aerosol. The difference between each dot and this baseline can thus be read as the non-local contribution to the aerosol concentration in Aosta–Downtown. Since ions chemical speciation started in 2017, only one year of overlap with the ALC is available at the moment (see Table 1). (b) Boxplot of the contribution of secondary (nitrate-rich and sulfate-rich) modes to the total PM_{10} concentration as a function of the day type. PMF-dataset "a" was used for both plots.

RC19. Page 25, Figure 13. Text is hard to read on the colored background. Consider using a different color map, or adding white background to the location labels.

AR19. White background has been added to the location labels (refer to AR11 for the new figures).

RC20. Pages 30-32. There is some redundancy in Figures 16-17-18-19. It is unclear why results from two locations are represented in the figure. In fact, the differences between the two sites are never discussed in detail in the text. It would be possible to remove the figure panels referring to Donnas without great loss of information.

AR20. By including both sites we intended to show the effect of the distance from the Po Valley on the receptor site. However, given this comment, we have removed the figure panels referring to Donnas. Figure 17 has been updated as follows (also considering RC3 by Reviewer#1):

Aosta–Downtown



Figure 17: Absolute (a,c) and relative (b,d) differences between simulated and observed PM_{10} concentrations at the surface in Aosta–Downtown. The mean bias error (MBE) and the normalised mean bias error (NMBE) for each case are reported in the plot titles. First row: FARM simulations as currently performed by ARPA. Second row: the PM_{10} concentrations from outside the boundaries of the domain were multiplied by a factor W=4.

RC21. Page 33, lines 2-3. "This is a clear indication that the external contribution (boundary conditions) is not optimally parametrised in the model". I am really not sure that I agree on this statement. If I understand correctly, FARM uses information from the whole COSMO-I2 domain, which completely includes the Po valley. Therefore, the problem does not lie in "external contributions" not being represented in the lateral boundary conditions. There might be problems with the lower boundary conditions (emission inventories), but I doubt these can be wrong by a factor of 4 over the whole Po valley. I am not even sure that model parametrisations matter here.

I would argue that the relatively coarse resolution of the weather model (2.8 km, marginally enough to resolve the valley) and of the transport model (only 16 vertical levels) play an important role. Please consider the topography cross-sections in Figure 20. Aosta seems to lie in a basin, likely non-existent in reality and introduced in the model by terrain smoothing. If the topography profile descended continuously from Aosta to Donnas and the Po Plain (as in reality), it would be possible to resolve the horizontal advection of aerosol-laden air from much lower altitudes on the plain, likely removing a large fraction of the negative concentration bias.

AR21. First of all, it must be noticed that FARM does not use "information from the whole COSMO-I2 domain, which completely includes the Po valley". We tried to better explain this in Sect. 3.1:

FARM is only run over a small domain (light blue rectangle in Fig. 1a,b), roughly corresponding to the Aosta Valley. A regional emission inventory ("local sources", updated to 2015) is supplied to the CTM over the same area to accurately assess the magnitude of the pollution load and its variability in time and space. Data from a national inventory and CTM model (QualeAria, here referred to as "boundary conditions", outer rectangle in Fig. 1a), taken along the border of the inner (light blue) rectangle, are also used to estimate the mass exchange from outside the borders of the FARM domain.

As a second point we added further analysis to check whether FARM performs better in a less complex terrain. To this purpose, we decided to include the data from the Ivrea station. This addition clearly showed that the same PM_{10} underestimation of the model is also obtained over flat terrain, demonstrating that this is not driven by inaccuracies of the simulated wind fields over the more complex terrain of Aosta. We think that this represents a remarkable improvement to our study, supporting the hypothesis that the model underestimations are mainly due to incorrect emission inventories/unaccounted physical processes (e.g., aqueous-phase chemistry) at a national level. This additional analysis required description of the measurement site, rationale, and comment of the results obtained (and relevant figures):

In this study, we mainly used data from four sites (...) and Ivrea (243 m a.s.l., urban background) (...), a site in the Po Plain (31 µg m^{-3} average PM₁₀ in 2017) located just outside the Aosta Valley, in the Italian Piedmont region (Fig. 1b). This was done to check if and how much inaccuracies of the model in reproducing the aerosol loads over the Aosta Valley are due to difficulties in simulating the wind field in such a complex terrain. In fact, the city centre of Ivrea (24000 inhabitants) is approximately in the middle between the measurement site (south of the city) and the nearest cell of our model domain (north of the city), the distance between these two points being only 2 km. The altitude of the cell (from the digital elevation model used in our simulations) is 241 m a.s.l., i.e. approximately the real one (...) The station of Ivrea features, among other instruments, a TCR Tecora Charlie/Sentinel PM₁₀ sampler. PM₁₀ concentrations are then determined by a gravimetric technique.

We have remarkably expanded Sect. 4.5 by adding the dataset from this station to the study, and comparing it to the corresponding FARM simulations:

The observed model-measurement discrepancies might originate from (1) an incomplete representation of the inventory sources (emission component), (2) inaccurate NWP modelling of the meteorological fields, and notably the wind (transport component), or a combination of (1) and (2). Some details on these aspects are provided in the following.

1. Emissions. Inaccuracies in the (local and national) emission inventories could degrade the comparison between simulations and observations. Based on results shown in Fig. 17(a,b), we decided to investigate the sensitivity of our simulations in Aosta–Downtown to the magnitude of the external contributions (boundary conditions). In particular, we used a simplified approach to speed up the calculations: assuming that the contribution from outside the regional domain and the local emissions add up without interacting, we simulated the surface PM_{10} concentrations turning on/off the boundary conditions (dark- and light-blue lines in Fig. 16) to roughly estimate the only contribution from sources outside the regional domain. Interestingly, the difference between the two FARM runs correlates well with the advection classes observed by the ALC (Fig. 18). This clear correlation between the simulations and the experimentally-determined atmospheric conditions is a first good indication that the NWP model used as input to the CTM yields reasonable meteorological inputs to FARM. Then, to further explore the sensitivity to the boundary conditions, we gradually increased, by a weighting factor W, the PM_{10} concentration from outside the boundaries of the domain trying to match the observed values. In Figs. 17(c, d) we show the results obtained with W=4. This exercise show that the overall mean bias error is much reduced compared to the original simulations (W=1, Figs. 17(a, b)), especially for the winter and autumn seasons, while slight overestimations are now visible for summer and spring. Also the annually-averaged MBE, NMBE and NMSD improve (Table 3), whilst the other statistical indicators remain stable or even slightly worsen.

Clearly, our simplified test has major limitations. (...)

2. Although the COSMO-I2 grid spacing is certainly in line with that of other state-of-the-art operational limited-area models, in our complex terrain it could be insufficient to appropriately resolve local meteorological phenomena triggered by the valley orography (Wagner et al., 2014), also considering that the actual model resolution is 6–8 times the grid cell (Skamarock, 2004). (...)

To disentangle the role of factors (1) and (2) discussed above, we evaluated the model performances in a flatter area of the domain, where simulations are expected to be less affected by the complex orography of the mountains. We therefore compared PM_{10} simulations and measurements in Ivrea (Fig. 16b). This test is also useful to operate the CTM model at the boundaries of the domain, with special focus on the emissions from the "boundary conditions". Model underestimation in both the warm and cold seasons is evident. Incidentally, it can be noticed that concentrations simulated without "boundary conditions" are nearly zero, since the considered cell is far from the strongest local sources and most part of the aerosol comes from outside the regional domain. As done for Aosta (...), Fig. 19(a,b) presents the absolute and relative differences between the model and the observations in Ivrea. The overall NMBE is -0.73, which rather well corresponds to the underestimation factor W=4 (or NMBE=-0.75) found for Aosta-Downtown and other sites in the valley. Since it is expected that in this flat area the NWP model is able to better resolve the circulation compared to the mountain valley, this result suggests poor CTM performances to be mostly related to underestimated emissions at the boundaries rather than to incorrect transport. In addition to this general underestimation, a large scatter between observations and simulations can be noticed in Fig. 16, the linear correlation index between simulations and observation in Ivrea being rather low (ρ =0.54). If such a large scatter, due to the erroneous boundary conditions, is already detectable at the border of the domain, it is likely that at least part of the RMSE reported in Table 3 for Aosta–Downtown can be attributed to inaccuracies in the national inventory. As already mentioned, an additional contribution to the observed RMSE could still be given by errors in modelling transport due to the coarse resolution of the NWP model, although we are not able to quantify the relative role of the two factors. To unravel this issue, high-resolution models (grid step 0.5 km, e.g. Golzio and Pelfini, 2018; Golzio et al., 2019) are being tested on the investigated area and their results will be addressed in future studies.

We report the new figures, previously mentioned in the text, here below.



Figure 16: Long-term (1 year) comparison between PM_{10} surface measurements and simulations (FARM) in Aosta–Downtown (a) and Ivrea (b).

lvrea



Figure 19: Absolute (a) and relative (b) differences between simulated and observed PM_{10} concentrations at the surface in Ivrea.

Concerning the impact of smoothed topography, please refer to AR10.

RC22. Page 33, line 10. "The reason for discrepancies is therefore likely in the emissions". Why not in deficiencies of the transport modelling, as I argued above?

AR22. See AR21.

RC23. Page 46, line 18: There seems to be a problem with the reference to Thunis et al, 2012 (the third author).

AR23. Thank you, the reference has been corrected.

References

- Diémoz, H., Barnaba, F., Magri, T., Pession, G., Dionisi, D., Pittavino, S., Tombolato, I. K. F., Campanelli, M., Della Ceca, L. S., Hervo, M., Di Liberto, L., Ferrero, L., and Gobbi, G. P.: Transport of Po Valley aerosol pollution to the northwestern Alps – Part 1: Phenomenology, Atmos. Chem. Phys., 19, 3065– 3095, doi:10.5194/acp-19-3065-2019, 2019.
- Giovannini, L., Antonacci, G., Zardi, D., Laiti, L., and Panziera, L.: Sensitivity of Simulated Wind Speed to Spatial Resolution over Complex Terrain, Energy Proced., 59, 323–329, doi:10.1016/j.egypro.2014.10. 384, 2014.
- Golzio, A. and Pelfini, M.: High resolution WRF over mountainous complex terrain: testing over the Ortles Cevedale area (Central Italian Alps), in: Conference: First National Congress, Italian Association of Atmospheric Sciences and Meteorology (AISAM), AISAM, 2018.
- Golzio, A., Ferrarese, S., Cassardo, C., Diolaiuti, G. A., and Pelfini, M.: Resolution improvement in WRF over complex mountainous terrain, in preparation for Boundary-Layer Meteorol., 2019.
- Norris, G. and Duvall, R.: EPA Positive Matrix Factorization (PMF) 5.0 Fundamentals and User Guide, U.S. Environmental Protection Agency, URL https://www.epa.gov/sites/production/files/2015-02/documents/pmf_5.0_user_guide.pdf, 2014.
- Schmidli, J., Böing, S., and Fuhrer, O.: Accuracy of Simulated Diurnal Valley Winds in the Swiss Alps: Influence of Grid Resolution, Topography Filtering, and Land Surface Datasets, Atmosphere, doi:10. 3390/atmos9050196, 2018.
- Seibert, P.: The riddles of foehn introduction to the historic articles by Hann and Ficker, Meteorol. Z., 21, 607–614, doi:10.1127/0941-2948/2012/0398, 2012.
- Skamarock, W. C.: Evaluating Mesoscale NWP Models Using Kinetic Energy Spectra, Mon. Weather Rev., 132, 3019–3032, doi:10.1175/MWR2830.1, 2004.
- Wagner, J. S., Gohm, A., and Rotach, M. W.: The Impact of Horizontal Model Grid Resolution on the Boundary Layer Structure over an Idealized Valley, Mon. Weather Rev., 142, 3446–3465, doi:10.1175/ MWR-D-14-00002.1, 2014.

Transport of Po Valley aerosol pollution to the northwestern Alps – Part 2: Long-term impact on air quality

Henri Diémoz¹, Gian Paolo Gobbi², Tiziana Magri¹, Giordano Pession¹, Sara Pittavino¹, Ivan K. F. Tombolato¹, Monica Campanelli², and Francesca Barnaba²

¹ARPA Valle d'Aosta, Saint-Christophe, Italy ²Institute of Atmospheric Science and Climate, CNR, Rome, Italy

Correspondence to: Henri Diémoz (h.diemoz@arpa.vda.it)

Abstract.

This work evaluates the impact of trans-regional aerosol transport from the polluted Po basin on particulate matter levels (PM_{10}) and physico-chemical characteristics in the northwestern Alps. To this purpose, we exploited a multi-sensor, multi-platform database over a <u>3-years 3-year</u> period (2015–2017) accompanied by a series of numerical simulations. The exper-

- 5 imental setup included operational (24/7) vertically-resolved vertically resolved aerosol profiles by an Automated LiDAR-Ceilometer (ALC), vertically-integrated vertically integrated aerosol properties by a sun/sky photometer, and surface measurements of aerosol mass concentration, size distribution and chemical composition. This experimental set of observations was then complemented by modelling tools, including Numerical Weather Prediction (NWP), Trajectory Statistical (TSM) and Chemical Transport (CTM) models, plus Positive Matrix Factorisation (PMF) on both the PM₁₀ chemical speciation analy-
- 10 ses and <u>particle</u> size distributions. In a first companion study(Diémoz et al., 2019), we showed and discussed through detailed case studies the 4-D phenomenology of recurrent episodes of aerosol transport from the polluted Po basin to the northwestern Italian Alps, and particularly to the Aosta Valley. Here we draw more general and statistically significant conclusions on the frequency of occurrence of this phenomenon, and on the quantitative impact of this regular, wind-driven, aerosol-rich "atmospheric tide" on PM₁₀ air quality levels in this alpine environment. <u>Combining vertically-resolved ALC measurements</u>
- 15 with wind informationBased on an original ALC-derived classification, we found that an advected aerosol layer is observed at the receptor site (Aosta) in 93% of days characterized by easterly winds (thermally-driven winds from the plain or synoptic circulation regimes), i.e., from the Po basin) and that the longer the time spent by air masses over the Po plain the higher this probability. On a seasonal basis, frequency of Frequency of these advected aerosol layers from the Po basin maximises in summer (70% of the days classified using the ALC profiles) and minimises in winter and spring (57 was found to be rather
- 20 stable over the seasons with about 50% of the classified days)days affected. Duration of these advection events ranges from few hours up to several days, while aerosol layer thickness ranges from 500 up to 4000 m. This phenomenon was found Our results confirm this phenomenon to be related to non-local emissions, to act at the regional scale and to largely impact both surface levels and column-integrated aerosol properties, with . In Aosta, PM₁₀ and AOD values respectively increasing increase respectively up to a factor of 3.5 and 4 in dates under the Po Valley influence. Similar variations in values observed
- 25 at different stations within the Aosta Valley also indicated the phenomenon to act at the regional scale and to be related to

non-local emissions. Pollution transport events were also shown to modify the mean chemical composition and typical size of particles in the target region. In fact, increase in secondary species, and mainly nitrate- and sulfate-rich components, were found to be effective proxies of the advections, with the transported aerosol responsible for at least 25% of the PM_{10} measured in the urban site of Aosta, and adding up to over 50 µg m⁻³ during specific episodes, thus exceeding alone the EU established

- 5 daily limit. This percentage is expected to be higher in the rural, pristine areas on the northwestern Alps, where chemical data were not available and trans-boundary contribution to might thus exceed the local one. Advected aerosols were also found to be on average finer, more light-scattering and more hygroscopic than the locally-produced ones. From a modelling point of view, our CTM simulations performed over a full year showed that the model is able to reproduce the phenomenon but, but markedly underestimates its impact on PM_{10} levels. As a sensitivity test, we employed the ALC-derived identification of aerosol ad-
- 10 vections to re-weight the emissions from outside the boundaries of the regional domain in order to match the observed PM_{10} field. This simplified exercise indicated that an increase of such "external" emissions by a factor of 4 in the model would is needed to halve the model PM_{10} maximum deviations and to significantly reduce the PM_{10} normalised mean bias forecasts error (MBE) from -10 to less than 2, the normalised mean standard deviation (NMSD) from over -50% to less than -10% and would halve the model maximum deviations from -35% to 5%).

15 1 Introduction

Mountain regions are often considered pristine areas, being typically far from large urban settlements and strong anthropogenic emission sources, and thus relatively unaffected by remarkable pollution footprints. However, atmospheric transport of pollutants from the neighbouring foreland is not uncommon, owing to the synoptic and regional circulation patterns. For example, in several regions of the world, thermally-driven flows represent a systematic characteristic of mountain weather

- 20 and climate (Hann, 1879; Thyer, 1966; Kastendeuch and Kaufmann, 1997; Egger et al., 2000; Ying et al., 2009; Serafin and Zardi, 2010; Schmidli, 2013; Zardi and Whiteman, 2013; Wagner et al., 2014a; Giovannini et al., 2017; Schmidli et al., 2018) favouring regular exchange of air masses between the plain and the highland sites (Weissmann et al., 2005; Gohm et al., 2009; Cong et al., 2015; Dhungel et al., 2018), with likely consequences on human health (Loomis et al., 2013; EEA, 2015; WHO, 2016), ecosystems (e.g., Carslaw et al., 2010; EMEP, 2016; Bourgeois et al., 2018; Burkhardt et al., 2018)
- 25 (e.g., Carslaw et al., 2010; EMEP, 2016; Bourgeois et al., 2018; Burkhardt et al., 2018; Allen et al., 2019; Rizzi et al., 2019), climate (Ramanathan et al., 2001; Clerici and Mélin, 2008; Philipona, 2013; Pepin et al., 2015; Zeng et al., 2015; Tudoroiu et al., 2016; Samset, 2018) and, not least, local economy, through loss of tourism revenues (de Freitas, 2003; Keiser et al., 2018). These phenomena have worldwide relevance, since nearly one quarter of the Earth's land mass can be classified as mountainous areas (Blyth, 2002).
- 30 Among them, the Alps are Alpine region (Fig. 1a) is of particular interest, due to both their its sensitive environment and their its proximity to the Po Valley (Fig. 1a; Tampieri et al., 1981; Seibert et al., 1998; Wotawa et al., 2000; Dosio et al., 2002; Campana et al., 2002; Campana et al., 2009; Finardi et al., 1981; Seibert et al., 1998; Wotawa et al., 2000; Dosio et al., 2002; Campana et al., 2009; Finardi et al., 2014). In fact, the Po basin represents a major pollution hotspot in Europe, with large emissions from highly populated



Figure 1. True colour corrected reflectance from MODIS Terra satellite (http://worldview.earthdata.nasa.gov) on 17 March 2017. (a) Italy, with indication of the Alpine and Po Valley regions, of the Aosta Valley FARM regional domain (northwest side, light blue rectangle), and the Po ValleyCOSMO-I2 domain (orange rectangle, approximately corresponding to the boundaries of the national inventory). (b) Zoom over the Aosta Valley. The circle markers represent the sites of 1) Aosta–Downtown; 2) <u>Aosta–Saint ChristopheAosta–Saint-Christophe</u>; 3) Donnas; 4) Ivrea. A thin aerosol layer over the Po basin starting to spread out into the Alpine valleys is visible in both figures.

urban areas (about 40% of the Italian population lives in this region, WMO, 2012) (about 40% of the Italian population lives in this region, WMO, 2012), intense anthropogenic activities, such as industry and agriculture, combined to a local topography promoting atmospheric stability (Chu et al., 2003; Barnaba and Gobbi, 2004; Schaap et al., 2004; Van Donkelaar et al., 2010; Fuzzi et al., (Chu et al., 2003; Barnaba and Gobbi, 2004; Schaap et al., 2010; Bigi and Ghermandi, 2014; Fuzzi

5 et al., 2015; EMEP, 2016; Belis et al., 2017; EEA, 2017). Despite the efforts to decrease the number of particulate matter (PM) exceedances, Italy is still failing to comply with the European air quality standards (EU Commission, 2008, 2018), the Po Valley being the major responsible for this situation.

In a first companion paper (Diémoz et al., 2019), we investigated the phenomenology of the aerosol-aerosol-rich air mass advections from the Po basin to the northwestern Alps, and specifically to the Aosta Valley (Fig. 1b). This was introduced and

- 10 thoroughly described by means of three specifically-selected case studies, each of them lasting several days and monitored by a large set of instruments. That investigation evidenced clear features of this phenomenology, and notably: 1) the aerosol transported to the northwestern Alps is clearly detectable and discernible from the locally-produced aerosol. Detection of such polluted aerosol layers over Aosta was primarily driven by remote-sensing profiling measurements (Automated LiDAR Ceilometers, ALCs), these showing recurrent arrival of thick and elevated aerosol-rich layers. Good correspondence was found
- 15 between the presence of these layers aloft, changes of column-averaged aerosol properties, and variations of PM surface concentrations and chemical composition; 2) the advected particles are small (accumulation mode), mainly of secondary origin,

weakly light-absorbing and highly hygroscopic compared to the locally-produced aerosol. Clouds are frequently observed to form within these layers during the night; 3) the air masses associated to with the observed elevated layers are found to originate from the Po basin and to be transported to the northwestern Alps by up-valley breezes diurnal wind systems and synoptic winds; 4) the chemical transport model (FARM, Flexible Air quality Regional Model) currently used by the local

- 5 Environment Protection Agency (ARPA Valle d'Aosta) was is able to reproduce this transport from a qualitative point of view and was usefully employed to interpret the observations. However, it fails at quantitatively estimating the PM mass contribution coming from outside the boundaries of the regional domain (approximately corresponding to the Valle d'Aosta region), likely due to deficiencies in the emission inventories therein and to unaccounted aerosol processes in the model (the main of which are likely e.g., those related to hygroscopicity/aqueous-phase chemistry triggered by high relative humidities).
- 10 The present work analyses the same phenomenon, but with a long-term perspective. Main aim of this study is to establish frequency of occurrence of aerosol transport from the Po Valley to the northwestern Alps and its impact on local air quality, depending on the interplay between frequency and severity of the episodes. In particular, this paper addresses the following questions:
 - 1. How frequently does advection from the Po plain to the northwestern Alps occur? What are the most common meteoro-
- 15

logical conditions favouring it?

- 2. What are the average properties of the advected aerosol both at the surface and in the layers aloft?
- 3. How much does transport from the Po basin impact the air quality EU-regulated metrics in the northwestern Alps?
- 4. Can we effectively predict the arrival of aerosol-polluted air masses to the northwestern Alps? How can we improve our air quality forecasts?
- To answer these questions, we took advantage of long-term (2015–2017) and almost uninterrupted series of measurements carried out in the Aosta region. The experimental dataset, thoroughly described by Diémoz et al. (2019), included measurements of vertically-resolved aerosol profiles by an ALC, vertically-integrated aerosol properties by a sun/sky photometer and surface measurements of the aerosol mass concentration, size distribution and chemical composition, most of these series encompassing a time period of several years. Indeed, a great number of intensive, short-term campaigns employing cutting-edge
- 25 research instruments and techniques were already performed on this topic was already performed in the Po basin, mainly focussing on its central, eastern and southern parts (Nyeki et al., 2002; Barnaba et al., 2007; Ferrero et al., 2010, 2014; Saarikoski et al., 2012; Landi et al., 2013; Decesari et al., 2014; Costabile et al., 2017; Bucci et al., 2018; Cugerone et al., 2018). However, continuous and multi-year datasets, especially from ground-based stations, are necessary to assess the influence of pollution transport on a longer term (e.g., Mélin and Zibordi, 2005; Clerici and Mélin, 2008; Kambezidis and Kaskaoutis, 2008; Mazzola
- 30 et al., 2010; Bigi and Ghermandi, 2014, 2016; Putaud et al., 2014; Arvani et al., 2016). In this context, local environmental agencies operate stable networks for continuous monitoring of air quality and meteorological parameters, and use standardised methodologies and universally recognised quality control procedures.

The study description is organised as follows: the investigated area is presented in Sect. 2, together with the experimental setup, further complemented by mathematical and the modelling tools (Sect. 3). Results are presented in Sect. 4 and include a) a first, original classification framework based on ALC data and meteorological variables; b) a long-term, statistical analysis of the phenomenon and its impacts on surface/column aerosol properties and c) comparison between observations and CTM simulations. Conclusions are drawn in Sect. 5.

2 Study region and experimental setup

5

We provide here a brief overview of the region of interest and of the experimental setup, this latter being described with full details in the companion paper (Diémoz et al., 2019).

- The Aosta Valley, located at the northwestern Italian border (Fig. 1a), is the smallest administrative region of the country, being only 80 km by 40 km wide and hosting 130000 inhabitants. The most remarkable geographical feature of the region, whose mean altitude exceeds 2000 m a.s.l., is the main valley, approximately oriented in a SE to NW direction and connecting the Po basin (about 300 m a.s.l., at the southeast side of the Aosta Valley region) to the Mont Blanc chain (4810 m a.s.l., separating Italy and France). Several tributary valleys depart from the main valley. The topography of the area triggers some of the most common weather regimes in the mountains, such as thermally-drivenwinds (from the east, during daytime), slope winds
- 15 perpendicular to and along the main valley axis, rain-shadow winds(, up-valley (daytime) and down-valley (nighttime) winds, up-slope (daytime) and down-slope (nighttime) winds, "Foehn", winds (Seibert, 2012) from the west), and frequent temperature inversions during wintertime anticyclonic days. Quite obviously, those meteorological phenomena are responsible of most of the variability of the atmospheric constituents (e.g., water vapour, Campanelli et al. (2018), and pollutants concentrations, Agnesod et al. (2003), and water vapour Campanelli et al. (2018)).
- The air quality network of the local regional Environmental Protection Agency is designed trying to capture the topographical and meteorological heterogeneity of the investigated area. In this study, we mainly used data from three four sites belonging to the this network (Fig. 1b): Aosta–Saint Christophe (560 Aosta–Downtown (580 m a.s.l., semi-rural), Aosta–Downtown (580 urban background), Aosta–Saint-Christophe (560 m a.s.l., urban background)and semi-rural), Donnas (316 m a.s.l., rural) and Ivrea (243 m a.s.l., urban background). The first two measuring stations, located 2.5 km apart, are situated respectively in the
- ²⁵ suburbs centre and in the centre suburbs of Aosta (45.7°N, 7.4°E), the main urban settlement of the region (36000 inhabitants). Car traffic, domestic heating and a steel mill are the main anthropogenic emission sources within the city, whose pollution levels are generally moderate (e.g., yearly average PM_{10} concentration of about 20 µg m⁻³, with summertime and wintertime averages of 13 and 31 µg m⁻³, respectively). Donnas is located at closer to the border with the Po basin (Fig. 1b) and is expected to be strongly impacted by pollution from outside the region. In fact, the PM_{10} concentration in this rural site is
- 30 about 18 μ g m⁻³ on a yearly average, i.e. only slightly lower than the concentration in Aosta. We also consider in our analysis the PM₁₀ records collected in the city of Ivrea, a site in the Po Plain (31 μ g m⁻³ average PM₁₀ in 2017) located just outside the Aosta Valley, in the Italian Piedmont region (Fig. 1b). This was done to check if and how much inaccuracies of the model in reproducing the aerosol loads over the Aosta Valley are due to difficulties in simulating the wind field in such a complex

terrain. In fact, the city centre of Ivrea (24000 inhabitants) is approximately in the middle between the measurement site (south of the city) and the nearest cell of our model domain (north of the city), the distance between these two points being only 2 km. The altitude of the cell (from the digital elevation model used in our simulations) is 241 m a.s.l., i.e. approximately the real one.

- 5 The Aosta–Saint Christophe atmospheric observatory Aosta–Downtown hosts a monitoring station for continuous measurements of the aerosol concentration and composition. PM_{10} and $PM_{2.5}$ daily concentrations are available from two Opsis SM200 Particulate Monitor instruments. An estimate of the PM_{10} hourly variability is furthermore provided by a Tapered Element Oscillating Microbalance (TEOM) 1400a monitor (Patashnick and Rupprecht, 1991), though not compensated for mass loss of semi-volatile compounds (Green et al., 2009), such as ammonium nitrate (e.g., Charron et al., 2004; Rosati et al., 2016).
- 10 Moreover, the collected PM₁₀ samples are chemically analysed on a daily basis. Ion Chromatography (AQUION/ICS-1000 modules) is employed for determining the concentration of water-soluble anions and cations (Cl⁻, NO₃⁻, SO₄²⁻, Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺), based on the CEN/TR 16269:2011 guideline. Elemental/organic carbon (EC/OC), using the thermal-optical transmission (TOT) method (Birch and Cary, 1996) and following the EUSAAR-2 protocol (Cavalli et al., 2010) according to the EN 16909:2017, are analysed alternatively to metal concentrations (Cr, Cu, Fe, Mn, Ni, Pb, Zn, As, Cd, Mo and Co), by
- 15 means of inductively coupled plasma mass spectrometry (ICP-MS) after acid mineralisation of the filter in aqueous solution (EN 14902:2005). In accordance with this laboratory schedule (i.e., metal analyses on two consecutive days, EC/OC on the following two days, followed by two sequences metal-metal-EC/OC), over a 10 days period EC/OC concentrations are routinely provided on 4 days on average and metal concentrations on 6 days. Particle-bound polycyclic aromatic hydrocarbons (PAHs) are detected continously in real-time by a photoelectric aerosol sensor (EcoChem PAS-2000). Gas-phase pollutants,
- 20 such as NO and NO_2 are also continuously monitored and were used in the present work to help identification of urban combustion sources (mainly traffic and residential heating).

The atmospheric observatory in Aosta–Saint-Christophe (WIGOS ID 0-380-5-1, Diémoz et al., 2011, 2014a) is located on the terrace of the ARPA building. At this site, advanced instrumentation is continuously operated, daily checked and maintained. Vertically-resolved measurements of aerosols and clouds are performed by a commercially-available Automated LiDAR

- 25 Ceilometer (ALC, Lufft CHM15k-Nimbus) running since 2015 in the framework of the Italian Alice-net (http://www.alice-net. eu/) and the European E-PROFILE (https://e-profile.eu) networks. The ALC emits in the atmosphere 8 μ J laser impulses at a single wavelength of 1064 nm with a frequency of 7 kHz and collects their backscatter by aerosol and molecules from ground up to an altitude of 15 km, with a vertical resolution of 15 m. This return signal is averaged over 15 s and saved by the firmware as range-, overlap- and baseline-corrected raw counts (β_{raw}). To convert the backscatter signal in SI units, a Rayleigh calibra-
- 30 tion (Fernald, 1984; Klett, 1985; Wiegner and Geiß, 2012) is necessary. This is accomplished by the operator based on data sampled during clear-sky nights. This allows to derive the particle backscatter coefficient (β_p) and the scattering ratio (SR, e.g. Zuev et al., 2017), which is the first primary, ALC-derived parameter used in this work to quantify the aerosol load. It is defined

as:

$$SR = \frac{\beta_p + \beta_m}{\beta_m} \tag{1}$$

 β_m being the molecular backscattering coefficient (thus SR=1 indicates a purely molecular atmosphere while SR increases with aerosol content).

At the same station, a POM-02 sun/sky photometer operates since 2012 as part of the European ESR-SKYNET network (http: //www.euroskyrad.net/). The instrument collects the direct light coming from the sun (every 1 min) and to retrieve the aerosol 5 optical depth (AOD, $\tau(\lambda)$) and its spectral dependence, calculated by fitting the Ångström (1929) law to the measurements in the 400–870 nm range:

$$\tau(\lambda) = b \, \lambda^{-a}$$

(2)

a being the Ångström exponent. Other aerosol optical and microphysical properties are retrieved from the light scattered from the sky in the almucantar plane (every 10 min) to retrieve the aerosol optical depth (AOD) and other aerosol optical 10 and microphysical properties at 11 wavelengths (315-2200 nm) using the SKYRAD pack version 4 code. The sun photometer is calibrated in situ with the improved Langley technique (Campanelli et al., 2007) and was recently successfully compared to other reference instruments (Kazadzis et al., 2018). Accurate cloud screening is achieved using the Cloud Screening of Sky Radiometer data (CSSR) algorithm by Khatri and Takamura (2009). Special care is needed to retrieve, among the other parameters, the aerosol single scattering albedo (SSA). In fact, it is known that the SSA retrieval by the SKYRAD pack version 15 4 is problematic and sometimes unnaturally close to unity (Hashimoto et al., 2012), irrespectively of the AOD. Although

improvements are expected in the next versions of SKYRAD.pack, data collected within the European ESR-SKYNET network are still processed with version 4. Therefore, only SSA lower than 0.99 at all wavelengths were retained in our analysis.

Finally, a Fidas 200s Optical Particle Counter (OPC) is used in Aosta-Saint-Christophe Aosta-Saint-Christophe to yield 20 an accurate estimation of the aerosol size distribution ($0.18-18 \mu m$) in proximity of the surface (Pletscher et al., 2016). This instrument, although not directly measuring the aerosol mass, obtained the certificate of equivalence to the gravimetric method by TÜV Rheinland Energy GmbH on the basis of a laboratory and a field test. A PM₁₀ comparison with the gravimetric technique was additionally performed in Aosta-Saint Christophe Aosta-Saint-Christophe and provided satisfactory results (29 days; slope 1.08 ± 0.04 ; intercept $-3.8 \pm 1.4 \text{ µg m}^{-3}$; $R^2 = 0.96$). Further instruments to monitor trace gases (Diémoz et al., 2014b) are employed at the station and the corresponding datasets will be investigated in future studies.

25

Aosta-Downtown hosts a monitoring station for continuous measurements of the aerosol concentration and composition, and daily concentrations are available from two Opsis SM200 Particulate Monitor instruments. An estimate of the hourly variability is furthermore provided by a Tapered Element Oscillating Microbalance (TEOM) 1400a monitor (Patashnick and Rupprecht, 1991) , though not compensated for mass loss of semi-volatile compounds (Green et al., 2009), such as ammonium nitrate (e.g., Charron et al., 200

- Moreover, the collected samples are chemically analysed on a daily basis. Ion Chromatography (AQUION/ICS-1000 modules) 30 is employed for determining the concentration of water-soluble anions and cations (,,,,,,,), based on the CEN/TR 16269:2011

guideline. Elemental/organic carbon (EC/OC), using the thermal-optical transmission (TOT) method (Birch and Cary, 1996) and following the EUSAAR-2 protocol (Cavalli et al., 2010) according to the EN 16909:2017, are analysed alternatively to metal concentrations (Cr, Cu, Fe, Mn, Ni, Pb, Zn, As, Cd, Mo and Co), by means of inductively coupled plasma mass spectrometry (ICP-MS) after acid mineralisation of the filter in aqueous solution (EN 14902:2005). Particle-bound polycyclic

5 aromatic hydrocarbons (PAHs) are detected continuously in real-time by a photoelectric aerosol sensor (EcoChem PAS-2000). Gas-phase pollutants, such as and are also continuously monitored and were used in the present work to help identification of urban combustion sources (mainly traffic and residential heating).

Some of the air quality parameters monitored in Aosta–Downtown are also measured at the Donnas station, i.e. PM_{10} daily concentrations (Opsis SM200) and nitrogen oxides (Horiba APNA-370 chemiluminescence monitor). Finally, the three

10 stations are equipped with instruments for tracking the standard meteorological parameters, such as temperature, pressure, relative humidity (RH) and surface wind velocity. A list of the instruments,

The station of Ivrea features, among other instruments, a TCR Tecora Charlie/Sentinel PM_{10} sampler. PM_{10} concentrations are then determined by a gravimetric technique.

A list of all measurements with relevant operating period and subset considered in this study, is presented in Table 1.

15 3 Modelling tools

Numerical models and statistical techniques were used to interpret and complement the observations described above. A numerical weather prediction model (COSMO, Consortium for Small-scale Modeling, www.cosmo-model.org) was employed to drive a chemical transport model (FARM, Flexible Air quality Regional Model) and a lagrangian model (LAGRANTO). These tools were thoroughly described by Diémoz et al. (2019) and are only briefly recalled here (Sect. 3.1), since they were

20 thoroughly described by Diémoz et al. (2019). Conversely, Trajectory Statistical Models (TSMs) and Positive Matrix Factorisation (PMF)were, adopted to interpret the long-term series of measurements used in this work, and are therefore are fully described in Sects. 3.2 and 3.3, respectively.

3.1 Numerical atmospheric models

COSMO is a non-hydrostatic, fully compressible atmospheric prediction model working on the meso- β and meso- γ scales (Bal-

- 25 dauf et al., 2011). Owing to the complex topography of the Aosta Valley, and the consequent need to resolve the atmospheric circulation at very small spatial scales, in the present work we used a nudged, high-resolution (2.8-The forecasts, 65 vertical levels, 2 runs/day) variant, called COSMO-I2 (or COSMO-IT), covering Italy. The COSMO forecasts, with inclusive of the complete set of parameters (including such as the 3-D wind velocity used here) , are disseminated daily for eight time steps (from 00 to 21 UTC)by the , are disseminated daily in two different configurations by the meteorological operative cen-
- 30 tre air force meteorological service (COMET): a lower-resolution version (COSMO-ME, 7 km horizontal grid and 45 levels vertical grid, 72 hours integration), covering central and southern Europe, and a nudged, higher-resolution version (COSMO-I2 or COSMO-IT, 2.8 km, 65 vertical levels, 2 runs/day), covering Italy (orange rectangle in Fig. 1a). Owing to the complex

Station	Elevation (m a.s.l.)	Measurement	Instrument	Data availability	Used in this study
Aosta-Downtown	580	PM_{10} hourly concentration	TEOM 1400a	1997–now	2015–2017
		$\rm PM_{10}$ and $\rm PM_{2.5}$ daily	Opsis SM200	2011–now	2015-2017
		concentrations			
		Water-soluble anion/cation	Dionex Ion Chromatography	2017–now	2017
		analyses on PM_{10} samples	System		
		EC/OC analyses on PM_{10}	Sunset thermo-optical analyser	2017–now ^a	2017
		samples			
		Metals on PM_{10} samples	Varian 820 MS	2000–now ^b	2017
		Total PAHs	EcoChem PAS-2000	1995–now	2017
		NO and NO2	Horiba APNA-370	1995–now	2017
		Standard meteorological	Vaisala WA15 (wind)	1995–now	2015-2017
		parameters			
Aosta-Saint-	560	Vertical profile of attenuated	CHM15k-Nimbus ceilometer	2015–now	2015-2017
Christophe		backscatter and derived			
(ARPA		products			
observatory)					
		Aerosol columnar properties	POM-02 sun/sky radiometer	2012–now ^c	2015–2017
		Surface particle size	Fidas 200s optical particle	2016–now	2016–2017
		distribution	counter		
Aosta-Saint-	545	Standard meteorological	Micros (wind)	1974–now	2015-2017
Christophe		parameters			
(weather station)					
Donnas	316	PM_{10} daily concentration	Opsis SM200	2010–now	2015–2017
		NO and NO2	Teledyne API200E	1995–now	2015–2017
Ivrea	243	PM_{10} daily concentration	TCR Tecora Charlie/	2006–now	2017
			Sentinel PM		

Table 1. Observation sites, measurements and instruments employed in this study.

^a The analysis is performed on 4 out of 10 days according to the laboratory schedule.

^b The analysis is performed on 6 out of 10 days according to the laboratory schedule.

^c Underwent major maintenance in the second half of 2016 and January 2017.

topography of the Aosta Valley, and the consequent need to resolve as much as possible the atmospheric circulation at small spatial scales, we used the latter version in the present work (cf. Sect. 4.5 for a discussion about possible effects of the finite model resolution in complex terrain).

FARM version 4.7 (Gariazzo et al., 2007; Silibello et al., 2008; Cesaroni et al., 2013; Calori et al., 2014) is a four-dimensional Eulerian model for simulating the transport, chemical conversion and deposition of atmospheric pollutants with 1 km spatial resolutiongrid, 1-hour temporal resolution and 16 different vertical levels (from the surface to 9290 m). FARM can be easily interfaced to most available diagnostic or prognostic NWP models, as done with COSMO in the present work. Pollutants emis-

- 5 sion from both area and point sources can be simulated by FARM, taking into account transformation of chemical species by gas-phase chemistry(Carter, 2000), dry and wet removalinto account. Particularly interesting for our study is the aerosol module (AERO3_NEW), coupled with the gas-phase chemical model and treating primary and secondary particle dynamics and their interactions with gas-phase species, thus accounting for nucleation, condensational growth and coagulation (Binkowski, 1999). Three particle size modes are simulated independently: the Aitken mode (diameter, $D < 0.1 \mu m$), the accumulation
- 10 mode (0.1 μm < D < 2.5 μm) and the coarse mode (D > 2.5 μm). A national (QualeAria,here referred to as "boundary conditions")and a regional FARM is only run over a small domain (light blue rectangle in Fig. 1a,b), roughly corresponding to the Aosta Valley. A regional emission inventory ("local sources", updated to 2015) emission inventories are is supplied to the CTM over the same area to accurately assess the magnitude of the pollution loads and their variability in both load and its variability in time and space. Data from a national inventory and CTM model (QualeAria, here referred to as "boundary conditional space).
- 15 conditions", outer rectangle in Fig. 1a), taken along the border of the inner (light blue) rectangle, are also used to estimate the mass exchange from outside the borders of the FARM domain.

3.2 Trajectory Statistical Models

The forecasted wind velocity profile from COSMO was here used as an input parameter into the publicly-available LA-GRANTO Lagrangian analysis tool, version 2.0 (Sprenger and Wernli, 2015), to numerically integrate the high-resolution 3-D

20 wind fields and to determine the origin of the air masses sampled by the ALC over Aosta-Saint ChristopheAosta-Saint-Christophe. We set up the program to start an ensemble of 8 trajectories in a circle of 1 km around the observing site and at 7 different altitudes from the ground to 4000 m a.s.l., for a total of 56 trajectories for every per run. A backward run time of 48 hours was considered sufficient, on average, to cover most of the domain of the meteorological model, while keeping minimising the numerical errorswith increasing running time contained.

25 3.3 Trajectory Statistical Models

Back-trajectories calculated with LAGRANTO were also then employed in Trajectory Statistical Models (TSMs) to provide a general picture of the geographical distribution of the probable aerosol sources. In this broadly used technique, the NWP domain is divided in <u>cells by a grid grid cells</u> (*i* and *j* indexesindices), and air parcels arriving to a specific receptor site are analysed. When a cell *ij* is crossed by a back-trajectory *l*, the tracer concentration c_l measured at the arrival point of that

30 trajectory (receptor) is considered. Finally, for each cell, a weighted average P_{ij} is calculated as follows to yield a map of the possible source areas (e.g., Kabashnikov et al., 2011):

$$P_{ij} = \frac{\sum_{l=1}^{L} F(c_l) \tau_{ij}(l)}{\sum_{l=1}^{L} \tau_{ij}(l)}$$
(3)

where L is the total number of trajectories and τ_{ij} the time spent by the trajectory l in the grid cell ij. F(c_l) is a function of the concentration at the receptor, and, depending on the chosen technique, can be the concentration itself (Concentration Weighted Trajectories (CWT) method, e.g, Hsu et al., 2003), the logarithm of the concentration (Concentration Field (CF)
method, Seibert et al., 1994) or the Heaviside step function H(c_l - c_T), where c_T is a concentration threshold (Potential Source Contribution Function (PSCF) method, e.g., Ashbaugh et al., 1985), generally the 75th percentile of the concentration series. In this last case, P_{ij} can be statistically interpreted as the conditional probability that concentrations above the threshold c_T at the receptor site are related to the passage of the relative back-trajectory through the location ij (e.g., Squizzato and Masiol, 2015). Additional iterative methods exist to reduce trailing effects and to better identify pollution hotspots (e.g., Stohl, 1996). Moreover, the obtained field P_{ij} can be further weighted as a function of the number of trajectories passing through each cell, thus reducing the impact of cells with limited statistics (Zeng and Hopke, 1989).

Any series of atmospheric measurements can be used as the concentration variable c_l in Eq. 3. In this work, we achieved especially meaningful results employing the scores from the PMF decomposition (Sect. 3.3), i.e. the contributions of the identified sources to the PM₁₀ daily concentration measured at Aosta–Downtown. The same approach, coupling PMF and

- 15 TSMs, was employed in other recent studies (e.g., Bressi et al., 2014; Waked et al., 2014; Zong et al., 2018). However, since only daily average information is available from the chemical speciation, the PMF output was repeated 8 times per day in order to allow correspondence with the 8 back-trajectories per day issued by COSMO and LAGRANTO. All methods (CWT, CF, PSCF) were explored and provided similar results. Therefore, since TSMs were used in this study to provide a general idea of the pollutant distribution and we did not aim at exhaustively discussing the differences among all existing techniques, only
- 20 results of the CF method were presented We present here (Fig. 13) results obtained with CF, having checked that application of CWT and PSCF methods provide similar outcomes. The COSMO-I2 domain was thus divided in 130x90 grid cells and 48h back-trajectories (interpolated to 5 minutes) were considered. Only points of the back-trajectories at altitudes lower than 2000 m a.s.l. were taken into account, i.e. this being the approximate average height of the mixing layer in the Po basin (Diémoz et al., 2019, and Fig. 20f in the present paper). Similarly, only trajectories ending at altitudes lower than 1500 m a.s.l. over
- 25 Aosta were examined, since the arriving air parcels should be able to impact on surface PM levels(, only trajectories ending close to the surface over Aosta were examined (i.e. altitudes < 1500 m a.s.l., the model surface altitude in the Aosta area being about 900 m a.s.l.). Weightings-To reduce statistical noise, every value P_{ij} of the resulting map was then multiplied by a weighting factor, w_{ij} , linearly varying from 0 (for N < 20 $N_{ij} \le 20$ end points in a cell) to 1 (for N > 200)were applied to the final concentration field to reduce statistical noise $N_{ij} \ge 200$), i.e. $w_{ij} = \min\{1, \frac{\max\{0, N_{ij} - 20\}}{180}\}$. To provide an idea of
- 30 the effect of this weighting procedure, TSM maps without any weighting (i.e., $w_{ij} = 1$) have been included in the Supplement (Fig. S7).

3.3 Positive Matrix Factorisation

The Positive Matrix Factorisation (PMF, Paatero and Tapper, 1994; Paatero, 1997) (PMF, Paatero and Tapper, 1994; Paatero, 1997) technique, as implemented in the US EPA PMF5.0 software (Norris and Duvall, 2014), was employed to study the 2017 series of aerosol chemical analyses and air quality measurements in Aosta–Downtown. The method allowed us to identify the possi-

5 ble emission sources of the observed aerosol, and notably to discriminate its local and non-local origin. PMF requires a long multivariate series of chemical analyses, matrix X (the *n* rows being the samples and *m* columns representing the chemical species) and decomposes it into the product of two positive-definite matrices, i.e. G ($n \times p$, matrix of factor contributions, *p* being the number of factors chosen for the decomposition) and F ($p \times m$, matrix of factor profiles) such that plus residuals (matrix E):

10 $\mathbf{X} = \mathbf{G}\mathbf{F} + \mathbf{E}$

(4)

A solution is found by minimizing the <u>so-called objective function Q</u>, i.e. the squared sum of the residuals (matrix E), weighted by their respective measurement uncertainties (this objective function is called Q) the measurement uncertainties. A total of 100 runs were performed in this work for each dataset to find an optimal solution. Here we considered three different series datasets X: a) the overall dataset of anion/cation analyses (data available almost every day in 2017, n = 360, "PMF-dataset a"),

- b) a subset of dataset a), selecting those measurements also having coincident EC/OC analyses (n = 132, "PMF-dataset b") and c) a subset of a), selecting those measurements also having coincident metal speciation (n = 209, "PMF-dataset c"). To make the decomposition of the three series comparable, an "unidentified" contribution, corresponding to the carbonaceous species only included in PMF-dataset b, was added to PMF-datasets a and c. This was done in the following way: we first estimated the total mass of crustal elements using the measured water-soluble calcium concentration as a proxy, with an empirical conversion
- 20 factor of 10 (e.g., Waked et al., 2014, a more rigorous estimate for the soil component, confirming this factor, was obtained from the PMF (e.g., Waked et al., 2014, note that a more rigorous estimate for the soil component was obtained from the PMF analysis itself and confirmed this factor); we then subtracted the sum of the available chemical components (including the estimated mass from soil components) from the total measured PM_{10} concentration, thus closing the mass balance. This difference was then added in PMF-datasets a and c as an unknown the "unidentified" source.
- Einally, NO, NO₂ and total PAHs measured at the same site (Aosta–Downtown) were furthermore included in the PMF to help identification of local pollution sources. The number of factors for each dataset was chosen based on the Q/Q_{exp} ratio and physical interpretability of the resulting factors (Sect. S4) and on the Q/Q_{exp} ratio. The latter is the ratio between the objective function obtained with the selected number of factors (Q, introduced before) and its expected value (Q_{exp}). Elevated (i.e., > 2) Q/Q_{exp} ratios could indicate that some samples and/or species are not well modelled and could be better explained by adding
- 30 another source (Norris and Duvall, 2014). The measured PM_{10} concentration was selected as the total variable to determine the contribution of each mode to the PM_{10} concentration.

PMF was additionally performed on volume size distributions measured by the Fidas OPC in Aosta–Saint Christophe Aosta–Saint-Christophe (Sect. 4.3.3), the *m* columns representing particle sizes, instead of chemical species.

4 Results

15

20

30

4.1 Daily classification schemes based on ALC measurements or meteorology

As Information on the vertical dimension obtained by the ALC was a key factor in identifying the events of aerosol pollution transport over the northwestern Alps (e.g., Diémoz et al., 2019). Therefore, a first step of our analysis was to set up a classification scheme of those advection dates based on ALC measurements. To this purpose, we used the longest ALC record available (2015–2017) and coupled it to meteorological measurements/forecasts. Since, to our knowledge, no previous ALC-based classification method is available in the scientific literature, we defined an original, daily-resolved classification scheme based on ceilometer measurements to group dates according to specific ALC-observed conditions. A graphical overview with examples for each class identified is given in Fig. 2. Overall, we identified six types (classes classes (A–F) of days:

- days "A": no aerosol layer or only low (i.e., < 500 m from the ground) aerosol layers visible for the whole day. These days are intended to represent unpolluted conditions or cases only affected by local sources, since aerosol transport from remote regions generally manifests as more elevated layers, as found by Diémoz et al. (2019);
 - days "B": only brief episodes of layers developing from the surface to altitudes > 500 m a.g.l. observed during the 24 hours. The origin of the aerosol layers of this intermediate class is uncertain, since they could either result from weak (not completely developed) advections from outside the investigated region or from puffs of locally-produced aerosol. For this reason, data in class "B" were used only as a transitional level, but not to draw definitive conclusions;
 - days "C": detection of an elevated, well-developed aerosol layer (usually, in the afternoon) and persistent during the night;
 - days "D": detection of an the aerosol layer from the previous day dissolving during the morning hours. No layers in the afternoon;
 - days "E": detection of a first aerosol layer from the previous day dissolving (or strongly reducing) in the morning and appearance of a separated structure in the afternoon;
 - days "F": thick aerosol layer persisting all day long.

Note that, although we also developed automatic recognition algorithms for classification purposes, we finally chose a classification based on visual inspection (for a total of 928 daily images <u>over the 3-year period</u>) to ensure the maximum quality of the flagging and avoid further sources of error.

We used the ALC classification above The ALC classification described above was used to derive the seasonal frequency distribution of the aerosol advections classescach aerosol advection class. Relevant results are shown in Fig. 3. Days not falling in those classes (e.g. complex-shaped layers, presence of low clouds or heavy rain hiding the aerosol layer, etc.) or including other kind of aerosol layers (as e.g., Saharan dust) were flagged as "Other" and were not considered for further analyses (this

accounting to for a maximum 26% of datesdays, in summer). The results reveal that clearly-detected aerosol advections (cases

13


Figure 2. Example of ALC images representative of each category (A-F) described in Sect. 4.1. Corresponding dates are 1 December 2015, 19 October 2016, 20 April 2016, 11 April 2015, 1 November 2017 and 27 January 2017. The dashed lines for categories C–F identify the sigmoid interpolation to the SR=3 envelope using the automated algorithm as explained in Sect. 4.2.



Figure 3. Frequency distribution of the aerosol layers observed in <u>Aosta–Saint Christophe Aosta–Saint-Christophe</u> based on the ALC classification described in Sect. 4.1. A=no layer, B=short episodes, C=layer detected in the afternoon, D=leaving layer, E=layer dissolving in the morning and a separated structure in the afternoon, F=persistent layer.

C to F) occur half (50%) of the days in summer and autumn and with a slightly lower frequency (45%) in winter and spring. A higher percentage of clear days (A) occurs in winter and spring, while persistent layers (F) are mostly found in winter-Conversely, while cases E are more frequent in summer.

- To assess how the ALC classification described above is related to the circulation patterns, we used the surface meteorological observations in Aosta-Saint Christophe Aosta-Saint-Christophe as drivers of a second, independent classification scheme as summarised in Table 2. Seasonal frequency distributions from this meteo-based scheme are shown-displayed in Fig. 4. These show that weather circulation types are remarkably variable during the year and help in interpreting the ALC-based results (Fig. 3). Winter In fact, winter is mostly characterised by stability wind calm (53% of the days), which sometimes favours stagnation and persistent aerosol layers (F). Plain-to-mountain breeze gradually increases diurnal winds gradually increase
- 10 from winter (only 11% of the days in that season, which reflects the high highest percentage of clear days A in that season(A)) to summer (68% of the days), when the thermally-driven regional circulation represents the main mechanism contributing to



Figure 4. Frequency distribution of circulation conditions in <u>Aosta–Saint Christophe Aosta–Saint-Christophe</u> based on the meteorological classification described in Table 2. Easterly winds (<u>Breeze diurnal wind systems</u> and <u>Upwindchannelled synoptic flow from the east</u>) are represented as orange slices, <u>stability wind calm</u> conditions in yellow and westerly winds (Foehn and <u>Downwindchannelled synoptic flow</u> from the west) in blue. Precipitation and unclassified days were not used in the study and are represented as grey slices and are not used further in the study.

Drimony	Secondary	Definition		
Fillinary	Secondary	Demitton		
classification	classification			
Easterly winds	Channelled synoptic flow from the east	$ v $ > 1 m s ⁻¹ , Δt >12 h, wind direction 0–180 $^{\circ}$		
		Night (19–8 UTC) ^a : calm wind ($ \boldsymbol{v} < 1 \text{ m s}^{-1}$)		
	Diurnal wind systems (east)	or low westerly wind ($ v < 1.5 \text{ m s}^{-1}$)		
		Day $(8-19 \text{ UTC})^{a}$: easterly wind $(v > 1.5 \text{ m s}^{-1}), \Delta t > 4 \text{ h}$		
Westerly winds	Foehn (west)	$ v > 4 \text{ m s}^{-1}, \Delta t > 4 \text{ h, wind direction} < 25^{\circ} \text{ or} > 225^{\circ} ^{\text{b}},$ RH< 40% no Foehn, $ v > 1 \text{ m s}^{-1}, \Delta t > 12 \text{ h, wind direction 180–360}^{\circ}$		
	Channelled synoptic flow from the west			
Wind calm	-	$ m{v} < 1~{ m ms^{-1}},\Delta t$ >20 h		
Other	Precipitation	Accumulated daily precipitation $> 1 \text{ mm}, \Delta t > 4 \text{ h}$		
	Unclassified	Every day that does not fall in the previous categories		

Table 2. Classification of weather regimes used in the study, based on wind speed (|v|), wind direction, daily duration of the condition (Δt) and other measured meteorological variables.

^a Both conditions must be met in order to discriminate such diurnal wind systems (inactive at night) from synoptic winds.

^b The directional range for the Foehn cases was determined experimentally by comparison with manual classification from a trained weather observer.

transport of polluted air masses (cases E). Foehn (rain shadow) winds are fundamental processes, especially in spring (22% of the days), leading to the removal of pollutants and frequent occurrence of clear days (A) in that season. Finally, upwind or downwind synoptic fluxes channelled synoptic flow from the east (or from the west) are also partly responsible for pushing polluted air masses respectively towards towards (or away from) the Aosta Valley, although they are markedly less frequent

5 compared to the breeze thermal winds mechanism.

The association between the ALC- and meteo-based classification schemes was explored using contingency tables (Fig. 5a). To this aim, the ALC classification (A–F) was further simplified and reduced to two main cases: presence of an arriving or stationary aerosol layer (classes C, E and F) and absence of any <u>elevated thick</u> aerosol layer (class A). Classes B (short and uncertain episodes) and D (layer leaving during the morning) were not used for the next considerations to avoid con-

- 10 funding conditions. Figure 5a shows that the presence of <u>an elevated a thick</u> aerosol layer is strongly connected to the wind provenancedirection, as expected. In fact, this scheme reveals that the layer appearance can be easily explained using the wind direction as the only information: when air masses come from the east, the probability of <u>identifying detecting</u> an aerosol layer over the Aosta Valley is of 93%, confirming the Po basin as an heavy aerosol source for the neighbouring regions, irrespectively of the day and period of the year. Conversely, this probability reduces to only 2% of the cases when the wind blows from
- 15 the west. In the period addressed, we thus detected few exceptions to the perfect correspondence (100% and 0%), which can however still be explained. For cases of easterly winds and no aerosol layer detected found (7%), possible reasons are:



Figure 5. (a) Contingency table showing occurrences of the aerosol layer seen by the ALC and wind provenancedirection. The blue boxes represent cases when the aerosol layer is not visible (day types A) and pink boxes cases when an arriving or stationary aerosol layer is revealed by the ALC (day types C, E and F). The numbers inside the boxes refers to the frequency and the number of occurrences for each couple of wind/layer classes. (b) Occurrence of the aerosol layer seen by the ALC and residence time of 48h back-trajectories in the Po plain.

- the up-valley breezediurnal wind, although clearly detected by the surface network, was too short of too short duration or too weak to transport the polluted air masses from the Po basin to Aosta-Saint-Christophe Aosta-Saint-Christophe (at least ~40 km must be travelled by the air masses along the main valley). This In our record, this was for example the case of 14 and 18 September 2015, and 17 June 2016;
- back-trajectories were indeed channelled along the central valley, however they did not come from the Po basin, but rather from the pre-Alpsother side of the Alps. This was the case, for instance, of 20 August 2015, in which air masses came from the Divedro Valley (north-east of the Aosta Valley) after crossing the Simplon pass (between Switzerland and Italy).

These exceptions also help understand why, in summer, about 70% of the days feature breeze systems (Fig. 4) while aerosol
layers are detected only about 50% of the days (Fig. 3). Indeed, some part (7%, Fig. 5a) of this 20% discrepancy can be attributed to the above events, the remaining 13% being likely associated with days with a complex aerosol structure (not classified in Fig. 2) or to days affected by low clouds and/or desert dust events. These days were therefore labelled as "Other" in Fig. 3.

There are few cases (2%, Fig. 5a) in our record in which, conversely, an aerosol layer was associated to with westerly, rather 15 than easterly, winds. These are as follows:

- on 5 February 2016 and 15 March 2016, the prevalent wind was from the west and the days were correctly identified as
 "DownwindChannelled synoptic flow from the west" and "Foehn", respectively. However, as soon as the wind turned and the valley-mountain breeze diurnal circulation started, the aerosol layer arrived;
- 5

- on 24 March 2016, a real case of transboundary/transalpine advection occurred and an aerosol-rich air mass was transported from the polluted French valleys on the other side of the Mont Blanc chain to the Aosta Valley. Although heavy pollution episodes can occur also on the French side of the Alps (e.g., Chazette et al., 2005; Brulfert et al., 2006; Bonvalot et al., 2016; Chemel et al., 2016; Largeron and Staquet, 2016; Sabatier et al., 2018), it is very rare to clearly spot an aerosol layer transported from that region to Aosta, since in those cases air masses coming from the west must have crossed the Alps and are almost always mixed with clean air from the uppermost layers, therefore only a dim aerosol backscatter signal is usually noticed from the ALC.

10

15

20

Overall, the good correspondence between the measured wind regimes and the aerosol layer detection by the ALC (Fig. 5a) suggests that the same association <u>ean_could</u> be employed to forecast the arrival of an advected aerosol layer based on the predicted wind fields. As an example of simple , forecasting capabilities of this phenomenon, Fig. 5b illustrates a contingency table relating the occurrence of an aerosol layer in <u>Aosta–Saint Christophe Aosta–Saint-Christophe</u> to the residence time over the Po Valley of the 48-hours back-trajectories arriving to the <u>Aosta–Saint Christophe Aosta–Saint-Christophe</u> observing site. Again, a direct, clear connection between both variables can be seen, with a 100% correspondence for residence times of air masses over the Po basin >10 hours.

Finally, it is worth mentioning that the proven high correlation between the circulation patterns (observed and/or simulated) and the presence of advected polluted layers in the Aosta area may be exploited in the future to reconstruct the impacts of aerosol transport on air quality over the Alpine region back to previous periods not covered by the ALC measurements (a 40-year-long meteorological database is available in the region).

4.2 Vertical and temporal characteristics of the advected aerosol layer

The 2015–2017 ALC time series in Aosta–Saint Christophe Aosta–Saint-Christophe is summarised in Fig. 6, showing monthly averages-1-hour resolved monthly average scattering ratios of the SR daily evolution. To draw the figure, the data were first

- 25 grouped in 1-hour bins (clouds were removed as well as periods with < 30 minutes of <u>A</u> data screening was preliminary performed by removing cloud periods with less than 30-minute measurements per hour), then monthly means were calculated (excluding points with and points < 1 week of measurements per month). The plot clearly shows that the maximum aerosol backscatter is generally found in the early morning and in the late afternoon, likely due to coupling between aerosol transport and hygroscopic effects (Diémoz et al., 2019), and not, as usual for plain sites, in correspondence of the midday development
- 30 of the mixing boundary layer. Also, the altitude of the layer varies with the season. Incidentally, months affected by exceptional events can be sharply distinguished, and namely : the frequent Saharan dust events in June and August 2017, and the forest fire plumes from Piedmont in October 2017, this latter again affecting the Aosta site in the afternoon, because of the breeze thermally-driven circulation. A similar climatology, showing the results in terms of average PM concentration profiles retrieved







Figure 7. (a) Aerosol layer top altitude for the classified days. Marker colour represents the type of the advection, while the marker shape represents the time of the day: morning (a.m.), afternoon (p.m.) or all day (applicable to cases F only). (b) Overall duration of the aerosol layer in a day (morning and afternoon durations were summed up in case E). The boxplots at the right represent synthetic information for each day type (same y-scale as the relative charts at the left). Missing measurements in summer 2015 are due to the replacement of the ALC laser module.

by the ALC is given in Fig. S1 of the Supplement. In that case, the conversion from backscatter to PM_{10} was obtained using the methodology described by Dionisi et al. (2018).

5

To quantitatively explore the spatial and temporal characteristics of the ALC-detected aerosol layer over the long term, we developed an automated procedure, described in detail in the Supplement (Sect. S2), to identify and fit with a sigmoid curve the space-time region of the ALC profiles affected by the aerosol advection (using $SR \ge 3$ as threshold value). This allowed to objectively determine, for example, the height and the duration of the these advections. The long-term statistics of these two variables is summarised in Fig. 7. The altitude of the polluted aerosol layer (Fig. 7a) displays, as expected,

a clear seasonal cycle, with minimum in winter and maximum, up to 4000 m a.g.l., in summer. The overall cycle mimics the variability of the convective boundary layer (CBL) height found in the Po basin by other studies (Barnaba et al., 2010; Decesari et al., 2014; Arvani et al., 2016), with somewhat higher values that reflect the modification of the boundary layer structure in mountainous areas (Teixeira et al., 2016)(De Wekker and Kossmann, 2015; Serafin et al., 2018). Notably, stronger,

- 5 multi-scale thermally-driven flows (e.g. the ones developing on the slopes of the valley) further redistribute the aerosol particles in the vertical direction, thus pushing the upper limit of the CBL to the ridge height. Average values of the different classes, represented as box plots in Fig. 7, are clearly impacted by the season of their maximum frequency over the year (Fig. 3). In fact, minimum altitude of the layer top was found in days F owing to their maximum occurrence in winter, while top altitude maximum was found in days E being these mostly summertime events. Great variability was also found for the duration of the
- 10 advection (Fig. 7b), ranging from few hours in the weakest events to 24 hours/day in the extreme cases (full day, by definition, for cases F). The coupling between the duration of the events , together with and their absolute aerosol load , are expected to be the major drivers of determines the impact of the investigated phenomenon on surface air quality. This is investigated , as discussed in the following paragraphs.

4.3 Impact of Po Valley advections on northwestern Alps surface-level PM loads and physico-chemical characteristics

- 15 To quantify the impact of the aerosol layers detected by the ALC on the air quality at ground level, we first investigated how the PM concentration daily evolution measured by the surface network changes depending on the ALC-identified classes (Sect. 4.3.1). We then used the Positive Matrix Factorisation of the multivariate dataset of chemical analyses to explore the chemical markers associated to with the transport from the Po basin (Sect. 4.3.2). The effectiveness of the identified markers and the coupling between chemistry and meteorology were also confirmed by the use of Trajectory Statistical Models. Finally,
- a link between aerosol chemical and physical properties was investigated in Sect. 4.3.3, where particle chemical composition is coupled to measurements of aerosol particle size. Finally, a preliminary assessment on of the effect of the investigated advections on surface pollutants other than particulate matter (e.g., NO_x partitioning) was also performed and is reported in the Supplement (Sect. S5).

4.3.1 Surface PM variability in relation to the ALC profiles classification scheme

- We started investigating the effect of aerosol transport on the Aosta Valley surface air quality by analysing the variations of the daily mean PM concentrations measured by the ARPA surface network as a function of the ALC classes introduced in Sect. 4.1. Figure 8a shows the relevant results, resolved by season. It is quite evident that PM_{10} concentrations in Aosta– Downtown are in fact highly correlated with the presence/strength of the aerosol layers seen by the ALC. PM_{10} on days with a persistent aerosol layer (class F) was found to be 2 to 3.5 times higher, on average, than on non-event days (class A, this
- 30 difference being statistically significant for all seasons, p-value < 0.05 from the Mann-Whitney test(Mann and Whitney, 1947))Mann and Whitney (1947) test. Similar behaviour was seen in PM_{2.5} concentrations measured at the same station (Fig. S2a) and in the PM_{2.5}/PM₁₀ ratio (Fig. S2b), this latter indicating that aerosol particles are smaller, on average, on days when the ALC detects a thick aerosol layer compared to non-event days. Causality between the presence of an elevated layer and



Figure 8. Daily-averaged surface concentrations (2015–2017) of PM_{10} (a), nitrogen oxides (b) and polycyclic aromatic hydrocarbons (c) in Aosta–Downtown as a function of the ALC classes and season. (d) PM_{10} surface concentration at the Donnas station. The EU-established PM_{10} daily limit of 50 µg m⁻³ and the NO₂ hourly limit of 200 µg m⁻³ are also drawn as references (horizontal dashed lines).



Figure 9. Comparison between daily PM_{10} anomalies (Eq. 5) registered in Aosta–Downtown and Donnas (40 km apart), relative to a monthly moving average. Circles colour represents the ALC classification, see legend, the "X" symbol represents the unclassified days. The 1:1 and the best fit line are also represented on the plot. The Pearson's correlation index between the two series is $\rho = 0.73$.

variations of the PM surface concentration, however, cannot be rigorously inferred at this point, since, in principle, common environmental conditions affecting both PM at ground and the aerosol in the layers aloft could originate the observed correlation. Nevertheless, it is interesting to note that this effect is less detectable for other pollutants measured by the network(Figs. 8b and e). In fact, concentrations of typical locally-produced pollutants such as NO_x and PAHs (Figs. 8b and c) do not change as much

5 as PM among the ALC classes, with only rare exceptions (e.g., class A, which is slightly influenced by foehn-Foehn winds, and class F, in which temperature inversions/low mixing layer heights may enhance the effect of local surface emissions). These results point towards excluding that the correlation (Fig. 8a) trivially originates from common weather conditions or from the uneven distribution of the ALC classes among the seasons.

Furthermore, large correlation was also found between the ALC classification obtained in Aosta-Saint Christophe only

- 10 available in Aosta–Saint-Christophe and the surface PM_{10} concentration recorded in Donnas (Fig. 8d). Since the two stations are located at 40 km distance, this result suggests that a major role is played by large-scale dynamics rather than by local sources and that the phenomena under consideration have at least a regional-scale extent. As a further test, we correlated the daily PM_{10} concentrations measured in Donnas to those measured in Aosta. To this purpose, we considered the anomaly $(\Delta PM_{10}, Eq. 5)$ with respect to the monthly moving average $(\langle PM_{10} \rangle_m)$, in order to avoid large, fictitious correlations due
- 15 to the same yearly cycle (maximum PM concentration in winter and minimum in summer), and only focus on the short-term dynamics, we considered the anomaly (Δ) with respect to the monthly moving average ($\langle PM_{10} \rangle_m$), i.e.:



Figure 10. (a) Daily PM_{10} cycle sampled by the OPC in <u>Aosta–Saint Christophe Aosta–Saint-Christophe</u> during non-event days (class A) and days with arriving and leaving aerosol layers (classes C and D), plotted together with the daily evolution of PAH as a proxy of the local sources (<u>dotted_dashed</u> line, right vertical axis, in ng m⁻³). (b) Same as (a) using the TEOM in Aosta–Downtown.

$$\Delta \mathbf{P}\mathbf{M}_{10}(t) = \mathbf{P}\mathbf{M}_{10}(t) - \langle \mathbf{P}\mathbf{M}_{10} \rangle_{\mathbf{m}}(t)$$
(5)

where t is a specific day. The scatterplot of these anomalies is shown in Fig. 9. It highlights the good correlation between the ΔPM_{10} at the two sites ($\rho = 0.73$ when considering all classes and $\rho = 0.76$ if only only considering advection cases, i.e. classes C to F, are considered). Using). Clustering of these results using the ALC classification to cluster these results (circle colours) further helps showing: 1) shows: the different ΔPM_{10} associated to with the different classes and 2) the good

5 (circle colours) further helps showing: 1) shows: the different ΔPM_{10} associated to with the different classes and 2) the good correspondence of this effect at both sites. Notably, in the most severe cases (E and F), the anomalies in Donnas are generally larger than in Aosta–Downtown (the best fit line being y = 1.1x + 0.8 for cases E and F only), which is compatible with this site being closer to the Po basin (Fig. 1).

The advected aerosol layers were also found to impact the daily evolution of surface PM concentrations. To investigate this
aspect, we used hourly and sub-hourly PM₁₀ data from both the Fidas 200s OPC in Aosta-Saint Christophe Aosta-Saint-Christophe and the TEOM in Aosta-Downtown. The Figure 10 shows the PM₁₀ daily cycles are presented for cases A (no advection), C and Donly, for sake of clarity, and for both Aosta-Saint Christophe, for both Aosta-Saint-Christophe (Fig. 10a) and Aosta-Downtown (Fig. 10b). Despite the instruments limitations described in Sect. 2, the figure shows that local effects play an important role on the PM₁₀ daily evolution, as visible from the morning and afternoon peaks. In fact, these maxima, common

to most pollutants, are the results of the coupled effect of varying local emissions and boundary layer height during the day. The daily cycle of PAH concentrations is added to the plot as a proxy of diurnal cycle from local sources (dotted dashed line, right y-axis). This cycle is very similar to the PM₁₀ one for case A. Conversely, the influence of the advections (C and D) is clearly visible on the corresponding PM₁₀ cycles. In fact, Fig. 10 shows the PM₁₀ concentration to markedly increase in the afternoon of days C (by 1.0 and 0.7 μ g m⁻³ h⁻¹ in Saint-Christophe Aosta-Saint-Christophe and Aosta-Downtown, respectively) and



Figure 11. Percentage of aerosol mass concentration carried by each mode for the three PMF-datasets considered in the analysis (PMF-dataset a: anion/cation only; PMF-dataset b: anion/cation together with EC/OC; PMF-dataset c: anion/cation together with metals), and their respective confidence intervals from the BS-DISP test. Details are provided in Sect. S4.

to markedly decrease in days D (by -1.3 and -0.7 μ g m⁻³ h⁻¹). This effect is similar at the two sites although less marked in Aosta–Downtown, especially at night. This is probably due to TEOM loss of volatile species in the advected aerosol (see also the PMF results on chemical analysis in the following section). Similar results (not shown) were obtained when splitting the data on a seasonal basis, which excludes that the observed behaviour originates from the seasonal cycle.

5 4.3.2 Chemical species within the advected aerosol layers

As a further step to adequately discriminate local and non-local sources, especially when dealing with data collected in an urban environment such as Aosta–Downtown, we took advantage of the one-year-long (2017) chemical aerosol characterisation dataset in the urban site of Aosta–Downtown and applied the PMF to the three datasets described in Sect. 3.3 (PMF-datasets a, b, c, i.e., anion/cation only, anion/cation with EC/OC and anion/cation with metals, respectively). The Results of this analysis

- 10 <u>show the</u> main factors shaping the composition of PM_{10} at the investigated site are <u>those</u> given in Fig.11, all details on this outcome being given in Sect. S4. The question addressed here is, <u>however</u>, how aerosol transport from the Po valley affects the chemical properties of PM_{10} sampled in Aosta. In the preliminary analysis of three cases studies reported in the companion paper, we found that the advected layers were rich of nitrates and sulfates (accounting, together with ammonium, for 30-40% of the total PM_{10} mass). This kind of secondary inorganic aerosol was indeed found in high concentrations in the Po basin by
- 15 previous studies (e.g., Putaud et al., 2002; Carbone et al., 2010; Putaud et al., 2010; Larsen et al., 2012; Saarikoski et al., 2012;



Figure 12. (a) Temporal chart of the contribution of secondary (nitrate-rich and sulfate-rich) modes to the total PM_{10} . The dashed line shaded area represents the estimated local production of secondary aerosol. The difference between each dot and the this baseline can thus be read as the non-local contribution to the aerosol concentration in Aosta–Downtown. Since ions chemical speciation started in 2017, only one year of overlap with the ALC is available at the moment (see Table 1). (b) Boxplot of the contribution of secondary (nitrate-rich and sulfate-rich) modes to the total PM_{10} concentration as a function of the day type. PMF-dataset "a" was used for both plots.

Gilardoni et al., 2014; Curci et al., 2015). Here we further tested on a longer dataset if the sum of the nitrate- and sulfate-rich factors (i.e. secondary aerosols) could in fact represent a good chemical marker of aerosol transport from the Po basin (e.g., Kukkonen et al., 2008; Tang et al., 2014). In this respect, it is worth highlighting that these PMF modes are not correlated with typical locally-produced pollutants such as NO_x and PAHs (this is revealed by the low percentage contribution of NO_x and

- 5 PAHs in nitrate-rich and sulfate-rich modes in Figs. S3–S5), which therefore excludes their local origin. We then combined the chemical information (the sum of the secondary components) with the ALC-classification. This was done for year 2017, which is the only overlapping period between anion/cation analyses and ALC profiles (see Table 1). Results are shown in Fig. 12. In particular, Fig. 12a shows the time evolution of the mass concentration carried by the secondary aerosol modes, in which each date is coloured according to the six ALC classes identified. It can be noticed that almost all peak values occur during days of
- 10 type E or F, i.e. when the advected layer is more persistent and able to strongly influence the <u>local</u> air quality at ground. The very good correspondence between the sum of the nitrate- and sulfate-rich modes contributions and the ALC classification, as



Figure 13. (a) Output of the Concentration Field <u>TSMTrajectory Statistical Model</u>, using the sum of the contributions from PMF nitrate- and sulfate-rich modes as concentration variable at the receptor (Aosta–Downtown). (b) <u>CF_Concentration Field</u> based on the traffic and heating mode from PMF. Trajectories are cut at the borders of the COSMO-I2 domain.

well as the poor correlation between the latter and the role of the other PMF-identified modes (not shown), suggests that the link between the secondary components and the aerosol layer advection is not trivially due to particular weather conditions affecting both variables. Rather, the coupling between the chemical and the ALC information indicates that aerosols of secondary origin dominate during the advection episodes from the Po basin.

A summary of the contribution of secondary modes to the total PM₁₀ as a function of the day type, on a statistical basis, is given in Fig. 12b. The Mann-Whitney test, applied on these data, confirms that the contribution of secondary pollutants is significantly lower for class A compared to B (p=0.0005, 3 · 10⁻⁸), C compared to D (p=6 · 10⁻⁸), D compared to E (p=9 · 10⁻⁷), and E compared to F (p=6 · 10⁻⁷). Figure 12b also allows to quantify the contribution of non-local sources using as baseline the results obtained for class A (i.e., no advection, dashed line shaded area in Fig. 12a). Note that this baseline
is very low on average, about 3 µg m⁻³, as expected from the unfavourable conditions for secondary aerosol production in the area under investigation (e.g., low expected emissions of ammonia, frequent winds, and from the unobserved correlation between secondary aerosol components and their gas-phase precursors). The non-local contribution is calculated as the average

difference between the mass from the secondary modes and this baseline, and amounts to about 5 $\mu g m^{-3}$ on a yearly basis,

- i.e. 24% of the total PM₁₀ concentration reconstructed from the PMF. MoreoverOn a seasonal basis, the absolute impact of air
 masses transport depends on the coupling between emissions (stronger in winter and weaker in summer) and weather regimes
 (e.g., thermal winds occurring more frequently in summer/autumn with respect to winter/spring, Sect. 4.1). This results in a
 - PM_{10} contribution of nearly 6 µg m⁻³ in winter and autumn, 4 µg m⁻³ in summer, and 3 µg m⁻³ in spring. In terms of relative contribution, this also depends on the "background" PM_{10} levels in Aosta. It is therefore highest in summer (32%), when

thermally-driven fluxes are more frequent and local emissions lower, and lowest in winter (16%), when thermal winds are less frequent and local emissions higher. Intermediate relative values are found in spring and summer (27% and 28%, respectively). In specific episodes, advections from the Po basin account for may still produce an increase of PM_{10} concentrations up to 50 μ g m⁻³in specific episodes, i.e. exceeding alone the EU daily threshold. The most important compounds contributing to the

- 5 this increase are nitrate, ammonium and and sulfate, i.e. the species characterising the nitrate- and sulfate-rich PMF modes. However, since the sulfate-rich mode additionally includes organic matter (OM, cf. Figs.S3–S5), the latter species is also relevant in the advection episodes, especially in summer, when the nitrate-rich mode has a lower-its lowest contribution. This finding agrees with previous works, identifying the Po basin as a source of organic aerosol (Putaud et al., 2002; Matta et al., 2003; Gilardoni et al., 2011; Perrone et al., 2012; Saarikoski et al., 2012; Sandrini et al., 2014; Bressi et al., 2016; Khan et al.,
- 10 2016; Costabile et al., 2017).

The Po Valley origin of the secondary components was also further proved coupling the chemical information to backtrajectories. Figure 13a shows the result of the TSM using the sum of the nitrate- and sulfate-rich modes as concentration variable at the receptor. It clearly reveals that trajectories crossing the Po basin have a much higher impact on secondary aerosol measured in Aosta compared to air parcels coming from the northern side of the Alps. Interestingly, this is not the case

15 using different source factors from PMF. For example, Fig. 13b reports the same map, but relative to the traffic and heating mode, which is clearly much more homogeneous compared to Fig. 13a and essentially reflects the density distribution of the trajectories. This behaviour highlights , as expected, the local origin of the PMF source factors other than the two (secondary aerosols) chosen as proxies of the advection (secondary aerosols). As sensitivity tests, similar maps, without any weighting applied, are reported in Fig. S7. No substantial differences can be noticed.

20 4.3.3 PMF of aerosol size distributions and links to chemical properties

Exploring the links between aerosol chemical and physical properties is useful to get insights into the atmospheric mechanisms leading to particles formation and transformation during their transport to the Aosta Valley. Therefore, we investigated correlations between the results of the chemical analyses on samples collected in Aosta–Downtown and particle size distributions (PSDs) measured by the Fidas OPC in Aosta–Saint Christophe Aosta–Saint-Christophe. To ensure comparability between these

- 25 two datasets, the particle volume distributions from the Fidas OPC (normally extending up to sizes of 18 μm) were weighted by a typical cut-off efficiency of PM₁₀ sampling head (Sect. S6). We then performed a PMF analysis of the hourly-averaged PSDs from the Fidas OPC (hereafter referred as size-PMF). Four principal modes were identified, as. Their relative contribution to the total particle volume is shown in Fig. 14a(their dependence on season, wind speed and direction is also provided in Fig. S9). The curves of the relative contributions of the four size-PMF modes to the total particle volume are approximately. These
- 30 modes are centred at 0.2, 0.5, 2 and 10 µm diameter, respectively (and will be thus referred to as 0.2, 0.5, 2 and 10-µm size-PMF modes in the following). A similar figure, showing their dependence on season, wind speed and direction, is also provided in Fig. S10. The number of factors (p) was chosen based on physical considerations: if p > 4, then the last mode (largest sizes) splits into sub-modes, which are not relevant for the present study; conversely, if fewer components (p < 4) are retained, they merge into unphysical modes (e.g., very large and very fine particles combined together).



Figure 14. (a) Modes identified by the PMF analysis applied on the PSDs measured by the Fidas OPC (size-PMF). (b–d) Comparison between the <u>PMF</u> scores time series obtained from <u>PMF</u>-analysis of chemical speciation (chem-PMF on PMF-dataset "a", showing mass, left-hand vertical axes, mass) and size distributions (size-PMF, showing volume, right-hand vertical axes, volume). The three panels show: (b) contributions from chem-PMF nitrate-rich (black) and size-PMF 0.5 μ m (green) modes; (c) contributions from the sum of the chem-PMF sulfate-rich and traffic and heating modes (black), and from the 0.2 μ m size-PMF mode (blue); (d) sum of contributions from chem-PMF road salting and soil modes (black), and from 2 and 10 μ m size-PMF modes (red). Correlation values (ρ) are also reported in each plot.

The scores from the size-PMF were then averaged on a daily basis to be compared to the chemical PMF ones (hereafter, chem-PMF; Sect. S4). Figure 14 compares time series of selected chem-PMF (dataset a) and size-PMF results. It shows that:

(a) the nitrate-rich mode from chem-PMF and the 0.5- μ m size-PMF mode correlate very strongly (ρ =0.81, Fig. 14b);

- (b) the sum of the sulfate-rich mode and traffic and heating mode correlate very strongly with the 0.2-µm size-PMF mode
- 5

10

15

(ρ =0.82, Fig. 14c)with the 0.2-size-PMF mode. Note that the correlation index decreases (ρ between 0.35 and 0.69) if the sulfate-rich mode and traffic and heating mode are compared separately with the 0.2 µm mode;

- (c) A moderate, statistically significant (p-value< $2.2 \cdot 10^{-16}$), Pearson's correlation index $\rho = 0.59$, Fig. 14d) correlation was found between the chem-PMF soil plus road salting modes and the size-PMF coarser modes (sum of 2 and 10 µm modes, Pearson's correlation index $\rho = 0.59$, p-value< $2.2 \cdot 10^{-16}$, Fig. 14d). This suggests that these latter likely arise from both indicate the same source, and likely non-exhaust traffic emissions, such as abrasion from brakes, tire wear and road (with typical sizes of 2–5 µm), and resuspension from the surface (up to >10 µm), as also found in other studies (e.g., Harrison et al., 2012). This agreement is remarkable when between the two independent datasets is remarkable considering that the particles were sampled at two different sites (Aosta–Saint Christophe Aosta–Saint-Christophe and Aosta– Downtown) with distinct environmental features, and that coarse particles usually travel for short ranges. It is also worth mentioning that the correlation between single modes from chem-PMF (road salting or soil) and size-PMF (2 or 10 µm) separately is weaker (ρ between 0.26 and 0.55) than the one resulting from their sum. Finally, from a preliminary analysis based on back-trajectories and desert dust forecasts (NMMB/BSC-Dust, http://ess.bsc.es/bsc-dust-daily-forecast), the 2
 - µm component seems to be additionally connected to deposition, and possibly resuspension, of mineral Saharan dust in Aosta, an effect already observed in other urban areas in central Italy (Barnaba et al., 2017).
- 20 A further interesting feature is the <u>clear</u> size separation of the 0.2 and 0.5 μm accumulation modes, which likely results from different aerosol formation mechanisms in the atmosphere: condensation/coagulation from primary emissions/aging on one side (0.2-μm mode, also known as "condensation mode"), and aqueous phase processes (e.g., in fog) on the other side (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2012; Costabile et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2 (forming "droplet mode" aerosol particles of about 0.5 μm diameter, e.g., Seinfeld and Pandis, 2006; Wang et al., 2 (forming "droplet mode" aerosol particles d
- 25 bile et al., 2017). Finally, the very good correlation between the fine particles and the nitrate- and sulfate-rich modes at the two stations is a further hint that this kind of particles are mostly of non-local origin.

Overall, the good agreement between the size- and chem-PMF results further strengthens their independent outputs.

4.4 Impact of Po Valley advections on northwestern Alps columnar aerosol properties

ALC measurements revealed the vertical extent and thickness of the advected polluted layers from the Po Valley. We therefore also investigated if and how much these layers impact the column-integrated aerosol properties (i.e., those sounded by groundbased sun photometers, but also by satellites). To this purpose, the ALC day type classification was applied to the measurements of the Aosta–Saint Christophe Aosta–Saint-Christophe sun photometer. The average AOD at 500 nmnm, binned by day type



Figure 15. (a) Average aerosol optical depth at 500 nm from the POM sun photometer for each day type (colour code as in Fig. 12) and season; (b) Ångström exponent calculated in the 400-870 nm nm band; (c) Yearly-averaged single scattering albedo at 500 nm. B days in spring were removed from panels (a) and (b) due to insufficient statistics (only 3 days).

and season, is shown in Fig. 15a. It can be noticed that a general increase of the AOD is found from classes A (about 0.04 on average) to F, for which AOD is more than 4 times higher (0.18). The advections are also found to affect the Ångström exponent (Eq. 2, Fig. 15b), a parameter used to qualitatively infer representative of the mean particle size. Results from this column-integrated perspective confirm that the transported particles are on average smaller than the locally-produced ones. In

5

fact, the mean Ångström exponents vary from 1.0 for days A in winter and autumn, up to 1.6 for days E-F, and show a general increase among the classes for every season. To confirm and fully understand this dependence of the Ångström exponents on the ALC classification, we also investigated the columnar volume-PSDs retrieved from the sun photometer almucantar scans during the Po Valley pollution transport events. This analysis (Fig. <u>\$10</u>\$11) confirms what we already found with the surface-



Figure 16. Long-term (1 year) comparison of between PM_{10} surface measurements and simulations (FARM) at the surface in Aosta– Downtown (a) and Donnas Ivrea (b). Simulations including and excluding the boundary conditions are both shown.

level PSDs from the Fidas OPC, i.e. that the advections increase the number of particles in all sizes, and particularly notably in the sub-micron fraction. This explains the higher Ångström exponents retrieved by the sun photometer during the advections. To again link <u>A further link between</u> aerosol physical and chemical properties <u>, is explored through</u> the variability of the single scattering albedo with day type is also shown in (Fig. 15c). Yearly-averaged data are shown in this case due to the

5 limited number of data points available for this parameter (the almucantar plane must be free of clouds to accurately retrieve the SSA, Sect. 2). Fig. Figure 15c reveals that SSA at 500 nm generally increases from class A to class F (0.92 to 0.98), which agrees with the fact that secondary, and thus more scattering, aerosol is transported with the advections. This information is particularly important for follow-up radiative transfer evaluations under the investigated conditions. In this respect, also note that further analysis more focussed on the impact of these advection on particle absorption properties is planned for the future.

10 4.5 Comparison between observations and CTM simulations

Difference between surface concentrations in Donnas simulated by FARM taking the boundary conditions into account (FARM_{tot}) and without taking them into account (FARM_{local}, only local sources), as a function of the advection class observed by the ALC.

Accurate simulations of air quality metrics are of utmost importance for Environmental Protection Agencies. Indeed, not only they are are they essential from a scientific/technical perspective (contributing to better understand the local and remote

15 pollution dynamics), but they also represent fundamental duty towards the public, being air quality forecasts disseminated every day. In this section, we compare long-term (1 year, 2017) daily averages of surface PM-PM₁₀ to the corresponding simulations



Figure 17. Differences Absolute (a,c) and relative (b,d) differences between simulated and observed PM_{10} concentrations at the surface in Aosta–Downtown. The mean bias error (MBE) and the normalised mean bias error (NMBE) for each case is are reported in the plot titles. First row: FARM simulations as currently performed in by ARPAfor the Donnas (a) and Aosta–Downtown (b) stations. Second row: the PM_{10} concentrations from outside the boundaries of the domain were multiplied by a factor W=4.

Aosta–Downtown

Table 3. Statistics of comparison between PM_{10} model forecasts and measurements at the site of Aosta–Downtown (mean bias error, MBE; root mean square error, RMSE; Pearson's correlation coefficient, ρ ; normalised mean bias error, NMBE; normalised mean standard deviation, NMSD). Results with both W = 1 and W = 4 weighting factors to the PM fraction arriving outside the boundaries of the regional domain are reported.

W	$MBE~(\mu gm^{-3})$	$\text{RMSE}~(\mu gm^{-3})$	NMBE (%)	NMSD (%)	ρ
1	-7	15	-35	-16	0.62
4	1	13	5	-8	0.61



Figure 18. Same as Fig. ??, Difference between PM_{10} surface concentrations in Aosta–Downtown simulated by FARM taking the boundary conditions into account (FARM_{tot}) and without taking them into account (FARM_{local}, only local sources), as a function of the advection class observed by the ALC.

by FARM at two sites of the regional domain (Aosta–Downtown and Donnas) in Aosta–Downtown (similar results are found for the other sites in the domain and are not reported here) and next to Ivrea. This exercise was also aimed at evaluating if and how the information gathered by the ALC can be used to understand deficiencies and thus improve the model performances. The one-year time series of both the experimental measured and simulated PM_{10} in the Aosta–Downtown and Ivrea sites

- 5 are displayed in Fig. 16. They Model and measurement show similarities (such as the same seasonal cycle, indicating that local emissions from the regional inventory and general atmospheric processes are correctly reproduced), but also divergences (especially PM_{10} underestimation by FARM, as already shown and discussed in the companion paper). Table 3 summarises some statistical indicators of the model-measurements comparison, such as and namely mean bias error (MBE), root mean square error (RMSE), Pearson's correlation coefficient (ρ)and normalised mean bias error (NMBE), normalised mean standard
- 10 deviation (NMSD), i.e. $(\sigma_M/\sigma_O 1)$, where σ_M and σ_O are the standard deviations of the model and observation) and

Pearson's correlation coefficient (ρ). The overall performances of FARM are comparable to the results obtained in other studies using CTMs (e.g., Thunis et al., 2012). In For the Aosta–Downtown case, we additionally explore the absolute (Fig. 17(a, b) we explore the a) and relative (Fig. 17b) differences between modelled and observed PM₁₀ in relation to the ALC-derived classes. It reveals These results reveal that the model underestimation for both Aosta–Downtown and Donnas mostly occurs in the cases

- 5 of strongest advections (F), with extreme model deviations as low as -40 μ g m⁻³ on days of class F. This is aclear indication that the external contribution (Fig. 17a), i.e., -50% or lower (Fig. 17b). The observed model-measurement discrepancies might originate from (1) an incomplete representation of the inventory sources (emission component), (2) inaccurate NWP modelling of the meteorological fields, and notably the wind (transport component), or a combination of (1) and (2). Some details on these aspects are provided in the following.
- 1. Emissions. Inaccuracies in the (local and national) emission inventories could degrade the comparison between simulations 10 and observations. Based on results shown in Fig. 17(a,b), we decided to investigate the sensitivity of our simulations in Aosta-Downtown to the magnitude of the external contributions (boundary conditions)is not optimally parametrised in the model. We made some sensitivity tests, using. In particular, we used a simplified approach to provide a general idea of the possible outcomes while speeding speed up the calculations. In particular, : assuming that the contribution from outside the regional domain and the local emissions add up without interacting, we simulated the surface PM_{10} 15 concentrations turning on/off the boundary conditions (dark- and light-blue lines in Fig. 16) to roughly estimate the only contribution from sources outside the regional domain. Interestingly, the difference between the two FARM runs correlates well with the advection classes observed by the ALC (Figs. ?? and Fig. 18). This clear correlation between the simulations and the experimentally-determined atmospheric conditions indicates is a first good indication that the 20 NWP model used as input to the CTM yields reasonable meteorological inputs to FARM. The reason for discrepancies is therefore likely in the emissions. To Then, to further explore the sensitivity to the boundary conditions, we then gradually increased, by a weighting factor W, the PM_{10} concentration from outside the boundaries of the domain trying to match the observed values. In FigFigs, 17(c, d) we show the results obtained with W=4. The This exercise show that the overall mean bias error is much reduced compared to the original simulations (W=1, Figs. 17(a, b)), especially for the winter 25 and autumn seasons, while slight overestimations are now visible for summer and spring. Also the annually-averaged MBE, NMBE and NMSD improve (Table 3), although whilst the other statistical indicators remain stable or even slightly worsen.
- 30
- Clearly, our simplified test has major limitations. For example, seasonally- and geographically-dependent (i.e., relative to the position on the regional domain border) factors W should be used to better correct deficiencies in the national inventory. Also, the high weighting factors needed to match the measurements in winter and autumn suggest not only underestimated emissions, but also missing aerosol production mechanisms (e.g., aqueous-phase particle production as in fog/cloud processing, Gilardoni et al. (2016)) during the cold season. In fact, this simplified exercise was only intended to roughly estimate the magnitude of the "outside PM_{10} tuning factor" necessary to match the observations. IndeedDespite this limitation, this numerical experiment demonstrates that the accuracy of the still shows quite convincingly that biases

in air quality forecasts may be improved can be reduced by revising the inventories , principally the national one, on the basis of experimental dataand particularly, as those from unconventional air-quality systems as ALC. (e.g., the ALCs used in this study);

2. Transport. Although the COSMO-I2 grid spacing is certainly in line with that of other state-of-the-art operational limited-area models, in our complex terrain it could be insufficient to appropriately resolve local meteorological phenomena 5 triggered by the valley or orgraphy (Wagner et al., 2014b), also considering that the actual model resolution is 6–8 times the grid cell (Skamarock, 2004). For example, Schmidli et al. (2018) show that at 2.2-km grid step the COSMO model poorly simulates valley winds, while at 1.1-km grid step the diurnal cycle of the valley winds is well represented. Similarly, Giovannini et al. (2014) show that 2-km step can be considered as the limit for a good representation of valley 10 winds in narrow Alpine valleys. Smoothed digital elevation model (DEM) used within COSMO and FARM could also play a direct role in the detected underestimation. In fact, as mentioned in Sect. 3.2, the model surface altitude of the Aosta urban area is 900 m a.s.l., whereas the actual altitude is about 580 m a.s.l. (Table 1). The adjacent cells are given an even higher altitude, owing to the fact that the valley floor and the neighbouring mountain slopes are not properly resolved at the current resolution. This results in an apparent lower elevation of the sites located in flatter and wider 15 areas, such as Aosta, while the real topography profile at the bottom of the valley presents a monotonic increase from the Po plain to Aosta. Therefore, it is expected that, just by better reproducing the orography, a higher resolution would allow to better resolve the horizontal advection of aerosol-laden air from much lower altitudes on the plain.

Most of these issues were extensively addressed in the companion paper (Diémoz et al., 2019). In that study, COSMO-I2 was shown to be capable of reproducing the mountain-plain wind patterns observed at the surface both on average (cf. Fig. S1 in the companion paper) and in specific case studies (Fig. S13 therein). Nevertheless, it was also found to slightly anticipate in time and overestimate the easterly diurnal winds in the first hours of the afternoon and to overestimate the nighttime drainage winds (katabatic winds, ventilating the urban atmosphere and reducing pollutant loads). These limitations were mostly attributed to the finite resolution of the model.

20

To disentangle the role of factors (1) and (2) discussed above, we evaluated the model performances in a flatter area of the
domain, where simulations are expected to be less affected by the complex orography of the mountains. We therefore compared PM₁₀ simulations and measurements in Ivrea (Fig. 16b). This test is also useful to operate the CTM model at the boundaries of the domain, with special focus on the emissions from the "boundary conditions". Model underestimation in both the warm and cold seasons is evident. Incidentally, it can be noticed that concentrations simulated without "boundary conditions" are nearly zero, since the considered cell is far from the strongest local sources and most part of the aerosol comes from outside
the regional domain. As done for Aosta (Fig. 17a, b), Fig. 19(a,b) presents the absolute and relative differences between the model and the observations in Ivrea. The overall NMBE is -0.73, which rather well corresponds to the underestimation factor W=4 (or NMBE=-0.75) found for Aosta–Downtown and other sites in the valley. Since it is expected that in this flat area the NWP model is able to better resolve the circulation compared to the mountain valley, this result suggests poor CTM performances to be mostly related to underestimated emissions at the boundaries rather than to incorrect transport. In addition

Ivrea



Figure 19. Average 3-D fields of concentration simulated by FARM extracted at the surface level Absolute (a, b, c) and on a vertical transect (d, e, f). (a) Average of all non-advection days (categorised as "A" by the experimental ALC-classification); relative (b) Average of advection days ("C", "E" differences between simulated and "F" from ALC-classification) using a weighting factor W = 1 for the observed PM₁₀ contribution coming from outside the boundaries of the regional domain; (c) Same as (b) using W = 4 as a weighting factor. (d), (e) and (f) are vertical transects along the main valley (i.e., along the dotted line in figures (a) to (c)) corresponding to the three cases above. The altitude of the three measuring sites shown in the figures is the one from the digital elevation model, which is a smoothed representation of the real topography. The white area in the second row of figures represents concentrations at the surface . The advections investigated in this study come from the east (right side of all figures) [yrea.]

to this general underestimation, a large scatter between observations and simulations can be noticed in Fig. 16, the linear correlation index between simulations and observation in Ivrea being rather low (ρ =0.54). If such a large scatter, due to the erroneous boundary conditions, is already detectable at the border of the domain, it is likely that at least part of the RMSE reported in Table 3 for Aosta–Downtown can be attributed to inaccuracies in the national inventory. As already mentioned, an additional contribution to the observed RMSE could still be given by errors in modelling transport due to the coarse resolution

of the NWP model, although we are not able to quantify the relative role of the two factors. To unravel this issue, high-resolution models (grid step 0.5 km, e.g. Golzio and Pelfini, 2018; Golzio et al., 2019) are being tested on the investigated area and their results will be addressed in future studies.

5

As a last step, we use the model to extend our observational findings to a wider domain, taking at the same time into account Finally, within the limits of the model performances discussed above, we use FARM to extend our observational findings to a wider domain compared to the few measuring sites. In Fig. 20, we provide some summarising plots of the 1-year simulations. These In particular, these show the 3-D simulated fields of PM_{10} concentrations in terms of both surface-level (a, b and c) and vertical transects along the main valley (d, e and f), averaged over all non-advection and advection days in 2017 (as determined by the ALC-classification). 2017. In this latter case, simulations with W = 1 (b,e) and W = 4 (c,f) are included.



Figure 20. Average 3-D fields of PM₁₀ concentration simulated by FARM extracted at the surface level (a, b, c) and on a vertical transect (d, e, f). (a) Average of all non-advection days (categorised as "A" by the experimental ALC-classification); (b) Average of advection days ("C", "E" and "F" from ALC-classification) using a weighting factor W = 1 for the PM₁₀ contribution coming from outside the boundaries of the regional domain; (c) Same as (b) using W = 4 as a weighting factor. (d), (e) and (f) are vertical transects along the main valley (i.e., along the dotted line in figures (a) to (c)) corresponding to the three cases above. The altitude of the three measuring sites shown in the figures is the one from the digital elevation model, which is a smoothed representation of the real topography. The white area in the second row of figures represents the surface. The advections investigated in this study come from the east (right side of all figures).

During non-advection cases the aerosol is supposed to be predominantly of local origin and major hotspots are the city of Aosta and the road along the main valley (Fig. 20a). Compared to this Compared to the "background condition" (ALC type A), the advection cases show higher PM_{10} concentrations and a different geographical distribution, with an increase while proceeding-increasing towards the entrance of the valley (Fig. 20b, e). Obviously, the absolute PM_{10} loads are higher when

5

the non-local contribution is multiplied by W = 4 (Fig. 20c, f), which, as discussed above, is the W value allowing providing a better matching with the PM_{10} concentrations actually measured in Aosta and Donnas Ivrea. Vertical profiles of simulated concentrations are also evidently impacted by the advections (e.g., Fig. 20d-f). A more complete set of maps is provided in Fig. \$11\$12, in which contribution of particulate produced inside the regional domain (local sources) and outside the domain (boundary conditions) have has been partitioned. Interesting feature in that view Fig. 20 is that the average maps of local PM_{10} do not change between advection/non-advection cases, while the (relative) concentration maps of aerosol coming from outside the domain show noticeable differences, thus confirming once again that the polluted air masses detected by the ALC in Aosta–Saint Christophe, on the base of which the classification is built, Aosta–Saint-Christophe are of non-local origin. In conclusion, the excellent correspondence between the model output and the experimental classification based on the ALC, i.e.

5 the remarkable difference of the concentrations and their vertical distribution between days of advection and non-advection, highlights both the <u>rather</u> good performances of the CTM model and the ability of the measurement dataset used to disclose the phenomenon.

5 Conclusions and perspectives

In this work we used long-term (2015–2017) multi-sensor observational datasets (Table 1) coupled to modelling tools to 10 describe pollution aerosol transport from the Po basin to the northwestern Alps. The deployment Key to this study was the deployment in Aosta of an Automated LiDAR Ceilometer (ALC) with its operational (24/7) capabilities in Aosta allowed. This allowed to: a) to collect and present the first-long term dataset of aerosol vertical distribution in the northwestern Alps; b) to provide observational evidence of the complex structure and dynamics of the atmosphere in the Alpine valleys; c) to stimulate the investigation of the phenomenon on a 4-D scale with complementing observational and modelling tools. The

15 analysis over the long-term allowed to complement the investigation of specific case studies provided in the companion paper (Diémoz et al., 2019) and answer some key scientific questions (as listed in the Introduction):

- 1. Driving meteorological factors and frequency of the aerosol pollution transport events. The newly-developed classification based on the ALC profiles (928 classified days, Fig. 3) permitted us to aggregate the days with similar aerosol vertical structures and examine the average properties of each group. Notably, we could understand the link between the wind flow regimes and the aerosol transport. The frequency of the elevated layers was observed to be on average 43–50% of the days, depending on the season (i.e. slightly less than 60% of the days falling in an ALC class different from "Other" in winter and spring to more than 70% in summer) and is clearly connected to eastern wind flows, such as plain-to-mountain, thermally-driven winds (blowing nearly 70% of the days in summer, Fig. 4). This was even clearer using Trajectory Statistical Models (Fig. 13) and contingency tables (Fig. 5) showing the clear connections between the arrival of an elevated aerosol layer over the northwestern Alps and the wind provenancedirection. Such layers were observed 93% of the days with easterly winds (Fig. 5a), with increasing probability of occurrence for increasing air masses residence time over the Po plain (aerosol layer detected >on over 80% of days when the trajectory residence times in the Po basin exceed 1 h, and up to 100% for residence times of days with air masses residence time >10 h, Fig. 5b);
- 2. Advected aerosol physical and chemical properties. The general spatio-temporal characteristics of the aerosol layers were statistically assessed based on the multi-year ALC series. The top altitudes of the layer ranges from 500 m to nearly 4000 m a.g.l., depending on season and according to the convective boundary layer height, and the . Notably, the high altitudes reached by the aerosol layer in summer are compatible with transport and deposition of other contaminants, such

20

as pesticides (Rizzi et al., 2019) and micro-plastics (Allen et al., 2019), on glaciers, snow fields and remote mountain catchments. Also, a dataset of aerosol layer heights may be valuable for follow-up studies on the radiative forcing of particulate matter in connection to the elevation-dependent warming in some mountain regions (Pepin et al., 2015).

5

The duration of the phenomenon ranges from few hours to several days in cases of atmospheric stability. The mean optical and microphysical properties of the aerosol layers were further sounded using a sun photometer. This showed that the columnar aerosol properties are all impacted by the pollution transport: aerosol optical depth (AOD at 500 nm) increases from 0.04 on non-event days up to 0.18 on event days, on average; relevant values of Ångström exponent (400-870 nm) and single scattering albedo (SSA at 500 nm) change from 1.0 to 1.6 and from 0.92 to 0.98, respectively, and depend on the duration/severity of the advection (Fig. 15). This indicates that the transported particles are on average smaller and more light-scattering than the locally-produced ones and agrees well with the results of measurements and analyses of aerosol properties at the surface. Positive matrix factorisation (PMF) of chemical properties at surface level allowed to disclose that particles advected from the Po Valley to the northwestern Alps are mostly of secondary origin (e.g., Fig. 12) and mainly composed by nitrates, sulfates and ammonium. Interestingly, we also found an organic component in the warm season, which confirms the Po Valley as an OM source. Moreover, particle size distributions were observed to showed increase in the fine fraction $(< 1 \ \mu m)$ fraction during event during Po Valley advection days. Correlation of chemical-PMF with independent PMF performed on aerosol size distribution also provided evidence of a clear link between chemical properties and aerosol size (correlations indexes correlation indices $\rho > 0.8$, Fig. 14), which proves that different aerosol formation mechanisms act in the atmosphere in different seasons. In particular, we found some evidence of aqueous processing of particles in winter (e.g., fog and/or cloud processing) through identification of a PMF "droplet mode" in the winter results:

3. Aerosol transport impact on air quality. At the surface levelbottom of the yalley, the advections were demonstrated to alter both the absolute concentrations of PM₁₀ (these being up to 3.5 times higher during event days compared to non-event days, Fig. 8), and their daily cycles, with hourly variations up to 30 µg m⁻³ (Fig. 10). PMF statistics on chemical data identified 5 to 7 different sources, depending on the available chemical characterisation, two of which (nitrate-rich mode in winter and sulfate-rich mode in summer) were attributed to the arrival of particles from outside the Aosta Valley as also confirmed by Trajectory Statistical Models (Fig. 13). Using their relevant scores as a proxy, we estimate an average 25% contribution of these transported nitrate and sulfate-rich components to the Aosta PM₁₀₇ reaching values. This impact varies on a seasonal basis. The relative contribution of non-local PM₁₀ is highest in summer (32%), when advection is most frequent and local PM₁₀ is lowest, while it is lowest in winter (16%), when advection is least frequent and local PM₁₀ is highest. In absolute terms the reverse occurs and the impact of transport is found to be highest in winter/autumn, reaching levels of 50 µg m⁻³ in some episodes and, more generally, favouring exceedances in proximity of the EU legislative limits. This (thus exceeding alone the EU legislative limits). This occurs due to the superposition of advected particles with higher background concentrations in the coldest part of the year, when emissions are highest and pollution is further enhanced by adverse weather conditions, i.e. low-level inversions.

20

10

locally worsened by the valley topography. Note that, although, in this study the impact of Po Valley advection on air quality was mostly quantified for the Aosta-Downtown station. In rural and remote sites of the region this non-local relative contribution is expected to be even much largerin rural and remote sites of the region larger, in relative terms. Increasing the number of stations collecting samples for chemical analyses would be an important follow-up to estimate the non-local contribution to PM_{10} in non-urban sites. The good correlation found between the PM_{10} concentration anomalies in the urban site of Aosta–Downtown and in the regional site of Donnas ($\rho > 0.7$, Fig. 9) is however a clear indication that aerosol loads all over the region are mainly influenced by trans-regional transport rather than by local emissions. It is also worth mentioning that the aerosol particle number (here only measured in the 0.18–18 um range. i.e., missing the finest particles) increased remarkably (up to >3000 particles cm⁻³) in some transport episodes, with possible implications on human health. Some preliminary results (Sect. S5) also indicate that these this regular air mass transport is also able to alter the oxidative capacity of the atmosphere (NO_x/NO ratio), although further investigation is needed to better quantify it.

In general, for both air quality and climate evaluations the identification of monitoring sites (and time periods) representative of "background conditions" is a critical issue to differentiate local and regional pollution levels (e.g., Yuan et al., 2018).

- 15 A global network of atmospheric "baseline" monitoring stations is for example maintained within the World Meteorological Organisation's (WMO) Global Atmosphere Watch (GAW) program (WMO, 2017), with the intention of providing regionally representative measurements relatively free of significant local pollution sources. Our observed (PM₁₀) daily cycle (Figs. 2, 6 and 10) indicates that caution should be paid when assuming mountain sites to be representative of background conditions. In particular, we show that the influence of nearby pollution sources in high altitude sites could be not only limited to the central hours of the day (when convection-driven growth of the nearby plain mixing layer is 20 known to uplift pollutants), but rather extend to night-time measurements via the mountain-valley circulation and particle growth mechanisms addressed in this study:
 - 4. Ability to predict the arrival and impacts of polluted air mass transport. Experimental data and their coupling with modelling tools well demonstrated the Po plain origin of the polluted layers and their increased probability of detection as a function of the air masses residence time over the origin region. Current chemical transport models, however, were found to reproduce well the phenomenon in qualitative terms, but to generally underestimate manifest both systematic underestimations of the absolute advected PM_{10} (biases) and large scatter to the measurements. Based on a-two 1-year long simulated datasetdatasets in Aosta-Downtown and Ivrea, we tested how the air quality forecasts could be improved by updating the emission inventories and using the ALC to constrain the modelled emissions. After the tested measurements-based tuning of some sensitivity tests, we tuned the national inventory to better match the measurements ("boundary conditions" increased by a factor of 4). In this case, the model underestimation was reduced from biases $< -40 \ \mu g \ m^{-3}$ to $-20 \ \mu g \ m^{-3}$ in the worst cases, with mean average biases $< 2 \ \mu g \ m^{-3}$ (Table 3), thus enhancing, on average, our ability to correctly predict the PM concentration and their relevant impacts (Fig. 20). Still, further improvements are necessary to enhance the model ability to better reproduce aerosol processes

5

10

25

in aqueous-phase chemistry (e.g., Gong et al., 2011)(e.g., aqueous-phase chemistry, Gong et al., 2011) and wind fields using higher-resolution NWP models.

In conclusion, we demonstrated that, despite considered a pristine environment, the northwestern Alps are regularly affected by a sort of "polluted aerosol tide", i.e. by a wind-driven transport of polluted aerosol layers from the Po plain. Overall, this calls for air pollution mitigation policies acting, at least, over the regional scale in this fragile environment.

Data availability. The ALC data are available upon request from the Alice-net (alicenet@isac.cnr.it) and E-PROFILE (http://data.ceda.ac.uk/ badc/eprofile/data/) networks. The sun photometer data can be downloaded from the EuroSkyRad network web site (http://www.euroskyrad. net/index.html) after authentication (credentials may be requested to M. Campanelli, m.campanelli@isac.cnr.it). The measurements from ARPA Valle d'Aosta air quality surface network are available at the web page http://www.arpa.vda.it/it/aria/la-qualit%C3%A0-dell-aria/ stazioni-di-monitoraggio/inquinanti-export-dati. The PM₁₀ measurements in Ivrea are available on the Piedmont regional government web

page http://www.regione.piemonte.it/ambiente/aria/rilev/ariaday/ariaweb-new/index.php/it/home. The weather data from the Aosta–Saint-Christophe and Saint-Denis stations can be retrieved from http://cf.regione.vda.it/richiesta_dati.php upon request to Centro Funzionale della Valle d'Aosta. The rest of the data can be asked to the corresponding author (h.diemoz@arpa.vda.it).

Author contributions. HD, FB and GPG conceived and designed the study, interpreted the results and wrote the paper. HD analysed the

15 data. TM supplied the meteorological observations and numerical weather predictions. GP performed the chemical transport simulations. SP carried out the EC/OC analyses and IKFT helped with the interpretation of the chemical speciation. MC supplied the POM calibration factors.

Competing interests. The authors declare that they have no conflict of interest.

5

10

Acknowledgements. The authors would like to thank: A. Brunier, G. Lupato, P. Proment, S. Vaccari, and M.C. Gibellino (ARPA Valle
d'Aosta) for carrying out the chemical analyses; M. Pignet, C. Tarricone, and M. Zublena (ARPA Valle d'Aosta) for providing the data from the air quality surface network; P. Lazzeri (APPA Trento) for helping with the PMF analysis. The authors would like to acknowledge the valuable contribution of the discussions in the working group meetings organised by COST Action ES1303 (TOPROF). They also gratefully acknowledge the Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland for the provision of the LAGRANTO software used in this publication, and ARPA Piemonte and Regione Piemonte for the PM₁₀ measurements at the Iyrea station.

References

- Agnesod, G., Moulin, P.-A., Pession, G., Villard, H., and Zublena, M.: Étude Air Espace Mont-Blanc Rapport Technique, Tech. rep., Espace Mont Blanc, http://www.arpa.vda.it/images/stories/ARPA/aria/progetti/progairmb_agnesod_2003.pdf, 2003.
- Allen, S., Allen, D., Phoenix, V. R., Le Roux, G., Durántez Jiménez, P., Simonneau, A., Binet, S., and Galop, D.: Atmospheric transport and
 deposition of microplastics in a remote mountain catchment, Nature Geoscience, 12, 339–344, https://doi.org/10.1038/s41561-019-03355, 2019.
 - Arvani, B., Bradley Pierce, R., Lyapustin, A. I., Wang, Y., Ghermandi, G., and Teggi, S.: Seasonal monitoring and estimation of regional aerosol distribution over Po valley, northern Italy, using a high-resolution MAIAC product, Atmos. Environ., 141, 106 – 121, https://doi.org/10.1016/j.atmosenv.2016.06.037, 2016.
- 10 Ashbaugh, L. L., Malm, W. C., and Sadeh, W. Z.: A residence time probability analysis of sulfur concentrations at grand Canyon National Park, Atmos. Environ., 19, 1263 – 1270, https://doi.org/10.1016/0004-6981(85)90256-2, 1985.
 - Baldauf, M., Seifert, A., Förstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational Convective-Scale Numerical Weather Prediction with the COSMO Model: Description and Sensitivities, Mon. Weather Rev., 139, 3887–3905, https://doi.org/10.1175/MWR-D-10-05013.1, 2011.
- 15 Barnaba, F. and Gobbi, G. P.: Aerosol seasonal variability over the Mediterranean region and relative impact of maritime, continental and Saharan dust particles over the basin from MODIS data in the year 2001, Atmos. Chem. Phys., 4, 2367–2391, https://doi.org/10.5194/acp-4-2367-2004, 2004.
 - Barnaba, F., Gobbi, G. P., and de Leeuw, G.: Aerosol stratification, optical properties and radiative forcing in Venice (Italy) during ADRIEX, Q. J. Roy. Meteorol. Soc., 133, 47–60, https://doi.org/10.1002/qj.91, 2007.
- 20 Barnaba, F., Putaud, J. P., Gruening, C., dell'Acqua, A., and Dos Santos, S.: Annual cycle in co-located in situ, total-column, and heightresolved aerosol observations in the Po Valley (Italy): Implications for ground-level particulate matter mass concentration estimation from remote sensing, J. Geophys. Res., 115, https://doi.org/10.1029/2009JD013002, 2010.
 - Barnaba, F., Bolignano, A., Di Liberto, L., Morelli, M., Lucarelli, F., Nava, S., Perrino, C., Canepari, S., Basart, S., Costabile, F., Dionisi, D., Ciampichetti, S., Sozzi, R., and Gobbi, G. P.: Desert dust contribution to PM10 loads in Italy: Methods and recommendations ad-
- 25 dressing the relevant European Commission Guidelines in support to the Air Quality Directive 2008/50, Atmos. Environ., 161, 288 305, https://doi.org/10.1016/j.atmosenv.2017.04.038, 2017.
 - Belis, C., Blond, N., Bouland, C., Carnevale, C., Clappier, A., Douros, J., Fragkou, E., Guariso, G., Miranda, A. I., Nahorski, Z., Pisoni, E., Ponche, J.-L., Thunis, P., Viaene, P., and Volta, M.: Strengths and Weaknesses of the Current EU Situation, pp. 69–83, Springer International Publishing, https://doi.org/10.1007/978-3-319-33349-6_4, 2017.
- 30 Bigi, A. and Ghermandi, G.: Long-term trend and variability of atmospheric PM₁₀ concentration in the Po Valley, Atmos. Chem. and Phys., 14, 4895–4907, https://doi.org/10.5194/acp-14-4895-2014, 2014.
 - Bigi, A. and Ghermandi, G.: Trends and variability of atmospheric PM_{2.5} and PM_{10-2.5} concentration in the Po Valley, Italy, Atmos. Chem. Phys., 16, 15777–15788, https://doi.org/10.5194/acp-16-15777-2016, 2016.
- Binkowski, F.: Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, vol. EPA/600/R 99/030, chap. The aerosol portion of Models-3 CMAO., DW Byun, and JKS Ching (Eds), 1999.
- Birch, M. E. and Cary, R. A.: Elemental Carbon-Based Method for Monitoring Occupational Exposures to Particulate Diesel Exhaust, Aerosol Sci. Tech., 25, 221–241, https://doi.org/10.1080/02786829608965393, 1996.

- Blyth, S.: Mountain watch: environmental change & sustainable developmental in mountains, United Nations Environment Programme, 2002.
- Bonvalot, L., Tuna, T., Fagault, Y., Jaffrezo, J.-L., Jacob, V., Chevrier, F., and Bard, E.: Estimating contributions from biomass burning, fossil fuel combustion, and biogenic carbon to carbonaceous aerosols in the Valley of Chamonix: a dual approach based on radiocarbon and levoglucosan, Atmos. Chem. Phys., 16, 13753–13772, https://doi.org/10.5194/acp-16-13753-2016, 2016.
- Bourgeois, I., Savarino, J., Caillon, N., Angot, H., Barbero, A., Delbart, F., Voisin, D., and Clément, J.-C.: Tracing the Fate of Atmospheric Nitrate in a Subalpine Watershed Using Δ17O, Environ. Sci. Tech., 52, 5561–5570, https://doi.org/10.1021/acs.est.7b02395, 2018.
 - Bressi, M., Sciare, J., Ghersi, V., Mihalopoulos, N., Petit, J.-E., Nicolas, J. B., Moukhtar, S., Rosso, A., Féron, A., Bonnaire, N., Poulakis, E., and Theodosi, C.: Sources and geographical origins of fine aerosols in Paris (France), Atmos. Chem. Phys., 14, 8813–8839,
- 10 https://doi.org/10.5194/acp-14-8813-2014, 2014.

- Bressi, M., Cavalli, F., Belis, C. A., Putaud, J.-P., Fröhlich, R., Martins dos Santos, S., Petralia, E., Prévôt, A. S. H., Berico, M., Malaguti, A., and Canonaco, F.: Variations in the chemical composition of the submicron aerosol and in the sources of the organic fraction at a regional background site of the Po Valley (Italy), Atmos. Chem. Phys., 16, 12875–12896, https://doi.org/10.5194/acp-16-12875-2016, 2016.
- Brulfert, G., Chemel, C., Chaxel, E., Chollet, J.-P., Jouve, B., and Villard, H.: Assessment of 2010 air quality in two Alpine valleys from modelling: Weather type and emission scenarios, Atmos. Environ., 40, 7893 7907, https://doi.org/10.1016/j.atmosenv.2006.07.021, 2006.
 - Bucci, S., Cristofanelli, P., Decesari, S., Marinoni, A., Sandrini, S., Größ, J., Wiedensohler, A., Di Marco, C. F., Nemitz, E., Cairo, F., Di Liberto, L., and Fierli, F.: Vertical distribution of aerosol optical properties in the Po Valley during the 2012 summer campaigns, Atmos. Chem. Phys., 18, 5371–5389, https://doi.org/10.5194/acp-18-5371-2018, 2018.
- 20 Burkhardt, J., Zinsmeister, D., Grantz, D. A., Vidic, S., Sutton, M. A., Hunsche, M., and Pariyar, S.: Camouflaged as degraded wax: hygroscopic aerosols contribute to leaf desiccation, tree mortality, and forest decline, Environ. Res. Lett., 13, 085001, https://doi.org/10.1088/1748-9326/aad346, 2018.
 - Calori, G., Silibello, C., and Marras, G.: FARM (Flexible Air quality Regional Model) Model formulation and user's Manual, Arianet, 4.7 edn., 2014.
- 25 Campana, M., Li, Y., Staehelin, J., Prevot, A. S., Bonasoni, P., Loetscher, H., and Peter, T.: The influence of south foehn on the ozone mixing ratios at the high alpine site Arosa, Atmos. Environ., 39, 2945 – 2955, https://doi.org/10.1016/j.atmosenv.2005.01.037, 2005.
 - Campanelli, M., Estellés, V., Tomasi, C., Nakajima, T., Malvestuto, V., and Martínez-Lozano, J. A.: Application of the SKYRAD Improved Langley plot method for the in situ calibration of CIMEL Sun-sky photometers, Appl. Opt., 46, 2688–2702, https://doi.org/10.1364/AO.46.002688, 2007.
- 30 Campanelli, M., Mascitelli, A., Sanò, P., Diémoz, H., Estellés, V., Federico, S., Iannarelli, A. M., Fratarcangeli, F., Mazzoni, A., Realini, E., Crespi, M., Bock, O., Martínez-Lozano, J. A., and Dietrich, S.: Precipitable water vapour content from ESR/SKYNET sun-sky radiometers: validation against GNSS/GPS and AERONET over three different sites in Europe, Atmos. Meas. Tech., 11, 81–94, https://doi.org/10.5194/amt-11-81-2018, 2018.
- Carbone, C., Decesari, S., Mircea, M., Giulianelli, L., Finessi, E., Rinaldi, M., Fuzzi, S., Marinoni, A., Duchi, R., Perrino, C., Sargolini, T.,
- 35 Vardè, M., Sprovieri, F., Gobbi, G., Angelini, F., and Facchini, M.: Size-resolved aerosol chemical composition over the Italian Peninsula during typical summer and winter conditions, Atmos. Environ., 44, 5269 – 5278, https://doi.org/10.1016/j.atmosenv.2010.08.008, 2010.

- Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth system, Atmos. Chem. Phys., 10, 1701–1737, https://doi.org/10.5194/acp-10-1701-2010, 2010.
- Carter, W.: Documentation of the SAPRC-99 Chemical Mechanism for VOC Reactivity Assessment. Final Report to California Air Resources
- 5 Board, Contract 92-329 and 95-308, SAPRC, Tech. rep., University of California, Riverside, CA, http://www.engr.ucr.edu/~carter/pubs/ s99doc.pdf, 2000.
 - Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, Atmos. Meas. Tech., 3, 79–89, https://doi.org/10.5194/amt-3-79-2010, 2010.

Cesaroni, G., Badaloni, C., Gariazzo, C., Stafoggia, M., Sozzi, R., Davoli, M., and Forastiere, F.: Long-term exposure to urban air pollution
 and mortality in a cohort of more than a million adults in Rome, Environ. Health Persp., 121, 324, https://doi.org/10.1289/ehp.1205862,

2013.

- Charron, A., Harrison, R. M., Moorcroft, S., and Booker, J.: Quantitative interpretation of divergence between PM10 and PM2.5 mass measurement by TEOM and gravimetric (Partisol) instruments, Atmos. Environ., 38, 415 423, https://doi.org/10.1016/j.atmosenv.2003.09.072, 2004.
- 15 Chazette, P., Couvert, P., Randriamiarisoa, H., Sanak, J., Bonsang, B., Moral, P., Berthier, S., Salanave, S., and Toussaint, F.: Three-dimensional survey of pollution during winter in French Alps valleys, Atmos. Environ., 39, 1035 – 1047, https://doi.org/10.1016/j.atmosenv.2004.10.014, 2005.
 - Chemel, C., Arduini, G., Staquet, C., Largeron, Y., Legain, D., Tzanos, D., and Paci, A.: Valley heat deficit as a bulk measure of wintertime particulate air pollution in the Arve River Valley, Atmos. Environ., 128, 208 215, https://doi.org/10.1016/j.atmosenv.2015.12.058, 2016.
- 20 Chu, D. A., Kaufman, Y. J., Zibordi, G., Chern, J. D., Mao, J., Li, C., and Holben, B. N.: Global monitoring of air pollution over land from the Earth Observing System-Terra Moderate Resolution Imaging Spectroradiometer (MODIS), J. Geophys. Res., 108, https://doi.org/10.1029/2002JD003179, 2003.
 - Clerici, M. and Mélin, F.: Aerosol direct radiative effect in the Po Valley region derived from AERONET measurements, Atmos. Chem. Phys., 8, 4925–4946, https://doi.org/10.5194/acp-8-4925-2008, 2008.
- 25 Cong, Z., Kawamura, K., Kang, S., and Fu, P.: Penetration of biomass-burning emissions from South Asia through the Himalayas: new insights from atmospheric organic acids, Scientific reports, 5, https://doi.org/10.1038/srep09580, 2015.
 - Costabile, F., Gilardoni, S., Barnaba, F., Di Ianni, A., Di Liberto, L., Dionisi, D., Manigrasso, M., Paglione, M., Poluzzi, V., Rinaldi, M., Facchini, M. C., and Gobbi, G. P.: Characteristics of brown carbon in the urban Po Valley atmosphere, Atmospheric Chemistry and Physics, 17, 313–326, https://doi.org/10.5194/acp-17-313-2017, https://www.atmos-chem-phys.net/17/313/2017/, 2017.
- 30 Cugerone, K., De Michele, C., Ghezzi, A., Gianelle, V., and Gilardoni, S.: On the functional form of particle number size distributions: influence of particle source and meteorological variables, Atmos. Chem. Phys., 18, 4831–4842, https://doi.org/10.5194/acp-18-4831-2018, 2018.
 - Curci, G., Ferrero, L., Tuccella, P., Barnaba, F., Angelini, F., Bolzacchini, E., Carbone, C., Denier van der Gon, H. A. C., Facchini, M. C., Gobbi, G. P., Kuenen, J. P. P., Landi, T. C., Perrino, C., Perrone, M. G., Sangiorgi, G., and Stocchi, P.: How much is particulate matter
- 35 near the ground influenced by upper-level processes within and above the PBL? A summertime case study in Milan (Italy) evidences the distinctive role of nitrate, Atmos. Chem. Phys., 15, 2629–2649, https://doi.org/10.5194/acp-15-2629-2015, 2015.
 - de Freitas, C. R.: Tourism climatology: evaluating environmental information for decision making and business planning in the recreation and tourism sector, Int. J. Biometeorol., 48, 45–54, https://doi.org/10.1007/s00484-003-0177-z, 2003.

- De Wekker, S. F. J. and Kossmann, M.: Convective Boundary Layer Heights Over Mountainous Terrain—A Review of Concepts, Front. Earth Sci., 3, 77, https://doi.org/10.3389/feart.2015.00077, 2015.
- Decesari, S., Allan, J., Plass-Duelmer, C., Williams, B. J., Paglione, M., Facchini, M. C., O'Dowd, C., Harrison, R. M., Gietl, J. K., Coe, H., Giulianelli, L., Gobbi, G. P., Lanconelli, C., Carbone, C., Worsnop, D., Lambe, A. T., Ahern, A. T., Moretti, F., Tagliavini, E., Elste,
- 5 T., Gilge, S., Zhang, Y., and Dall'Osto, M.: Measurements of the aerosol chemical composition and mixing state in the Po Valley using multiple spectroscopic techniques, Atmos. Chem. Phys., 14, 12 109–12 132, https://doi.org/10.5194/acp-14-12109-2014, 2014.
 - Dhungel, S., Kathayat, B., Mahata, K., and Panday, A.: Transport of regional pollutants through a remote trans-Himalayan valley in Nepal, Atmos. Chem. Phys., 18, 1203–1216, https://doi.org/10.5194/acp-18-1203-2018, 2018.
 - Diémoz, H., Siani, A. M., Casale, G. R., di Sarra, A., Serpillo, B., Petkov, B., Scaglione, S., Bonino, A., Facta, S., Fedele, F., Grifoni,
- 10 D., Verdi, L., and Zipoli, G.: First national intercomparison of solar ultraviolet radiometers in Italy, Atmos. Meas. Tech., 4, 1689–1703, https://doi.org/10.5194/amt-4-1689-2011, 2011.
 - Diémoz, H., Campanelli, M., and Estellés, V.: One Year of Measurements with a POM-02 Sky Radiometer at an Alpine EuroSkyRad Station, J. Meteorol. Soc. Jpn., 92A, 1–16, https://doi.org/10.2151/jmsj.2014-A01, 2014a.
 - Diémoz, H., Siani, A. M., Redondas, A., Savastiouk, V., McElroy, C. T., Navarro-Comas, M., and Hase, F.: Improved retrieval of
- 15 nitrogen dioxide (NO₂) column densities by means of MKIV Brewer spectrophotometers, Atmos. Meas. Tech., 7, 4009–4022, https://doi.org/10.5194/amt-7-4009-2014, 2014b.
 - Diémoz, H., Barnaba, F., Magri, T., Pession, G., Dionisi, D., Pittavino, S., Tombolato, I. K. F., Campanelli, M., Della Ceca, L. S., Hervo, M., Di Liberto, L., Ferrero, L., and Gobbi, G. P.: Transport of Po Valley aerosol pollution to the northwestern Alps – Part 1: Phenomenology, Atmos. Chem. Phys., 19, 3065–3095, https://doi.org/10.5194/acp-19-3065-2019, 2019.
- 20 Dionisi, D., Barnaba, F., Diémoz, H., Di Liberto, L., and Gobbi, G. P.: A multiwavelength numerical model in support of quantitative retrievals of aerosol properties from automated lidar ceilometers and test applications for AOT and PM₁₀ estimation, Atmos. Meas. Tech., 11, 6013–6042, https://doi.org/10.5194/amt-11-6013-2018, 2018.
 - Dosio, A., Galmarini, S., and Graziani, G.: Simulation of the circulation and related photochemical ozone dispersion in the Po plains (northern Italy): Comparison with the observations of a measuring campaign, J. Geophys. Res., 107, LOP 2–1–LOP 2–24,
- 25 https://doi.org/10.1029/2000JD000046, 2002.
 - EEA: Air Quality in Europe 2015 Report, Tech. rep., https://doi.org/10.2800/62459, 2015.

EEA: Air Quality in Europe - 2017 Report, Tech. rep., https://doi.org/10.2800/850018, 2017.

30 0493(2000)128<1106:DWITHK>2.0.CO;2, 2000.

- EMEP: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, Tech. rep., MSC-W, CCC and CEIP, 2016.
- EU Commission: Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe, Official Journal of the European Union, pp. L152/1–44, 2008.
- 35 EU Commission: European Commission Press release: Air quality: Commission takes action to protect citizens from air pollution, Brussels, 17 May 2018, Press Release Database, http://europa.eu/rapid/press-release_IP-18-3450_en.htm, 2018.
 - Fernald, F. G.: Analysis of atmospheric lidar observations: some comments, Appl. Opt., 23, 652–653, https://doi.org/10.1364/AO.23.000652, 1984.

Egger, J., Bajrachaya, S., Egger, U., Heinrich, R., Reuder, J., Shayka, P., Wendt, H., and Wirth, V.: Diurnal Winds in the Himalayan Kali Gandaki Valley. Part I: Observations, Mon. Weather Rev., 128, 1106–1122, https://doi.org/10.1175/1520-0402(2000)128 <1106.DWITHK> 2.0 CO:2. 2000

- Ferrero, L., Perrone, M. G., Petraccone, S., Sangiorgi, G., Ferrini, B. S., Lo Porto, C., Lazzati, Z., Cocchi, D., Bruno, F., Greco, F., Riccio, A., and Bolzacchini, E.: Vertically-resolved particle size distribution within and above the mixing layer over the Milan metropolitan area, Atmos. Chem. Phys., 10, 3915–3932, https://doi.org/10.5194/acp-10-3915-2010, 2010.
- Ferrero, L., Castelli, M., Ferrini, B. S., Moscatelli, M., Perrone, M. G., Sangiorgi, G., D'Angelo, L., Rovelli, G., Moroni, B., Scardazza, F.,
 Močnik, G., Bolzacchini, E., Petitta, M., and Cappelletti, D.: Impact of black carbon aerosol over Italian basin valleys: high-resolution measurements along vertical profiles, radiative forcing and heating rate, Atmos. Chem. Phys., 14, 9641–9664, https://doi.org/10.5194/acp-

14-9641-2014, 2014.

Finardi, S., Silibello, C., D'Allura, A., and Radice, P.: Analysis of pollutants exchange between the Po Valley and the surrounding European region, Urban Clim., 10, 682 – 702, https://doi.org/10.1016/j.uclim.2014.02.002, source apportionment and modelling of urban air

10 pollution, 2014.

20

- Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs, Atmos. Chem. Phys., 15, 8217–8299, https://doi.org/10.5194/acp-15-8217-2015, 2015.
- 15 Gariazzo, C., Silibello, C., Finardi, S., Radice, P., Piersanti, A., Calori, G., Cecinato, A., Perrino, C., Nussio, F., Cagnoli, M., et al.: A gas/aerosol air pollutants study over the urban area of Rome using a comprehensive chemical transport model, Atmos. Environ., 41, 7286–7303, https://doi.org/10.1016/j.atmosenv.2007.05.018, 2007.

Gilardoni, S., Vignati, E., Cavalli, F., Putaud, J. P., Larsen, B. R., Karl, M., Stenström, K., Genberg, J., Henne, S., and Dentener, F.: Better constraints on sources of carbonaceous aerosols using a combined ¹⁴C – macro tracer analysis in a European rural background site, Atmos. Chem. Phys., 11, 5685–5700, https://doi.org/10.5194/acp-11-5685-2011, 2011.

Gilardoni, S., Massoli, P., Giulianelli, L., Rinaldi, M., Paglione, M., Pollini, F., Lanconelli, C., Poluzzi, V., Carbone, S., Hillamo, R., Russell, L. M., Facchini, M. C., and Fuzzi, S.: Fog scavenging of organic and inorganic aerosol in the Po Valley, Atmos. Chem. Phys., 14, 6967–6981, https://doi.org/10.5194/acp-14-6967-2014, 2014.

Gilardoni, S., Massoli, P., Paglione, M., Giulianelli, L., Carbone, C., Rinaldi, M., Decesari, S., Sandrini, S., Costabile, F., Gobbi, G. P.,

- 25 Pietrogrande, M. C., Visentin, M., Scotto, F., Fuzzi, S., and Facchini, M. C.: Direct observation of aqueous secondary organic aerosol from biomass-burning emissions, P. Natl. A. Sci., 113, 10013–10018, https://doi.org/10.1073/pnas.1602212113, 2016.
 - Giovannini, L., Antonacci, G., Zardi, D., Laiti, L., and Panziera, L.: Sensitivity of Simulated Wind Speed to Spatial Resolution over Complex Terrain, Energy Proced., 59, 323–329, https://doi.org/10.1016/j.egypro.2014.10.384, 2014.

Giovannini, L., Laiti, L., Serafin, S., and Zardi, D.: The thermally driven diurnal wind system of the Adige Valley in the Italian Alps, Q. J.

- 30 Roy. Meteor. Soc., 143, 2389–2402, https://doi.org/10.1002/qj.3092, 2017.
 - Gohm, A., Harnisch, F., Vergeiner, J., Obleitner, F., Schnitzhofer, R., Hansel, A., Fix, A., Neininger, B., Emeis, S., and Schäfer, K.: Air Pollution Transport in an Alpine Valley: Results From Airborne and Ground-Based Observations, Bound.-Lay. Meteorol., 131, 441–463, https://doi.org/10.1007/s10546-009-9371-9, 2009.

Golzio, A. and Pelfini, M.: High resolution WRF over mountainous complex terrain: testing over the Ortles Cevedale area (Central Italian

35 Alps), in: Conference: First National Congress, Italian Association of Atmospheric Sciences and Meteorology (AISAM), AISAM, 2018. Golzio, A., Ferrarese, S., Cassardo, C., Diolaiuti, G. A., and Pelfini, M.: Resolution improvement in WRF over complex mountainous terrain, in preparation for Boundary-Layer Meteorol., 2019.

- Gong, W., Stroud, C., and Zhang, L.: Cloud Processing of Gases and Aerosols in Air Quality Modeling, Atmosphere, 2, 567–616, https://doi.org/10.3390/atmos2040567, 2011.
- Green, D. C., Fuller, G. W., and Baker, T.: Development and validation of the volatile correction model for PM10 An empirical method for adjusting TEOM measurements for their loss of volatile particulate matter, Atmos. Environ., 43, 2132 – 2141,
- 5 https://doi.org/10.1016/j.atmosenv.2009.01.024, 2009.

Hann, J. v.: Zur Theorie der Berg- und Talwinde, Z. Öst. Meteorol., 14, 444-448, 1879.

- Harrison, R. M., Jones, A. M., Gietl, J., Yin, J., and Green, D. C.: Estimation of the Contributions of Brake Dust, Tire Wear, and Resuspension to Nonexhaust Traffic Particles Derived from Atmospheric Measurements, Environ. Sci. Tech., 46, 6523–6529, https://doi.org/10.1021/es300894r, 2012.
- 10 Hashimoto, M., Nakajima, T., Dubovik, O., Campanelli, M., Che, H., Khatri, P., Takamura, T., and Pandithurai, G.: Development of a new data-processing method for SKYNET sky radiometer observations, Atmos. Meas. Tech., 5, 2723–2737, https://doi.org/10.5194/amt-5-2723-2012, 2012.
 - Hsu, Y.-K., Holsen, T. M., and Hopke, P. K.: Comparison of hybrid receptor models to locate PCB sources in Chicago, Atmos. Environ., 37, 545 562, https://doi.org/10.1016/S1352-2310(02)00886-5, 2003.
- 15 Kabashnikov, V. P., Chaikovsky, Anatoli, P., Kucsera, T. L., and Metelskaya, N. S.: Estimated accuracy of three common trajectory statistical methods, Atmos. Environ., 45, 5425 – 5430, https://doi.org/10.1016/j.atmosenv.2011.07.006, 2011.
 - Kaiser, A.: Origin of polluted air masses in the Alps. An overview and first results for MONARPOP, Environ. Pollut., 157, 3232 3237, https://doi.org/10.1016/j.envpol.2009.05.042, 2009.
 - Kambezidis, H. and Kaskaoutis, D.: Aerosol climatology over four AERONET sites: An overview, Atmos. Environ., 42, 1892 1906,
- 20 https://doi.org/10.1016/j.atmosenv.2007.11.013, 2008.
 - Kastendeuch, P. P. and Kaufmann, P.: Classification of summer wind fields over complex terrain, Int. J. Climatol., 17, 521–534, https://doi.org/10.1002/(SICI)1097-0088(199704)17:5<521::AID-JOC143>3.0.CO;2-Q, 1997.
 - Kazadzis, S., Kouremeti, N., Diémoz, H., Gröbner, J., Forgan, B. W., Campanelli, M., Estellés, V., Lantz, K., Michalsky, J., Carlund, T., Cuevas, E., Toledano, C., Becker, R., Nyeki, S., Kosmopoulos, P. G., Tatsiankou, V., Vuilleumier, L., Denn, F. M., Ohkawara, N., Ijima,
- 25 O., Goloub, P., Raptis, P. I., Milner, M., Behrens, K., Barreto, A., Martucci, G., Hall, E., Wendell, J., Fabbri, B. E., and Wehrli, C.: Results from the Fourth WMO Filter Radiometer Comparison for aerosol optical depth measurements, Atmos. Chem. Phys., 18, 3185–3201, https://doi.org/10.5194/acp-18-3185-2018, 2018.
 - Keiser, D., Lade, G., and Rudik, I.: Air pollution and visitation at U.S. national parks, Sci. Adv., 4, https://doi.org/10.1126/sciadv.aat1613, 2018.
- 30 Khan, M., Masiol, M., Formenton, G., Di Gilio, A., de Gennaro, G., Agostinelli, C., and Pavoni, B.: Carbonaceous PM2.5 and secondary organic aerosol across the Veneto region (NE Italy), Sci. Total Environ., 542, 172 181, https://doi.org/10.1016/j.scitotenv.2015.10.103, 2016.
 - Khatri, P. and Takamura, T.: An Algorithm to Screen Cloud-Affected Data for Sky Radiometer Data Analysis, J. Meteorol. Soc. Jpn., 87, 189–204, https://doi.org/10.2151/jmsj.87.189, 2009.
- 35 Klett, J. D.: Lidar inversion with variable backscatter/extinction ratios, Appl. Opt., 24, 1638–1643, https://doi.org/10.1364/AO.24.001638, 1985.
Kukkonen, J., Sokhi, R., Luhana, L., Härkönen, J., Salmi, T., Sofiev, M., and Karppinen, A.: Evaluation and application of a statistical model for assessment of long-range transported proportion of PM2.5 in the United Kingdom and in Finland, Atmos. Environ., 42, 3980 – 3991, https://doi.org/10.1016/j.atmosenv.2007.02.036, fifth International Conference on Urban Air Quality, 2008.

Landi, T., Curci, G., Carbone, C., Menut, L., Bessagnet, B., Giulianelli, L., Paglione, M., and Facchini, M.: Simulation of size-

- 5 segregated aerosol chemical composition over northern Italy in clear sky and wind calm conditions, Atmos. Res., 125-126, 1 11, https://doi.org/10.1016/j.atmosres.2013.01.009, 2013.
 - Largeron, Y. and Staquet, C.: Persistent inversion dynamics and wintertime PM10 air pollution in Alpine valleys, Atmos. Environ., 135, 92 108, https://doi.org/10.1016/j.atmosenv.2016.03.045, 2016.
 - Larsen, B., Gilardoni, S., Stenström, K., Niedzialek, J., Jimenez, J., and Belis, C.: Sources for PM air pollution in the Po Plain, Italy: II.
- Probabilistic uncertainty characterization and sensitivity analysis of secondary and primary sources, Atmos. Environ., 50, 203 213, https://doi.org/10.1016/j.atmosenv.2011.12.038, 2012.
 - Loomis, D., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Baan, R., Mattock, H., and Straif, K.: The carcinogenicity of outdoor air pollution, Lancet Oncology, 14, 1262, https://doi.org/10.1016/S1470-2045(13)70487-X show, 2013.

Mann, H. B. and Whitney, D. R.: On a Test of Whether one of Two Random Variables is Stochastically Larger than the Other, Ann. Math.

- 15 Stat., 18, 50–60, 1947.
 - Matta, E., Facchini, M. C., Decesari, S., Mircea, M., Cavalli, F., Fuzzi, S., Putaud, J.-P., and Dell'Acqua, A.: Mass closure on the chemical species in size-segregated atmospheric aerosol collected in an urban area of the Po Valley, Italy, Atmos. Chem. Phys., 3, 623–637, https://doi.org/10.5194/acp-3-623-2003, 2003.
- Mazzola, M., Lanconelli, C., Lupi, A., Busetto, M., Vitale, V., and Tomasi, C.: Columnar aerosol optical properties in the Po Valley, Italy,
 from MFRSR data, J. Geophys. Res., 115, https://doi.org/10.1029/2009JD013310, 2010.
 - Mélin, F. and Zibordi, G.: Aerosol variability in the Po Valley analyzed from automated optical measurements, Geophy. Res. Lett., 32, https://doi.org/10.1029/2004GL021787, 2005.
 - Norris, G. and Duvall, R.: EPA Positive Matrix Factorization (PMF) 5.0 Fundamentals and User Guide, U.S. Environmental Protection Agency, https://www.epa.gov/sites/production/files/2015-02/documents/pmf_5.0_user_guide.pdf, 2014.
- 25 Nyeki, S., Eleftheriadis, K., Baltensperger, U., Colbeck, I., Fiebig, M., Fix, A., Kiemle, C., Lazaridis, M., and Petzold, A.: Airborne Lidar and in-situ Aerosol Observations of an Elevated Layer, Leeward of the European Alps and Apennines, Geophys. Res. Lett., 29, 33–1–33–4, https://doi.org/10.1029/2002GL014897, 2002.
 - Paatero, P.: Least squares formulation of robust non-negative factor analysis, Chemometr. Intell. Lab., 37, 23 35, https://doi.org/10.1016/S0169-7439(96)00044-5, 1997.
- 30 Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, Environmetrics, 5, 111–126, https://doi.org/10.1002/env.3170050203, 1994.
 - Patashnick, H. and Rupprecht, E. G.: Continuous PM-10 Measurements Using the Tapered Element Oscillating Microbalance, J. Air Waste Manage., 41, 1079–1083, https://doi.org/10.1080/10473289.1991.10466903, 1991.
 - Pepin, N., Bradley, R. S., Diaz, H. F., Baraer, M., Caceres, E. B., Forsythe, N., Fowler, H., Greenwood, G., Hashmi, M. Z., Liu,
- 35 X. D., Miller, J. R., Ning, L., Ohmura, A., Palazzi, E., Rangwala, I., Schöner, W., Severskiy, I., Shahgedanova, M., Wang, M. B., Williamson, S. N., and Yang, D. Q.: Elevation-dependent warming in mountain regions of the world, Nat. Clim. Change, 5, 424–430, https://doi.org/10.1038/nclimate2563, 2015.

Perrone, M., Larsen, B., Ferrero, L., Sangiorgi, G., De Gennaro, G., Udisti, R., Zangrando, R., Gambaro, A., and Bolzacchini, E.: Sources of high PM2.5 concentrations in Milan, Northern Italy: Molecular marker data and CMB modelling, Sci. Total Environ., 414, 343 – 355, https://doi.org/10.1016/j.scitotenv.2011.11.026, 2012.

Philipona, R.: Greenhouse warming and solar brightening in and around the Alps, Int. J. Climatol., 33, 1530–1537, https://doi.org/10.1002/joc.3531, 2013.

Pletscher, K., Weiss, M., and Moelter, L.: Simultaneous determination of PM fractions, particle number and particle size distribution in high time resolution applying one and the same optical measurement technique, Gefahrst. Reinhalt. L., 76, 425–436, http://www.gefahrstoffe. de/gest/article.php?data[article_id]=86622, 2016.

Putaud, J. P., Van Dingenen, R., and Raes, F.: Submicron aerosol mass balance at urban and semirural sites in the Milan area (Italy), J.
 Geophys. Res., 107, LOP 11–1–LOP 11–10, https://doi.org/10.1029/2000JD000111, 2002.

- Putaud, J.-P., Dingenen, R. V., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R., Hansson, H., Harrison, R., Herrmann, H., Hitzenberger, R., Hüglin, C., Jones, A., Kasper-Giebl, A., Kiss, G., Kousa, A., Kuhlbusch, T., Löschau, G., Maenhaut, W., Molnar, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A., and Raes, F.: A European aerosol phenomenology – 3:
- 15 Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, Atmos. Environ., 44, 1308 – 1320, https://doi.org/10.1016/j.atmosenv.2009.12.011, 2010.
 - Putaud, J. P., Cavalli, F., Martins dos Santos, S., and Dell'Acqua, A.: Long-term trends in aerosol optical characteristics in the Po Valley, Italy, Atmos. Chem. Phys., 14, 9129–9136, https://doi.org/10.5194/acp-14-9129-2014, 2014.
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Aerosols, Climate, and the Hydrological Cycle, Science, 294, 2119–2124,
- 20 https://doi.org/10.1126/science.1064034, 2001.

5

- Rizzi, C., Finizio, A., Maggi, V., and Villa, S.: Spatial-temporal analysis and risk characterisation of pesticides in Alpine glacial streams, Environ. Pollut., 248, 659 – 666, https://doi.org/10.1016/j.envpol.2019.02.067, 2019.
 - Rosati, B., Gysel, M., Rubach, F., Mentel, T. F., Goger, B., Poulain, L., Schlag, P., Miettinen, P., Pajunoja, A., Virtanen, A., Klein Baltink, H., Henzing, J. S. B., Größ, J., Gobbi, G. P., Wiedensohler, A., Kiendler-Scharr, A., Decesari, S., Facchini, M. C., Weingartner, E., and
- 25 Baltensperger, U.: Vertical profiling of aerosol hygroscopic properties in the planetary boundary layer during the PEGASOS campaigns, Atmos. Chem. Phys., 16, 7295–7315, https://doi.org/10.5194/acp-16-7295-2016, 2016.
 - Saarikoski, S., Carbone, S., Decesari, S., Giulianelli, L., Angelini, F., Canagaratna, M., Ng, N. L., Trimborn, A., Facchini, M. C., Fuzzi, S., Hillamo, R., and Worsnop, D.: Chemical characterization of springtime submicrometer aerosol in Po Valley, Italy, Atmos. Chem. Phys., 12, 8401–8421, https://doi.org/10.5194/acp-12-8401-2012, 2012.
- 30 Sabatier, T., Paci, A., Canut, G., Largeron, Y., Dabas, A., Donier, J.-M., and Douffet, T.: Wintertime Local Wind Dynamics from Scanning Doppler Lidar and Air Quality in the Arve River Valley, Atmosphere, 9, https://doi.org/10.3390/atmos9040118, 2018.

Samset, B. H.: How cleaner air changes the climate, Science, 360, 148–150, https://doi.org/10.1126/science.aat1723, 2018.

- Sandrini, S., Fuzzi, S., Piazzalunga, A., Prati, P., Bonasoni, P., Cavalli, F., Bove, M. C., Calvello, M., Cappelletti, D., Colombi, C., Contini, D., de Gennaro, G., Di Gilio, A., Fermo, P., Ferrero, L., Gianelle, V., Giugliano, M., Ielpo, P., Lonati, G., Marinoni, A., Massabò, D., Molteni,
- U., Moroni, B., Pavese, G., Perrino, C., Perrone, M. G., Perrone, M. R., Putaud, J.-P., Sargolini, T., Vecchi, R., and Gilardoni, S.: Spatial and seasonal variability of carbonaceous aerosol across Italy, Atmos. Environ., 99, 587 – 598, https://doi.org/10.1016/j.atmosenv.2014.10.032, 2014.

- Schaap, M., van Loon, M., ten Brink, H. M., Dentener, F. J., and Builtjes, P. J. H.: Secondary inorganic aerosol simulations for Europe with special attention to nitrate, Atmos. Chem. Phys., 4, 857–874, https://doi.org/10.5194/acp-4-857-2004, 2004.
- Schmidli, J.: Daytime Heat Transfer Processes over Mountainous Terrain, J. Atmos. Sci., 70, 4041–4066, https://doi.org/10.1175/JAS-D-13-083.1, 2013.
- 5 Schmidli, J., Böing, S., and Fuhrer, O.: Accuracy of Simulated Diurnal Valley Winds in the Swiss Alps: Influence of Grid Resolution, Topography Filtering, and Land Surface Datasets, Atmosphere, https://doi.org/10.3390/atmos9050196, 2018.
 - Seibert, P.: The riddles of foehn introduction to the historic articles by Hann and Ficker, Meteorol. Z., 21, 607–614, https://doi.org/10.1127/0941-2948/2012/0398, 2012.
 - Seibert, P., Kromp-Kolb, H., Baltensperger, U., Jost, D. T., and Schwikowski, M.: Trajectory Analysis of High-Alpine Air Pollution Data, pp. 595–596, Springer US, Boston, MA, https://doi.org/10.1007/978-1-4615-1817-4_65, 1994.
- Seibert, P., Kromp-kolb, H., Kasper, A., Kalina, M., Puxbaum, H., Jost, D. T., Schwikowski, M., and Baltensperger, U.: Transport of polluted boundary layer air from the Po Valley to high-alpine sites, Atmos. Environ., 32, 3953 – 3965, https://doi.org/10.1016/S1352-2310(97)00174-X, 1998.
- Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to Climate Change 2nd ed., John Wiley & Sons,

15 2006.

10

- Serafin, S. and Zardi, D.: Daytime Heat Transfer Processes Related to Slope Flows and Turbulent Convection in an Idealized Mountain Valley, J. Atmos. Sci., 67, 3739–3756, https://doi.org/10.1175/2010JAS3428.1, 2010.
- Serafin, S., Adler, B., Cuxart, J., De Wekker, S. F. J., Gohm, A., Grisogono, B., Kalthoff, N., Kirshbaum, D. J., Rotach, M. W., Schmidli, J., Stiperski, I., Večenaj, Ž., and Zardi, D.: Exchange Processes in the Atmospheric Boundary Layer Over Mountainous Terrain, Atmosphere,
- 20 9, https://doi.org/10.3390/atmos9030102, 2018.
- Silibello, C., Calori, G., Brusasca, G., Giudici, A., Angelino, E., Fossati, G., Peroni, E., and Buganza, E.: Modelling of PM 10 concentrations over Milano urban area using two aerosol modules, Environ. Modell. Softw., 23, 333–343, https://doi.org/10.1016/j.envsoft.2007.04.002, 2008.

Skamarock, W. C.: Evaluating Mesoscale NWP Models Using Kinetic Energy Spectra, Mon. Weather Rev., 132, 3019–3032, https://doi.org/10.1175/MWR2830.1, 2004.

- Sprenger, M. and Wernli, H.: The LAGRANTO Lagrangian analysis tool version 2.0, Geosci. Model Dev., 8, 2569–2586, https://doi.org/10.5194/gmd-8-2569-2015, 2015.
- Squizzato, S. and Masiol, M.: Application of meteorology-based methods to determine local and external contributions to particulate matter pollution: A case study in Venice (Italy), Atmos. Environ., 119, 69 81, https://doi.org/10.1016/j.atmosenv.2015.08.026, 2015.
- 30 Stohl, A.: Trajectory statistics-A new method to establish source-receptor relationships of air pollutants and its application to the transport of particulate sulfate in Europe, Atmos. Environ., 30, 579 – 587, https://doi.org/10.1016/1352-2310(95)00314-2, 1996.
 - Tampieri, F., Trombetti, F., and Scarani, C.: Summer daily circulation in the Po Valley, Italy, Geophys. Astro. Fluid, 17, 97–112, https://doi.org/10.1080/03091928108243675, 1981.
 - Tang, L., Haeger-Eugensson, M., Sjöberg, K., Wichmann, J., Molnár, P., and Sallsten, G.: Estimation of the long-range transport contribution
- 35 from secondary inorganic components to urban background PM10 concentrations in south-western Sweden during 1986–2010, Atmos. Environ., 89, 93 – 101, https://doi.org/10.1016/j.atmosenv.2014.02.018, 2014.
 - Teixeira, M. A. C., Kirshbaum, D. J., Ólafsson, H., Sheridan, P. F., and Stiperski, I., eds.: The Atmosphere over Mountainous Regions, Frontiers Research Topics, 2016.

- Thunis, P., Pederzoli, A., and Pernigotti, D.: Performance criteria to evaluate air quality modeling applications, Atmos. Environ., 59, 476 482, https://doi.org/10.1016/j.atmosenv.2012.05.043, 2012.
- Thyer, N. H.: A theoretical explanation of mountain and valley winds by a numerical method, Archiv für Meteorologie, Geophysik und Bioklimatologie, Serie A, 15, 318–348, https://doi.org/10.1007/BF02247220, 1966.
- 5 Tudoroiu, M., Eccel, E., Gioli, B., Gianelle, D., Schume, H., Genesio, L., and Miglietta, F.: Negative elevation-dependent warming trend in the Eastern Alps, Environ. Res. Lett., 11, 044 021, https://doi.org/10.1088/1748-9326/11/4/044021, 2016.
 - Van Donkelaar, A., Martin, R. V., Brauer, M., Kahn, R., Levy, R., Verduzco, C., and Villeneuve, P. J.: Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: development and application, Environ. Health Persp., 118, 847, https://doi.org/10.1289/ehp.0901623, 2010.
- 10 Wagner, J. S., Gohm, A., and Rotach, M. W.: The impact of valley geometry on daytime thermally driven flows and vertical transport processes, Q. J. Roy. Meteor. Soc., 141, 1780–1794, https://doi.org/10.1002/qj.2481, 2014a.
 - Wagner, J. S., Gohm, A., and Rotach, M. W.: The Impact of Horizontal Model Grid Resolution on the Boundary Layer Structure over an Idealized Valley, Mon. Weather Rev., 142, 3446–3465, https://doi.org/10.1175/MWR-D-14-00002.1, 2014b.
 - Waked, A., Favez, O., Alleman, L. Y., Piot, C., Petit, J.-E., Delaunay, T., Verlinden, E., Golly, B., Besombes, J.-L., Jaffrezo, J.-L., and
- 15 Leoz-Garziandia, E.: Source apportionment of PM₁₀ in a north-western Europe regional urban background site (Lens, France) using positive matrix factorization and including primary biogenic emissions, Atmos. Chem. Phys., 14, 3325–3346, https://doi.org/10.5194/acp-14-3325-2014, 2014.
 - Wang, X., Wang, W., Yang, L., Gao, X., Nie, W., Yu, Y., Xu, P., Zhou, Y., and Wang, Z.: The secondary formation of inorganic aerosols in the droplet mode through heterogeneous aqueous reactions under haze conditions, Atmos. Environ., 63, 68 – 76, https://doi.org/10.1016/j.atmosenv.2012.09.029, 2012.
- Weissmann, M., Braun, F. J., Gantner, L., Mayr, G. J., Rahm, S., and Reitebuch, O.: The Alpine Mountain–Plain Circulation: Airborne
 - Doppler Lidar Measurements and Numerical Simulations, Mon. Weather Rev., 133, 3095–3109, https://doi.org/10.1175/MWR3012.1, 2005.

WHO: Ambient air pollution: a global assessment of exposure and burden of disease, Tech. rep., 2016.

- 25 Wiegner, M. and Gei
 ß, A.: Aerosol profiling with the Jenoptik ceilometer CHM15kx, Atmos. Meas. Tech., 5, 1953–1964, https://doi.org/10.5194/amt-5-1953-2012, 2012.
 - WMO: WMO/IGAC Impacts of megacities on air pollution and climate, Tech. rep., World Meteorological Organization, https://library.wmo. int/pmb_ged/gaw_205.pdf, 2012.

WMO: WMO Global Atmosphere Watch (GAW) Implementation Plan: 2016-2023, Tech. rep., World Meteorological Organization, https:

30 //library.wmo.int/doc_num.php?explnum_id=3395, 2017.

20

Wotawa, G., Kröger, H., and Stohl, A.: Transport of ozone towards the Alps – results from trajectory analyses and photochemical model studies, Atmos. Environ., 34, 1367 – 1377, https://doi.org/10.1016/S1352-2310(99)00363-5, 2000.

Ying, C., Chunsheng, Z., Qiang, Z., Zhaoze, D., Mengyu, H., and Xincheng, M.: Aircraft study of Mountain Chimney Effect of Beijing, China, J. Geophys. Res., 114, https://doi.org/10.1029/2008JD010610, 2009.

35 Yuan, Y., Ries, L., Petermeier, H., Steinbacher, M., Gómez-Peláez, A. J., Leuenberger, M. C., Schumacher, M., Trickl, T., Couret, C., Meinhardt, F., and Menzel, A.: Adaptive selection of diurnal minimum variation: a statistical strategy to obtain representative atmospheric CO₂ data and its application to European elevated mountain stations, Atmos. Meas. Tech., 11, 1501–1514, https://doi.org/10.5194/amt-11-1501-2018, 2018.

- Zardi, D. and Whiteman, C. D.: Diurnal Mountain Wind Systems, pp. 35–119, Springer Netherlands, Dordrecht, https://doi.org/10.1007/978-94-007-4098-3_2, 2013.
- Zeng, Y. and Hopke, P.: A study of the sources of acid precipitation in Ontario, Canada, Atmos. Environ., 23, 1499 1509, https://doi.org/10.1016/0004-6981(89)90409-5, 1989.
- 5 Zeng, Z., Chen, A., Ciais, P., Li, Y., Li, L. Z. X., Vautard, R., Zhou, L., Yang, H., Huang, M., and Piao, S.: Regional air pollution brightening reverses the greenhouse gases induced warming-elevation relationship, Geophys. Res. Lett., 42, 4563–4572, https://doi.org/10.1002/2015GL064410, 2015.
 - Zong, Z., Wang, X., Tian, C., Chen, Y., Fu, S., Qu, L., Ji, L., Li, J., and Zhang, G.: PMF and PSCF based source apportionment of PM2.5 at a regional background site in North China, Atmos. Res., 203, 207 215, https://doi.org/10.1016/j.atmosres.2017.12.013, 2018.
- 10 Zuev, V. V., Burlakov, V. D., Nevzorov, A. V., Pravdin, V. L., Savelieva, E. S., and Gerasimov, V. V.: 30-year lidar observations of the stratospheric aerosol layer state over Tomsk (Western Siberia, Russia), Atmos. Chem. Phys., 17, 3067–3081, https://doi.org/10.5194/acp-17-3067-2017, 2017.

Ångström, A.: On the Atmospheric Transmission of Sun Radiation and on Dust in the Air, Geografiska Annaler, 11, 156–166, 1929.