# Impact of halogen chemistry on summertime air quality in coastal and continental Europe: application of CMAQ model and implication for regulation

Qinyi Li<sup>1</sup>, Rafael Borge<sup>2</sup>, Golam Sarwar<sup>3</sup>, David de la Paz<sup>2</sup>, Brett Gantt<sup>4</sup>, Jessica Domingo<sup>2</sup>, Carlos A. Cuevas<sup>1</sup>, and Alfonso Saiz-Lopez<sup>1\*</sup>

2 Environmental Modelling Laboratory, Department of Chemical & Environmental Engineering, Universidad Politécnica de Madrid (UPM), Madrid, Spain

3 National Exposure Research Laboratory, Environmental Protection Agency, Research Triangle Park, NC 27711, United States

4 Office of Air Quality Planning and Standards, Environmental Protection Agency, Research Triangle Park, NC 27711, United States

\*Correspondence to: Alfonso Saiz-Lopez (a.saiz@csic.es)

#### Abstract:

10

Halogen (Cl, Br, and I) chemistry has been reported to influence the formation of secondary air 15 pollutants. Previous studies mostly focused on the impact of chlorine species on air quality over large spatial scales. Very little attention has been paid to the effect of the combined halogen chemistry on air quality over Europe and its implications for control policy. In the present study, we utilize a widely-used regional model, the Community Multiscale Air Quality Modeling System (CMAQ), incorporated with the latest halogen sources and chemistry, to simulate the 20 abundance of halogen species over Europe and to examine the role of halogens in the formation of secondary air pollution. The results suggest that the CMAQ model is able to reproduce the level of O<sub>3</sub>, NO<sub>2</sub>, and halogen species over Europe. Chlorine chemistry slightly increases the levels of OH, HO<sub>2</sub>, NO<sub>3</sub>, O<sub>3</sub>, and NO<sub>2</sub> and substantially enhances the level of Cl radical. 25 Combined halogen chemistry induces complex effects on OH (ranging from -0.023 pptv to 0.030 pptv) and HO<sub>2</sub> (in the range of -3.7 to 0.73 pptv), significantly reduces the concentrations of NO<sub>3</sub> (as much as 20 pptv) and O<sub>3</sub> (as much as 10 ppbv), and decreases NO<sub>2</sub> in the highly polluted regions (as much as 1.7 ppbv) and increases NO<sub>2</sub> (up to 0.20 ppbv) in other areas. The maximum effects of halogen chemistry occur over oceanic and coastal regions, but some noticeable impacts also occur over continental Europe. Halogen chemistry affects the number of 30 days exceeding the European Union target threshold for the protection of human being and vegetation from ambient O<sub>3</sub>. In light of the significant impact of halogen chemistry on air quality, we recommend that halogen chemistry be considered for inclusion in air quality policy assessments, particularly in coastal cities.

<sup>5 1</sup> Department of Atmospheric Chemistry and Climate, Institute of Physical Chemistry Rocasolano, CSIC, Madrid 28006, Spain

#### 35 **1. Introduction**

Halogen (Cl, Br, and I) species and related processes have been known to deplete stratospheric ozone (O<sub>3</sub>) for several decades (Molina and Rowland, 1974; Farman et al., 1985). In the troposphere, the chemistry of halogens in the troposphere has been described in detail in recent reviews and references therein (Saiz-Lopez and von Glasow, 2012; Simpson et al., 2015), so it is just briefly outlined here. Halogen species affect the concentration of air pollutants, e.g., directly destroying O<sub>3</sub> (R1), indirectly decreasing O<sub>3</sub> production by reducing NO<sub>2</sub> (R2 and R3), influencing the NO/NO<sub>2</sub> ratio (R2 and R4) and the HO<sub>2</sub>/OH ratio (R5 and R6). The budgets of NO<sub>x</sub> (NO+NO<sub>2</sub>) and HO<sub>x</sub> (OH+HO<sub>2</sub>) also affect the formation of O<sub>3</sub> (e.g., Sillman, 1999; Li et al., 2018).

50

40

$$X (Cl, Br, I) + O_3 \rightarrow XO + O_2$$
(R1)

$$XO + NO_2 \rightarrow XONO_2$$
 (R2)

$$XONO_2 + H_2O(l) \rightarrow HOX + HNO_3(l)$$
(R3)

$$XO + NO \rightarrow X + NO_2$$
 (R4)

$$XO + HO_2 \rightarrow HOX + O_2$$
 (R5)

$$HOX + hv \rightarrow OH + X$$
 (R6)

Chlorine radical (Cl) initiates the oxidation of hydrocarbons (R7), including methane (CH<sub>4</sub>) and non-methane volatile organic compounds (NMVOC), in a similar way to OH radical, reducing the lifetime of CH<sub>4</sub> and NMVOC and leading to the formation of  $O_3$  in the presence of  $NO_x$ (Thornton et al., 2010).

$$RH + Cl \rightarrow HCl + RO_2 \tag{R7}$$

The combined effect of halogen chemistry on air quality, therefore, is complicated and depends heavily on local conditions, e.g., atmospheric compositions, oxidative capacity, etc. (Sherwen et al., 2016; Muñiz-Unamunzaga et al., 2018). Evaluation of the complex role of halogen chemistry in air quality requires the employment of advanced, high-resolution chemical transport models.

60

A number of modeling studies have been conducted to investigate the impact of individual halogen species on air quality. The chemistry of chlorine, mainly that of nitryl chloride (ClNO<sub>2</sub>), has been reported to increase the oxidation capacity and the formation of  $O_3$  in recent studies

(Sarwar et al., 2012, 2014; Li et al., 2016; Sherwen et al., 2017; Sommariva et al., 2018). Bromine (Fernandez et al., 2014) and iodine (Saiz-Lopez et al., 2014) chemistry are reported to decrease the concentration of O<sub>3</sub> over the oceanic and terrestrial regions.

Only a few regional modeling studies have explored the combined influence of the halogen chemistry on air quality. The first modeling study with combined halogen (Cl, Br, and I) chemistry was conducted by Sarwar et al. (2015) who used a hemispheric version of the Community Multiscale Air Quality (CMAQ) model (Ching and Byun, 1999; Byun and Schere, 2006; Mathur et al., 2017) and reported a decrease of surface O<sub>3</sub> by ~15% to ~48% over the Northern Hemisphere by Br and I. Gantt et al. (2017) then utilized the CMAQ model to explore the role of halogen chemistry at a regional scale over the continental United States (US). While these studies focused on Northern Hemisphere and the continental US, Muñiz-Unamunzaga et al. (2018) applied the full-halogen chemistry version of CMAQ with a resolution of 4 km and reported up to 5 ppbv decrease of O<sub>3</sub> in the city of Los Angeles, California, US. Sherwen et al. (2017) used a global model, GEOS-Chem, in a regional configuration (with a grid size of 0.25°  $\times$  0.315°, ~25km x ~25km) and predicted a large decrease of O<sub>3</sub>, on average 13.5 pptv (25%) and as much as 28.9 pptv (45%) in Europe. Sarwar et al. (2019) further updated the halogen chemistry in CMAQ model and reported a reduction of 3 to 12 ppbv of annually average O<sub>3</sub> over seawater and 3 to 6 ppbv over coastal and 3 ppbv over inland area by Br and I. These previous regional studies using various models (or versions of models) in different areas reported a large range of the halogen impact on O<sub>3</sub> highlighting the uncertainty in this research field.

85

65

70

75

80

The regulation of air quality and the control of air pollutants emission in Europe started in the early 1970s and over forty years of effort has successfully improved air quality throughout Europe (EEA, 2018a). Nonetheless, poor air quality persist in major cities like Madrid, Paris, and London (EEA, 2018a); this shows the need for continuied air quality management and effective policy. Because the influence of halogens on air quality is uncertain and potentially has an impact on air quality management decisions, we have conducted regional simulations using the latest version of the CMAQ model implemented with comprehensive halogen sources and chemistry (Sarwar et al., 2019) to examine the overall effect of halogen species on air quality model

predictions (Sommariva et al., 2018), we used a CMAQ model domain with 12 km horizontal resolution (higher than the previous studies on halogen impact covering Europe) to simulate the levels of halogen species over Europe, examine the effect on the oxidation capacity and the concentration of air pollutants, and explore the potential implications for air quality policy related to NO<sub>2</sub> and O<sub>3</sub>.

95

105

## 2. Method and Materials

#### 100 **2.1 Data**

The meteorological inputs for the CMAQ model were obtained from the Weather Research and Forecasting model (WRF 3.7.1) (Skamarock and Klemp, 2008; Borge et al., 2008a) as an offline input. The WRF model was initialized from global reanalyses from the National Centers for Environmental Prediction (NCEP) Global Forecast System (GFS) with a spatial resolution of  $1^{\circ} \times 1^{\circ}$  and a temporal resolution of 6 h (available online at http://rda.ucar.edu/datasets/ds083.2/) which was updated daily from NCEP global analyses with 0.5° resolution (available online at http://www.nco.ncep.noaa.gov/pmb/products/sst/). NCEP's ADP global upper-air (NCAR archive ds351.0) and global surface observations (NCAR archive ds461.0) were used to drive

Anthropogenic emissions for the year 2016 were taken from the 0.1° × 0.1° gridded EMEP inventory (EMEP/CEIP, 2014). It should be noted that no anthropogenic chlorine sources are included in our emission inventory. The temporal profiles and vertical distribution needed to resolve the emissions were those used in the EuroDelta experiment (van Loon et al., 2007). Biogenic emissions were estimated using the Model of Emissions of Gases and Aerosols from Nature (MEGANv2.10) (Guenther et al., 2012). All emissions were gridded to our model domain, temporally allocated and chemically speciated using the Sparse Matrix Operator Kernel Emissions (SMOKE) model, version 3.6.5 (UNC, 2015; Borge et al., 2008b).

the simulation with a Newtonian relaxation technique in the WRF model.

In addition, we used measurement data of  $NO_2$  and  $O_3$  from over 400 background stations (traffic and industrial stations are not included) across Europe from database *AirBase* (public air

120 quality database system of the European Environment Agency, 2018) to compare the results of our simulation with observations (Fig. 1). Among these stations, 315 are located in inland areas (208 for NO<sub>2</sub> and 266 for O<sub>3</sub>), and 125 are located in the coastal areas (83 for NO<sub>2</sub> and 93 for  $O_3$ ).



125 Figure 1. Geographic representation of the modeling domain and air quality stations used for model evaluation

## 2.2 Modeling system

130

The CMAQ model is widely used and includes comprehensive representations of many essential atmospheric processes. The skill of the model in reproducing observed air quality has been demonstrated in many previous studies (Foley et al., 2010; Appel et al., 2013, 2017; Mathur et al., 2017), including applications over Europe (Borge et al., 2008a; Appel et al., 2012; Solazzo et al., 2017). CMAQ version 5.2 (www.epa.gov/cmaq; doi:10.5281/zenodo.1167892) containing the Carbon Bond chemical mechanism (Appel et al., 2017) with halogen chemistry was used in this study. We updated the chlorine chemistry of Sarwar et al. (2012) and implemented it in 135 CMAQ version 5.2 (Table S1). The heterogeneous hydrolysis of dinitrogen pentoxide (N<sub>2</sub>O<sub>5</sub>) can produce  $CINO_2$  and nitric acid (HNO<sub>3</sub>) in the presence of particulate chloride. In the absence of particulate chloride, heterogeneous hydrolysis of N<sub>2</sub>O<sub>5</sub> produces only HNO<sub>3</sub>. Sarwar et al. (2015) implemented the initial bromine and iodine chemistry in CMAQ which has recently been updated by Sarwar et al. (2019). The updates include revising the gas-phase bromine and iodine 140 reactions, incorporating several heterogeneous reactions of bromine and iodine species, incorporating several aqueous-phase bromine reactions, and revising the iodine and bromine emissions. We combine the updated chlorine, bromine, and iodine chemistry with the Carbon Bond chemical mechanism and use it in this study.

#### 145 **2.3 Simulation setup**

A detailed description of physics and other model options (Table S2) can be found in (de la Paz et al., 2016). The CMAQ model is applied over a domain covering the entirety of Europe (Fig. 1) with 12 km horizontal resolution. The vertical extent of the model extended from the surface to 100 mbar and contained 35 layers with an average surface layer thickness of approximately 20 m. The CMAQ chemical transport model is configured to use the Piecewise Parabolic Method to describe advection processes, the Asymmetric Convective Model (version 2) to describe vertical diffusion processes, and the multiscale method to describe horizontal diffusion processes. Gasphase chemistry, aqueous chemistry, aerosol processes, and dry and wet deposition were also included. The Rosenbrock solver was used for gas-phase chemistry.

- The study was completed for the month of July 2016 with a spin-up period of 30 days. We performed three simulations to isolate the effect of halogen chemistry on air quality (in brackets the name of the scenario used hereafter):
  - (1) Base model without halogen chemistry (BASE),
  - (2) BASE with chlorine chemistry (CL), and
- 160 (3) CL with Br and I chemistry (HAL).

The BASE model simulation includes the Carbon Bond 2015 chemical mechanism but does not contain any halogen chemistry, and only the HNO<sub>3</sub> is produced from the heterogeneous hydrolysis of  $N_2O_5$ . The CL simulation contains the Carbon Bond chemical mechanism with chlorine chemistry and considers ClNO<sub>2</sub> and HNO<sub>3</sub> production from the heterogeneous uptake of  $N_2O_5$  on the aerosol surface. The HAL simulation contains the Carbon Bond chemical mechanism with full halogen chemistry and produces ClNO<sub>2</sub> and HNO<sub>3</sub> from the heterogeneous

165

uptake of N<sub>2</sub>O<sub>5</sub> on the aerosol surface.

Boundary conditions for the model were derived from the hemispheric CMAQ simulations (Mathur et al., 2017). Three different annual simulations were conducted using the hemispheric
CMAQ model for 2016: the first simulation used the Carbon Bond chemical mechanism without any halogen chemistry, the second simulation used the Carbon Bond chemical mechanism and the chlorine chemistry, and the third simulation used the Carbon Bond chemical mechanism and the full halogen chemistry. Results from the corresponding hemispheric CMAQ simulation were used to generate boundary conditions for the BASE, CL, and HAL simulations. Therefore, the difference between CL and BASE simulations represents the impact of the chlorine chemistry on air quality and the difference between HAL and BASE simulations represents the effect of halogen chemistry on air quality.

3. Results and Discussions

#### 180 **3.1 Evaluation of model performance**

The performance of the CMAQ model in simulating air quality over Europe is evaluated using observation data collected from >400 measurement stations. We separate the stations into coastal (within 24 km from the coast) and continental stations (Fig. 1). Table 1 presents the statistics of the model performance for  $O_3$  and  $NO_2$  for BASE and HAL simulations.

The BASE and HAL simulations generally reproduce the concentration levels and the temporal variations of  $O_3$  and  $NO_2$  both at coastal and continental stations. The correlation coefficients between simulations and observations (Fig. S1) show that CMAQ satisfactorily reproduces the variation of  $O_3$  and  $NO_2$  over most of Europe especially the coastal regions (> 0.6 for  $O_3$  and > 0.4 for  $NO_2$ ). The BASE simulation under-predicts  $O_3$  compared to observations both at coastal and continental stations (Table 1), possibly due to the uncertainty of NMVOC emission inventory (Sherwen et al., 2017) and the underestimated  $NO_x$  (Table 1). The HAL simulation slightly improves the correlation coefficient of  $O_3$  but decreases the average level of  $O_3$ compared to the BASE case. Diurnal variation plots (Fig. S2) suggest that both BASE and HAL simulations can reproduce the temporal patterns of O<sub>3</sub>.

The BASE simulation under-predicts NO<sub>2</sub> compared to observations both at coastal and continental stations (Table 1). Such an under-estimation of NO<sub>2</sub> can occur for many reasons including (1) positive artifacts of NO<sub>2</sub> monitors (Jung et al., 2017), (2) under-estimation of NO<sub>x</sub> in the emission inventory, and (3) rapid transformation of NO<sub>2</sub> into HNO<sub>3</sub> in the model compared to the real atmosphere. Model performance is reasonable as the NO<sub>2</sub> underestimation is relatively small. The HAL case predicted very similar NO<sub>2</sub> concentrations (Table 1).

Overall, the evaluation of CMAQ results over Europe demonstrates that the model is capable of reproducing the levels of atmospheric chemical species and can be used to investigate the impact of halogen chemistry on air quality over Europe. It also suggests that the incorporation of halogen chemistry changes the model performance for  $O_3$  concentrations by a small margin without a noticeable impact on the model performance for  $NO_2$ .

205

Tuble T. Suustear Summary of Model performance								
Statistics	O <sub>3</sub> (µg/m <sup>3</sup> )				NO <sub>2</sub> (μg/m <sup>3</sup> )			
	Coastal		Inland		Coastal		Inland	
	Base	HAL	Base	HAL	Base	HAL	Base	HAL
MB	-0.9	-6.8	-2.3	-6.6	-3.7	-3.7	-3.1	-3.1
ME	16.9	17.5	19.7	20.2	5.8	5.8	5.7	5.6
RMSE	22.6	23.1	25.8	26.4	7.7	7.7	7.2	7.2
r	0.65	0.67	0.60	0.61	0.44	0.44	0.42	0.42
IOA	0.64	0.62	0.62	0.61	0.31	0.31	0.18	0.19

Table 1. Statistical summary of model performance

Note: MB = Mean Bias, RMSE = Root Mean Square Error, r = correlation coefficient, IOA = Index of Agreement.

#### 210 **3.2 Simulated halogen species**

Average surface concentrations of the inorganic halogen species predicted in the HAL simulation over the ocean are summarized in Table 2. HCl is the dominant chlorine species with an average level of 247.9 pptv representing over 96% to the total inorganic chlorine (Cl<sub>y</sub>) while the average ClNO<sub>2</sub> and HOCl is 4.9 pptv (1.9%) and 3.8 pptv (1.5%), respectively, and the remaining species contributing less than 1%. The Br<sub>y</sub> species are relatively evenly partitioned, with HOBr (1.06 pptv, 27.0%), BrCl (0.71 pptv, 18.2%), BrNO<sub>3</sub> (0.67 pptv, 16.9), HBr (0.66

pptv, 16.8%), and Br<sub>2</sub> (0.33 pptv, 8.4%) being the abundant species while the remaining species contribute <5%. HOI (5.1 pptv, 52.0%), INO<sub>3</sub> (2.8 pptv, 28.7%), IO (1.0 pptv, 10.3%) contribute over 90% of I<sub>y</sub> over the ocean, while the remaining species contribute ~10%. The predicted average concentrations of the critical halogen radicals, Cl, BrO, and IO, are 2.0x10<sup>-4</sup> pptv, 0.03 pptv, and 0.4 pptv, respectively, over the ocean in the Europe.

**Concentration Percentage** Concentration Percentage Concentration Percentage Species Species **Species** (ppt) (%) (ppt) (%) (ppt) (%) 27.0 96.1 HOI HCl 247.9 HOBr 1.06 5.1 52.0 4.9 1.9 BrCl 18.2 2.8 CINO<sub>2</sub> 0.71 INO<sub>3</sub> 28.7 3.8 HOCI 1.5 BrNO<sub>3</sub> 0.67 16.9 Ю 1.0 10.3 1.2 0.5 16.8 0.4 4.0 CINO<sub>3</sub> HBr 0.66 I 0.25 0.1 BrO 0.38 9.7 0.3 3.1 ClO  $I_2O_3$ 0.02 0.008 0.33 8.4 HI 0.1 1.0 Cl<sub>2</sub> Br<sub>2</sub> 0.0002 0.0001 0.09 0.05 0.5 Cl BrNO<sub>2</sub> 2.4 I2 Br 0.03 0.7 INO 0.03 0.3 0.01 0.1  $I_2O_2$  $INO_2 \\$ 0.01 0.1 0.004 < 0.1  $I_2O_4$ **Total Cl** 258.1 100 3.9 100 9.8 100 Total Br Total I

Table 2. Simulated average concentrations of inorganic halogen species over the ocean



225

230

Figure 2. Monthly average ClNO<sub>2</sub>, HCl, BrO, and IO concentration in the HAL simulation.

The spatial distributions of key halogen species are shown in Fig 2. The HAL simulation with full halogen chemistry simulates generally higher CINO<sub>2</sub> levels (with the highest monthly average value of 113.0 pptv) along the coast of the Mediterranean Sea and the North Sea with some influence into continental Europe, especially in Germany. The simulated HCl shows a similar pattern to that of CINO<sub>2</sub> but with much higher concentration (>10 times higher). The predicted BrO levels over Europe are low (average value ~0.17 pptv) with the largest predicted value occurring within the Arctic circle while GEOS-Chem predicted >1.0 pptv level of BrO in Mediterranean Sea (Sherwen et al., 2017). The predicted IO peaks over the Mediterranean region with a maximum value of 6.9 pptv.

Direct measurements of halogen species are very scarce and not available for the period covered in the present study (July 2016). Since a direct comparison is not possible, here we present a comparison of the simulated concentrations with observations from previous studies (Table 3), to provide an approximate assessment of the representation of halogen species in the HAL simulation of the CMAQ model over Europe.

Table 3. The comparison of observed and simulated halogen species

Location	Species	Observation *	Simulation #	
Hessen, Germany <sup>a</sup>	ClNO <sub>2</sub>	800.0	273.4	
London, United Kingdom <sup>b</sup>	ClNO <sub>2</sub>	724.0	801.5	
Weybourne, United Kingdom <sup>c</sup>	ClNO <sub>2</sub>	65	373	
Weybourne, United Kingdom <sup>d</sup>	ClNO <sub>2</sub>	946	373	
Weybourne, United Kingdom <sup>e</sup>	ClNO <sub>2</sub>	1100 (summer) 75.6 (autumn) 733 (winter)	373	
Leicester, United Kingdom <sup>e</sup>	CINO <sub>2</sub>	274 (spring) 74.2 (summer) 248 (winter)	274	
Penlee Point , United Kingdom <sup>e</sup>	ClNO <sub>2</sub>	922	319	
Mace Head, Ireland <sup>f</sup>	BrO	6.5	10.1	
Brittany, France <sup>g</sup>	BrO	7.5	0.4	
Dead Sea <sup>h</sup>	BrO	100.0	0.2	
Mace Head, Ireland <sup>i</sup>	ΙΟ	4.0~50.0	3.9	
Brittany, France <sup>j</sup>	ΙΟ	7.7~30.0	1.1	
Dagebull, Germany <sup>k</sup>	ΙΟ	2.0	9.0	
Atlantic Ocean <sup>1</sup> (Prados-Roman et al., 2015)	Ю	0.4 to 0.5 (daytime average)	0.4 to 2.0 (daytime average)	

\*: Maximum value (pptv). #: Maximum value (pptv) from the HAL simulation.

- a: Phillips et al. 2012.
  b: Bannan et al., 2015.
  c: Banna et al., 2017.
  d: Sherwen et al., 2017.
- e: Sommariva et al., 2018.
- f: Saiz-Lopez et al., 2004. g: Mahajan et al. 2009.

g. Manajan et al. 2007.
h: Matveev et al., 2001; Holla et al., 2015.
i: Allan et al., 2000; Commane et al., 2011.
j: Britter et al., 2005; Furneaux et al., 2010.
k: Peters et al., 2005.

l: Prados-Roman et al., 2015

Numerous CINO<sub>2</sub> measurements have been reported around the globe which show that CINO<sub>2</sub> is ubiquitous in the boundary layer with maximum values ranging from hundreds to thousands pptv in polluted coastal (Osthoff et al., 2008; Wang et al., 2016) and continental regions (Tham et al., 2016; Thornton et al., 2010). A few campaigns have been conducted in Europe. Phillips et al. (2012) reported a maximum value of 800 pptv for ClNO<sub>2</sub> in Hessen, Germany where CMAQ 245 predicts a concentration of 273 pptv. Bannan et al. (2015) observed a peak value of 724 pptv in London where CMAQ predicts a concentration of 802 pptv. Simulations with the GEOS-Chem model (Sherwen et al., 2017) reported maximum values of 110 pptv and 140 pptv at Hessen and London, respectively. Several field campaigns have been conducted in Weybourne in the past 250 few years to measure CINO<sub>2</sub>. Sherwen et al. (2017) reported a peak concentration of 946 pptv. Bannan et al. (2019) reported a peak value of 65 pptv, and Sommariva et al. (2018) reported a peak value of 1100 pptv in summer, 75.6 pptv in autumn and 733 pptv in winter. CMAQ simulated a maximum of 373 pptv at that location while GEOS-Chem predicted 458 pptv. Sommariva et al. (2018) also reported measurements of CINO<sub>2</sub> at Leicester with a maximum 255 value of 274 pptv in spring, 74.2 pptv in summer, 248 pptv in winter, and that at Penlee Point a peak value of 922 pptv. CMAQ predicted a maximum of 274 pptv at Leicester and 319 pptv at Penlee Point. Eger et al. (2019) conducted shipborne observation of CINO<sub>2</sub> in the Mediterranean Sea and reported up to 600 pptv ClNO<sub>2</sub> during their campaign, which is similar to the prediction of the present study.

- The observed levels of HCl in Europe range from <100 pptv to 5000 pptv (Hossaini et al., 2016 and the reference therein). The CMAQ model predicted monthly average concentrations of HCl 6.3 to 1249 pptv, which is similar to the observation ranges. GEOS-Chem (Sherwen et al., 2017) predicted a maximum of 12 pptv for HCl, which is significantly lower than the measurements in Europe.
- BrO measurements have been reported at ground-based sites and during the ship cruises which generally demonstrate a range of 0.5 to 2.0 pptv maximum values for land measurements and 3.0 to 3.6 pptv for ship measurements (Saiz-Lopez and von Glasow, 2012). BrO observations have been reported at several coastal sites in Europe. BrO level of up to 6.5 pptv (Saiz-Lopez et al., 2004) and 7.5 pptv (Mahajan et al., 2009) were reported in Mace Head and Brittany,

270 respectively. CMAQ predicts 10.1 pptv and 0.4 pptv at those locations, which are lower than the measurements. Sherwen et al. (2017) also predicted small values with a maximum of 0.8 pptv in Mace Head and 0.5 pptv in Brittany. An extremely high level of BrO, ~100 pptv, was observed over the Dead Sea (Matveev et al., 2001; Holla et al., 2015). CMAQ is not able to reproduce such a high level of BrO due to the lower bromide content in typical ocean water (which was used in the present study for the Dead Sea) compared to the exceptionally high bromide content 275 in Dead Sea (Tas et al., 2006; Sarwar et al., 2015).

Global measurements show that the IO levels observed by ground-based campaigns were generally between 0.2 and 2.4 pptv while those by ship measurement were ~3.5 pptv (Saiz-Lopez and von Glasow, 2012). Observations of IO have also been conducted in Europe. Maximum IO levels of 4.0~50.0 pptv were measured at Mace Head (Allan et al., 2000; 280 Commane et al., 2011). CMAQ predicts a value of 3.9 pptv at Mace Head while GEOS-Chem predicted a value of 0.6 pptv (Sherwen et al., 2017). In Brittany, up to 7.7~30.0 pptv of IO were observed by Bitter et al. (2005) and Furneaux et al. (2010). CMAQ predicts 1.1 pptv of IO at Brittany and Sherwen et al. (2017) predicted 0.07 pptv. A maximum IO concentration of 2.0 285 pptv was reported in Dagebull (Peters et al., 2005), and CMAQ predicts 9.0 pptv at that site, while GEOS-Chem predicted 1.8 pptv (Sherwen et al., 2017). Prados-Roman et al. (2015) reported the level of IO during a ship-based campaign in the range of <0.4 to >1.4 pptv (daytime average) around the globe and 0.4 to 0.5 pptv (daytime average) in the south of Spain and the west of Africa (over the Atlantic), and the present study predicted 0.4 to 2.0 pptv (daytime 290 average) of IO over those areas.



#### 3.3 Influence of halogen chemistry on the atmospheric oxidation capacity

Figure 3. Monthly average OH and HO<sub>2</sub> concentration in the BASE simulation, and changes due to chlorine (CL) and full halogen chemistry (HAL).

Fig. 3 shows the monthly average concentrations of the OH and HO<sub>2</sub> radicals predicted by the BASE simulation and the impact of chlorine chemistry (CL-BASE), and the full halogen chemistry (HAL-BASE), on the simulated OH and HO<sub>2</sub> levels. In the BASE simulation, the highest OH concentration levels are predicted over the oceans especially along ship tracks, with 300

a maximum value of 0.38 pptv (~9.5  $\times$  10<sup>6</sup> molecule cm<sup>-3</sup>). The chlorine chemistry slightly increased the OH level over most of the domain by up to 0.0041 pptv ( $1.0 \times 10^5$  molecule cm<sup>-3</sup>). The impact of the halogen chemistry has competing effects on OH concentrations with a maximum increase of 0.030 pptv ( $7.5 \times 10^5$  molecule cm<sup>-3</sup>) and a reduction of as much as 0.023 pptv ( $5.8 \times 10^5$  molecule cm<sup>-3</sup>). The net change of OH due to halogen was near zero. The BASE simulation predicts the highest values of HO<sub>2</sub> over the Mediterranean Sea with a maximum value of 21 pptv. The chlorine chemistry increases the HO<sub>2</sub> level in the areas of elevated ClNO<sub>2</sub> 305 predictions (Fig. 2 and Fig. 3). The further addition of halogen chemistry lowers the HO<sub>2</sub> as much as 3.7 pptv and increases HO<sub>2</sub> up to 0.73 pptv compared to the BASE simulation. The overall difference of HO<sub>2</sub> because of halogens was -0.59 pptv in the European domain. The effect of halogen chemistry on OH and HO<sub>2</sub> is the combined effect of the following four pathways: (1) conversion of HO<sub>2</sub> to OH via XO (R5 and R6), in which HO<sub>2</sub> decreases and OH 310 increases; (2) reduction of O<sub>3</sub> (R1 and Figure 7) and the reduced production of OH by O<sub>3</sub> photolysis, in which both the OH and HO<sub>2</sub> decrease; (3) increase of NO<sub>2</sub> (R4 and Figure 7) and the enhanced consumption of OH by the reaction with NO<sub>2</sub>, in which both OH and HO<sub>2</sub> decrease; and (4) increased oxidation of VOCs due to halogens in which both OH and HO<sub>2</sub> increase. Pathway (1) leads to the increase of OH and decrease of HO<sub>2</sub> over coastal area, Mediterranean 315 Sea and Baltic Sea (except the ship tracks). Pathway (2) results in a decrease of OH and  $HO_2$ over the remote ocean. Pathway (2), (3), and (4) results in a decrease of OH and an increase or decrease of HO<sub>2</sub> along the ship tracks in Mediterranean Sea.

Sarwar et al. (2015) reported a small overall decrease of OH (1%) and a significant decrease of 320 HO<sub>2</sub> (11%) in the Northern Hemisphere due to the bromine and iodine chemistry. Their results suggest a considerable reduction of the HO<sub>2</sub>/OH ratio which is consistent with the present study. Muñiz-Unamunzaga et al. (2018) found a slight increase of diurnal OH (1-2%) and a noticeable decrease of HO<sub>2</sub> (4%) leading to a decrease of HO<sub>2</sub>/OH in Los Angeles, California. Sherwen et al. (2017) suggested that OH was reduced across their European domain due to the halogen chemistry, and concluded that the shift of HO<sub>2</sub> to OH by XO could not compensate for the 325 decrease of OH due to the loss of O<sub>3</sub>. Another GEOS-Chem study, however, predicted an increase of OH over the Mediterranean Sea (Stone et al., 2018). The discrepancy among the previous studies and between those works and the present one is difficult to deduce and requires

further investigation. Several possible causes could lead to different simulated levels of halogens and their impact on oxidants, including the different mechanism of producing and recycling halogen species (Sarwar et al., 2019), spatial resolution (Sommariva et al., 2018), emission inventory (Wang et al., 2019), and different spatio-temporal scale of interest (Stone et al., 2018).



Figure 4. Monthly average NO<sub>3</sub> and Cl radical concentrations in the BASE simulation, and changes induced by chlorine (CL) and full halogen chemistry (HAL).

335

340

345

Fig. 4 presents the monthly average prediction of NO<sub>3</sub> and Cl radicals in the BASE scenario and the influence of chlorine (CL-BASE) and halogen chemistry (HAL-BASE) on the levels of NO3 and Cl. The BASE simulation predicted relatively high NO<sub>3</sub> concentrations over the Mediterranean Sea along the busy shipping tracks. Although concentrations as high as 71 pptv are found, the majority of oceanic regions have concentrations in the range of 10 to 40 pptv and between 0 and 4 pptv over land. The chlorine chemistry slightly increases the NO<sub>3</sub> radical due to the increase of the O<sub>3</sub> (see section 3.4). In contrast, the halogen chemistry considerably reduces NO<sub>3</sub> concentrations by as much as 20 pptv at nighttime over the Mediterranean Sea. Muñiz-Unamunzaga et al. (2018) reported a 20-50% (2 to 4 pptv) decrease of NO<sub>3</sub> radical in Los Angeles, California when considering the halogen chemistry.

350

In the BASE simulation, the Cl concentration was negligible because there was no relevant chlorine source incorporated in the CMAQ model. The CL simulation contains the production of ClNO<sub>2</sub> and its subsequent photolysis which increases the Cl concentration of as high as  $7.0 \times 10^{-10}$ <sup>4</sup> pptv (~ $1.75 \times 10^4$  atom cm<sup>-3</sup>). The HAL simulation predicted a very similar magnitude and spatial distribution of chlorine concentration. Sherwen et al. (2017) reported Cl concentrations less than  $1.4 \times 10^4$  atom cm<sup>-3</sup> (~5.6 × 10<sup>-4</sup> pptv) over Europe, comparable to our prediction. Hossaini et al. (2016) reported more than  $1.0 \times 10^4$  atom cm<sup>-3</sup> (~4.0 × 10<sup>-4</sup> pptv) of chlorine over Asia, Europe and North America, with a maximum of  $8.5 \times 10^4$  atom cm<sup>-3</sup> (~ $3.4 \times 10^{-3}$  pptv), 355 using a global chemical transport model (TOMCAT) that incorporated chlorine sources from sea salt dechlorination, coal and biomass burning, oxidation of natural and anthropogenic chlorocarbon, and heterogeneous reactions on sea salt and sulfate aerosol. In their study, Hossaini et al. (2016) used the Reactive Emission Inventory of Chlorine (Keene et al., 1999) which Wang et al. (2019) reported to be unrealistic for present day applications. 360

The current and the previous studies simulated a broad range of the surface Cl concentrations although they were all within the scope of the reported observed (observation-based calculation) values of  $10^3$  to  $10^5$  atom cm<sup>-3</sup> (~  $4.0 \times 10^{-5}$  to  $4.0 \times 10^{-3}$  pptv) according to the review of Saiz-Lopez and von Glasow (2012). In light of the considerable variation of observed and model predicted Cl level, further study may be needed to comprehensively evaluate the significant role of Cl in the troposphere.



Figure 5. Monthly average of daily-maximum concentrations of OH and HO<sub>2</sub> in the BASE simulation, and changes due to chlorine (CL) and full halogen chemistry (HAL).



375

Figure 6. Monthly average of daily-maximum concentrations of NO<sub>3</sub> and Cl radical in the BASE simulation, and changes induced by chlorine (CL) and full halogen chemistry (HAL).

Fig. 5. and Fig. 6 demonstrate the monthly average of the daily-maximum concentrations of OH, HO<sub>2</sub>, NO<sub>3</sub>, and Cl in the BASE simulation and also the impact of chlorine (CL-BASE) and the halogen chemistry (HAL-BASE). The maximum values of OH, HO<sub>2</sub>, and Cl were predicted during the daytime but they peak at different hours with Cl in the early morning and OH and HO<sub>2</sub> later in the day, while the highest levels of NO<sub>3</sub> radical were simulated during night-time.

The monthly average of daily-maxima of OH, HO<sub>2</sub>, NO<sub>3</sub>, and Cl (Fig. 5 and Fig. 6) have similar

spatial pattern and higher concentrations (or changes of concentration) compared to those of the
monthly averages (Fig. 3 and Fig. 4). The monthly average of daily-maximum OH (HO<sub>2</sub> and NO<sub>3</sub>) radical is 3.7 (2.5 and 3.4) times of the monthly average OH (HO<sub>2</sub> and NO<sub>3</sub>) concentration in the BASE simulations, while both the monthly average of daily-maximum and monthly average Cl shows negligible values. The monthly average of daily-maximum changes of OH (HO<sub>2</sub>) concentration due to the chlorine and halogen chemistry has magnitude of -0.10 to 0.09
pptv (-4.0 to 0.8 pptv), which is wider than (the same as) that of the monthly averages, i.e., -0.020 to 0.03 pptv (-4.0 to 0.8 pptv). For the NO<sub>3</sub> (Cl) radicals, the magnitude of changes in monthly average is -20 to 1.0 pptv (0.0 to 0.0008 pptv) while that in monthly average of daily-maximums is -55 to 1.0 pptv (0.0 to 0.0055 pptv).

We also examine the diurnal variations of the four radicals in the BASE and HAL scenarios (Fig.
S3 and S4). Halogens have small effect on the diurnal pattern of OH. HO<sub>2</sub> is reduced by halogens especially in the mid-day. NO<sub>3</sub> radical is strongly decreased throughout the night after the addition of halogens. Cl atom is released by the halogen chemistry in the early morning. The significant effects of halogen chemistry on the diurnal variation of OH, HO<sub>2</sub>, NO<sub>3</sub>, and Cl radicals highlight the role of halogen chemistry in regulating the atmospheric oxidation capacity throughout the day with the highest effect on Cl in the early morning, maximum effects on OH and HO<sub>2</sub> in daytime, and largest effect on NO<sub>3</sub> at night.



### 400 **3.4 Impact of halogen chemistry on regulated gaseous air pollutants**

Figure 7. Monthly average NO<sub>2</sub> and O<sub>3</sub> concentration in the BASE simulation, and changes induced by chlorine (CL) and full halogen chemistry (HAL).

405

The monthly average modeled  $NO_2$  and  $O_3$ , two major gaseous air pollutants in Europe, and the effect of chlorine and halogen chemistry on the two regulated gaseous species were shown in Fig. 7. The BASE simulation produced many hot spots of  $NO_2$  over Europe, in the vicinity of the major cities and the ship trajectories. The chlorine chemistry slightly increases the level of  $NO_2$ 

(by up to 0.038 ppbv) in the majority of the domain since (1) the production and the subsequent photolysis of ClNO<sub>2</sub> recycles the NO<sub>x</sub> extending its lifetime, which increases both NO<sub>2</sub> and NO 410 (Fig. 7 and S5) and (2) the increased O<sub>3</sub> level (Fig. 7) enhances the transformation of NO to NO<sub>2</sub>, which increases NO<sub>2</sub> and decreases NO (Fig. 7 and S2). Some grid cells show a decrease of NO<sub>2</sub> (as much as 0.09 ppbv) because the enhanced oxidative capacity (Section 3.3) promotes the cleansing of NO<sub>2</sub> via OH forming HNO<sub>3</sub>. The full halogen chemistry enhances NO<sub>2</sub> (up to 0.20 ppbv) over the North Sea and the Mediterranean Sea and decreases NO<sub>2</sub> (as much as 1.7 ppbv) 415 in the most polluted hot spots. The increase of NO<sub>2</sub> occurs by the reactions of XO with NO. Meanwhile, in the most polluted regions, the NO-NO<sub>2</sub> balance is predominantly controlled by the reactions of NO with HO<sub>2</sub> and O<sub>3</sub>. With the decrease of HO<sub>2</sub> and O<sub>3</sub> due to the halogen chemistry, the transformation of NO to NO<sub>2</sub> is reduced which leads to decreasing NO<sub>2</sub> and increasing NO.

420 The monthly average O<sub>3</sub> concentration over Europe from the BASE simulation was relatively high (up to 57 ppbv) especially over the southern Europe, where higher temperature and more intensive radiation promote the formation of this secondary pollutant. Chlorine chemistry increases O<sub>3</sub> levels over the land of Europe with a maximum increment of 0.22 ppbv and decreases O<sub>3</sub> over the oceanic area as much as 0.76 ppbv. The full halogen chemistry decreases  $O_3$  throughout the domain with a maximum reduction of 10 ppby. On average, the halogen 425 chemistry reduces O<sub>3</sub> concentration by more than 3.0 ppbv in coastal Europe and by over 2.0 ppbv over western and central Europe (nearly one thousand kilometers from the ocean). Our model simulation highlights the fact that halogen chemistry has a large impact on O<sub>3</sub> concentrations over the oceanic areas and a moderate impact on O<sub>3</sub> over coastal and continental regions of Europe. 430

Muñiz-Unamunzaga et al. (2018) reported a decrease of -2.0 ppbv O<sub>3</sub> in the inland areas of the western US (several hundreds of kilometers from the ocean) and a reduction of 2.5 to 5.0 ppbv O<sub>3</sub> in the coastal regions due to the full-halogen chemistry. Sarwar et al. (2015) suggested that the inclusion of halogen processes reduced O<sub>3</sub> concentrations by 2.0 to 4.0 ppbv over most of the terrestrial regions in the North Hemisphere, and over 6.0 ppbv in some coastal areas. Sherwen et al. (2017) used a revised version of GEOS-Chem (Sherwen et al., 2016) with

halogen chemistry to show substantial reductions in  $O_3$  over Europe with an average reduction of 13.5 ppbv in the domain and a maximum of 28.9 ppbv in some locations.

### 440 **3.5 Implications for policy assessment**

The current air quality management in Europe has two main objectives: (1) to protect human health and (2) to protect the environment. While many plans and measures have prioritized Particulate Matter (PM) or NO<sub>2</sub>, policies to reduce O<sub>3</sub> concentrations are still needed (EEA, 2018a). The World Health Organization (WHO) Air Quality Guidelines value for O<sub>3</sub> (maximum daily 8-hour mean of 100  $\mu$ g·m<sup>-3</sup>) was exceeded in 96% of all the reporting stations in Europe, 445 although this is especially true for the areas near the Mediterranean Sea. According to the European Environment Agency latest report, 12% of the EU-28 urban population is exposed to O<sub>3</sub> concentrations above the European Union target value threshold (maximum daily 8-hour mean of 120 µg·m<sup>-3</sup> not to be exceeded on more than 25 days/year, as set out by the Directive 2008/50/EC) in 2016. Apart from significant potential health effects (Jerrett et al., 2009; Malley 450 et al., 2017),  $O_3$  is also known to have a negative impact on vegetation (Mills et al., 2011). The target value for the protection of vegetation (18,000 µg·m<sup>-3</sup>·h accumulated over May to July), based on the Accumulated Ozone exposure over a Threshold of 40 ppbv index (AOT40 index), was exceeded in about 31 % of all agricultural land in all European countries. The critical level for this pollutant (10,000 µg·m<sup>-3</sup>·h accumulated over April to September) was exceeded in 60% 455 of the total forest area of the continent in 2016 (EEA, 2018a). Here we use the simulation results (BASE and HAL) in July to calculate the accumulated O<sub>3</sub> and the difference between two scenarios. We note that the accumulated  $O_3$  is considerably reduced (~15% along the coast) after the addition of halogens (Fig. 8).



Figure 8. Accumulated O<sub>3</sub> in July in the BASE and HAL simulations, and absolute and relative changes between the two simulations.

We find that halogen chemistry strongly affects ambient O<sub>3</sub> concentration and may need to be considered in the formulation of plans and strategies for O<sub>3</sub> non-attainment areas. We see differences between BASE and HAL simulations (over land in July 2016) as high as 12% and 36% for the number of days with daily maximum 8 h O<sub>3</sub> over 120 µg·m<sup>-3</sup> and the monthly average daily maximum 8 h O<sub>3</sub> level, respectively (Fig. S6 and Fig. S7). Furthermore, we notice strong regional differences, mainly between coastal and inland areas. The considerable effect of halogen chemistry on air quality implies the need to improve the robustness and accuracy of modeling tools to design customized policies to control O<sub>3</sub>.

In Section 3.3 and 3.4, we have also discussed the effect of halogen chemistry on the partitioning of OH/HO<sub>2</sub> and NO/NO<sub>2</sub>. The budgets of HO<sub>x</sub> and NO<sub>x</sub> are key parameters to accurately simulate the formation of O<sub>3</sub> and its response to the reductions of the precursors, namely NO<sub>x</sub> and VOCs (e.g., Li et al., 2018). Air quality models are predominantly used to formulate air pollution control policy by examining the responses of O<sub>3</sub> levels to various

reduction rates of  $NO_x$  and/or VOCs. The models which do not include the comprehensive halogen chemistry can potentially lead to different  $O_3$  concentration responses to  $NO_x$  and/or VOC emission changes in Europe.

This study also demonstrates that chlorine chemistry affects the formation of O<sub>3</sub>. The current policy is only designed to control the long-lived chlorinated species (Hossaini et al., 2015), but not reactive chlorine species, e.g., HCl, chloride, and short-lived chlorocarbons, from the coal burning, biomass burning, and industrial activities. The coal-fired power plants in EU (EEA, 2018b; Kuklinska et al., 2015) can potentially provide chlorine sources, making the implications

#### 485

### 4. Conclusion

of halogen chemistry even more relevant.

We applied the CMAQ model with comprehensive halogen chemistry (Cl, Br and I) to conduct high-resolution simulations for examining the impact of halogen chemistry on air quality over Europe.

490 The comparison of model results with observations from over 400 monitoring sites indicates that the CMAQ model is capable of reproducing the concentrations and temporal variations of air pollutants over Europe and can be employed to study the impact of halogen chemistry in Europe. The comparison of predicted halogen species concentrations with measurements suggests that the CMAQ model is able to predict observed levels of chlorine and iodine species although it underestimates bromine species.

The chlorine chemistry enhances the atmospheric oxidation capacity by significantly increasing the level of Cl radical and affects the levels of OH, HO<sub>2</sub>, NO<sub>3</sub>, O<sub>3</sub>, and NO<sub>2</sub>. The combined halogen chemistry marginally increases the level of OH and reduces HO<sub>2</sub>, NO<sub>3</sub>, and O<sub>3</sub>. The impact of halogen chemistry on ambient concentration of NO<sub>2</sub> is smaller but non-negligible.

500 Halogen chemistry significantly influences the atmospheric oxidation capacity throughout the day by imposing the highest effect on Cl in the early morning, maximum effects on OH and HO<sub>2</sub>

in daytime, and largest effect on  $NO_3$  at night. Halogen chemistry can have a strong influence on atmospheric composition over oceanic and coastal regions but also some noticeable impacts over continental Europe. This study highlights the potential benefit of incorporating halogen chemistry into air quality models for policy development.

Although the incorporation of the halogen chemistry may improve the capabilities of 3D Eulerian chemical transport models, we acknowledge that large uncertainties still exist in the assessment of halogen chemistry impact due to emission inventories (e.g., chlorine emission inventory; Wang et al., 2019), model configuration (e.g., grid size; Sommariva et al., 2018), chemical mechanism (e.g., photolysis rate of iodine oxides, recycling rate of halogen species on aerosol; Simpson et al., 2015), etc. Further field, laboratory, and theoretical studies are needed to constraint modeling studies for evaluating the impacts of halogen chemistry on air quality and for assessing air quality policy implications.

515 **Data availability.** The data used and demonstrated in this study are available upon the request to the corresponding author.

*Author contributions*. AS-L designed research. GS and BG conducted the CMAQ modeling. RB, DP, JD, GS, BG, QL, and AS-L analyzed the results. QL, RB, GS, and AS-L wrote the paper with contributions from all authors.

520 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgment.* This study has received funding from the European Research Council Executive Agency under the European Union's Horizon 2020 Research and Innovation programme (Project 'ERC-2016-COG 726349 CLIMAHAL').

# DISCLAIMER

505

510

525 The views expressed in this paper are those of the authors and do not necessarily represent the views or policies of the U.S. EPA.

### **Reference:**

530

555

Allan, B. J., McFiggans, G., Plane, J. M. C. and Coe, H.: Observations of iodine monoxide in the remote marine boundary layer, J. Geophys. Res. Atmos., 105(D11), 14363–14369, doi:10.1029/1999JD901188, 2000.

Appel, K. W., Chemel, C., Roselle, S. J., Francis, X. V, Hu, R.-M., Sokhi, R. S., Rao, S. T. and Galmarini, S.: Examination of the Community Multiscale Air Quality (CMAQ) model performance over the North American and European domains, Atmos. Environ., 53, 142–155, doi:https://doi.org/10.1016/j.atmosenv.2011.11.016, 2012.

- Appel, K. W., Pouliot, G. A., Simon, H., Sarwar, G., Pye, H. O. T., Napelenok, S. L., Akhtar, F. and Roselle, S. J.: Evaluation of dust and trace metal estimates from the Community Multiscale Air Quality (CMAQ) model version 5.0, Geosci. Model Dev., 6(4), 883–899, doi:10.5194/gmd-6-883-2013, 2013.
- Appel, K. W., Napelenok, S. L., Foley, K. M., Pye, H. O. T., Hogrefe, C., Luecken, D. J.,
  Bash, J. O., Roselle, S. J., Pleim, J. E., Foroutan, H., Hutzell, W. T., Pouliot, G. A.,
  Sarwar, G., Fahey, K. M., Gantt, B., Gilliam, R. C., Heath, N. K., Kang, D., Mathur, R.,
  Schwede, D. B., Spero, T. L., Wong, D. C. and Young, J. O.: Description and evaluation
  of the Community Multiscale Air Quality (CMAQ) modeling system version 5.1, Geosci.
  Model Dev., 10(4), 1703–1732, doi:10.5194/gmd-10-1703-2017, 2017.
- Bannan, T. J., Murray Booth, A., Bacak, A., Muller, J. B. A., Leather, K. E., Le Breton, 545 M., Jones, B., Young, D., Coe, H., Allan, J., Visser, S., Slowik, J. G., Furger, M., Prévôt, A. S. H., Lee, J., Dunmore, R. E., Hopkins, J. R., Hamilton, J. F., Lewis, A. C., Whalley, L. K., Sharp, T., Stone, D., Heard, D. E., Fleming, Z. L., Leigh, R., Shallcross, D. E. and Percival, C. J.: The first UK measurements of nitryl chloride using a chemical ionization mass spectrometer in central London in the summer of 2012, and an investigation of the 550 role of Cl oxidation, J. Geophys. Res., 120(11), 5638-5657, atom doi:10.1002/2014JD022629, 2015.

Bitter, M., Ball, S. M., Povey, I. M. and Jones, R. L.: A broadband cavity ringdown spectrometer for in-situ measurements of atmospheric trace gases, Atmos. Chem. Phys., 5(9), 2547–2560, doi:10.5194/acp-5-2547-2005, 2005.

Borge, R., Alexandrov, V., José del Vas, J., Lumbreras, J. and Rodríguez, E.: A comprehensive sensitivity analysis of the WRF model for air quality applications over the Iberian Peninsula, Atmos. Environ., 42(37), 8560–8574, doi:https://doi.org/10.1016/j.atmosenv.2008.08.032, 2008a.

Borge, R., Lumbreras, J. and Rodríguez, E.: Development of a high-resolution emission

inventory for Spain using the SMOKE modelling system: A case study for the years 2000 and 2010, Environ. Model. Softw., 23(8), 1026–1044, doi:https://doi.org/10.1016/j.envsoft.2007.11.002, 2008b.

Byun, D. and Schere, K. L.: Review of the Governing Equations, Computational Algorithms, and Other Components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, Appl. Mech. Rev., 59(2), 51–77 [online] Available from: http://dx.doi.org/10.1115/1.2128636, 2006.

Ching, J. and Byun, D.: Science Algorithms of the EPA MODELS-3 Community Multiscale Air Quality (CMAQ) Modelling System, Washington, D.C., 1999.

570 Commane, R., Seitz, K., Bale, C. S. E., Bloss, W. J., Buxmann, J., Ingham, T., Platt, U., Pöhler, D. and Heard, D. E.: Iodine monoxide at a clean marine coastal site: observations of high frequency variations and inhomogeneous distributions, Atmos. Chem. Phys., 11(13), 6721–6733, doi:10.5194/acp-11-6721-2011, 2011.

EEA (European Environmental Agency): Air Quality in Europe – 2018 report. [online]
 Available from: https://www.eea.europa.eu/publications/air-quality-in-europe-2018, 2018a.

EEA (European Environmental Agency): European Union emission inventory report 1990–2016 under the UNECE Convention on Long-range Transboundary Air Pollution (LRTAP)., 2018b.

- 580 Eger, P. G., Friedrich, N., Schuladen, J., Shenolikar, J., Fischer, H., Tadic, I., Harder, H., Martinez, M., Rohloff, R., Tauer, S., Fachinger, F., Drewnick, F., Brooks, J., Darbyshire, E., Sciare, J., Pikridas, M., Lelieveld, J. and Crowley, J. N.: Shipborne measurements of ClNO2 in the Mediterranean Sea and around the Arabian Peninsula during summer, Atmos. Chem. Phys. Discuss., (June), 1–37, doi:10.5194/acp-2019-531, 2019.
- 585 EMEP/CEIP: Present state of emissions as used in EMEP models, (Available at: <u>http://www.ceip.at/webdab\_emepdatabase/emissions\_emepmodels)</u>, 2014.

Farman, J. C., Gardiner, B. G. and Shanklin, J. D.: Large losses of total ozone in Antarctica reveal seasonal ClOx/NOx interaction, Nature, 315(6016), 207–210, doi:10.1038/315207a0, 1985.

Fernandez, R. P., Salawitch, R. J., Kinnison, D. E., Lamarque, J.-F. F. and Saiz-Lopez,
 A.: Bromine partitioning in the tropical tropopause layer: Implications for stratospheric injection, Atmos. Chem. Phys., 14(24), 13391–13410, doi:10.5194/acp-14-13391-2014, 2014.

Foley, K. M., Roselle, S. J., Appel, K. W., Bhave, P. V., Pleim, J. E., Otte, T. L., Mathur,

- 595 R., Sarwar, G., Young, J. O., Gilliam, R. C., Nolte, C. G., Kelly, J. T., Gilliland, A. B. and Bash, J. O.: Incremental testing of the Community Multiscale Air Quality (CMAQ) modeling system version 4.7, Geosci. Model Dev., 3(1), 205–226, doi:10.5194/gmd-3-205-2010, 2010.
- Furneaux, K. L., Whalley, L. K., Heard, D. E., Atkinson, H. M., Bloss, W. J., Flynn, M.
  J., Gallagher, M. W., Ingham, T., Kramer, L., Lee, J. D., Leigh, R., McFiggans, G. B.,
  Mahajan, A. S., Monks, P. S., Oetjen, H., Plane, J. M. C. and Whitehead, J. D.:
  Measurements of iodine monoxide at a semi polluted coastal location, Atmos. Chem.
  Phys., 10(8), 3645–3663, doi:10.5194/acp-10-3645-2010, 2010.
- Gantt, B., Sarwar, G., Xing, J., Simon, H., Schwede, D., Hutzell, W. T., Mathur, R. and
   Saiz-Lopez, A.: The Impact of Iodide-Mediated Ozone Deposition and Halogen
   Chemistry on Surface Ozone Concentrations Across the Continental United States,
   Environ. Sci. Technol., 51(3), 1458–1466, doi:10.1021/acs.est.6b03556, 2017.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K. and Wang, X.: The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic emissions, Geosci. Model Dev., 5(6), 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.

610

Holla, R., Schmitt, S., Frieß, U., Pöhler, D., Zingler, J., Corsmeier, U. and Platt, U.: Vertical distribution of BrO in the boundary layer at the Dead Sea, Environ. Chem., 12(4), 438–460 [online] Available from: https://doi.org/10.1071/EN14224, 2015.

Hossaini, R., Chipperfield, M. P., Saiz-Lopez, A., Harrison, J. J., Von Glasow, R., Sommariva, R., Atlas, E., Navarro, M., Montzka, S. A., Feng, W., Dhomse, S., Harth, C., Mühle, J., Lunder, C., O'Doherty, S., Young, D., Reimann, S., Vollmer, M. K., Krummel, P. B. and Bernath, P. F.: Growth in stratospheric chlorine from short-lived chemicals not controlled by the Montreal Protocol, Geophys. Res. Lett., 42(11), 4573– 4580, doi:10.1002/2015GL063783, 2015.

Hossaini, R., Chipperfield, M. P., Saiz-Lopez, A., Fernandez, R., Monks, S., Feng, W., Brauer, P. and Von Glasow, R.: A global model of tropospheric chlorine chemistry: Organic versus inorganic sources and impact on methane oxidation, J. Geophys. Res., 121(23), 14,271-14,297, doi:10.1002/2016JD025756, 2016.

Jerrett, M., Burnett, R. T., Pope, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle,
E. and Thun, M.: Long-Term Ozone Exposure and Mortality, N. Engl. J. Med., 360(11),
1085–1095, doi:10.1056/NEJMoa0803894, 2009.

Jung, J., Lee, J., Kim, B. and Oh, S.: Seasonal variations in the NO2 artifact from

chemiluminescence measurements with a molybdenum converter at a suburban site in
 Korea (downwind of the Asian continental outflow) during 2015–2016. Atmos. Environ.,
 165, 290-300, 2017.

Keene, W., Khalil, M. A. K., Erickson, D., McCulloch, A., Graedel, T. E., Lobert, J. M., Aucott, M. L., Gong, S. L., Harper, D. B., Kleiman, G., Midgley, P., Moore, R. M., Seuzaret, C., Sturges, W. T., Benkovitz, C. M., Koropalov, V., Barrie, L. A., and Li, Y. F.: Composite global emissions of reactive chlorine from anthropogenic and natural sources: Reactive Chlorine Emissions Inventory, J. Geophys. Res.-Atmos., 104, 8429–8440, 1999.

635

650

Kuklinska, K., Wolska, L. and Namiesnik, J.: Air quality policy in the U.S. and the EU – a review, Atmos. Pollut. Res., 6(1), 129–137, doi:10.5094/APR.2015.015, 2015.

de la Paz, D., Borge, R. and Martilli, A.: Assessment of a high resolution annual WRF-BEP/CMAQ simulation for the urban area of Madrid (Spain), Atmos. Environ., 144, 282–296, doi:https://doi.org/10.1016/j.atmosenv.2016.08.082, 2016.

Li, Q., Zhang, L., Wang, T., Tham, Y. J., Ahmadov, R., Xue, L., Zhang, Q. and Zheng, J.: Impacts of heterogeneous uptake of dinitrogen pentoxide and chlorine activation on ozone and reactive nitrogen partitioning: improvement and application of the WRF-Chem model in southern China, Atmos. Chem. Phys., 16(23), 14875–14890, doi:10.5194/acp-16-14875-2016, 2016.

Li, Q., Zhang, L., Wang, T., Wang, Z., Fu, X. and Zhang, Q.: "New" Reactive Nitrogen Chemistry Reshapes the Relationship of Ozone to Its Precursors, Environ. Sci. Technol., 52(5), 2810–2818, doi:10.1021/acs.est.7b05771, 2018.

van Loon, M., Vautard, R., Schaap, M., Bergström, R., Bessagnet, B., Brandt, J., Builtjes,
P. J. H., Christensen, J. H., Cuvelier, C., Graff, A., Jonson, J. E., Krol, M., Langner, J.,
Roberts, P., Rouil, L., Stern, R., Tarrasón, L., Thunis, P., Vignati, E., White, L. and
Wind, P.: Evaluation of long-term ozone simulations from seven regional air quality
models and their ensemble, Atmos. Environ., 41(10), 2083–2097,
doi:https://doi.org/10.1016/j.atmosenv.2006.10.073, 2007.

Mahajan, A. S., Oetjen, H., Lee, J. D., Saiz-Lopez, A., McFiggans, G. B. and Plane, J. M. C.: High bromine oxide concentrations in the semi-polluted boundary layer, Atmos. Environ., 43(25), 3811–3818, doi:10.1016/j.atmosenv.2009.05.033, 2009.

Malley, C. S., Henze, D. K., Kuylenstierna, J. C. I., Vallack, H. W., Davila, Y.,
 Anenberg, S. C., Turner, M. C. and Ashmore, M. R.: Updated Global Estimates of
 Respiratory Mortality in Adults ≥30Years of Age Attributable to Long-Term Ozone

Exposure, Environ. Health Perspect., 125(8), 87021, doi:10.1289/EHP1390, 2017.

- Mathur, R., Xing, J., Gilliam, R., Sarwar, G., Hogrefe, C., Pleim, J., Pouliot, G., Roselle,
  S., Spero, T. L., Wong, D. C. and Young, J.: Extending the Community Multiscale Air Quality (CMAQ) Modeling System to Hemispheric Scales: Overview of Process Considerations and Initial Applications, Atmos. Chem. Phys., 17, 12449–12474, doi:10.5194/acp-17-12449-2017, 2017.
- Matveev, V., Peleg, M., Rosen, D., Tov-Alper, D. S., Hebestreit, K., Stutz, J., Platt, U.,
  Blake, D. and Luria, M.: Bromine oxide—ozone interaction over the Dead Sea, J. Geophys. Res. Atmos., 106(D10), 10375–10387, doi:10.1029/2000JD900611, 2001.
- Mills, G., Pleijel, H., Braun, S., Büker, P., Bermejo, V., Calvo, E., Danielsson, H., Emberson, L., Fernández, I. G., Grünhage, L., Harmens, H., Hayes, F., Karlsson, P.-E. and Simpson, D.: New stomatal flux-based critical levels for ozone effects on vegetation, Atmos. Environ., 45(28), 5064–5068, doi:https://doi.org/10.1016/j.atmosenv.2011.06.009, 2011.

Molina, M. J. and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone, Nature, 249(5460), 810–812, doi:10.1038/249810a0, 1974.

- Muñiz-Unamunzaga, M., Borge, R., Sarwar, G., Gantt, B., de la Paz, D., Cuevas, C. A. and Saiz-Lopez, A.: The influence of ocean halogen and sulfur emissions in the air quality of a coastal megacity: The case of Los Angeles, Sci. Total Environ., 610–611, 1536–1545, doi:10.1016/j.scitotenv.2017.06.098, 2018.
- Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M.,
  Sommariva, R., Bates, T. S., Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H.,
  Burkholder, J. B., Talukdar, R. K., Meagher, J., Fehsenfeld, F. C. and Brown, S. S.: High
  levels of nitryl chloride in the polluted subtropical marine boundary layer, Nat. Geosci.,
  1(5), 324–328, doi:10.1038/ngeo177, 2008.
- Peters, C., Pechtl, S., Stutz, J., Hebestreit, K., Hönninger, G., Heumann, K. G., Schwarz,
  A., Winterlik, J. and Platt, U.: Reactive and organic halogen species in three different European coastal environments, Atmos. Chem. Phys., 5(12), 3357–3375, doi:10.5194/acp-5-3357-2005, 2005.

695

Phillips, G. J., Tang, M. J., Thieser, J., Brickwedde, B., Schuster, G., Bohn, B., Lelieveld,
J. and Crowley, J. N.: Significant concentrations of nitryl chloride observed in rural continental Europe associated with the influence of sea salt chloride and anthropogenic emissions, Geophys. Res. Lett., 39(10), 1–5, doi:10.1029/2012GL051912, 2012.

Saiz-Lopez, A. and von Glasow, R.: Reactive halogen chemistry in the troposphere, Chem. Soc. Rev., 41(19), 6448–6472, doi:10.1039/c2cs35208g, 2012.

Saiz-Lopez, A., Plane, J. M. C. and Shillito, J. A.: Bromine oxide in the mid-latitude marine boundary layer, Geophys. Res. Lett., 31(3), 4–7, doi:10.1029/2003GL018956, 2004.

Saiz-Lopez, A., Fernandez, R. P., Ordóñez, C., Kinnison, D. E., Martín, J. C. G., Lamarque, J. F. and Tilmes, S.: Iodine chemistry in the troposphere and its effect on ozone, Atmos. Chem. Phys., 14(23), 13119–13143, doi:10.1177/0964663912467814, 2014.

705

715

730

Sarwar, G., Simon, H., Bhave, P. and Yarwood, G.: Examining the impact of heterogeneous nitryl chloride production on air quality across the United States, Atmos. Chem. Phys., 12(14), 6455–6473, doi:10.5194/acp-12-6455-2012, 2012.

Sarwar, G., Simon, H., Xing, J. and Mathur, R.: Importance of tropospheric CINO2 chemistry across the Northern Hemisphere, Geophys. Res. Lett., 41(11), 4050–4058, doi:10.1002/2014GL059962, 2014.

Sarwar, G., Gantt, B., Schwede, D., Foley, K., Mathur, R. and Saiz-Lopez, A.: Impact of Enhanced Ozone Deposition and Halogen Chemistry on Tropospheric Ozone over the Northern Hemisphere, Environ. Sci. Technol., 49(15), 9203–9211, doi:10.1021/acs.est.5b01657, 2015.

Sarwar, G., Gantt, B., Foley, K., Fahey, K., Spero T. L., Kang, D., Mathur, R., Hosein F., Xing, J., Sherwen, T., Saiz-Lopez, A.: Influence of bromine and iodine chemistry on annual, seasonal, diurnal, and background ozone: CMAQ simulations over the Northern Hemisphere, Atmospheric Environment, 213, 395-404, 2019.

- Sherwen, T., Schmidt, J. A., Evans, M. J., Carpenter, L. J., Großmann, K., Eastham, S. D., Jacob, D. J., Dix, B., Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S. and Ordóñez, C.: Global impacts of tropospheric halogens (Cl, Br, I) on oxidants and composition in GEOS-Chem, Atmos. Chem. Phys., 16(18), 12239–12271, doi:10.5194/acp-16-12239-2016, 2016.
- Sherwen, T., Evans, M. J., Sommariva, R., Hollis, L. D. J., Ball, S. M., Monks, P. S., Reed, C., Carpenter, L. J., Lee, J. D., Forster, G., Bandy, B., Reeves, C. E. and Bloss, W. J.: Effects of halogens on European air-quality, Faraday Discuss., 200, 75–100, doi:10.1039/c7fd00026j, 2017.

Sillman, S.: The relation between ozone, NO(x) and hydrocarbons in urban and polluted rural environments, Atmos. Environ., 33(12), 1821–1845, doi:10.1016/S1352-

2310(98)00345-8, 1999.

740

745

Simpson, W. R., Brown, S. S., Saiz-Lopez, A., Thornton, J. A. and Von Glasow, R.: Tropospheric Halogen Chemistry: Sources, Cycling, and Impacts, Chem. Rev., 115(10), 4035–4062, doi:10.1021/cr5006638, 2015.

735 Skamarock, W. C. and Klemp, J. B.: A time-split nonhydrostatic atmospheric model for weather research and forecasting applications, J. Comput. Phys., 227(7), 3465–3485, doi:https://doi.org/10.1016/j.jcp.2007.01.037, 2008.

Solazzo, E., Hogrefe, C., Colette, A., Garcia-Vivanco, M. and Galmarini, S.: Advanced error diagnostics of the CMAQ and Chimere modelling systems within the AQMEII3 model evaluation framework, Atmos. Chem. Phys., 17(17), 10435–10465, doi:10.5194/acp-17-10435-2017, 2017.

Sommariva, R., Hollis, L. D. J., Sherwen, T., Baker, A. R., Ball, S. M., Bandy, B. J., Bell, T. G., Chowdhury, M. N., Cordell, R. L., Evans, M. J., Lee, J. D., Reed, C., Reeves, C. E., Roberts, J. M., Yang, M. and Monks, P. S.: Seasonal and geographical variability of nitryl chloride and its precursors in Northern Europe, Atmos. Sci. Lett., 19(8), 1–10, doi:10.1002/asl.844, 2018.

Stone, D., Sherwen, T., Evans, M. J., Vaughan, S., Ingham, T., Whalley, L. K., Edwards, P. M., Read, K. A., Lee, J. D., Moller, S. J., Carpenter, L. J., Lewis, A. C. and Heard, D. E.: Impacts of bromine and iodine chemistry on tropospheric OH and HO2: Comparing observations with box and global model perspectives, Atmos. Chem. Phys., 18(5), 3541–3561, doi:10.5194/acp-18-3541-2018, 2018.

Tas, E., Peleg, M., Pedersen, D. U., Matveev, V., Biazar, A. P. and Luria, M.: Measurement-based modeling of bromine chemistry in the boundary layer: 1. Bromine chemistry at the Dead Sea, Atmos. Chem. Phys., 6(12), 5589–5604, doi:10.5194/acp-6-5589-2006, 2006.

- Tham, Y. J., Wang, Z., Li, Q., Yun, H., Wang, W., Wang, X., Xue, L., Lu, K., Ma, N., Bohn, B., Li, X., Kecorius, S., Größ, J., Shao, M., Wiedensohler, A., Zhang, Y. and Wang, T.: Significant concentrations of nitryl chloride sustained in the morning: Investigations of the causes and impacts on ozone production in a polluted region of northern China, Atmos. Chem. Phys., 16(23), 14959–14977, doi:10.5194/acp-16-14959-2016, 2016.
  - Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., D ube, W. P., Wolfe, G. M., Quinn, P. K., Middlebrook, A. M., Alexander, B. and Brown, S. S.: A large atomic chlorine source inferred from mid-continental reactive nitrogen chemistry, Nature, 464(7286), 271–274, doi:10.1038/nature08905, 2010.

UNC (University of North Carolina at Chapel Hill): SMOKE v3.6.5 User's Manual. 765 Institute for the Environment. Available at: https://www.cmascenter.org/smoke/documentation/3.6.5/html, 2015. Wang, T., Tham, Y. J., Xue, L., Li, Q., Zha, Q., Wang, Z., Poon, S. C. N., Dubé, W. P., Blake, D. R., Louie, P. K. K., Luk, C. W. Y., Tsui, W. and Brown, S. S.: Observations of 770 nitryl chloride and modeling its source and effect on ozone in the planetary boundary of southern China, J. Geophys. Res. Atmos., 121(5), 2476-2489, layer

doi:10.1002/2015JD024556, 2016.

Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B.,
Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A.,
Huey, G. L. and Liao, H.: The role of chlorine in global tropospheric chemistry, Atmos. Chem.
Phys., 19(6), 3981–4003, doi:10.5194/acp-19-3981-2019, 2019.