

The sensitivity of PM_{2.5} acidity to meteorological parameters and chemical composition changes: 10-year records from six Canadian monitoring sites

Ye Tao¹, Jennifer G. Murphy²

5 ¹Department of Physical and Environmental Sciences, University of Toronto Scarborough, Toronto, ON, M1C 1A4, Canada

²Department of Chemistry, University of Toronto, Toronto, ON, M5S 3H6, Canada

Corresponding to: Jennifer G. Murphy (jmurphy@chem.utoronto.ca)

Abstract. Aerosol pH is difficult to measure directly but can be calculated if the chemical composition is known with sufficient accuracy and precision to calculate the aerosol water content and the H⁺ concentration through the equilibrium among acids and their conjugate bases. In practical terms, simultaneous measurements of at least one semi-volatile constituents, e.g. NH₃ or HNO₃, are required to provide a constraint on the calculation of pH. Long-term records of aerosol pH are scarce due to the limited monitoring of NH₃ in conjunction with PM_{2.5}. In this study, 10-year (2007-2016) records of pH of PM_{2.5} at six eastern Canadian sites were calculated using the E-AIM II model with the input of gaseous NH₃, gaseous HNO₃ and major water-soluble inorganic ions in PM_{2.5} provided by Canada's National Air Pollution Surveillance (NAPS) Program. Clear seasonal cycles of aerosol pH were found with lower pH (~2) in summer and higher pH (~3) in winter consistently across all six sites, while the day-to-day variations of aerosol pH were higher in winter compared to summer. Tests of the sensitivity of aerosol pH to meteorological parameters demonstrate that the changes in ambient temperature largely drive the seasonal cycle of aerosol pH. The sensitivity of pH to chemical composition shows that pH has different responses to the changes in chemical composition in different seasons. During summertime, aerosol pH was mainly determined by temperature with limited impact from changes in NH_x or sulfate concentrations. However, in wintertime, both meteorological parameters and chemical composition contribute to the variations in aerosol pH, resulting in the larger variation during wintertime. This study reveals that the sensitivity of aerosol pH to chemical composition is distinctly different under different meteorological conditions and needs to be carefully examined for any particular region.

10
15
20

Key words: PM_{2.5}, aerosol pH, NH_x phase partitioning, meteorological parameters, E-AIM

25 1. Introduction

Aerosol acidity is among the most important parameters for atmospheric particulate chemistry as it has a large impact on both gas/particle partitioning of semi-volatile ionizable components and rates of many reactions occurring in aerosol liquid water (Seinfeld and Pandis, 2006; Losey et al., 2016; Guo et al., 2017a; Nah et al., 2018; Freedman et al., 2019). Laboratory and

field studies have also shown that enhanced acidity in aerosol can increase the formation of secondary organic aerosol (Surratt et al., 2007; Zhang et al., 2007b), one of the major components responsible for particulate air pollution (Zhang et al., 2007a; Huang et al., 2014). Studies have shown that highly acidic particles can have adverse health effects related to respiratory diseases (Utell et al., 1983; Dockery et al., 1996). More acidic particles can enhance the solubility of several trace metals in fine particulate matter (PM_{2.5}) to induce oxidative potential inside the human body as an indirect health effect of acidic particles (Fang et al., 2017).

pH is the parameter frequently used to describe the acidity of aerosol aqueous phase and is calculated as the negative logarithm of hydrogen ion activity. Few methods can directly measure the pH of aerosol (Rindelaub et al., 2016; Craig et al., 2018; Wei et al., 2018), so aerosol acidity is usually presented using indirect proxies such as ion balance (cation-to-anion ratio), neutralization ratio ($[\text{NH}_4^+]/(2[\text{SO}_4^{2-}]+[\text{NO}_3^-])$) and strong acidity ($\Sigma\text{cation}-\Sigma\text{anion}$) (Yao et al., 2006; Du et al., 2010; Zhou et al., 2017). However, these indirect methods can lead to substantial uncertainty in acidity assessment (Hennigan et al., 2015; Guo et al., 2016; Murphy et al., 2017). Calculation of pH through thermodynamic modelling such as E-AIM (Clegg et al., 1998) and ISORROPIA II (Fountoukis and Nenes, 2007) with the input of reliable chemical compositions of the aerosol and at least one semi-volatile gases and meteorological data has been shown to be a more rigorous approach to calculate the pH of aerosol liquid water. Murphy et al. (2017) and Song et al. (2018) both showed that the constraint from phase partitioning of NH₃/NH₄⁺ should be included in calculations using aerosol thermodynamic models to get reliable pH calculations, which indicates that NH₃ observation can greatly improve the reliability of the aerosol acidity assessment.

However, long-term observations of NH₃ with corresponding particulate matter composition measurements are very scarce. Some of the few long-term monitoring networks are NAMN in the UK, SEARCH in the US and NAPS in Canada. In the UK, 30 of the NAMN (National Ammonia Monitoring Network) sites have been providing monthly-integrated NH₃ and particulate inorganic ionic composition data since 2000 (Tang et al., 2018). SEARCH (Southeastern Aerosol Research and CHaracterization Network) is one long-term monitoring programs of NH₃ in the United States, which has both 24-hour average gas phase NH₃ and particle phase ionic species measurements at eight sites located in Southeastern United States with the sampling frequency of every three days. In Canada, the National Air Pollution Surveillance (NAPS) program is the only program employing the long-term measurement of 24-hour average NH₃ concentrations with a sampling frequency of every three days at twelve monitoring sites where PM_{2.5} constituents are also measured.

Several studies have examined the response of pH to changes in chemical composition with conceptual modelling. Weber et al. (2016) showed that in summertime, aerosol tends to remain highly acidic with the reduction of sulfate and the increase of ammonia-to-sulfate ratio. The modelling results of Guo et al. (2017b) showed that for wintertime meteorological conditions aerosol pH has similar sensitivity to NH₃ concentrations in China and the eastern US under fixed temperature and relative

humidity. These two sensitivity tests of aerosol pH suggest that aerosol pH seems to have weak response to chemical composition changes. However, these studies both carried out conceptual modelling or calculations under fixed meteorological parameters. The time series of aerosol pH in six regions within the United States calculated by Lawal et al. (2018) showed that pH has seasonal variation that depends on the region, suggesting that changes in meteorological conditions can also contribute to pH changes. Thus, the relative contributions of changes in meteorological conditions and chemical composition to variability in pH are still unclear and require systematic studies. As a result, in this study, we focus on the long-term variations of aerosol pH at six eastern Canadian sites (four urban and two rural) along with conceptual modellings of the sensitivity of aerosol pH to meteorological parameters and chemical composition changes to assess the major factors determining aerosol pH, and its variability, in different sites and seasons.

2. Data sources and analysis method

2.1 NAPS database and chosen study sites

All the PM_{2.5} composition and trace gas data used in this study were downloaded from the NAPS program website (www.ec.gc.ca/rnsps-naps/). The NAPS program provides long-term monitoring of air quality-relevant pollutants at various sites across Canada. Altogether, there are twelve sites with comprehensive gas phase and particle phase composition measurements, including four coastal sites (Halifax, Saint John, Burnaby and Abbotsford), one central site (Edmonton) and seven eastern sites (Toronto, Simcoe, Hamilton, Windsor, Ottawa, Montreal and St Anicet). 24-hour ambient samples of gaseous components (including NH₃ and HNO₃) and the dominant inorganic components of PM_{2.5} were taken from 00:00-24:00 generally every three days.

Six sites in the provinces of Ontario and Quebec were chosen for this study (Figure 1), including Toronto (the largest city in Canada), Ottawa, two other metropolitan cities (Windsor and Montreal) and two sites in rural regions (Simcoe and St Anicet). The details of these monitoring sites including their longitudes and latitudes, NAPS monitoring station codes, and duration of the data record used for this study can be found in Table S1. These sites were chosen because 1) they have long-term (at least 8 year) records of gaseous and PM_{2.5} chemical components concentrations; 2) they are located in the central region of North America with little influence of sea salt (Vet et al., 2014) and therefore the acidity of PM_{2.5} is mainly dominated by the chemistry of sulfate, total nitrate and total ammonia (Young et al., 2013; Allen et al., 2015; Hennigan et al., 2015).

Measurements of 24 h-average concentrations between 2007 and 2016 of alkaline and acidic gaseous components and major water-soluble inorganic ions in PM_{2.5} were used in this study for pH calculation. Citric acid-coated denuders were used to collect gas phase NH₃ samples. Na₂CO₃-coated denuders were used to collect gas phase HNO₃ samples. After alkaline and acidic gases were removed by passing through the two denuders, PM_{2.5} samples were captured by a filter pack consisting of a

front Teflon filter and a back-up Nylon filter. The use of nylon filter is to capture volatilized nitrate during sampling as the artefact of the evaporation ammonium nitrate and correct the data accordingly (Yu et al., 2006; Babich et al., 2011). All the extracts were measured by ion chromatography. Field blanks were routinely performed as the background corrections. The detailed protocol for ambient constituent measurements for NAPS database can be seen in Dabek-Zlotorzynska et al. (2011).
5 In total, there are 1067, 851, 711, 840, 713 and 742 sets of valid data used in the sites of Toronto, Ottawa, Windsor, Montreal, Simcoe and St Anicet, respectively.

Relevant historical meteorological data including hourly average temperature, relative humidity and atmospheric pressure were downloaded from <http://climate.weather.gc.ca>. The closest stations with complete data recordings were chosen as the meteorological data source. 24-hour average values of meteorological parameters during each sampling time period were
10 calculated and used as the input for E-AIM thermodynamic calculation. Specifically, the St Clair meteorological station was chosen as the meteorological data source for Toronto NAPS site (0.9 km from sampling site), the Delhi CS station for Simcoe site (21 km), the Windsor A station for Windsor site (9 km), the Ottawa CDA RCS station for Ottawa site (6 km), the McTavish station for Montreal site (16 km), and the St-Anicet 1 station for St Anicet site (0.1 km).

2.4 E-AIM thermodynamic modelling

The Extended AIM thermodynamic model (<http://www.aim.env.uea.ac.uk/aim/aim.php>) was applied for the calculation of aerosol acidity as E-AIM relies on few assumptions during the calculation of aerosol inorganic components behavior and it can also calculate the activity coefficient of each ion, which is crucial for aerosol pH calculation (Wexler, 2002). Specifically, model II, which mainly focuses on the thermodynamic calculations of H⁺-NH₄⁺-SO₄²⁻-NO₃⁻-H₂O system (Clegg et al., 1998), was used in this study. Model II was chosen rather than model III or IV because it can deal with a wider range of relative
20 humidity and temperature, which gives more reliable information of the seasonal variation of aerosol acidity. The SNA (sulfate, nitrate and ammonium) components in PM_{2.5} generally contributed more than 80% of total measured charges in the particles. Temperature, relative humidity, total ammonia (summation of gas phase ammonia and ammonium in PM_{2.5}), total nitrate (summation of gas phase nitrate acid and nitrate in PM_{2.5}) and sulfate were used as the inputs for this model. Partitioning of both NH₃/NH₄⁺ and HNO₃/NO₃⁻ allows for a strong constraint on the aerosol pH calculation (Murphy et al., 2017). Particles
25 were forced to be metastable without any formation of salts or ice precipitation. Aerosol pH was calculated according to the formula (Robinson and Stokes, 2002):

$$\text{pH} = -\log(a_{\text{H}^+}) = -\log(f_{\text{H}^+} \times \gamma_{\text{H}^+} \times 55.509) \quad [1]$$

where a_{H^+} is the activity of hydrogen ion in aerosol liquid water, f_{H^+} is the hydrogen ion's mole-fraction based activity coefficient, γ_{H^+} is the mole-fraction of hydrogen ion, and 55.509 is the conversion factor from mole-based activity coefficient

to molality-based one. Model IV was also used to assess the influence from non-volatile cations.

3. Results and discussion

3.1 Time series of pH in six Canadian sites

The 10-year time series of PM_{2.5} pH values at the six sites is illustrated in Figure 2. To our knowledge, this is the first
5 long-term study of aerosol pH in Canada and provides one of the longest records for the evaluation of trends anywhere in the
world. The reliability of the pH calculations depends on several assumptions, including that daily average values are
appropriate for the calculations, and that the gas and particle phases are equilibrated. The limitations of using E-AIM II model
also include the lack of consideration of non-volatile cations, aerosol liquid water contributed by organic species or possible
10 phase separation etc. One rigorous method to evaluate the reliability of the calculated pH is to compare the input (measured)
and output (modelled) gas-particle partitioning of semi-volatile species. The comparisons of modelled and measured gas phase
fractions of total ammonia, denoted as $\text{Frac}(\text{NH}_3) = n(\text{NH}_3) / [n(\text{NH}_3) + n(\text{NH}_4^+)]$, are plotted in Figure S2, where the linear
regressions were all close to 1:1 line with high R^2 (>0.88), indicating that the E-AIM modelling results are consistent with the
observed phase partitioning characteristics of $\text{NH}_3/\text{NH}_4^+$. The use of the phase partitioning of both $\text{NH}_3/\text{NH}_4^+$ and $\text{HNO}_3/\text{NO}_3^-$
to constrain the aerosol pH helps to avoid significant over- or underpredictions (Murphy et al., 2017).

15 In Figure 2, all the time series of aerosol pH show a strong seasonal oscillation pattern between 1.5 to 4.0. By coloring
the pH values by ambient temperature, we can clearly see that particles were generally more acidic when ambient temperature
was higher. This variation pattern is consistent in both spatial (six sites) and temporal (10-year) terms. In the long-term, the
seasonal variation of aerosol pH fits well with the variation of ambient temperature. However, larger variation of aerosol pH
can be identified during wintertime. In some cases, the aerosol is very acidic, with $\text{pH} < 2$, even when temperature < 0 °C. This
20 variation pattern is more clearly illustrated in the box and whisker plots of pH in each month in six sites shown in Figure 3.
The pH values are lowest in the summer (July and August averages < 2) and highest in the winter (January and February
averages > 2.5). The aerosol was consistently more acidic and had smaller pH variation in warmer months, and showed higher
average (and median) pH values but exhibited larger variation in wintertime.

To assess the potential influence of non-volatile cations including Na^+ , K^+ , Mg^{2+} and Ca^{2+} to the seasonal variation
25 pattern of aerosol pH, aerosol pH was also calculated using E-AIM model IV (Friese and Ebel, 2010) with the input of Na^+
concentration representing the contributions of all non-volatile cations ($[\text{Na}^+] + [\text{K}^+] + 2[\text{Mg}^{2+}] + 2[\text{Ca}^{2+}]$). The pH calculation
results are illustrated in Figure S3. Because E-AIM IV deals with narrower range of meteorological conditions ($T \geq 263.15$ K,
 $\text{RH} \geq 60\%$), E-AIM IV calculations were only possible for a limited part of the data sets. However, shown in Figure S3, E-
AIM IV modelling results show similar time series compared to E-AIM II results. These results confirm that the aerosol pH

seasonal variation pattern was not significantly affected by the non-volatile cations. The involvement of non-volatile cations typically increases pH values by less than 0.1, and the change is less than 0.4 for more than 90% of data (shown in Figure S4). Using Na^+ to replace other non-volatile cations does not account for the precipitation of CaSO_4 , which has the same effect on aerosol pH of reducing sulfate. The substitution also does not perfectly reflect the impact on aerosol liquid water content, which may indirectly affect the pH. In general, the gas fraction of NH_x calculated from the output of the thermodynamic model matches very closely with the measured gas fraction of NH_x (Figure S2), suggesting that we are not missing substantial contributions to the ion balance in the particles by only considering ammonium sulfate.

The factors influencing aerosol pH can be roughly classified into two categories: meteorological parameters (including ambient temperature and relative humidity) and chemical composition (including gas phase and particle phase components). The following discussion will focus on the conceptual modelling of aerosol pH sensitivity to meteorological conditions in comparison to the impacts of changes in chemical composition.

3.2 Seasonal cycle of aerosol pH

To assess the sensitivity of aerosol pH to meteorological conditions and chemical composition changes individually, we first examine the pH of aerosol with chemical composition held fixed at representative values under different meteorological conditions. Then we study the aerosol pH sensitivity to chemical composition changes under fixed typical meteorological conditions of each of the four seasons. Figure 4 displays the aerosol pH calculated under every combination of relative humidity from 30% to 95% and temperature from $-10\text{ }^\circ\text{C}$ to $30\text{ }^\circ\text{C}$ with fixed chemical composition. The chosen concentrations for sulfate, $\text{TNO}_3(=\text{HNO}_3+\text{pNO}_3)$ and NH_x are 15 nmol m^{-3} , 30 nmol m^{-3} and 180 nmol m^{-3} NH_x , respectively, which are close to the ten-year average concentrations of chemical composition in Toronto atmosphere (which were 17.6 nmol m^{-3} sulfate, 34.2 nmol m^{-3} TNO_3 and 188 nmol m^{-3} NH_x).

It can be seen in Figure 4 that particles tend to become more acidic at higher temperature and lower relative humidity and become more neutralized when temperature is lower or RH is higher. The square symbols on Figure 4 are the monthly average values of ambient temperature and relative humidity in Toronto from 2007-2016 with standard deviations as error bars. The seasonal cycles of meteorological conditions in five other sites are listed in Table S2, and generally show similar seasonal cycling of RH and temperature. Even excluding changes in chemical composition, we find from Figure 4 that we can explain the summer minimum and winter maximum in pH from Figure 3. Temperature is the main factor driving the seasonal variation of aerosol pH while the changes due to RH variation are much smaller. This result suggests the central role of meteorological conditions, especially temperature, in the determination of aerosol pH seasonal cycle in mid- and high-latitude regions with $\text{NH}_x \gg$ sulfate in chemical composition.

The temperature dependence of aerosol pH can be theoretically derived from the phase partitioning $\text{NH}_3/\text{NH}_4^+$ based on the equilibrium: $\text{NH}_3(\text{g}) \leftrightarrow \text{NH}_3(\text{aq})$ and $\text{NH}_4^+ \leftrightarrow \text{NH}_3(\text{aq}) + \text{H}^+$, which is governed by K_H and pK_a , respectively (Hennigan et al., 2015). Under equilibrium, aerosol pH can be theoretically derived from the observed gas/particle concentration ratio of NH_x and aerosol liquid water [ALW] as:

$$\text{pH} = \log([\text{NH}_3]/[\text{NH}_4^+]) + \log[\text{ALW}] + pK_a + \log K_H \quad [2]$$

, which Hennigan et al. (2015) showed having good agreement with E-AIM modelling results. Because both K_H and pK_a have strong temperature dependencies (Chameides, 1984; Bell et al., 2007), aerosol pH is going to be temperature-dependent even if liquid water content or $\text{NH}_3/\text{NH}_4^+$ partitioning behavior does not change. The partial derivative of aerosol pH dependence on temperature will give $\partial \text{pH} / \partial T = \partial (pK_a + \log K_H) / \partial T \approx -0.05 \text{ (K}^{-1}\text{)}$, which corresponds to 0.1 unit increase (decrease) of aerosol pH if temperature decreases (increases) by 2 °C. The pH gradient shown in Figure 4 also illustrated that approximately 10 °C increase of temperature is required for 0.5 unit decrease in aerosol pH under the same chemical composition and RH, which is consistent with the temperature sensitivity derived through NH_x phase partitioning method.

It is possible that the sensitivity of aerosol pH to chemical composition is different in different seasons. As a result, the sensitivity of aerosol pH to composition changes in each season is further studied through the conceptual modelling. In Figure 5(a)-(d) the pH calculated under different combinations of NH_x and sulfate concentrations under typical temperature and RH conditions in spring (April, $T=10 \text{ °C}$, $\text{RH}=60 \%$), summer (July and August, $T=25 \text{ °C}$, $\text{RH}=70 \%$), fall (October, $T=10 \text{ °C}$, $\text{RH}=75 \%$) and winter (January and February, $T=-5 \text{ °C}$, $\text{RH}=75 \%$) in the Toronto atmosphere are shown. Total nitrate concentrations were set to be two times sulfate concentrations. This assumption is representative of the Toronto atmosphere as ten-year average of sulfate concentration (17.6 nmol m^{-3}) is approximately half of total nitrate molar concentration (34.2 nmol m^{-3}) though for any given day the ratio could be different. The corresponding calculation results of particle phase fraction of nitrate $\epsilon(\text{NO}_3^-)$ in four seasons is shown in Figure S5. During summer ammonium nitrate formation is unfavorable, while in wintertime ammonium nitrate can form very efficiently. As a result, in summertime, the aerosol liquid water content was mainly contributed by ammonium sulfate while in wintertime it was affected by both sulfate and nitrate. The effect of nitrate formation to aerosol pH will be further discussed in section 3.4.

The markers on Figure 5 represent the median and the range (from 10th percentile to 90th percentile) of NH_x and sulfate concentrations in four seasons in six monitoring sites from 2007 to 2016. The three more westerly sites (Toronto, Windsor and Simcoe) tend to have higher sulfate in all seasons and Toronto has slightly higher NH_x . However overall all six sites occupy regions of similar chemical sensitivity in the plots. The conceptual modelling of aerosol pH under summertime condition shown in Figure 5(b) suggests that the relatively low variability in summertime pH shown in Figure 3 results from the aerosol

pH being insensitive to chemical composition changes under the corresponding meteorological conditions. Specifically, in summertime, pH was very constrained to the small range around 2 with little influence from the variation of chemical composition, even though the sulfate loadings were the most variable in the summer. Judging from the concentration ranges of sulfate and NH_x shown in all six sites, in summertime even decreasing sulfate concentration by one order of magnitude do not significantly change the pH values, and this effect became more obvious with higher NH_x :sulfate molar ratio.

By comparison, the pH calculated under spring and fall temperature (10 °C) shown in Figure 5(a) and 5(c) showed that pH became more sensitive to NH_x variation but still remain quite insensitive to sulfate concentration changes within the relevant concentration range. In spring, the median concentrations of NH_x in the sites of Toronto, Simcoe and Windsor were higher compared to the other three sites, leading to average April pH values in Toronto, Simcoe and Windsor (2.40~2.69) that were consistently higher than the other three sites (2.15~2.31). In October, NH_x and sulfate concentrations had wider ranges of concentration, but we can conclude from Figure 5(c) that the pH changes due to chemical composition were mainly driven by the variation of NH_x . The effect of relative humidity changes can be seen from the comparison between Figure 5(a) and 5(c), where it shows that with fixed chemical composition, aerosol will be slightly less acidic at higher RH.

By comparison, under wintertime meteorological conditions, aerosol pH was higher in the region NH_x :sulfate>2. Apart from the effect of lower temperature, lower concentration of NH_x and lower NH_x to sulfate molar ratio also made aerosol pH much more sensitive to chemical composition changes than the other seasons because it will make the $\log([\text{NH}_3]/[\text{NH}_4^+])$ part in equation 2 more sensitive to chemical component changes. This could contribute to the larger variation of wintertime aerosol pH shown in Figures 2 and 3. In Simcoe, Windsor and St Anicet, the sites with lower NH_x :sulfate molar ratios, wintertime pH values all exhibited significant variations. Under these conditions, not only the uncertainty in the SNA concentration measurements, but also the potential contributions from organic matter such as organic ammonium salts (Schlag et al., 2017; Tao and Murphy, 2018) or organosulfates (Vogel et al., 2016; Glasius et al., 2018), which current thermodynamic models do not fully consider, can contribute to uncertainty in aerosol pH calculation. These conceptual modelling results also provide insight into the challenges posed in the direct measurement of aerosol pH off-line as it is both strongly impacted by ambient environmental conditions and the NH_3 concentration equilibrated with the particle phase.

One key assumption in the above conceptual modelling is forcing total nitrate to sulfate molar ratio to be 2. To test the effect of this assumption on the aerosol pH, we calculate the pH of aerosol on each sampling day assuming total nitrate to sulfate molar ratio to be 2 and compare with the pH calculated with the measured total nitrate. The results show that this assumption has negligible influence on aerosol pH for samples from April to November but has larger influence on wintertime aerosol, indicating that wintertime aerosol is more sensitive to nitrate concentration (the statistical summary is shown in Figure S6). The influence of nitrate to wintertime aerosol pH will be further discussed in section 3.4.

3.3 Long-term trends of pH in summertime

The above time series analysis and conceptual modellings of aerosol pH suggest that in the study region during summertime, aerosol pH is strongly impacted by ambient temperature with much weaker response to the other factors while during wintertime aerosol pH is both affected by temperature and chemical composition. Figure 6 shows the summertime (data from July and August) median aerosol pH in each year in the six monitoring sites. Summertime average temperature was also plotted on right axes in each graph. The clear anti-correlation indicates that interannual variability in summer temperatures causes interannual variability in aerosol pH. Another application of this result can be found in Battaglia et al. (2017), who showed that urban aerosol in U.S. cities tended to be more acidic than rural aerosol due to urban heat island effect, consistent with the important role of temperature to aerosol pH determination.

The study conducted by Weber et al. (2016) found that in summertime, aerosol remained highly acidic even with significant reduction of sulfate in the US. A similar conclusion could be drawn for conditions in Canada, where we found the role of temperature to be more significant than variations in chemical composition for trends in aerosol pH. Significant changes in the particle phase sulfate have been observed during the summers of 2007-2016. Figure 7 displays the summertime average (data from July and August) concentrations of PM_{2.5} sulfate at the six sites. Linear regressions are used to show the annual reductions of fine particulate sulfate. Significant decreasing trends (p-value of linear regressions <0.05) of sulfate have been observed in five out of six sites (excluding the St Anicet site where large inter-annual variation was found). Specifically, sulfate concentrations have experienced 61%, 56%, 62% and 67% reductions in Toronto, Simcoe, Windsor and Ottawa, respectively, over the decade from 2007 to 2016. There is no statistically significant long-term trend in the NH_x values at any of these sites. Summertime aerosol pH is not decreasing with the decreasing trend of sulfate but is mainly influenced by the inter-annual variation of meteorological conditions.

3.4 pH trends in wintertime

To assess the effect of chemical composition changes on aerosol pH in winter meteorological conditions, wintertime average concentrations of sulfate and NH_x in each year in each site were plotted on Figure 8(a) with the color representing the corresponding average pH values. The background color in Figure 8(a) represents pH calculated under different combinations of sulfate and NH_x concentrations with fixed temperature (-5 °C), RH (75%) and sulfate to TNO₃ molar ratio (1:2), the same as Figure 5(d) but with a different scale and focused on a smaller range of chemical conditions. The figure clearly shows that changes in chemical composition under wintertime meteorological conditions can lead to significant changes in aerosol pH. Changes of 20 nmol m⁻³ in either NH_x or sulfate can change aerosol pH by more than 0.5 units for some cases. Also, although most of the wintertime average pH values calculated for the actual chemical composition at each site (colors of the markers)

generally agree with the pH calculated as the conceptual model (colors of the same position in the background), some of the data points were slightly more acidic than the background value, and this is likely the result of our simplified assumption of fixed sulfate to TNO₃ molar ratio during the calculation in the conceptual modelling.

Figure 8(b) shows the effects that temperature, relative humidity and TNO₃ concentrations can have on the pH of aerosol with 10 nmol m⁻³ sulfate under wintertime meteorological conditions. Generally, lower temperature and higher relative humidity will make aerosol pH higher. However, the effect of the addition of total nitrate is more complicated. Based on equation [2], the added TNO₃ concentration can impact aerosol pH in two opposite ways. First, because ammonium nitrate is more hygroscopic than ammonium sulfate (Gysel et al., 2007), the particulate nitrate formation will raise the liquid water content [ALW] in aerosol, increasing the aerosol pH; however, nitrate formation will also scavenge NH₃ from gas phase, and the smaller value of [NH₃]/[NH₄⁺] will make aerosol more acidic. These two factors altogether contribute to the arc-shaped curve of the relationship between aerosol pH and nitrate shown in Figure 8(b). Figure S7 shows the box and whisker plots of wintertime average molar ratio of TNO₃ to sulfate concentration in each year in each sampling site, where it shows that the molar concentration of TNO₃ was generally higher than 2 times of sulfate concentration, which could contribute to wintertime aerosol being more acidic than the value calculated forcing sulfate:TNO₃=1:2. As a result, during wintertime, both chemical composition and meteorological conditions can have significant impacts on aerosol pH without a clear dominant factor. As a corollary, the uncertainty in measurements of particle composition, the influence of non-volatile cations and the potential contribution from organic acids or organosulfates (the error caused by misrepresenting organosulfates as inorganic sulfate) will also have a more significant impact on the determination of aerosol pH in winter by potentially over- or underestimating the sulfate or nitrate that is balancing the ammonium.

4. Conclusion

The pH values of fine particulate matter at six eastern Canadian sites, including Toronto, Windsor, Simcoe, Ottawa, Montreal, and St Anicet from 2007 to 2016 were calculated by E-AIM model constrained by input from the NAPS database and meteorological data. Strong seasonal cycles of aerosol pH were found in all six sites over the ten years. A consistent pattern of aerosol pH fluctuation from acidic (pH <2) in summer to higher pH around 3 in winter was found at all six sites. Ambient temperature proved to be mainly responsible for the seasonal cycle of aerosol pH. The sensitivity tests of pH to chemical composition shows aerosol pH has different sensitivity to chemical composition changes in different seasons. During summertime, aerosol pH was mainly controlled by ambient temperature while the fluctuation of NH_x and sulfate concentrations did not contribute significantly to the inter-annual variation of aerosol pH. In comparison, in wintertime, aerosol pH can be greatly influenced by both chemical compositions and meteorological conditions. When temperatures are low, aerosol pH at these six sites are very sensitive to all the changes in NH_x, sulfate and TNO₃ concentrations, leading to the higher variation of

aerosol pH in wintertime compared to that in summertime. This study focused on a number of sites with relatively low ambient mass loadings of aerosol inorganic constituents dominated by sulfate-nitrate-ammonium system. Regions in which PM_{2.5} is strongly influenced by dust or marine sources may have different sensitivities for chemical and meteorological factors. However, for places with high particulate pollution and different concentration level of NH₃, similar approaches can also be applied to assess the aerosol pH sensitivity to chemical composition and meteorological parameters to assess the generalizability of our findings.

Data availability

All the data and model used in this study are publicly available, including chemical composition data (<http://maps-cartes.ec.gc.ca/rnspa-naps/data.aspx>), meteorological data (<http://climate.weather.gc.ca/>) and E-AIM modelling (<http://www.aim.env.uea.ac.uk/aim/aim.php>). The time series of 24-hour average meteorological data, chemical composition and aerosol pH calculated with E-AIM II and IV used in this study is available on the supplement excel file of this paper. Further relevant information can be obtained upon request on corresponding author.

Author contributions

YT performed all the data processing and modelling calculation. YT and JGM analyzed data, drew the conclusions and wrote the manuscript together.

Acknowledgement

The authors thanks Enyu Xiong for the help of organizing data. Ye Tao is supported by Ontario Connaught Scholarship from the University of Toronto. The authors acknowledge Environmental Canada National Air Pollution Surveillance Network and Analysis and Air Quality Section of Environmental Canada who contributed to the analysis of PM chemical speciation.

References

- Allen, H. M., Draper, D. C., Ayres, B. R., Ault, A., Bondy, A., Takahama, S., Modini, R. L., Baumann, K., Edgerton, E., Knote, C., Laskin, A., Wang, B., and Fry, J. L.: Influence of crustal dust and sea spray supermicron particle concentrations and acidity on inorganic NO₃⁻ aerosol during the 2013 Southern Oxidant and Aerosol Study, *Atmospheric Chemistry and Physics*, 15, 10669-10685, 10.5194/acp-15-10669-2015, 2015.
- Babich, P., Davey, M., Allen, G., and Koutrakis, P.: Method Comparisons for Particulate Nitrate, Elemental Carbon, and PM_{2.5} Mass in Seven U.S. Cities, *Journal of the Air & Waste Management Association*, 50, 1095-1105, 10.1080/10473289.2000.10464152, 2011.
- Battaglia, M. A., Douglas, S., and Hennigan, C. J.: Effect of the Urban Heat Island on Aerosol pH, *Environmental science & technology*, 51, 13095-13103, 10.1021/acs.est.7b02786, 2017.
- Bell, T. G., Johnson, M. T., Jickells, T. D., and Liss, P. S.: Ammonia/ammonium dissociation coefficient in seawater: A significant numerical correction, *Environmental Chemistry*, 4, 10.1071/en07032, 2007.
- Chameides, W. L.: The photochemistry of a remote marine stratiform cloud, *Journal of Geophysical Research*, 89,

- 10.1029/JD089iD03p04739, 1984.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic Model of the System $\text{H}^+ - \text{NH}_4^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{H}_2\text{O}$ at Tropospheric Temperatures, *The Journal of Physical Chemistry A*, 102, 2137-2154, 10.1021/jp973042r, 1998.
- Craig, R. L., Peterson, P. K., Nandy, L., Lei, Z., Hossain, M. A., Camarena, S., Dodson, R. A., Cook, R. D., Dutcher, C. S., and Ault, A. P.: Direct Determination of Aerosol pH: Size-Resolved Measurements of Submicrometer and Supermicrometer Aqueous Particles, *Analytical chemistry*, 90, 11232-11239, 10.1021/acs.analchem.8b00586, 2018.
- Dabek-Zlotorzynska, E., Dann, T. F., Kalyani Martinelango, P., Celio, V., Brook, J. R., Mathieu, D., Ding, L., and Austin, C. C.: Canadian National Air Pollution Surveillance (NAPS) $\text{PM}_{2.5}$ speciation program: Methodology and $\text{PM}_{2.5}$ chemical composition for the years 2003–2008, *Atmospheric Environment*, 45, 673-686, 10.1016/j.atmosenv.2010.10.024, 2011.
- Dockery, D. W., Cunningham, J., Damokosh, A. I., Neas, L. M., Spengler, J. D., Koutrakis, P., Ware, J. H., Raizenne, M., and Speizer, F. E.: Health effects of acid aerosols on North American children: respiratory symptoms, *Environmental health perspectives*, 104, 500-505, 10.1289/ehp.96104500, 1996.
- Du, H., Kong, L., Cheng, T., Chen, J., Yang, X., Zhang, R., Han, Z., Yan, Z., and Ma, Y.: Insights into Ammonium Particle-to-Gas Conversion: Non-sulfate Ammonium Coupling with Nitrate and Chloride, *Aerosol and Air Quality Research*, 10, 589-595, 10.4209/aaqr.2010.04.0034, 2010.
- Fang, T., Guo, H., Zeng, L., Verma, V., Nenes, A., and Weber, R. J.: Highly acidic ambient particles, soluble metals and oxidative potential: A link between sulfate and aerosol toxicity, *Environmental science & technology*, 51, 2611-2620, 10.1021/acs.est.6b06151, 2017.
- Fountoukis, C., and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for $\text{K}^+ - \text{Ca}^{2+} - \text{Mg}^{2+} - \text{NH}_4^+ - \text{Na}^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{Cl}^- - \text{H}_2\text{O}$ aerosols, *Atmospheric Chemistry and Physics*, 7, 4639-4659, 10.5194/acp-7-4639-2007, 2007.
- Freedman, M. A., Ott, E. E., and Marak, K. E.: Role of pH in Aerosol Processes and Measurement Challenges, *The journal of physical chemistry. A*, 123, 1275-1284, 10.1021/acs.jpca.8b10676, 2019.
- Friese, E., and Ebel, A.: Temperature dependent thermodynamic model of the system $\text{H}^+ - \text{NH}_4^+ - \text{Na}^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{Cl}^- - \text{H}_2\text{O}$, *The journal of physical chemistry. A*, 114, 11595-11631, 10.1021/jp101041j, 2010.
- Glasius, M., Bering, M. S., Yee, L. D., de Sa, S. S., Isaacman-VanWertz, G., Wernis, R. A., Barbosa, H. M. J., Alexander, M. L., Palm, B. B., Hu, W., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Shrivastava, M., Martin, S. T., and Goldstein, A. H.: Organosulfates in aerosols downwind of an urban region in central Amazon, *Environmental science. Processes & impacts*, 20, 1546-1558, 10.1039/c8em00413g, 2018.
- Guo, H., Sullivan, A. P., Campuzano-Jost, P., Schroder, J. C., Lopez-Hilfiker, F. D., Dibb, J. E., Jimenez, J. L., Thornton, J. A., Brown, S. S., Nenes, A., and Weber, R. J.: Fine particle pH and the partitioning of nitric acid during winter in the northeastern United States, *Journal of Geophysical Research: Atmospheres*, 121, 30355-30376, 10.1002/2016jd025311, 2016.
- Guo, H., Liu, J., Froyd, K. D., Roberts, J. M., Veres, P. R., Hayes, P. L., Jimenez, J. L., Nenes, A., and Weber, R. J.: Fine particle pH and gas-particle phase partitioning of inorganic species in Pasadena, California, during the 2010 CalNex campaign, *Atmospheric Chemistry and Physics*, 17, 5703-5719, 10.5194/acp-17-5703-2017, 2017a.
- Guo, H., Weber, R. J., and Nenes, A.: High levels of ammonia do not raise fine particle pH sufficiently to yield nitrogen oxide-dominated sulfate production, *Scientific reports*, 7, 12109, 10.1038/s41598-017-11704-0, 2017b.
- Gysel, M., Crosier, J., Topping, D. O., Whitehead, J. D., Bower, K. N., Cubison, M. J., Williams, P. I., Flynn, M. J., McFiggans, G. B., and Coe, H.: Closure study between chemical composition and hygroscopic growth of aerosol particles during TORCH2, *Atmospheric Chemistry and Physics*, 7, 6131-6144, 10.5194/acp-7-6131-2007, 2007.
- Hennigan, C. J., Izumi, J., Sullivan, A. P., Weber, R. J., and Nenes, A.: A critical evaluation of proxy methods used to estimate the acidity of atmospheric particles, *Atmospheric Chemistry and Physics*, 15, 2775-2790, 10.5194/acp-15-2775-2015, 2015.
- Huang, R. J., Zhang, Y., Bozzetti, C., Ho, K. F., Cao, J. J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., El Haddad, I., and Prevot, A. S.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218-222, 10.1038/nature13774, 2014.

- Lawal, A. S., Guan, X., Liu, C., Henneman, L. R. F., Vasilakos, P., Bhogineni, V., Weber, R. J., Nenes, A., and Russell, A. G.: Linked Response of Aerosol Acidity and Ammonia to SO₂ and NO_x Emissions Reductions in the US, *Environmental science & technology*, 52, 9861-9873, 10.1021/acs.est.8b00711, 2018.
- Liu, H. J., Zhao, C. S., Nekat, B., Ma, N., Wiedensohler, A., van Pinxteren, D., Spindler, G., Müller, K., and Herrmann, H.: Aerosol hygroscopicity derived from size-segregated chemical composition and its parameterization in the North China Plain, *Atmospheric Chemistry and Physics*, 14, 2525-2539, 10.5194/acp-14-2525-2014, 2014.
- Losey, D. J., Parker, R. G., and Freedman, M. A.: pH Dependence of Liquid-Liquid Phase Separation in Organic Aerosol, *The journal of physical chemistry letters*, 7, 3861-3865, 10.1021/acs.jpcllett.6b01621, 2016.
- Murphy, J. G., Gregoire, P., Tevlin, A., Wentworth, G., Ellis, R., Markovic, M., and VandenBoer, T.: Observational constraints on particle acidity using measurements and modelling of particles and gases, *Faraday Discuss.*, 200, 379-395, 10.1039/c7fd00086c, 2017.
- Nah, T., Guo, H., Sullivan, A. P., Chen, Y., Tanner, D. J., Nenes, A., Russell, A., Ng, N. L., Huey, L. G., and Weber, R. J.: Characterization of aerosol composition, aerosol acidity, and organic acid partitioning at an agriculturally intensive rural southeastern US site, *Atmospheric Chemistry and Physics*, 18, 11471-11491, 10.5194/acp-18-11471-2018, 2018.
- Rindelaub, J. D., Craig, R. L., Nandy, L., Bondy, A. L., Dutcher, C. S., Shepson, P. B., and Ault, A. P.: Direct Measurement of pH in Individual Particles via Raman Microspectroscopy and Variation in Acidity with Relative Humidity, *The journal of physical chemistry. A*, 120, 911-917, 10.1021/acs.jpca.5b12699, 2016.
- Robinson, R. A., and Stokes, R. H.: *Electrolyte solutions*, Courier Corporation, 2002.
- Schlag, P., Rubach, F., Mentel, T. F., Reimer, D., Canonaco, F., Henzing, J. S., Moerman, M., Otjes, R., Prevot, A. S. H., Rohrer, F., Rosati, B., Tillmann, R., Weingartner, E., and Kiendler-Scharr, A.: Ambient and laboratory observations of organic ammonium salts in PM₁, *Faraday discussions*, 200, 331-351, 10.1039/c7fd00027h, 2017.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, in, John Wiley & Sons, Inc., 2006.
- Song, S., Gao, M., Xu, W., Shao, J., Shi, G., Wang, S., Wang, Y., Sun, Y., and McElroy, M. B.: Fine-particle pH for Beijing winter haze as inferred from different thermodynamic equilibrium models, *Atmospheric Chemistry and Physics*, 18, 7423-7438, 10.5194/acp-18-7423-2018, 2018.
- Surratt, J. D., Lewandowski, M., Offenberg, J. H., Jaoui, M., Kleindienst, T. E., Edney, E. O., and Seinfeld, J. H.: Effect of Acidity on Secondary Organic Aerosol Formation from Isoprene, *Environmental science & technology*, 41, 5363-5369, 10.1021/es0704176, 2007.
- Tang, Y. S., Braban, C. F., Dragosits, U., Dore, A. J., Simmons, I., van Dijk, N., Poskitt, J., Dos Santos Pereira, G., Keenan, P. O., Conolly, C., Vincent, K., Smith, R. I., Heal, M. R., and Sutton, M. A.: Drivers for spatial, temporal and long-term trends in atmospheric ammonia and ammonium in the UK, *Atmospheric Chemistry and Physics*, 18, 705-733, 10.5194/acp-18-705-2018, 2018.
- Tao, Y., and Murphy, J. G.: Evidence for the importance of semi-volatile organic ammonium salts in ambient particulate matter, *Environmental science & technology*, 53, 108-116, 10.1021/acs.est.8b03800, 2018.
- Utell, M. J., Morrow, P. E., Speers, D. M., Darling, J., and Hyde, R. W.: Airway Responses to Sulfate and Sulfuric Acid Aerosols in Asthmatics, *American Review of Respiratory Disease*, 128, 444-450, 10.1164/arrd.1983.128.3.444, 1983.
- Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C.-U., Aas, W., Baker, A., Bowersox, V. C., Dentener, F., Galy-Lacaux, C., Hou, A., Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher, T., Mahowald, N. M., Nickovic, S., Rao, P. S. P., and Reid, N. W.: A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus, *Atmospheric Environment*, 93, 3-100, 10.1016/j.atmosenv.2013.10.060, 2014.
- Vogel, A. L., Schneider, J., Muller-Tautges, C., Phillips, G. J., Pohlker, M. L., Rose, D., Zuth, C., Makkonen, U., Hakola, H., Crowley, J. N., Andreae, M. O., Poschl, U., and Hoffmann, T.: Aerosol Chemistry Resolved by Mass Spectrometry: Linking Field Measurements of Cloud Condensation Nuclei Activity to Organic Aerosol Composition, *Environmental science & technology*, 50, 10823-10832, 10.1021/acs.est.6b01675, 2016.
- Weber, R. J., Guo, H., Russell, A. G., and Nenes, A.: High aerosol acidity despite declining atmospheric sulfate concentrations

- over the past 15 years, *Nature Geoscience*, 9, 282-285, 10.1038/ngeo2665, 2016.
- Wei, H., Vejerano, E. P., Leng, W., Huang, Q., Willner, M. R., Marr, L. C., and Vikesland, P. J.: Aerosol microdroplets exhibit a stable pH gradient, *Proceedings of the National Academy of Sciences of the United States of America*, 115, 7272-7277, 10.1073/pnas.1720488115, 2018.
- 5 Wexler, A. S.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *Journal of Geophysical Research*, 107, 1-14, 10.1029/2001jd000451, 2002.
- Yao, X., Yan Ling, T., Fang, M., and Chan, C. K.: Comparison of thermodynamic predictions for in situ pH in $PM_{2.5}$, *Atmospheric Environment*, 40, 2835-2844, 10.1016/j.atmosenv.2006.01.006, 2006.
- Young, A. H., Keene, W. C., Pszenny, A. A. P., Sander, R., Thornton, J. A., Riedel, T. P., and Maben, J. R.: Phase partitioning
10 of soluble trace gases with size-resolved aerosols in near-surface continental air over northern Colorado, USA, during winter, *Journal of Geophysical Research: Atmospheres*, 118, 9414-9427, 10.1002/jgrd.50655, 2013.
- Yu, X. Y., Lee, T., Ayres, B., Kreidenweis, S. M., Malm, W., and Collett, J. L.: Loss of fine particle ammonium from denuded nylon filters, *Atmospheric Environment*, 40, 4797-4807, 10.1016/j.atmosenv.2006.03.061, 2006.
- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A.
15 M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophysical Research Letters*, 34, L13801, 10.1029/2007gl029979, 2007a.
- 20 Zhang, Q., Jimenez, J. L., Worsnop, D. R., and Canagaratna, M.: A Case Study of Urban Particle Acidity and Its Influence on Secondary Organic Aerosol, *Environmental science & technology*, 41, 3213-3219, 10.1021/es061812j, 2007b.
- Zhou, M., Zhang, Y., Han, Y., Wu, J., Du, X., Xu, H., Feng, Y., and Han, S.: Spatial and temporal characteristics of $PM_{2.5}$ acidity during autumn in marine and coastal area of Bohai Sea, China, based on two-site contrast, *Atmospheric Research*, 202, 196-204, 10.1016/j.atmosres.2017.11.014, 2017.

25




Figure 1. Six NAPS monitoring sites chosen for this study: Toronto, Ottawa, Windsor, Montreal, Simcoe and St Anicet.




Figure 2. The time series of PM_{2.5} pH at six sites colored by ambient temperature.




Figure 3. The box and whisker plot of aerosol pH in each month in six sites. In each box, the top, middle and bottom lines represent 75th, 50th and 25th percentile of statistical data, the upper and lower whiskers represent 90th and 10th percentile, and the square mark represent the mean value.




Figure 4. Plot of E-AIM calculated pH under different combinations of relative humidity (from 30%-95%) and temperature (from -10 °C to 30 °C) with the fixed input chemical compositions (15 nmol m^{-3} sulfate, 30 nmol m^{-3} TNO_3 and 180 nmol m^{-3} NH_x). The square plots on the graph represent monthly average of meteorological parameters in Toronto from 2007-2016 with standard deviations as error bars.




Figure 5. Aerosol pH calculated with fixed meteorological parameters (temperature and RH) under different combinations of NH_x and sulfate concentrations. Total nitrate concentrations are set to be 2 times that of sulfate. The markers on the graph represent median summertime concentrations of NH_x and sulfate in six sites from year 2007 to 2016 with error bars indicating 10th percentile to 90th percentile. The fixed meteorological parameters chosen in Figure (a)-(d) are representative of the conditions in spring (April), summer (July and August), fall (October) and winter (January and February) shown in Figure 4.

5




Figure 6. Median of aerosol pH in summertime (data on July and August) from year 2007 to 2016 in six monitoring sites. The error bars with median pH values represent the range from 25% to 75% percentile. Average ambient temperature in summertime (July and August) in each site is also plotted on right axis to illustrate the reverse trend of temperature with pH inter-annual variation.




Figure 7. 10-year trends of summertime average concentrations of sulfate and NH_x in each site. Linear regressions of sulfate were illustrated on each graph to represent the reduction of particulate sulfate from 2007 to 2016.




Figure 8. (a) The plot of aerosol pH under wintertime meteorological conditions (-5°C , $\text{RH}=75\%$) under different combinations of sulfate and NH_x with more focused region compared to Figure 5(d). The markers on the graph represents the average wintertime (data from January and February) sulfate and NH_x concentrations in each year in each sampling site with color indicating average pH value. (b) pH of aerosol with 10 nmol/m^3 sulfate under different combinations of temperature and relative humidity as the function of TNO_3 in the system.

5