

Interactive comment on “Evaluating the Relative Importance of Northern African Mineral Dust Sources Using Remote Sensing” by Natalie L. Bakker et al.

Ian Ashpole (Referee)

ian.ashpole@smu.ca

Received and published: 10 May 2019

General comments

This is an enjoyable and insightful paper that combines multiple different satellite datasets to enhance our knowledge of the geomorphology of North African dust sources and estimate how much dust is emitted from different regions. As well as being of scientific value in themselves, the results are of significance for our understanding of the fertilization link between Saharan dust and the Amazon and this case is strongly made in the paper introduction, discussion and conclusion. However, I do feel that the methods need greater clarification/explanation, and that in some places a

C1

mention of the methodological/data limitations is needed. In my opinion, this paper will be well worthy of publication once a few issues are addressed.

My review is structured as follows: I first explain my major comment about the paper. I then outline a series of specific comments, section-by-section, which are less minor (some of these will likely be addressed in the process of addressing my major comment), before suggesting a few small technical corrections at the end.

Major Comment

My only major question/comment refers to your methods and is more a question of greater explanation/clarification needed, as opposed to me questioning the validity of what you have done (the method seems great to me, if I understand it correctly).

Basically, I'm not entirely clear how you get to the final results that are presented in Tables 1 and 2. From several re-reads of the methods, it is my understanding that you only classify the source geomorphology (Section 2.3) of the 20 dust events that you study in detail in Section 2.2 (using VIIRS)? Yet, in Table 1, you give the classification of 3512 sources. Does this mean that the 20 events studied originated from 3512 specific point sources in total; or does this number come from all of the events identified in the SEVIRI data (Section 2.1)? If the latter, how do you know the geomorphology of the point sources that contribute to the events that were *not* studied using the VIIRS data in Section 2.2 and Sentinel data in Section 2.3? (A related point: you don't state anywhere in Section 2.1 that you identify the actual sources of events identified using SEVIRI data, just that the events were "identified". If you did backtrack these events to sources, you should say so.)

I *think* the answer to the above is in the first couple of sentences of Section 2.6, but I am not 100% sure: "Analysis of the dust source locations from the 10 largest dust storms in each dust season showed that a single geomorphological source dominated

C2

in most areas. On the basis of this observation, 16 individual dust source regions were identified and mapped.” Does this mean, that all (or most of) of the events identified with SEVIRI (Section 2.1) originated from within the 16 broader regions that are outlined in Figure 3? And because you had studied in detail events that originated from these sources in the course of Sections 2.2 and 2.3, you were able to deduce the geomorphology of the whole broader area? So, for example, if an event detected in SEVIRI that was not one of the largest 20 originated from within the boundaries of the Bodélé, it was assumed to have a source geomorphology of paleolake? If I *am* correct about this, then I am confused as to how you get stats in Table 1 for geomorphology types of “Sand deposits” through to “Anthropogenic”, since all of the 16 broad regions are classified as “Alluvial deposits” and/or “Paleolakes”*. But then, some of the pins in Figure 3 (which are the sources of the 20 largest events) are colour coded to represent “Sand deposits” etc. So, do the figures in Table 1 actually only come from these top 20 events? Hopefully you can see why I am confused!

*Relatedly: 4 of these broad regions weren’t analysed as part of Sections 2.2 and 2.3 as none of the 20 biggest events originated from them: At what stage, then, did you classify their geomorphology? I think this is implied in “mapping” stage of the quoted text from the start of Section 2.6, but how you did this mapping isn’t clear.

Specific comments

(*Suggested text modifications for each section are in italics*)

General: It would be helpful to clarify/be consistent with your use of the term “source” and variants (“point source”, “source region”, etc). In some cases you use “source” to refer to the specific point sources and features identified in Sections 2.2 and 2.3, and in other cases you use it to refer to the 16 broader source regions from Figure 3. Sometimes you use “point source” and “source regions” to refer to these, but not always.

C3

Abstract

Arguably, the first 3 sentences set this up to be a paper about fertilization, but this isn’t really its focus – Amazon fertilization can only really be commented on from your (albeit quite significant) results. Can I suggest shifting these sentences to the end of the abstract/de-emphasizing the fertilization angle?

1 Introduction

P1, L20: Mineral dust is likely important to more parts of the earth system than just climate (i.e. biogeochemical cycles, as you emphasise this paper). You could broaden this opening statement a little to reflect this.

P2, L14: “While the major dust source areas within the Sahara and the Sahel have been identified. . .less is known about the geomorphology of these sources”. True, but some studies (e.g. Ashpole and Washington 2013, Crouvi et al 2012) have considered this (AW13 identifies several alluvial features as dust sources in the central and western Sahara) and could be mentioned here?

P2, L28: “. . .as well as the total dust mass emitted per class and dust source region are calculated”. Should “calculated” be changed to “estimated”, since the “calculations” contain a terms with a lot of potential error/unknowns?

2.1. Detecting dust storms

Did you detect the sources of all dust events that were identified? You don’t explicitly say this, but it seems that you must have done for the later methods to make sense?

P5, L6-7: “By visually interpreting images, all dust events of the 2015-2017 boreal winter dust seasons were identified”. Did you have some subjective threshold of what was and wasn’t a dust event? From experience of working with these data I know that there are sometimes minor dust plumes that are hard to reliably trace to a specific source area. Similarly, how about when detection/interpretation/tracking was complicated by the presence of clouds? I appreciate that you do the best job you can with the data at

C4

hand and that no analysis is perfect, but I think it is important to note somewhere if you actually just identified, for example, all dust events that could be tracked to a source with confidence, since this will introduce an element of uncertainty (albeit probably small) into your results.

2.2 Identifying dust sources

P5, L14-15 (and also Figure 2): “Dust sources were identified manually using visual interpretation, by tracking the individual plumes back to their first point of occurrence (illustrated in Figure 2)”. I think this needs a little more explanation/clarification: The VIIRS data just show you a dust plume at a single point in time. How can you be *sure* what the first point of occurrence is? Is it taken as the most upwind point of the plume (with the upwind direction deduced from watching plume evolution in the SEVIRI data)? Additionally, in the example of Figure 2, you seem to have attributed the plume to 5 specific source locations, but what about the spaces in between these points, which also seem to be a shade of pink? Or is the whole upwind edge actually taken as the source location and the black points are just illustrative? OR, is this stage just narrowing down the suspected first points of occurrence so that you know what area to study for likely emissive features in a later step? Further explanation/clarification would be helpful here!

P5, L16: It would be helpful to clarify whether the selection of the 10 largest dust storms per season was qualitative, or whether some other quantitative means was used to pick them (such as area covered, based on the step outlined in Section 2.4?).

2.3 Classifying geomorphology of dust sources

General: Is it possible to include example images (or a case study) of sources meeting each classification as an additional figure (or supporting material)? I think this would help non-expert readers to understand (and visualise) the different source types.

P6, L11-13: “To account for the spatial resolution differences between VIIRS and

C5

Sentinel-2, point sources were classified as the most prominent feature and/or most common geomorphic class within 750 m of the point source.” I find the wording a little confusing here. Does the second “point source” refer to the original point identified in the VIIRS data?

2.4 Determining dust plume area

General:

1) What happens when dust plumes detected & measured in VIIRS are the result of several smaller plumes from different sources merging? How do you apportion the plume between the different sources? Or is this never an issue?

2) A problem with using VIIRS to determine dust plume area as opposed to SEVIRI is that it only gives one snapshot per day of dust plumes. Given that the size of plumes will change due to transport and advection, this must introduce a bias in your results, since if VIIRS observed the plumes at a different time of day they would be smaller or larger? This has knock-on effects for average plume AOD calculations and dust mass calculations. You are limited in what you can do by the data at hand, but this seems important to acknowledge. Relatedly, what time of day is the VIIRS observation actually made, and how does this relate to the modal emission time for the region, which I think is during the morning in this season (based on the work of e.g. Schepanski et al. 2009 & others)?

P6, L23-24: “dust plumes could be outlined and their size calculated”. For clarification, this enables you to treat each individual dust plume separately and assign that dust plume area to certain source?

2.5 Measuring average aerosol optical depth per plume

General: You may have clarified this in addressing my major comment, but do you get the average AOD of every dust plume identified in Section 2.1, or just the subset studied in Sections 2.2 – 2.3?

C6

General: How do you actually go about calculating the average AOD per plume? Is it a case of overlapping the VIIRS and MODIS data, getting all AOD values that overlap with flagged dust plumes from Section 2.4, and calculating the arithmetic mean (or median) for each individual dust plume? How do you deal with the resolution mismatch between the MODIS data (10 km) and VIIRS data (0.75 km)?

P6, L31: “MODIS monitors AOD on a daily basis”. Is there any temporal offset between the MODIS and VIIRS data, or are the sensors carried on the same satellite? This seems important to me because if the plume area/mean AOD are obtained at different points in the plume lifetime, its location may have changed, which would be problematic if using VIIRS pixels to get AOD values from MODIS data. . .

2.6 Calculating dust mass per dust source

Suggested modification for section title: “Estimating emitted dust mass per dust source region”? Terms in your calculation contain a lot of potential error/uncertainty (plume size, mean AOD, extinction coefficients. . .), so this is really more of a rough estimate. Also, you seem to consider dust mass for broader source regions as opposed to individual point sources, which is what the term “source” also occasionally refers to in this paper.

P7, L9-17: This paragraph seems like it could be better placed in its own sub-section, explaining how you pull the methods of Section 2.1 – 2.5 together to create a set of dust plumes per broad source region, as that step (if I understand your methods correctly) seems like it needs greater emphasis. However, this may not be the case depending on how you address my major comment.

P7, L9-10: “Analysis. . . showed that a single geomorphological source dominated in most areas”. Do you mean single geomorphological source class?

P7, L10: “On the basis of this observation, 16 individual dust source regions were identified and mapped”. What does “and mapped” mean? Does this refer to digital

C7

boundary creation or some such to help with categorising the source area of all of the individual plumes detected in Section 2.1? I think clarification/clearer explanation is needed in this section.

P7, L14-15: “Each dust plume was then assigned a geomorphic class based on the region it came from”. I’m confused. By “each dust plume”, are you referring to all the plumes detected in Section 2.1? If so, you didn’t state in that section that you identified sources. I am guessing this must be the case because (I think?) that you also have the size and mean AOD for all these plumes. . .

P7, L15-17: “The accuracy of this methodology was evaluated by randomly selecting three additional days, analysing the geomorphic class of each source, and determining what percentage of these dust sources coincided with that defined by the map of predominant dust geomorphologies.” Suggested addition for clarity: “. . . analysing the geomorphic class of each detected dust source. . .” Additionally: does it make sense to move your statements about the accuracy assessment results from the end of Section 3 (P9, L9-11) to here? They don’t add anything to the overall paper results, and the current wording here leaves me asking the question “well. . . how accurate did your methodology turn out to be?!”

P7, L19: “. . . whereby dust mass is calculated. . .” Again, can I suggest calculated be substituted for estimated? Can I also suggest a statement be made about the reliability of these results, given all of the (presumably quite uncertain and highly variable) parameters in Kaufman’s equations, as well as the fact that you are using a plume average AOD and plume size calculation which are taken from one snapshot through the dust event’s lifetime (and these values could therefore be quite different if calculated using observations at a different time of day).

P8, L3: “although it should be noted that the equation has an uncertainty of $\pm 30\%$ for AOD values between 0.2 and 0.4.” Ben-Ami et al. also note that “in cases when total AOD > 0.4, the error may be larger” – this should be added.

C8

3 Results

General: In Table 2, you only seem to attribute dust mass to events originating from within the 16 broad source regions outlined in Figure 3. This figure shows that a small number of the point sources identified during the 20 largest dust storms that you analysed in Section 2.2 fell outside of these regions (e.g. in Egypt) and, presumably, at least a small portion of the events identified in SEVIRI data (Section 2.1) did too. If I'm right about your methodology and you calculated dust mass for all plumes, this means your "Total" mass stats (and statements about this) are missing emission from these events? Even if this is a minor proportion of the total mass emitted from the 16 main source regions, it still seems worthy of mention.

P8, L19: Can you give mean AOD for the other source regions, in addition to the Bodélé Depression? It would be interesting to see whether there are significant differences between sources. For example, some source areas (such as Sudan and Ennedi, both alluvial) seem to be frequently active, but emit a comparatively small amount of dust (and the converse is true for, e.g. Algeria – paleolake). It would be interesting to know if this is a function of differences in mean AOD or plume size. Given your comments about differences in the biogeochemical make up of dust from different source geomorphologies, it seems plausible that this might in some way affect the observed AOD values from specific source types too. Alternatively, it could say something about differences in erosivity/erodibility in these regions. Either way, worthy of comment as this is useful knowledge for remote sensing/dust modelling communities (as well as others!)

P9, L4-5 & Table 2: A reminder (possibly just in Table 2 caption) that South Air and Erg Chech dust mass was apportioned 50/50 to the paleolake and alluvial deposit categories would be useful, as it took me a while to work out how exactly you got to the 64% & 36% numbers.

P9, L11-12: "The ten largest dust storms per season emitted 37.6% of the total dust

C9

mass." What proportion of the total detected dust storms did the ten largest dust storms per season account for? This would be useful to know to contextualise the 37.6% figure.

Technical corrections (typos, citations, etc)

P5, L4: add a reference to Figure 2 (as this visualises pink dust)?

P5, L5: "...and blue the 10.8 μm channel". I suggest the following addition for clarity (since this is not a BT): "...and blue the brightness temperature (BT) of the 10.8 μm channel".

P5, L8: "[the high temporal resolution of SEVIRI] makes it suitable for observing dust storms". I suggest the following addition: "[the high temporal resolution of SEVIRI] makes it suitable for observing the evolution of dust storms". This is the strength of the higher temporal resolution data; you can still "observe" dust storms (or dust plumes, at least) with lower temporal resolution data.

P5, L8-9: "...its low spatial resolution means it is less suited to determining the location and nature of the sources". I suggest the following addition: "...its low spatial resolution means it is less suited to determining the precise location and nature of the sources". The strength of the higher spatial resolution is the precision it affords, since SEVIRI can give the correct location to a scale of 3 km, no?

P8, L13: "During the 2015-2017 winter dust season between 82.3 - 127.0 Tg of dust was emitted". You should include a reference to Table 2 here.

P12, L15: "N and P macronutrients". Expand abbreviations/chemical symbols?

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-253>, 2019.

C10