

Interactive comment on “Controls on the water vapor isotopic composition near the surface of tropical oceans and role of boundary layer mixing processes” by Camille Risi et al.

Camille Risi et al.

crimd@lmd.jussieu.fr

Received and published: 4 July 2019

Printer-friendly version

Discussion paper



Response to reviewers

July 4, 2019

Summary:

We thank both reviewers for their comments. As detailed in the point-by-point response below, we have taken them into account. We have already implemented most of the necessary changes in a revised manuscript, which we are motivated to re-submit in the next few weeks. However, we still need more time to work on 4 aspects that are mentioned by the reviewers, because they need substantial new analyses:

1. quantify the effect of horizontal advection, add an appendix on the method for this quantification, and discuss this effect in the main text.
2. quantify the effect of rain evaporation, add this quantification in the appendix that is already devoted to this subject, and discuss this effect in the main text.
3. quantify the effect of assuming a Rayleigh curve for δD with constant α_{eff} , and discuss this effect in the main text.
4. better document and illustrate the spatio-temporal variability in free tropospheric profiles.

[Printer-friendly version](#)

[Discussion paper](#)



We have already started to work on these aspects, as detailed below, but we have not finished yet. We will use the revision time to finish the work and address more completely the reviewers' comments on these specific issues.

1 Reviewer 1

We thank reviewer 1 for his/her comments.

This paper presents an analytical steady state vertical mixing model to investigate the controls on the water vapor isotopic composition in the subcloud layer over the tropical oceans. It is a nicely simple model that considers the most important processes, which are surface evaporation and vertical mixing and predicts the subcloud layer water vapour isotope composition from a combination of mass balance equations for all isotope species. I enjoyed reading this paper very much, I particularly like the approach chosen for testing this analytical model, which combines model simulation data and ship-based observations. The ideas presented in this paper are exciting and very valuable for upcoming large field campaigns in which isotope observations are planned in different parts of the lower troposphere.

I thus recommend minor revisions with the following minor points:

1) In the abstract it should be clearly stated that the proposed analytical model is a steady state formulation, which neglects horizontal gradients and thus the impact of horizontal advection.

[Printer-friendly version](#)

[Discussion paper](#)



We agree: we modify as follows: “We propose a steady-state analytical model ... The model relies on the hypothesis that δD profiles are steeper than mixing lines, and we neglect the effects of rain evaporation and horizontal advection on δD_0 ”

2) P. 1, L. 10: “When the air mixing into the SCL is lower in altitude it is moister”: I think this is true most of the time and certainly in a climatological sense, but of course when including differential advection elevated moist layers can occur. I guess adding “it is generally moister” would make me very happy.

We modify as suggested.

3) P. 3, L. 4: Shouldn't cloud top cooling be mentioned here as well?

We modify as: “driven by cloud-top radiative cooling, mixing and evaporative cooling of droplets”

4) P. 5, L. 4: Neglecting the large-scale horizontal gradients in air properties, particularly in the trade wind regions seems to me like a strong assumption. Given the sensitivity of dD to SST and the considerable SST gradient across the North Atlantic, I find that this caveat could be discussed a bit more explicitly here.

We agree that the neglect of horizontal advection is an important caveat that we need to discuss more, and if possible, quantify. There are several comments along this line from you and from reviewer 2.

We are working on a way to rigorously estimate the effect of horizontal advection of

[Printer-friendly version](#)[Discussion paper](#)

isotopic gradients on our results. To do so, we extend our Eq. 9: we get:

$$R_0 = \frac{R_{oce}}{\alpha_{eq}} \cdot \frac{1}{h_0 + \alpha_K \cdot (1 - h_0) \cdot \left(\frac{1 - r_{orig}^{\alpha_{eff}}}{1 - r_{orig}} + \phi \cdot (1 - \beta) \right)} \quad (1)$$

where $\phi = \frac{F_{adv} \cdot q_0}{E}$, $\beta = \frac{R_{adv}}{R_0}$, F_{adv} is the air flux coming from advection in kg/m²/s, R_{adv} is the isotopic ratio of the advected air assuming an upstream advection scheme. All these variables can be diagnosed from the model outputs.

We have not finished yet, we will continue to work on it during the revision time. If this method works well, we will add an additional appendix in the paper, analogous to the appendix on rain evaporation. In the main text, we will add the results from this quantification. Discussions will be added on how horizontal advection affects our results.

Coming back to the effect of horizontal gradients in the North Atlantic, with the above-mentioned equation, we hope to get a map that quantifies the effect of horizontal advection. If this map supports your comment, we will add a sentence on the North Atlantic that reflects your comment.

5) P. 5, L. 20: “qs is the saturation specific humidity at SST”

corrected

6) P. 6, L26: In the closure section and the discussion of the free tropospheric profile the role of horizontal advection is again neglected. This is maybe a good assumption in the tropic but it should still be mentioned explicitly.

We neglect horizontal advection in the box model described in section 2.1. However,

our closure assumption does not need to neglect horizontal advection. Horizontal advection effects can be implicitly taken into account through the α_{eff} : we now clarify this: “Effects of horizontal advection and rain evaporation are encapsulated into α_{eff} .” As detailed below as a response to another comment, we now give specific examples of how horizontal advection may affect α_{eff} .

It is possible that horizontal advection distorts the δD profile from a Rayleigh curve. We will quantify the effect of assuming a Rayleigh curve compared to the full simulated profile by LMDZ, as detailed further down (equation 2 of this response). This will allow us to address this issue.

7) P. 8, L26: “Depending on microphysical details that are too complex to be addressed here”, Graf et al. 2019 could be referenced here

We add this reference.

8) P. 9. Fig. 3: Which value was chosen for the SST? This could be mentioned in the caption as well as a reference to which equilibrium fractionation factor was used

We add this information in the caption: “For this illustrative purpose, we assume SST=30°C, $h_0 = 0.8$ and $\delta D_{oce} = 0$.”. We also add it to Fig. 4.

9) P. 12, L. 2: “However, if the end member is defined below 500 hPa (e.g. 600 hPa), results are not always reasonable”, why is this so?

Now we explain this in the text: “However, the end member should be defined above

Printer-friendly version

Discussion paper



500 hPa to ensure that it is well above boundary layer processes. If the end member is defined below 500 hPa (e.g. 600 hPa), there are a few cases where q increases with altitude ($q_f > q_0$) due to horizontal advection or convective detrainment from nearby moister regions. Meanwhile, δD decreases monotonically, leading to unrealistic values for α_{eff} ”.

10) P. 12, L. 7: In my opinion, this makes it difficult to interpret rorig. But probably there are conditions when rorig and thus zorig are more physically meaningful than others. Could the authors maybe add a list with explicit and quantitatively expressed conditions in which they would argue that the assumptions involved in Eq. 9 are satisfied?

We will work on “quantitatively expressed conditions” for the validity of assumption in Eq. 9 during the revision time. Off course, the difficulty in this work is the “quantitatively”.

- As explained above (equation 1 of this response), we will quantify the effect of horizontal advection on δD_0 , r_{orig} and z_{orig} .
- In addition, we will make use of the equation that is already in appendix B to quantify the effect of rain evaporation. We already have all the necessary LMDZ outputs to diagnose the η and α_{re} parameters.

11) P. 13, L. 4-5: Is there a literature reference that the authors could indicated for this calculation of zi from observations?

We now give several literature references and more explanation on this calculation

method: “The temperature inversion is an abrupt increase in temperature that caps the boundary layer. Therefore, a method to automatically estimate its altitude is to detect a maximum in the vertical gradient of potential temperature (Stull (1988); Oke (1988); Sorbjan (1989); Garratt (1994); Siebert et al. (2000)). This method is sensitive to the resolution of vertical profiles (Siebert et al. (2000); Seidel et al. (2010)). Therefore, we adapted this method in order to yield z_i values that best agree with what we would estimate from visual inspection of individual temperature profiles. In LMDZ, we calculate z_i as the first level at which the vertical potential temperature gradient exceeds 3 times the moist-adiabatic lapse rate. In observations, we calculate z_i as the first level at which the vertical potential temperature gradient exceeds 5 times the moist-adiabatic lapse rate, because radio-soundings are noisier than simulated profiles. ”

12) P. 13, L. 15: I did not immediately understand what was meant by composites belonging to a given interval of ω_{500} , I was expecting a map. A reference to the results figure referred to here would have helped me.

We now explain better how the composites are calculated, and we add a reference to Fig. 10 as an example: “The type of clouds and mixing processes depends strongly on the large-scale velocity at 500 hPa (ω_{500} , map show in Fig. 6a), with shallow clouds in subsiding regions and deeper clouds in ascending regions (Fig. 1). Therefore, it is convenient to plot variables as composites as a function of ω_{500} (Bony et al. (2004)). To make such plots, we divide the ω_{500} range from -30 to 50 hPa/d into intervals of 5 hPa/d. In each given interval, we average all seasonal-mean values at all locations over tropical oceans that belong to a this interval (e.g. Fig. 10a will be an example).

The cloud cover strongly correlates with the inversion strength, which can be quantified by the Estimated Inversion Strength (EIS, Wood and Bretherton (2006), map shown in Fig. 6b). We thus also plot variables as composites as a function of EIS. To make such

[Printer-friendly version](#)[Discussion paper](#)

plots, we divide the EIS range from -1 K to 9 K into intervals of 0.5K. In each given interval, we average all seasonal-mean values at all locations over tropical oceans that belong to this interval I (e.g. Fig. 10b will be an example). “

13) P. 13, L. 22: By how much (range of variability) were the four factors varied?

The four factors were varied from a control value to their simulated value.

We now clarify this in the text with a modified paragraph: “To understand what controls the δD_0 spatio-temporal variations, δD_0 is decomposed into 4 contributions based on Eq. (9). First, we define $r_{orig,basic} = 0.6$, $\alpha_{eff,basic} = 1.09$, $SST_{basic}=25^\circ\text{C}$, $h_{0,basic} = 0.8$ as a basic state. We call $\delta D_{eq9}(r_{orig}, \alpha_{eff}, SST, h_0)$ the function giving δD_0 as a function of r_{orig} , α_{eff} , SST and h_0 following the Eq. (9). The relative contributions of r_{orig} , α_{eff} , SST and h_0 to δD_0 variations are estimated as $\delta D_{eq9}(r_{orig}, \alpha_{eff,basic}, SST_{basic}, h_{0,basic})$, $\delta D_{eq9}(r_{orig,basic}, \alpha_{eff}, SST_{basic}, h_{0,basic})$, $\delta D_{eq9}(r_{orig,basic}, \alpha_{eff,basic}, SST, h_{0,basic})$ and $\delta D_{eq9}(r_{orig,basic}, \alpha_{eff,basic}, SST_{basic}, h_0)$ respectively.”.

14) P. 14, Section 4.1: This section seemed very technical for me. I also see it more as a methodological aspect than a result. I would recommend to either shift it to a technical appendix (since the paper is quite long) or to the methods section.

We will work during the revision time on moving this section to the method section or appendix. This may depend on what we do to address the comments on better documenting the spatio-temporal variability in free tropospheric δD profiles.

Printer-friendly version

Discussion paper



15) P. 15, Fig. 7: Mention that these are different random (?) grid points in the caption. I would have liked a more general evaluation also describing the temporal and spatial variability in the vertical profiles simulation by LMDZ.

We now explain how these points were selected: “These profiles were selected automatically as the first day and tropical ocean location (scanning all latitudes from South to North and longitudes from 180W to 180E) for which the RMS difference between the LMDZ profile and the Rayleigh line ($RMS_{Rayleigh}$) and the RMS difference between the LMDZ profile and the Rayleigh line (RMS_{mixing}) satisfy the following conditions: (a) $|RMS_{Rayleigh} - 10| < 1permil$ and $|RMS_{mixing} - 25| < 1permil$, (b) $|RMS_{Rayleigh} - 25| < 1permil$ and $|RMS_{Rayleigh} - 10| < 1$ and (c) $|RMS_{Rayleigh} - 25| < 1permil$ and $|RMS_{Rayleigh} - 25| < 1permil$.”.

Regarding the “more general evaluation also describing the temporal and spatial variability in the vertical profiles simulation by LMDZ”: this comment echoes one from the second reviewer. We will think about how to address it during the revision time.

16) P. 15, L. 3: could the authors mention the region where they think that α_{eff} may also reflect horizontal advection effects?

Although we will work on quantifying the effect of horizontal advection on δD_0 , i.e. in the SCL, the effect of horizontal advection on α_{eff} is a completely different subject that is beyond the scope of this paper. Therefore, we only reply based on the literature: “The pattern of α_{eff} may also reflect horizontal advection effects, where strong isotopic gradients align with winds (e.g. from the Eastern to the Western Pacific, Dee et al. (2018)). “

[Printer-friendly version](#)[Discussion paper](#)

17) P.16: I find it interesting that the mixing and Rayleigh lines have large biases in front of the eastern continental boundaries where the inversion is strongest and, where there is a strong decoupling between the FT and BL. In particular in these regions, I would expect horizontal advection to play a key role, (e.g. the SAL layer in front of the eastern North African Coast, see Lacour et al. 2017, ACP). Maybe the authors find a good way to shortly note this in the text.

We now add this comment: “For example, we note that mixing and Rayleigh lines have large biases in front of the eastern continental boundaries where the inversion is strongest, leading to a strong decoupling between the FT and the boundary layer. Horizontal advection is expected to play a key role in these regions (e.g. the Saharian layer in front of the eastern North African Coast, Lacour et al. (2017)). “

18)P. 17, L. 2: Maybe one could add oceanic upwelling and atmospheric deep convection. Jumping from upwelling to deep convection in the same sentence, I was not sure whether deep convection in the ocean or the atmosphere was meant here.

We modify as suggested.

19) P. 17, L. 4: “Decreases as omega500 is more strongly ascending or descending”
-> “with increasing vertical winds (omega500) of both signs”

We modify as suggested.

[Printer-friendly version](#)[Discussion paper](#)

20) P. 17, L. 11: “in more ascending regions” -> a reference to Fig. 8d would have helped me here.

We add this reference

21) P. 17, L. 25: “The fact that the effect...” I had difficulties to understand this sentence.

We remove this sentence that was not so useful.

22) P. 17, L. 33: h_0 (62%) is the largest explained fraction of all the variables considered and should thus be put first. This could be a hint that large-scale horizontal advection plays an important role at the synoptic timescale in these regions.

We now move h_0 first in the sentence.

We will write a comment on horizontal advection once we have quantified its effects.

23) P. 19, Fig. 10: The bin sizes (number of data points per bin should be added).

We now add this information in the figures, and we write in the caption: “The number of samples in each bin is indicated on a logarithmic scale on the right-hand-side as bars.”

If I understood correctly from the caption, the authors used the seasonal averaged fields from LMDZ. Why not making these composites using the 6-hourly outputs? For me there is a timescale discrepancy between the processes (mixing, evaporation) that the authors look at and the averaging timescale of the used fields.

Printer-friendly version

Discussion paper



The composites are based on seasonal-mean EIS or ω_{500} because this allows a better link with the large-scale dynamical regime. Mixing and evaporation are processes that act at short time scales, but their relationship to the large-scale circulation is best constrained by energetics at time scales longer than synoptic. Let's consider ω_{500} , for example. It relates to convective activity and other diabatic processes through the conservation equation of moist static energy. Adiabatic cooling by large-scale ascent balances latent heating by convection (Yanai et al. (1973)), or adiabatic heating by large-scale subsidence balances radiative cooling (Emanuel et al. (1994)). The stationarity in the conservation of moist static equation is most valid at scales longer than synoptic, otherwise, the storage term becomes important (Masunaga and Sumi (2017)). This is why in Bony et al. (2004) and subsequent papers (e.g. Bony et al. (2013)) based on ω_{500} , monthly-mean ω_{500} is used, which yields similar results to seasonal-mean.

A similar rationale applies to EIS. This is why many papers on EIS use seasonal-mean values, notably the paper defining this quantity (Wood and Bretherton (2006)).

We now explain this in section 3.5: “Note that such composites are done on seasonal-mean ω_{500} because cloud processes and their associated diabatic heating are tied to the large-scale circulation through energetic constraints (Yanai et al. (1973); Emanuel et al. (1994)) that are best valid at longer time scales (otherwise, the energy storage term may become significant, e.g. Masunaga and Sumi (2017)). This is why ω_{500} is generally averaged over a month or longer (e.g. Bony et al. (2004); Bony et al. (2013))”

and for EIS: “Using seasonal-mean values is consistent with Wood and Bretherton (2006) and with the better link at longer time scales between cloud processes and the large-scale dynamical regime.”

[Printer-friendly version](#)[Discussion paper](#)

In addition, we do not look at the diurnal variations: the stationarity assumption in our simple model would be violated. We now explain this in the abstract: “the steady-state assumption restricts the application of this model to time scales longer than daily.” and in section 2.1: “We assume that the SCL is at steady state. For example, its depth is constant. Since the SCL properties may exhibit a diurnal cycle (Duynkerke et al. (2004)), this hypothesis restricts the application of this model to time scales longer than daily. “

24) P. 27, L. 28: a reference to a more technical paper such as Aemisegger et al. 2012 AMT, would be nice here.

We add this reference

Small technical comments:

- 1) P. 2, L. 22 : “suffers **from** a low bias”
- 2) P. 3, L. 28: “capturing **the** second-order...”
- 3) P. 8, L. 26: no parenthesis after B)
- 4) P. 12, L.7: “based” -> “biased”
- 5) P. 15, L . 1: Figure 8d
- 6) P. 15, L. 4: “Values **of** alphaeff...”
- 7) P. 15, L. 5: using a fractionation coefficient alpha eq **as** a function of temperature”
- 8) P. 17, L. 14: space missing between rorig and (
- 9) P. 18, L. 1: “Overall, **the** results...”

We correct all these mistakes.

[Printer-friendly version](#)

[Discussion paper](#)



10) In general, the authors do not consistently use B15 for Benetti et al. (2015)

We now use B15 consistently.

11) P. 21, L. 2: “with **the** strongest inversion”

12) P. 27, L. 27: measurement errors

13) P. 28, L. 2: “if **we** measure...”

14) P. 29, L. 14: very precise

15) P. 30, L. 11: **from** which altitude the air comes

We correct all these mistakes.

2 Reviewer 2

We thank reviewer 2 for his/her comments.

This paper presents a simple box model solving the water isotope budget in the sub-cloud layer to quantify the relative contributions of sea surface temperature, relative humidity, mid-tropospheric depletion, and the fraction of moisture from the free troposphere (r_{orig}) on the variability of δD in near-surface water vapor (δD_0). The contribution of r_{orig} is further separated into contributions of specific humidity at the surface, and the height (z_{orig}), relative humidity and temperature from which the free tropospheric air originates. z_{orig} is found to be an important factor explaining the seasonal-spatial and daily variations of δD_0 . This means that measurements of δD_0 , if precise enough, can potentially be used to estimate z_{orig} and distinguish between different mixing processes in the atmosphere.

[Printer-friendly version](#)

[Discussion paper](#)



The paper is interesting and well written, and it nicely demonstrates the use of measuring water vapor isotopes on short time scales. The box model's theoretical framework is described in detail and its drawbacks are clearly identified by the authors. I only have a few comments about the methods, the rest are mainly ideas for clarifying the paper. I recommend that the paper be published after minor revisions.

General comments

1) I like the method for quantifying the contributions of different factors by linear regression. I see how this works when the contributing factors have the same units as the variable of interest, which was the case in the previous studies that used this method and are cited in this paper (Risi et al. 2010, Oueslati et al., 2016). Here the different factors all have different units, and the slope therefore depends on the units, or how much the components vary. I assume this was accounted for somehow, as the slopes in the tables are all unitless, but it is not clear from the text, and makes me a bit skeptical about the results. More explanation on that would be useful.

The contributing factors have the same units as the variable of interest, i.e. permil. We now clarify this in the text: "To understand what controls the δD_0 spatio-temporal variations, δD_0 is decomposed into 4 contributions based on Eq. (9). First, we define $r_{orig,basic} = 0.6$, $\alpha_{eff,basic} = 1.09$, $SST_{basic} = 25^\circ\text{C}$, $h_{0,basic} = 0.8$ as a basic state. We call $\delta D_{eq9}(r_{orig}, \alpha_{eff}, SST, h_0)$ the function giving δD_0 as a function of r_{orig} , α_{eff} , SST and h_0 following the Eq. (9). The relative contributions of r_{orig} , α_{eff} , SST and h_0 to δD_0 variations are estimated as $\delta D_{eq9}(r_{orig}, \alpha_{eff,basic}, SST_{basic}, h_{0,basic})$, $\delta D_{eq9}(r_{orig,basic}, \alpha_{eff}, SST_{basic}, h_{0,basic})$, $\delta D_{eq9}(r_{orig,basic}, \alpha_{eff,basic}, SST, h_{0,basic})$ and $\delta D_{eq9}(r_{orig,basic}, \alpha_{eff,basic}, SST_{basic}, h_0)$ respectively. All these components have the same units as δD_0 (permil). "

[Printer-friendly version](#)
[Discussion paper](#)


2) As stated in the paper, the methods rely on the assumption that the δD profile follows a Rayleigh-like line, and that there is no effect of rain evaporation. Figures 7 and 8 show that the δD profile is often closer to a mixing line than a Rayleigh line, and the large contribution of r_{orig} mainly comes from ascending regions, where clouds are most likely precipitating. It would be nice to see some quantification of how this impacts the results. A possible way to do this is to remove days/locations where the RMSE of the mixing line is smaller than the RMSE of the Rayleigh line and where there is precipitation, then repeat the analysis for these new fields and add the results in brackets in Tables 1, 2 and as dotted lines in Figures 10, 12.

We are working on quantifying the effect of rain evaporation on our results using the equation in the appendix B and diagnostics of η and α_{re} from LMDZ. We do not have the results yet. Depending on our results, we will decide what is the best way to show this quantification in the paper.

3) The paper presents the new box model as an extension of the model by Benetti et al. (2015), which is technically true, but can be a bit misleading because its application is different. Rather than predicting δD_0 from z_{orig} , it predicts z_{orig} from δD_0 and therefore requires δD_0 to be known. This means it cannot be applied to initialize Rayleigh models like the model by Benetti et al. (2015), which assumes constant z_{orig} . This could be written more clearly (e.g., from the abstract it seems like the model can be used to predict δD_0 , which is only possible if z_{orig} is known).

The model can be used both ways, either to predict δD_0 as a function of z_{orig} , or to estimate z_{orig} from δD_0 . If someone wants to initialize Rayleigh models with it, one can

[Printer-friendly version](#)[Discussion paper](#)

make assumptions on z_{orig} . We write: “We propose a steady-state analytical model to predict δD_0 as a function of ... and the altitude from which the free tropospheric air originates (z_{orig}). “

4) Changing some of the colors and colormaps could make the figures easier to understand. For example, I think the contributions of different factors and how they add up in Figures 9 and 11 would be more intuitive with a perceptually uniform colormap going from light to dark colors. Also, the red and pink lines in Figures 3, 4, 7, 10, 12, 15 look very similar to each other. It would be good to use a different color for one of them.

- I read about color scales and I changed the color scales from rainbow to single hue in all maps.

- We now change the pink into purple in all the Figures.

Specific comments

P1 L13: [D]/[H] instead of [HDO]/[H2O]

We modify.

P2 L22: high bias instead of low bias?

Corrected.

P2 L30: Please introduce the abbreviation for LCL

Done.

P3 L4: pointed **out** the important role

Corrected.

P3 L15: “We do not call it entrained”: The word entrained/entrainment still appears a few times in the text (e.g. in Fig. 2, the title of section 4.4)

We modify all the occurrences of “entrainment”, except when it really refers to entrainment.

P3 L23: during **a** field campaign, global outputs **of** an isotope-enabled GCM.

Corrected.

P3 L24: “at the global scale”: Really? There are no global maps. Are the numbers in Tables 1, 2 and the lines in Figures 10, 12 from global output, or from the region shown on the maps?

We now modify by “in the Tropics”. We precise in the captions for these Tables and Figures: “All seasons and locations over tropical oceans ($30^{\circ}N - 30^{\circ}S$, ocean fraction > 80%) are considered.”

P3 L28: capturing **the** second-order parameter d-excess

Corrected.

P3 L32: “MJ79 already performs quite well for d-excess”: Pfahl and Wernli (2009) would probably disagree.

Printer-friendly version

Discussion paper



Now we modify as: “since MJ79 the effect of convective mixing is larger on d-excess than on δD (Risi et al. (2010); Benetti et al. (2014)).” which does not contradict Pfahl and Wernli (2009).

P5 L23: $r \rightarrow r_{orig}$

Corrected.

P6 L20: measurements

Corrected.

P6 L20: “Therefore, variations of δD_0 that are mediated by q_0 or h_0 do not interest us”: But δD in the FT is prescribed as a function of q (confusing).

We remove this confusing sentence and we write: “We attempt to express neither h_0 as a function of q_0 as in B15, and nor the q profile as a function of q_0 ”.

P8 L11: Refer to l'Hopital's rule?

Now we add: “(l'Hopital's rule was used to calculate this limit).”

P8 L21: follows as mixing line

Corrected.

P9: Fig.3: $\alpha_{eff} = \alpha_{eq}$ instead of $\alpha_{eff} = 1/\alpha_{eq}$

Corrected.

P11 L25: “Only profiles during the ascending phase of the balloons are considered”: (Why?)

We now justify this choice: “Only profiles during the ascending phase of the balloon are considered, because the descent phase is often located far away from the initial launch point (McGrath et al. (2006); Seidel et al. (2011)). “

P11 L27 (title): write somewhere that these results are based on LMDZ output (not observations)

Now we write: “Here we explain how z_{orig} is estimated based on LMDZ outputs.”. Later in the sub-section, we write: “When estimating z_{orig} from observations, we follow the same methodology except that...”.

P12 Fig. 5: Describe abbreviations (LCL, EIS, SCL) in caption.

Done

P12 L2: “if the end member is defined below 500hPa (e.g. 600hPa) results are not always reasonable”: In what sense? Why?

Now we write: “However, the end member should be defined above 500 hPa to ensure that it is well above boundary layer processes. If the end member is defined below 500 hPa (e.g. 600 hPa), there are cases where q increases with altitude ($q_f > q_0$) due to horizontal advection or convective detrainment from nearby moister regions;

meanwhile, δD decreases monotonically, leading to unrealistic values for α_{eff} .”

P15 Fig. 7: What meteorological conditions do these examples represent? Would it be possible to show all (/more) simulated profiles in the background, e.g. in some transparent color, to get a better feeling for the variability? Also, I suggest adding markers to highlight where the levels are.

- Now we explain in the caption how these examples were selected (see response to rev 1).

- We will give more information on the type of meteorological conditions during these examples.

- This comments joins that from rev 1 who asks for a better documentation of the spatio-temporal variability among profiles. We will work on this question during the revision time to find the most adequate (and also concise) way to document and illustrate this variability.

- Now we add markers to highlight model levels.

P15 L1: Figure 8d instead of 8c.

Corrected

P15 L5: α_{eq} as a function of temperature

Corrected

P16 Fig. 8: in boreal winters of all years

Corrected

P17 L22: “in the cold upwelling regions”: for example where?

Now we add: “cold upwelling regions, for example off Peru or Namibia”

P17 L23: probably reflects

Corrected (here and also elsewhere)

P17 L24: “the effect of r_{orig} can be seen on the composites as a function of EIS and not as a function of ω_{500} ”: I don’t see this, please elaborate.

We modify as: “We note that higher r_{orig} in regions of stronger EIS contributes to the decrease of δD_0 with EIS (slightly decreasing green curve in Fig. 10b), but it does not contribute to the decrease of δD_0 with ω_{500} (flat green curve in Fig. 10a).”

P17 L30: followed by h_0 (23%), r_{orig} (16%), ...

Corrected

P18 Fig. 9: Are the correlations significant everywhere? Otherwise, add hatching where not significant?

In Fig. 9, we do not show the correlations, but rather the contributions on a δD_0 scale. When a map shows nearly constant values, it means that the contribution to δD_0 spatial

variations is small. When a map shows patterns that are similar to the simulated δD_0 , it means that the contribution to δD_0 spatial variations is large. We add this explanation to the caption: “When a map shows patterns that are similar to the simulated δD_0 , it means that the contribution to δD_0 spatial variations is large.”

The spatial-seasonal correlations are shown in Table 1. We now write between brackets when correlations are not statistically significant at 99%. We write in the caption: “The threshold for the correlation coefficient to be statistically significant at 99 % is 0.15 or lower in all cases. We write correlation coefficient and slope values between brackets when they are not significant at 99%.”

P19 Fig.10: ω_{500} (hPa/d)

Corrected

P20 Tab. 2: q_0 seems to be important in Fig. 12, but the slope is 0.0 here, h_0 seems to be unimportant in Fig. 12 but slope is 0.91 here. Why is that?

We now explain this: “Note that this effect can be seen only in most stable regions, but when considering all subsiding regions, the contribution is near zero (Table 2). “

P20 L1: “it would translate into a lower z_{orig} .”: Why?

We now explain this better at several places: section 3.3: “For example, in case of deep convection with depleting rain evaporation, a larger r_{orig} is necessary to match the depleted δD_0 , and a lower z_{orig} is necessary to match this large r_{orig} .” section 4.3: “if the large r_{orig} was purely an artifact of the neglect of rain evaporation, it would translate totally into a lower z_{orig} , since a lower z_{orig} is necessary to match a larger r_{orig} ”

P22 Fig.12: ω_{500} (hPa/d)

Corrected

P25 L6: the cruises goes

Corrected

P25 L8: “when considering only the 6 data points when $z_{orig} < 2000\text{m}$ ”: Rationale behind this?

We now clarify what we mean: “Remarkably, there are 6 days when z_{orig} coincides with z_i with a root means square error of 31 and correlation coefficient of 0.996 (Fig. 15c). This indicates that the air exactly comes from the inversion layer. When recalling that z_{orig} and z_i are estimated from completely independent observations, the coincidence is remarkable and lends support to the fact that on these days, our z_{orig} estimate is physical. However, there remains 9 days when z_{orig} is much higher than z_i . This may reflect more penetrative downdrafts as we approach deeper convective regimes. But it may also be an artifact of our neglect of horizontal advection. For example, on these days which are characterized by lower h_0 , neglecting the advection of enriched water vapor from nearby regions with higher h_0 could be mis-interpreted as lower r_{orig} and thus higher z_{orig} .”

P25 L14: ... at the seasonal-spatial and daily scale is the proportion of the water vapor in the SCL that is originates from above

Corrected

Printer-friendly version

Discussion paper



P26 Fig. 15: $r \rightarrow r_{orig}$

Corrected

P27 L1: there \rightarrow they

Corrected

P27 L13: the **temporal** variability of α_{eff} . Is it possible to estimate the uncertainty from the spatial variability of α_{eff} as well (in the vertical, i.e. how much the δ profile differs from a Rayleigh line with constant α_{eff})?

During the revision time, we will estimate this source of error. If the δD profile doesn't follow a Rayleigh line with constant α_{eff} , an analytical solution is not guaranteed, but a numerical solution can be found as long as δD doesn't follow a mixing line. In the general case:

$$R_0 = \frac{\left(1 - \frac{q(z_{orig})}{q_0}\right) \cdot R_{oce}/\alpha_{eq} + \alpha_K \cdot (1 - h_0) \cdot \frac{q(z_{orig})}{q_0} \cdot R(z_{orig})}{\left(1 - \frac{q(z_{orig})}{q_0}\right) \cdot h_0 + \alpha_K \cdot (1 - h_0)} \quad (2)$$

We can thus numerically estimate the value for z_{orig} that yields the simulated R_0 , given the simulated vertical profiles of q and R . We will compare this result with that obtained when assuming a Rayleigh line with constant α_{eff} .

P27 L21: estimating z_{orig} from δD_0 measurements on a daily basis (?)

Interactive
comment

Printer-friendly version

Discussion paper



Now we write: “estimating z_{orig} from daily δD_0 measurements cannot be useful unless we measure δD profiles on a daily basis as well.”

P28 L2: and if **we** measure

Corrected

P28 L3: swap trade-wind cumulus and strato-cumulus clouds

Corrected

P29 L14: very **precised** estimates

Corrected

P29 L18: the altitude from which the air **is** originates, and is not **to biased** by

Corrected

References

- Benetti, M., Reverdin, G., Pierre, C., Merlivat, L., Risi, C., and Vimeux., F. (2014). Deuterium excess in marine water vapor: Dependency on relative humidity and surface wind speed during evaporation. *J. Geophys. Res.*, 119:584–593, DOI: 10.1002/2013JD020535.
- Bony, S., Bellon, G., Klocke, D., Sherwood, S., Fermepin, S., and Denvil, S. (2013). Robust direct effect of carbon dioxide on tropical circulation and regional precipitation. *Nature Geoscience*, 6(6):447–451.

Printer-friendly version

Discussion paper



- Bony, S., Dufresne, J.-L., Le Treut, H., Morcrette, J.-J., and Senior, C. (2004). On dynamic and thermodynamic components of cloud changes. *Climate Dynamics*, 22:71–86.
- Dee, S. G., Nusbaumer, J., Bailey, A., Russell, J. M., Lee, J.-E., Konecky, B., Buening, N. H., and Noone, D. C. (2018). Tracking the strength of the walker circulation with stable isotopes in water vapor. *Journal of Geophysical Research: Atmospheres*, 123(14):7254–7270.
- Duykerke, P. G., de Roode, S. R., van Zanten, M. C., Calvo, J., Cuxart, J., Cheinet, S., Chlond, A., Grenier, H., Jonker, P. J., Köhler, M., et al. (2004). Observations and numerical simulations of the diurnal cycle of the eurocs stratocumulus case. *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130(604):3269–3296.
- Emanuel, K., Neelin, D., and Bretherton, C. (1994). On large-scale circulations in convecting atmospheres. *Quarterly Journal of the Royal Meteorological Society*, 120:1111–1143.
- Garratt, J. R. (1994). The atmospheric boundary layer. *Earth-Science Reviews*, 37(1-2):89–134.
- Lacour, J.-L., Flamant, C., Risi, C., Clerbaux, C., and Coheur, P.-F. (2017). Importance of the saharan heat low in controlling the north atlantic free tropospheric humidity budget deduced fromiasi δd observations. *Atmospheric Chemistry and Physics*, 17:9645–9663.
- Masunaga, H. and Sumi, Y. (2017). A toy model of tropical convection with a moisture storage closure. *Journal of Advances in Modeling Earth Systems*, 9(1):647–667.
- McGrath, R., Semmler, T., Sweeney, C., and Wang, S. (2006). Impact of balloon drift errors in radiosonde data on climate statistics. *Journal of climate*, 19(14):3430–3442.
- Oke, T. R. (1988). *Boundary layer climates*. Halsted press, New York.
- Risi, C., Landais, A., Bony, S., Masson-Delmotte, V., Jouzel, J., and Vimeux, F. (2010). Understanding the 17O-excess glacial-interglacial variations in Vostok precipitation. *J. Geophys. Res.*, 115, D10112:doi:10.1029/2008JD011535.
- Seidel, D. J., Ao, C. O., and Li, K. (2010). Estimating climatological planetary boundary layer height from radiosonde observations: comparison of methods and uncertainty analysis. *J. Geophys. Res.*, 115, D16113.
- Seidel, D. J., Sun, B., Pethey, M., and Reale, A. (2011). Global radiosonde balloon drift statistics. *Journal of Geophysical Research: Atmospheres*, 116(D7).
- Siebert, P., Beyrich, F., Gryning, S. E., Joffre, S., and Rasmussen, A. (2000). Review and intercomparison of operational methods for the determination of the mixing height. *Atmos. Environ.*, 34:1001–1027.

[Printer-friendly version](#)[Discussion paper](#)

- Sorbjan, Z. (1989). *Structure of the atmospheric boundary layer*. Prentice Hall, Englewood Cliffs, N. J.
- Stull, R. B. (1988). *An intruduction to boundary layer meteorology*. Dordrect Kluwer.
- Wood, R. and Bretherton, C. S. (2006). On the relationship between stratiform low cloud cover and lower-tropospheric stability. *Journal of climate*, 19(24):6425–6432.
- Yanai, M., Esbensen, S., and Chu, J.-H. (1973). Determination of bulk properties of tropical cloud clusters from large-scale heat and moisture budgets. *Journal of the Atmospheric Sciences*, 30(4):611–627.

[Printer-friendly version](#)[Discussion paper](#)