

# Measurement of aerosol properties during wintertime in Beijing

Misti Levy Zamora<sup>1,3,†</sup>, Jianfei Peng<sup>1,2,†</sup>, Min Hu<sup>2,\*</sup>, Song Guo<sup>1,2</sup>, Wilmarie Marrero-Ortiz<sup>1</sup>, Dongjie Shang<sup>2</sup>, Jing Zheng<sup>2</sup>, Zhuofei Du<sup>2</sup>, Zhijun Wu<sup>2</sup> & Renyi Zhang<sup>1,2,\*</sup>

5 <sup>1</sup>Department of Atmospheric Sciences and Department of Chemistry, Center for Atmospheric Chemistry and the Environment, Texas A&M University, College Station, TX 77843, USA

<sup>2</sup>State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

<sup>3</sup>Department of Environmental Health and Engineering, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe St., Baltimore, Maryland 21205, USA

10

† These authors contribute to this paper equally.

\*Correspondence to: Min Hu (minhu@pku.edu.cn); Renyi Zhang (Renyi-Zhang@tamu.edu)




Figure S1. A comparison of the mixing layer depth (m) determined by the HYSPLIT model and the mass concentration ( $\mu\text{g m}^{-3}$ ) in (A) Autumn 2013 and (B) Winter 2015.