

1 **Supporting Information for**

2

3 **Chemical Composition and Hydrolysis of Organic Nitrate Aerosol formed from Hydroxyl and**
4 **Nitrate Radical Oxidation of α -pinene and β -pinene**

5

6 Masayuki Takeuchi¹ and Nga L. Ng^{2,3*}

7


8 ¹School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, Georgia, 30332,
9 USA

10 ²School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, Georgia,
11 30332, USA

12 ³School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, 30332,
13 USA

14 *Corresponding author: ng@chbe.gatech.edu


15



16

17 Figure S1. FIGAERO-HR-ToF-I-CIMS time-series data of select organic nitrate aerosols (i.e.,
 18 $C_{10}H_{13,15,17}NO_{5-8}$). Shown are the data from Exp. 3 (α -pinene+OH), though very similar trends are observed
 19 for β -pinene+OH from Exp. 6.


20



21


22 Figure S2. Comparison of FIGAERO-HR-ToF-I-CIMS mass spectra of organic nitrate aerosols (i.e.,
 23 $C_{10}H_{11,13,15,17,19,21}NO_{4-11}$) with ambient measurement data (Lee et al., 2016; Huang et al., 2019). All mass
 24 spectra are normalized to the maximum signal.

25




27

28 Figure S3. Concentrations of NO_3 and $\text{NO}_{3,\text{Org}}$ derived from two independent methods. “Excess NH_4 ”
29 refers to a method to apportion the inorganic nitrate contribution to NO_3 based on the increase in NH_4 and
30 “ $\text{NO}^+/\text{NO}_2^+$ ” refers to the method based on its ratio (Farmer et al., 2010).



31
 32 Figure S4. HR-ToF-AMS time-series data of SO₄ and SMPS aerosol volume concentration in Exp. 3 (low
 33 RH, dry AS), Exp. 4 (high RH, dry AS), and Exp. 5 (high RH, wet AS). A significant difference in the
 34 volume concentration among the experiments indicates the deliquesced nature in Exp. 5 (high RH, wet AS)
 35 and the effloresced nature of seed particles in Exp. 4 (high RH, dry AS).


36



37

38 Figure S5. Time-series data of NO_3/Org in Exp. 15 normalized to that in Exp. 14 based on the method
39 present in prior study (Boyd et al., 2015).

40



41

42 Figure S6. Average thermogram of organic aerosol (i.e., organic nitrate and non-nitrated organic aerosol)

43 at the peak of SOA growth in Exp. 3-5. Signals are normalized to the maximum signal for each

44 experiment.

45 Table S1. Summary of R_{AN} and R_{ON} applied to derive the concentration of $NO_{3,Org}$. R_{AN} from 300 nm
46 ammonium nitrate particle at the nearest date is used unless noted otherwise. R_{ON}/R_{AN} used for high RH
47 experiments are taken from the same experiment but under low RH experiments.

ID	R_{AN}	R_{ON}	R_{ON}/R_{AN}
1	2.87	6.25	2.18
2	2.87	6.25	2.18
3	2.80	5.75	2.05
4	2.80	5.75	2.05
5	2.88	5.91	2.05
6	3.06	5.02	1.64
7	3.06	5.02	1.64
8	3.24	8.64	2.67
9	2.87	7.66	2.67
10	2.70	8.24	3.05
11	2.80	8.55	3.05
12	2.80	8.60	3.07
13	2.88	8.85	3.07
14	2.87	7.13	2.48
15	2.56 ^a	6.36	2.48

48 ^a R_{AN} is taken from 30-min average of NO^+/NO_2^+ during seed only period as the R_{AN} from the calibration at
49 the nearest date is slightly higher.

50

51

52 Table S2. Fraction of pON based on the number of carbon in each system.

System	C _{<9} pON	C ₁₀ pON	C ₁₁₋₁₉ pON	C ₂₀ pON
α -pinene+OH	47 %	52 %	1 %	0 %
β -pinene+OH	58 %	41 %	1 %	0 %
α -pinene+NO ₃	11 %	26 %	8 %	54 %
β -pinene+NO ₃	10 %	58 %	4 %	28 %

53

References

- 54
55
56 Boyd, C. M., Sanchez, J., Xu, L., Eugene, A. J., Nah, T., Tuet, W. Y., Guzman, M. I., and Ng, N. L.:
57 Secondary organic aerosol formation from the beta-pinene+NO₃ system: effect of humidity and peroxy
58 radical fate, *Atmos Chem Phys*, 15, 7497-7522, 10.5194/acp-15-7497-2015, 2015.
- 59 Farmer, D. K., Matsunaga, A., Docherty, K. S., Surratt, J. D., Seinfeld, J. H., Ziemann, P. J., and Jimenez,
60 J. L.: Response of an aerosol mass spectrometer to organonitrates and organosulfates and implications for
61 atmospheric chemistry, *P Natl Acad Sci USA*, 107, 6670-6675, 10.1073/pnas.0912340107, 2010.
- 62 Huang, W., Saathoff, H., Shen, X. L., Ramisetty, R., Leisner, T., and Mohr, C.: Chemical Characterization
63 of Highly Functionalized Organonitrates Contributing to Night-Time Organic Aerosol Mass Loadings and
64 Particle Growth, *Environ Sci Technol*, 53, 1165-1174, 10.1021/acs.est.8b05826, 2019.
- 65 Lee, B. H., Mohr, C., Lopez-Hilfiker, F. D., Lutz, A., Hallquist, M., Lee, L., Romer, P., Cohen, R. C., Iyer,
66 S., Kurten, T., Hu, W. W., Day, D. A., Campuzano-Jost, P., Jimenez, J. L., Xu, L., Ng, N. L., Guo, H. Y.,
67 Weber, R. J., Wild, R. J., Brown, S. S., Koss, A., de Gouw, J., Olson, K., Goldstein, A. H., Seco, R., Kim,
68 S., McAvey, K., Shepson, P. B., Starn, T., Baumann, K., Edgerton, E. S., Liu, J. M., Shilling, J. E., Miller,
69 D. O., Brune, W., Schobesberger, S., D'Ambro, E. L., and Thornton, J. A.: Highly functionalized organic
70 nitrates in the southeast United States: Contribution to secondary organic aerosol and reactive nitrogen
71 budgets, *P Natl Acad Sci USA*, 113, 1516-1521, 10.1073/pnas.1508108113, 2016.

72