Reply to reviewers' comments on "History of desert dust deposition recorded in the Elbrus ice core" We would like to thank both reviewers for their comments that help us to improve and clarify the manuscript.

Please note that this is a companion paper of another manuscript submitted to ACP <u>https://www.atmos-chem-phys-discuss.net/acp-2019-402/</u> Preunkert et al., "The Elbrus (Caucasus, Russia) ice core glaciochemistry to reconstruct anthropogenic emissions in central Europe: The case of sulfate."

Some additional text was added to this manuscript as suggested by the reviewer of the Preunkert et al. paper (see section 4.3). Figure 5 was changed and an additional Figure 7 was added following the recommendation of one of the reviewers as well.

10 Reply to

5

25

Anonymous Referee #1

Received and published: 5 June 2019

The manuscript by Kutuzov et al. presents a calcium-dust proxy record from an ice core drilled on Mt. Elbrus, Caucasus,

- 15 spanning the time period 1774-2013 CE. The discussion on the dust proxy include separating the background signal and the main dust events, evaluating the frequency and amplitude of dust events, establishing a relation with the potential dust sources by means of analysis of atmospheric circulation patterns and climate indices. I found the dataset very interesting per se, which warrants publication in ACP. The discussion and interpretation of the record is quite detailed and includes very interesting findings, following an approach established by the same group of authors. However, I also found that one
- 20 relevant issue, related to the almost 10-fold increase in snow accumulation with potential implications on the interpretation of the proxy record, is not discussed. Therefore I recommend a major review of the manuscript. General comments

After reading the manuscript, I am not sure whether the trends mostly reflect changes in deposition/accumulation processes rather than changes in dust emissions. This aspect is not discussed in the manuscript, while I believe it is central for the interpretation of the record. I elaborate on this consideration in the lines below.

- In Figure 1a one can be appreciate how the summer half-year thickness, expressed in meters of water equivalent, increases significantly since the beginning of the record. This implies increased snow accumulation, in addition to the expected ice compaction with depth. In fact the "companion" discussion paper by Preunkert et al., reports a "decrease of the net annual snow accumulation from 1.5 mwe (0.8 mwe in summer and 0.7 mwe in winter) near the surface to 0.18 mwe (0.15 mwe in
- 30 summer and 0.03 mwe in winter) at 157 m depth". Therefore we see an almost 10-fold increase in snow accumulation rates along the core. The authors discussed the related potential issues in determining the the ability to detect the frequency of dust events; in order to overcome this issue, they adopted a strategy with finer sampling in the bottom sections of the core. While this precaution is an effective measure to that aim, it does not respond to the issue of whether the increased accumulation rates reflect increased precipitation and wet scavenging, in other words a larger or more frequent sampling of the

atmospheric dust loading during precipitation events. As a result, it cannot be safely concluded which effect primarily (or maybe both) determines the observed trends in the dust proxy. This kind of reasoning is partly grounded in the long-standing debate on whether for instance dust concentrations or deposition fluxes are a better proxy for atmospheric dust / dust variations (e.g. Fischer et al., 2007; Mahowald et al., 2011).

5 I recommend that these issues are throughly discussed in the manuscript, and the interpretations and conclusions weighted accordingly.

Taken into account. We thank reviewer for these general comments. One important issue was raised by the reviewer. The significant change in the accumulation rate indeed may influence the results of ice cores interpretation. However, in this manuscript we do not show or discuss any accumulation changes. The presented figures and data show only the thickness of

- 10 annual layers. In order obtain an accumulation rate the layer thickness must be corrected for the compression which occurred since it was deposited (e.g. Paterson and Waddington, 1984). This effect at Elbrus can be clearly seen at Figure 9 in (Mikhalenko et al., 2015) which shows the annual layer thickness and the Nye model fit. The monotonic decrease of annual layer thickness is an effect of layer thinning. A separate paper dedicated to the accumulation rate change in Elbrus should be submitted soon. The accumulation is calculated using the dating and available reference horizons together with
- 15 depth age modelling. When accounted for the ice layer thinning the accumulation variations are within 20-30% and there is no linear trend in accumulation change over the whole period. Therefore we do not expect any significant influence of the accumulation change on Ca2+ concentrations giving the reasonable sampling resolution. The observed trends in Ca2+ concentration cannot be explained by the changes in accumulation.

We can see that this misunderstanding was due to inconsistent wording in figure captions and in the text of two companion 20 manuscripts. We added paragraph to explain this issue.

"It show be noted that Fig. 3 show the thickness of layers and does not represent the linear change in accumulation rate. In order obtain an accumulation rate the layer thickness must be corrected for the compression which occurred since it was deposited (e.g. Paterson and Waddington, 1984) which is out of the scope of this paper."

25 Specific comments

p. 2 / lines 3-4: Please explain how

Taken into account. Text revised. "The discrepancies between models are partly explained by very limited observations of dust variability over the past and therefore limited possibilities to evaluate the model's reproducibility of the dust cycle." 2/4-5: please provide some references

30 Done. References added (e.g. Gautam et al., 2009; Chudnovsky et al., 2017; Li and Sokolik, 2018).

2/8: it would seem more precise to say that many of the archives reported in the cited manuscript show a doubling of the respective dust signals

Taken into account. Text revised.

2/9-13: Given the level of detail reported here with reference to the cited paper, it may be worth reporting other studies as well (e.g. Ginoux et al., 2012; older papers assessing the issue at the global level)

Taken into account. Text revised, reference added.

2/25: Two references are listed as Kutuzov et al. 2015 a,b. Please delete the nonrelevant one. In addition, remove from the reference list the discussion paper (Kutuzov et al., 2013).

Done

5

3/9: Make sure that the special character is properly displayed. In addition, in the legends of Figures 1 and 2, a resolution of 0.5 x 0.5 degree is reported. Which one is correct?

Both the resolutions are correct. The first one is related to NCEP/NCAR Reanalysis, the second one simply show a resolution with which the Figures 1 and 2 were produced.

3/12: Rather than aerosols, the HYSPLIT analysis reported here shows that "Elbrus glaciers receive AIR MASSES from sources . . ."

Text revised

3/14-21: Please clarify whether the density plots in Figure 2 are based solely on the the back-trajectories passing close to the ground (and what about Figure 1?). In addition, please explain how did you define the well-mixed boundary layer.

- 15 ground (and what about Figure 1?). In addition, please explain how did you define the well-mixed boundary layer. Text revised. "Density plots were calculated only for 10 day backward trajectories which descended below mixed layer depth. The depth is calculated by HYSPLIT_4 (using NCEP/NCAR Reanalysis data) for each point of backward trajectory as the height of the first exceeding of potential air temperature over surface air temperature by 2 K in the point (Draxler & Hess, 1998)."
- 20 3/25: Please report the geographical coordinates

Done

3/26: ranged from . . . AT 10 m depth . . .

Done

3/30: Could you report in a few words the main aspects of that methodology?

- 25 Done. "Cores were subsampled and decontaminated at -15°C using the pre-cleaned electric plane tool methodology described in (Preunkert & Legrand, 2013). In brief, in a first step, ice samples were cut with a band saw. After that, all surfaces of the cut samples were cleaned under a clean air bench by using a pre-cleaned electric plane tool over which the ice was slid. To control the decontamination efficiency process blank ice samples, consisting of ultrapure frozen MilliQ water were preceded regularly."
- 30 3/30-31: Please specify whether the sampling is continuous along the core

Done. Text revised.

4/8-9: Define what is meant by decontamination blank

Done. Text revised. "To control the decontamination efficiency process blank ice samples, consisting of ultrapure frozen MilliQ water were preceded regularly."

4/18-19: Please rephrase

Done. Text slightly revised

5/7-8: Please report briefly the methodology of the cited paper, i.e. how one can identify dust events on the basis of Ca2+ and acidity records.

5 *The section with this part of the text was removed. This criteria is explained further in the manuscript in section 4.1.* 5/20-22: It would be interesting to estimate/report the uncertainty arising from this assumption, and propagate it to the dust proxy.

Rejected. The sea salt Ca+2 fraction of the summer Ca2+ concentration is $1\pm0.7\%$. We show that we can neglect its influence.

10 5/22-25: Can you provide an estimate of how much this uncertainty could amount to?

Rejected. We really cannot say more than what is stated. "That percentage is clearly an upper limit since, in precipitation deposited at continental free tropospheric sites (e.g. Legrand, 2002), Na+ is not only related to sea-salt due to the presence of leachable sodium in alumino-silicate particles but also Na+ from halide evaporates present in the deserts".
This upper limit (1 and 1.5%) is low enough to completely neglect the sea-salt contribution to the calcium level.

15 5/31 - 6/3: It is not clear what you mean by "disturb" in both sentences. Please rephrase.

Taken into account. Text revised.

6/8: What do yo mean by "warm periods"? Warm years/decades? Warm seasons?

Warm seasons. Text revised.

References

20 Fischer, H., Siggaard-Andersen, M. L., Ruth, U., Rothlisberger, R., and Wolff, E. W.: Glacial/Interglacial changes in mineral dust and sea-salt records in polar ice cores: sources, transport, and deposition, Rev. Geophys., 45, RG1002, doi:10.1029/2005RG000192, 2007.

Ginoux, P., Prospero, J., Gill, T., Hsu, N., and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS deep blue aerosol products, Rev. Geophys., 50, RG3005, doi:10.1029/2012RG000388, 2012.

Mahowald, N., Albani, S., Engelstaedter, S., Winckler, G., and Goman, M.: Model insight into glacial-interglacial paleodust records, Quat. Sci. Rev., 30, 832–854, 2011

References

25

Paterson, W. S. B. and Waddington, E. D.: Past precipitation rates derived from ice core measurements: methods and data analysis, Rev. Geophys., 22, 123–130, 1984
 Mikhalenko, V., Sokratov, S., Kutuzov, S., Ginot, P., Legrand, M., Preunkert, S., Lavrentiev, I., Kozachek, A., Ekaykin, A.,

Faïn, X., Lim, S., Schotterer, U., Lipenkov, V., and Toropov, P.: Investigation of a deep ice core from the Elbrus western plateau, the Caucasus, Russia, The Cryosphere, 9, 2253-2270, https://doi.org/10.5194/tc-9-2253-2015, 2015.

History of desert dust deposition recorded in the Elbrus ice core.

Stanislav Kutuzov¹, Michel Legrand^{2,3}, Suzanne Preunkert^{2,3}, Patrick Ginot^{2,3,6}, Vladimir Mikhalenko¹, Karim Shukurov⁴, Alexey Poliukhov⁵, Pavel Toropov^{1,5}

5

10

¹ Institute of Geography, Russian Academy of Sciences, Moscow, 119017, Russia

² Université Grenoble Alpes, CNRS, Institut des Géosciences de l'Environnement (IGE), Grenoble, 38402, France

³ CNRS, Institut des Géosciences de l'Environnement (IGE), Grenoble, 38402, France

⁴ A.M. Obukhov Institute of Atmospheric Physics of Russian Academy of Sciences, Moscow, 119017, Russia

⁵ Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia

⁶Observatoire des Sciences de l'Univers de Grenoble, IRD/UGA/CNRS, Grenoble, 38400, France

Correspondence to: Stanislav Kutuzov (kutuzov@igras.ru)

Abstract. Ice cores are one of the most valuable paleoarchives. Records from ice cores provide information not only about the amount of dust in the atmosphere but also about dust sources and their changes in the past. A 182 m long ice core was

- 15 recovered from the western plateau of Mt. Elbrus (5115 m.a.s.l.) in 2009. This record was extended with a shallow ice core drilling in 2013. Here we analyse Ca²⁺ concentrations, a commonly used proxy of dust, recorded in the Elbrus ice core over the time period of 1774-2013 CE. The Ca²⁺ record reveals a quasi-decadal variability with a generally increasing trend. Using multiple regression analysis, we found a statistically significant spatial correlation of the Elbrus Ca²⁺ summer concentrations with precipitation and soil moisture content in the Levant region (specifically Syria and Iraq). The Ca²⁺ record
- also correlates with drought indices in North Africa (r=0.69 p<0.001) and Middle East regions (r=0.71 p<0.001). Dust concentrations prominently increase in the ice core over the past 200 years confirming that the recent droughts in the Fertile Crescent between 1998-2012 CE were the most severe aridity for at least the past two centuries. For the most recent 33 years (1979-2012 CE), significant correlations between Ca²⁺ and Pacific circulation indices (Pacific Decadal Oscillation, Southern Oscillation Index and Niño 4) may indicate that the increased frequency of extreme El Niño and La Niña events due to a
- warming climate extended their influence to the Middle East. It was shown that the increase of Ca^{2+} concentration in the ice core cannot be attributed to human activities, such as coal combustion and cement production.

1 Introduction

30

planetary radiation balance, atmospheric chemistry, biosphere and human health (Middleton, 2017). Despite the significance of atmospheric dust, knowledge of its regional variability and long term trends over past centuries is still poor (Mahowald et al., 2010). Dust concentration in the atmosphere depends on specific meteorological conditions, which may also be influenced by large scale circulation patterns (e.g. ENSO, NAO). Long term trends are controlled by changes in precipitation and vegetation cover in dust source regions, with the vegetation cover being dependent on both natural (climatic changes)

Dust is the most important aerosol emitted to the atmosphere by mass (Knippertz & Stuut, 2014). Atmospheric dust affects

and anthropogenic (land cover change) causes. The complexity of dust emission, atmospheric transport and deposition mechanisms result in large uncertainties in atmospheric dust models (Mahowald et al., 2007). The discrepancies between models are partly explained by very limited observations of dust variability over the past and therefore limited possibilities to evaluate the model's reproducibility of the dust cycle. However, reliable information on multiannual dust variability back to

- 5 1980 is now available from satellite data <u>(e.g. Gautam et al., 2009; Chudnovsky et al., 2017; Li and Sokolik, 2018)</u>.
 Analysis of recent aerosol variations and trends over different land and ocean regions show that despite significant trends over some major continental source regions, average values demonstrate little change in the past three decades (1980-2009) because opposite trends cancel each other out in the global average (e.g. Chin et al., 2014). Recent broad-scale assessments of changes of dust emissions based on sedimentary achieves show that global dust emissions doubled a doubling of the dust
- 10 deposition in many sedimentary achieves since the mid-18th century which was attributed to anthropogenic land use and short term climate variability (Hooper & Marx, 2018). Globally anthropogenic dust sources account for 25% of emission but this value vary and can be considerably higher, up to 75% in some regions (Ginoux et al., 2012). Climate-aerosol model simulations with the ability to separate natural and anthropogenic dust sources show that there was a 25% increase in dust emissions between the 19th century and today. These changes are attributed to climate change (56%) and anthropogenic land
- 15 cover change (40%) although the model underestimates dust concentrations in Asia, Middle East, and the US (Stanelle et al., 2014).

Records of past changes in dust concentrations are essential to better constrain connections between dust emissions and both natural and anthropogenic environmental changes. In this respect, proxy data <u>are fundamentalis vital</u>. Ice cores are natural archives of past concentrations of various impurities present in the atmosphere, including dust (e.g. Legrand & Mayewski,

- 20 1997). The particle concentrations in ice cores provide information not only on the concentrations of atmospheric dust in the past but also give insight into the strengths of the dust sources and their changes through the time. Polar ice cores from Greenland and Antarctica reconstruct the changes of dust content in the atmosphere over hundreds thousands of years at a hemispherical scale (e.g., De Angelis et al., 1997; Delmonte et al., 2002; Legrand, 1987; Petit et al., 1999; Ruth et al., 2003). In contrast, data from ice cores drilled at mid-latitude mountain glaciers proximal to arid areas reconstruct local- to regional-
- 25 scale dust aerosol emission histories over shorter timescales (e.g. Grigholm et al., 2015, 2017; Kaspari et al., 2009; Osterberg et al., 2008; Thompson, 2000; Bohleber et al., 2018).

Mineral dust from North Africa and deserts of the Middle East is regularly deposited on glaciers in the Caucasus Mountains (Kutuzov et al., 2013; Kutuzov et al., 2015). Due to its high elevation (over 5000 m.a.s.l.) and proximity to arid and semiarid areas, the Caucasus are-is a natural trap for desert dust. The absence of melt water infiltration Conditions-near the

30

summit of Elbrus ensures the preservation of a reasonably long climatic record in an ice core not affected by melt water infiltration. High accumulation rates assure high temporal resolution of the ice-core data (Mikhalenko et al., 2015). In addition to a study focusing on the long-term trend of black carbon in the Elbrus ice (Lim et al., 2017), several studies of other chemical species are currently in progress, including studies of calcium and of sulfate. Data are discussed in two companion papers including the glaciochemistry of the deep Elbrus ice core drilled in 2009 (Preunkert et al., this issue).

Here, we report changes in Ca^{2+} concentrations recorded in the Elbrus ice core between 1774-2012 CE and connections with natural variability, climatic and land use changes in the dust source regions.

2. Location, climatology, and backward air-mass trajectories

- 5 The Caucasus are situated between the Black and Caspian seas, and generally trend east-southeast, with the Greater Caucasus range often considered as the divide between Europe and Asia. The 2020 glaciers in the Caucasus cover an area of $1193 \pm 27 \text{ km}^2$ (Tielidze & Wheate, 2018). Elbrus mountain glaciers contain about 10% of Caucasus ice volume and cover an area of 112.6 km² (Kutuzov, et al., 2015) (Fig. 1). Glaciers cover an altitudinal range from 2800 to 5642 m a.s.l. with the coldest conditions present above 5200 m a.s.l. where mean summer air temperature stays below 0°C.
- To characterize possible sources of aerosols deposited on glaciers, we calculated three-dimensional backward trajectories of air parcels (elementary air particles) arriving at the Elbrus plateau (5100 m a.s.l.) using the NOAA HYSPLIT_4 trajectory model (Draxler & Hess, 1998; Stein et al., 2015) and NCEP/NCAR Reanalysis data on 2.5×2.5 <u>degree</u> grids (Kistler et al., 2001) for the 1948-2013 period. 10 day backward trajectories were calculated for every 6 hours for the whole period-, resulting in a total of ~100k modelled backward trajectories. Over this time period, <u>air particles aerosol</u> transport is defined
- 15 by the westerlies. Elbrus glaciers receive-<u>air particles highly likely polluted with aerosols aerosols</u> from sources located in Turkey and the Mediterranean, Eastern and Central Europe, the Middle East, North Africa and Southern Russia (Fig. 1b). To identify potential dust source contributions we also analysed vertical distribution of the backward trajectories. An objective criterion was chosen to extract locations of possible dust entrainment along the trajectories. The criterion is met when the air parcel is close to the ground, i.e. within the well-mixed boundary layer, allowing the uptake of mineral aerosols
- (Sodemann et al., 2006). Density plots were calculated only for 10 day backward trajectories which descended below mixed layer depth. The depth is calculated by HYSPLIT 4 (using NCEP/NCAR Reanalysis data) for each point of backward trajectory as the height exceeding potential air temperature over surface air temperature by 2 K (Draxler & Hess, 1998). Density plots using NCEP/NCAR Reanalysis data were calculated for 10 day backward trajectories which fell within the boundary layer. The majority of the backward trajectories show a south-west origin with the highest frequency over the Middle East, castern Mediterranean and North Africa in all seasons. In winter (December-February) air masses tend to come
- from more remote locations while summer (June August) reveals possible transport from the Caspian Sea region and Southern Russia (Fig. 2).

3. Methodology

3.1. Ice core analysis

A 181.8 m long ice core was recovered at the western plateau of Elbrus, in the central Caucasus (43°20'53.9" N, 42°25'36.0"E; 5115 m asl) during August-September 2009. Drilling was performed in a dry borehole with a lightweight 5 electromechanical drilling system. Borehole temperatures ranged from -17 °C to-at 10 m depth to -2.4 °C at 181 m (Mikhalenko et al., 2015). Ice cores were packed in insulated core boxes and shipped in a frozen state to the cold laboratory of the Institute des Géosciences de l'Environnement (IGE) in Grenoble, France for analyses. A total of 3724 samples were prepared from the surface (2009) down to 168.6 m of the Elbrus core. Cores were subsampled and decontaminated at -15°C using the pre-cleaned electric plane tool methodology described in (Preunkert & Legrand, 2013). In brief, in a first step, ice 10 samples were cut with a band saw. After that, all surfaces of the cut samples were cleaned under a clean air bench by using a pre-cleaned electric plane tool over which the ice was slid. Sampling was continuous along the core. To control the decontamination efficiency process blank ice samples, consisting of ultrapure frozen MilliO water were preceded regularly. Cores were subsampled and decontaminated under a clean bench located in a cold room using the methodology described in (Preunkert & Legrand, 2013). Due to the glacier compression with depth, we applied a variable sampling resolution of 10 cm 15 from 0 157 m, and then a sampling resolution of every 2 cm below 157 m depth. As a result (Fig. 3), the temporal resolution remains relatively consistent throughout the core, with 12 samples per summer over the 1950 2010 CE time

period to 14 samples each summer over the 1900-1950 CE time period. In addition, a 20.5 m long ice core was extracted in June 2013 at the same location to expand the existing record from 2009 to 2012. In total 515 samples of the firn core were analysed (85 samples per year).

We determined cations (Na⁺, K⁺, Mg²⁺, Ca²⁺, and NH₄⁺) using a Dionex ICS-1000 chromatograph equipped with a CS12 separator column. For anions, a Dionex 600 equipped with an AS11 separator column was used with an eluent mixture of H₂O, NaOH at 2.5 and 100 mM, and CH₃OH. A gradient pump system allows the determination of inorganic species (Cl⁻, NO₃⁻, and SO₄²⁻) as well as short-chain carboxylates (Legrand et al., 2013). For all investigated ions, ice decontamination blanks were insignificant compared to respective concentrations in the ice cores.

25 **3.2. Dating the** ice core

Seasonal ice-core stratigraphy of chemical parameters and ice-core dating based on annual layer counting of the deep Elbrus ice core is described in detail in (Mikhalenko et al., 2015). The seasonal oscillations of NH_4^+ and succinic acid allow dating the core with seasonal resolution. Based on the ammonium and succinic acid profiles, each annual layer was divided into two parts corresponding to snow deposition under winter conditions and spring-summer-autumn conditions (Preunkert et al., 2015).

2000; Legrand et al., 2013; Mikhalenko et al., 2015). The annual counting was confirmed with 1 year uncertainty over the last one hundred years by reference horizons from a tritium peak (1963) and the Katmai 1912 horizon (Mikhalenko et al., 2015). Though the annual counting becomes less straight forward prior to 1860, Mikhalenko et al. (2015) reported an ice age

of 1825 at 156.6 m depth (i.e., 122.3 meter water equivalent, mwe). This time scale is consistent with the presence of a volcanic horizon at approximately 1833-1840 such as which can be attributed to the eruption of the Coseguina volcano (in 1835). More recently, Preunkert et al. (this issue) extended the annual counting down to 168.5 m depth (i.e., 131.6 mwe) using the ammonium and succinate records demonstrating that the seasonally-resolved record extends back to 1774.

5 Due to the glacier compression with depth, we applied a variable sampling resolution of 10 cm from 0 – 157 m, and then a sampling resolution of every 2 cm below 157 m depth. As a result (Fig. 3), the temporal resolution remains relatively consistent throughout the core, with 12 samples per summer over the 1950-2010 CE time period to 14 samples each summer over the 1900-1950 CE time period.

<u>It show be noted that Fig. 2 show the thickness of annual layers and does not represent the linear change in accumulation</u>
 <u>rate. In order obtain an accumulation rate the layer thickness must be corrected for the compression which occurred since it</u>
 was deposited (e.g. Paterson and Waddington, 1984) which is out of the scope of this paper.

In addition, a 20.5 m long ice core was extracted in June 2013 at the same location to expand the existing record from 2009 to 2012. In total 515 samples of the firn core were analysed (85 samples per year).

For the 2013 ice core the succinic acid data were not available. We therefore used a combination of NH_4^+ , BC and $\delta^{18}O$ 15 profiles for annual counting and seasonal dissection. These results suggest that the 2013 firn core extends back to 2007, as confirmed by the similar patterns of NH_4^+ , Ca^{2+} and $\delta^{18}O$ between the first 3.8 mwe of the 2009 core and the 7.2-10.7 mwe interval of the 2013 core (Lim et al., 2017).

3.3. Data presentation

The amount of dust in ice cores depends on many factors and corresponds to the presence of dust particles in the atmosphere. Primarily, dust emissions are influenced by the characteristics of the dust source (soil type, geomorphology, soil moisture) as well as by meteorological conditions (surface winds). Once dust clouds are uplifted to the mid troposphere their transport depends on the main circulation patterns. In mountainous areas with high snow accumulation, wet deposition defines annual and seasonal aerosol concentrations. During spring and summer the majority of air masses arrive to the Elbrus site from arid areas with calcareous soils. We consider that all of the Ca²⁺ in the Elbrus ice core is of natural origin and therefore Ca²⁺

25 concentrations can be considered as a proxy for dust.

Using ammonium and succinate stratigraphy to separate the winter and summer seasons (Sect. 3.2), we determined half-year summer and winter means from 1774 to 2010 (Fig. 4). In Fig. 5, we report the seasonal cycle of calcium and ammonium averaged over two different periods of the 20th century (1900-1950 and 1950-2010). As discussed in the next section, calcium present in the Elbrus ice has two distinct origins: (1) sporadic arrival of desert dust plume events at the site and (2)

30 "background" concentrations. Combining the Ca²⁺ and acidity records, Preunkert et al., (this issue) identified samples which are likely influenced by desert dust events (616 samples on a total of 2524 samples in summer, 67 samples on a total of 1150 samples in winter). For Fig. 4, 5, and 6, we report total calcium and "background" (also denoted Ca²⁺ red., see details in section 4.1) concentrations. The difference between Ca²⁺ and Ca²⁺ red is related to dust concentration (denoted Ca²⁺ dust). To discuss the long-term winter and summer trends of calcium in relation to past climatic conditions, we also reported both total concentrations and those calculated after removal of samples suspected to contain input from large dust deposition events (Ca²⁺_{red}, Values; Fig. 6).

3.3. Climatic data analysis

- 5 In order to assess which factors drive the variations in dust content in the Elbrus ice core, we compared the dust concentrations with various climatic parameters in the potential dust sources. For climate data which may potentially influence the dust emission (temperature, precipitation, wind speed, soil moisture) we used ERA-Interim Reanalysis fields for the period of 1979-2013 (Dee et al., 2011). Since the variables used in this study are continues and normally distributed the Pearson's correlation coefficient was used. The statistical significance of the correlation was checked using Student's t-
- 10 distribution for probability. The correlations between Ca^{2+} concentrations in the ice core and each climatological parameter fields were computed.

The Standardised Precipitation-Evapotranspiration Index (SPEI) was used as a drought proxy (Vicente-Serrano et al., 2010). It was shown that this index is preferable among the other drought indices to investigate the relation of dust emission to droughts (Achakulwisut et al., 2018). We considered SPEI calculated over 1, 2, 3, 6 and 12 month. The time series of SPEI

- 15 were obtained by averaging over the regions of interest. To identify the possible factors which influence the dryness in the dust sources and dust transport to the Caucasus Mountains, we also investigated the Ca²⁺ record in relation with various circulation indices. The following indices were analyzed: North Atlantic Oscillation (NAO), Pacific Decadal Oscillation (PDO), Atlantic Multidecadal Oscillation (AMO), Tripole Index for the Interdecadal Pacific Oscillation TPI (IPO) and El Niño-Southern
- 20 Oscillation (ENSO) indexes, Southern Oscillation Index (SOI), Niño 3, Niño 3.4, Niño 4, Niño 1-2 (https://www.esrl.noaa.gov/psd/gcos_wgsp/Timeseries/). Correlations were calculated for periods of 30 years, using 5 years sliding windows. All series were detrended prior to analysis. The following indices were analyzed: PDO, SOI, AMO, NAO, <u>TPI (IPO) Niño 3, Niño 3.4, Niño 4, Niño 1-2 ().</u>

4. Results and discussion

- 25 The amount of dust in ice cores depends on many factors and corresponds to the presence of dust particles in the atmosphere. Primarily, dust emissions are influenced by the characteristics of the dust source (soil type, geomorphology, soil moisture) as well as by meteorological conditions (surface winds). Once dust clouds are uplifted to the mid troposphere their transport depends on the main circulation patterns. In mountainous areas with high snow accumulation, wet deposition defines annual and seasonal aerosol concentrations. During spring and summer the majority of air masses arrive to the Elbrus site from arid
- 30 areas with calcareous soils. We consider that all of the Ca^{2+} in the Elbrus ice core is of natural origin and therefore Ca^{2+} concentrations can be considered as a proxy for dust.

The total dust deposited at Mt. Elbrus may have three different natural components: (i) dust from local sources (nunataks, rock outcrops), (ii) sporadic arrival of large aeolian desert dust events and (iii) large scale background continuous terrigenous aerosol emissions. Apart from regions strongly impacted by sea-salt aerosols, the presence of calcium in aerosols in continental atmospheres is expected to mainly originate from exposed continental sediments. Even in polar regions

- 5 (Antarctica or Greenland), calcium in ice mainly comes from dust, where only a small fraction is related to sea-salt emitted from the ocean (De Angelis et al., 1997; Legrand, 1987). Assuming that Na⁺ present in the <u>ELB-Elbrus</u> melted ice samples is only related to sea-salt emissions, and considering the [Na⁺]/[Ca²⁺] ratio of 0.038 in seawater, we find that only 1.0 ± 0.7 % in summer (1.40 ± 1.0 % in winter) of total Ca²⁺ may be attributable to sea-salt emissions. That percentage is clearly an upper limit since, in precipitation deposited at continental free tropospheric sites (e.g. Legrand, 2002), Na⁺ is not only related to
- 10 sea-salt due to the presence of leachable sodium in alumino-silicate particles but also Na⁺ from halide evaporates present in the deserts.

The volcanic rocks of Elbrus near the drilling site do not contain calcite and we can therefore assume that Ca^{2+} present in the Elbrus ice archive information on past dust emissions, including continuous background emissions and large dust plumes reaching the site. In the following sections, we attempt to separate these two possible calcium sources.

15 4.1. Calcium concentrations during desert dust events and background conditions

The majority of the calculated backward trajectories show a south-west origin with the highest frequency over the Middle East, eastern Mediterranean and North Africa in all seasons. In winter (December-February) air masses tend to come from more remote locations while summer (June-August) reveals possible transport from the Caspian Sea region and Southern Russia (Fig. 23).

- 20 Large dust plumes originating from the Middle East and less frequently from the Sahara reach the Caucasus (Kutuzov et al., 2013). As seen in the Alps, these events disturb-impacts the chemistry of snow deposits to create calcium-rich alkaline snow layers (Wagenbach et al., 1996). Deposition of these plumes increases the concentrations of numerous chemical species in Alpine ice because either they were present in dust at the emission stage or, being acidic, they interacted with alkaline material during transport (Usher et al., 2003). Deposition of light-absorbing impurities (in particular, black carbon and dust)
- 25 plays an important role in changes of the snow and glaciers and enhance the response of the mountain cryosphere to climate changes via snow-albedo feedbacks (Ginot et al., 2014; Gabbi et al., 2015; Di Mauro et al., 2017; Skiles et al., 2018). In this work, as well as in Preunkert et al. (this issue) Elbrus samples are considered to be impacted by dust events if they
- contain more than 120 ppb of calcium and are also below the 25% quartile of a robust spline through the calculated raw acidity profile. These selection criteria result in 616 dust-<u>deposition</u> summer samples (from a total of 2524) and 67 winter
- 30 samples (from a total of 1150). Similar results were obtained when changing the calcium concentration criteria from 120 ppb to 100 or 140 ppb.

Using ammonium and succinate stratigraphy to separate the winter and summer seasons (Sect. 3.2), we determined half-year summer and winter means from 1774 to 2010 (Fig. 4). In Fig. 5, we report the seasonal cycle of calcium and ammonium averaged over two different periods of the 20th century (1900-1950 and 1950-2010). For Fig. 4, 5, and 6, we report total ealeium and "background" (also denoted Ca^{2+}_{red} , see details in section 4.1) concentrations. The difference between Ca^{2+} and Ca^{2+}_{red} is related to dust concentration (denoted Ca^{2+}_{dust}). To discuss the long term winter and summer trends of calcium in relation to past climatic conditions, we also reported both total concentrations and those calculated after removal of samples suspected to contain input from large dust deposition events (Ca^{2+}_{red} , Values; Fig. 6).

The mean concentration of Ca^{2+} was 145 ppb with a maximum of 5506 ppb. Most of the dust from long distance transport is deposited on Elbrus during warm <u>periodsseasons</u>. The average background summer Ca^{2+} concentration was 103 ppb compared to 44 ppb in winter layers. Most of the large dust deposition events also <u>occursoccur</u> during the warm season which further increases the difference between summer (178 ppb) and winter (54 ppb) layers (Table 1).

As seen in Fig. 6, most of long-term calcium trend is detected in summer with: (1) more frequent arrivals of large dust events after 1950 and (2) an increase of 100 ppb of the calcium background level after 1950. The maximum annual concentrations were found in 1999 and 2000 annual layers (980 and 850 ppb). There was a pronounced period of increased dust content in the 1960s with a following decrease in late 1970s. Exceptional peaks compared to the background occur during the

following years: 1802, 1957, 1863, 1917, 1933, 1963, 1984, 1989, 1999, 2000, 2008 and 2009.

4.2. Increasing frequency of desert dust events

5

15

20

Warm season layers contain an increase in Ca²⁺ throughout the entire investigated time period. The most pronounced increase in peak Ca²⁺ concentrations occurred after 1950s (Fig. 6). It is important to emphasize that this appearance of more frequent calcium peaks after 1950 compared to the preceding decades cannot be attributed to a decreasing time resolution of the record with a smoothing of events with depth (Fig. <u>32</u>). More specifically, the enhanced occurrence of summer calcium peaks between 1960 and 1970 compared 1950-1960 does not correspond to a significant decrease of the temporal resolution (from 9 samples over the 1960-1970 years compared to 11-12 samples over the 1950-1960 years).

In recent decades (1950-2010), over which time period more dust events were detected in the Elbrus ice, a clear spring maximum of the dust fraction is observed (Fig. 5). These dust peaks are consistent with the timing of the arrival of large dust plumes at the site (Kutuzov et al., 2013). Recent major long-range dust outbreak event over North Africa, eastern Mediterranean, and Caucasus occurred on 22 and 23 March 2018 [Solomos et al., 2018] and resulted in a significant dust deposition on glaciers.

The calcium peaks in Elbrus ice containing dust are accompanied by an increase of magnesium (Fig. 7). Interestingly, the mean $[Mg^{2+}]/[Ca^{2+}]$ ratio in Elbrus samples containing dust (0.035, Fig. 7) is similar to those in atmospheric dust aerosols

30 mean [Mg²⁺]/[Ca²⁺] ratio in Elbrus samples containing dust (0.035, Fig. 7) is similar to those in atmospheric dust aerosols from the Sahara or Middle East. For instance, Koçak et al. (2012) reported dust event related aerosol concentrations of sodium, magnesium, and calcium from two Eastern Mediterranean sites, i.e. from Erdemli (Turkey) with dust arriving from Middle East and from Heraklion (Crete) with dust from Sahara. It is important to emphasize that, as for the Elbrus ice core,

the atmospheric concentrations of these cations correspond to their water-soluble fraction (not the total fraction), which were measured with Ion Chromatography. In the case of Erdemli during Middle East dust events, Kocak et al. (2012) reported atmospheric concentrations of 7085 ng m⁻³ for Ca²⁺ and 423 ng m⁻³ for Mg²⁺ (Table 2). Since Erdemli is located at 22 m above sea level and 10 m away from the sea, in addition to the leachable fraction of magnesium from dust, a fraction of

- 5 magnesium would here come from sea-salt. To correct concentration from the sea-salt contribution, we have used the Na⁺ concentration (1148 ng m⁻³) and assumed a mean $[Na^+]/[Ca^{2+}]$ ratio in dust of 0.08 as observed in Elbrus ice samples containing dust (not shown). Thus, neglecting the sea-salt calcium contribution, we estimate a dust sodium contribution of 567 ng m-3 (0.08 x 7085 ng m⁻³). With that, and using the $[Mg^{2+}]/[Na^+]$ ratio in seawater (0.12), we estimate that 70 ng m⁻³ of magnesium are originated from sea-salt and calculate a $[Mg^{2+}]/[Ca^{2+}]$ ratio for dust aerosol close to 0.05 (Table 2). A
- 10
- similar value is obtained for aerosol at Heraklion during a Saharan dust event (0.043, Table 2). The content of Mg^{2+} in Elbrus samples impacted by dust is therefore very consistent with what is observed in atmospheric aerosol from the Eastern Mediterranean region during dust events. As discussed by Preunkert et al. (this issue), the composition of dust events reaching Elbrus in the summer is consistent with what is found in atmospheric aerosols collected at Mediterranean sites during large dust events. For instance, the [Mg²⁺]/[Ca²⁺] ratio in Elbrus samples containing dust (0.035) is similar to those in 15 atmospheric dust aerosols from the Sahara or Middle East.
 - In recent decades (1950 2010), over which time period more dust events were detected in the Elbrus ice, a clear spring maximum of the dust fraction (in April) is observed (Fig. 5). These April dust peaks are consistent with the timing of the arrival of large dust plumes at the site (Kutuzov et al., 2013).

4.3. Enhanced background concentrations

- Figure 6 demonstrates an increase in the calcium background concentrations after 1950. This increase may be influenced by 20 human activities, such as coal combustion and cement production, thereby contributing to the background calcium trend detected in the Elbrus ice. Similar impacts of anthropogenic emissions on various species in natural dust emissions, including calcium, was reported by (Kalderon-Asael et al., 2009) demonstrating that, under strong stratification in the lower atmosphere in Israel, part of the atmospheric calcium may be anthropogenic nature. At the scale of Europe, Lee and Pacyna (1999) estimated that 0.8 Tg of anthropogenic calcium are emitted per year, coal combustion contributing for 60% and 25
- cement for 30% of total. To date, these anthropogenic calcium emissions remain however one order of magnitude weaker than dust calcium emissions from North-East Africa (12 Tg yr⁻¹) or West Asia (12.7 Tg yr⁻¹) (Zhang et al., 2015). Particles emitted during both coal combustion and cement production are rich in calcium (calcite). We therefore may expect

a weaker $[Mg^{2+}]/[Ca^{2+}]$ ratio in particles emitted by these anthropogenic processes compared to that in natural dust particles.

Examination of Mg^{2+} and Ca^{2+} in Elbrus ice layers that not impacted by dust events helps to determine a possible 30 contribution of anthropogenic activities to the increasing background calcium trend in Elbrus ice. As seen in Fig. 78, the $[Mg^{2+}_{red.}]/[Ca^{2+}_{red.}]$ ratio in Elbrus summer ice is of 0.069 over the 1960-2010 compared to 0.126 over the 1774-1920. If we attribute this recent decrease of $[Mg^{2+}_{red}]/[Ca^{2+}_{red}]$ to a growing impact over the recent decades of calcium from cement (neglecting the presence of soluble magnesium in cement), the mean increase of $Mg_{red.}^{2+}$ level (10.8 ppb over the 1774-1920 years and 17.6 ppb over the 1960-2010 years) would lead to a $Ca_{red.}^{2+}$ increase of 55 ppb from 1774-1920 to 1960-2010. This value is half of the observed $Ca_{red.}^{2+}$ increase between the two time periods (71 ppb over 1774-1920 and 183 ppb over 1960-2010). These calculations suggest that half of the observed increase of background calcium concentration after 1950 are

5

attributable to anthropogenic activities. As seen in Fig. 89, even prior to 1950 the $[Mg^{2+}_{red.}]/[Ca^{2+}_{red.}]$ ratio sometimes dropped to values as low as 0.1 or less (around 1820, 1850, 1865, or 1908). Although the time period 1940-1950 has a low ratio (0.09) the calcium concentrations moderately increased (27 ppb; less than a third of the overall increase after 1950). Figure 8-9 suggests that a decrease in the $[Mg^{2+}_{red.}]/[Ca^{2+}_{red.}]$ ratio is not always associated with enhanced calcium levels but that enhanced magnesium levels also lead to an enhancement of the $[Mg^{2+}_{red.}]/[Ca^{2+}_{red.}]$ ratio. For instance, over 1853-1861, a

10 mean $[Mg^{2+}_{red.}]/[Ca^{2+}_{red.}]$ ratio of 0.22 occurs at the same time as a relatively high $Mg^{2+}_{red.}$ level (16 ppb instead of 10.8 ppb over the 1774-1920 time period).

Another indirect argument against a dominant contribution from the growing impact of cement use on the background calcium levels at Elbrus derives from an earlier study of calcium and aluminium at the Col du Dome site, Mont Blanc, France (De Angelis & Gaudichet, 1991). The authors demonstrate that both the increase of the frequency of dust arrival in

15 the 1980s and the background dust levels occurred without any coinciding decrease of the aluminium to calcium ratio. Given the low aluminium content of cement compared to desert dust, these observations suggest that the Col du Dome site is not significantly impacted by growing use of cement. We may expect that this impact is even weaker at the Elbrus site.

4.4. Climatic factors

Two major dust sources contribute mineral particles to the Elbrus glaciers, the Sahara and the Middle East. It was established
that majority of the small scale dust sources in the Middle East are located in northern Mesopotamia (northern Syria – northwestern Iraq) and the Syrian Desert (Kutuzov et al., 2013). The Levant is a major source of atmospheric dust (Middleton, 1986) with natural, anthropogenic and hydrological (intermittent streams and lakes) sources. The area between the Tigris and the Euphrates in Iraq contains natural desert dust sources while the Neinava region in Iraq was recently identified as the most active dust source in the Middle East (Moridnejad et al., 2015). In the northern Sahara, strong sporadic dust storms originating in the Libyan desert and the foothills of the Ahaggar Mountains in eastern Algeria may reach the Caucasus in the spring (Kutuzov et al., 2013).

In order to assess which factors drive the variations in dust content in the Elbrus ice core, we compared the dust concentrations with various elimatic parameters in the potential dust sources. A statistically significant spatial correlation occurs between Ca^{2+} and precipitation as well as between soil moisture content in the Middle East (Fig. 910). Unlike in the

30

¹ Middle East, no spatial correlation was found between Ca^{2+} and the amount of precipitation in North Africa. It should be noted, however, that the observation network is very sparse in the arid areas and there are much larger uncertainties in Global Precipitation Climatology Centre (GPCC) datasets in North Africa.

The Ca²⁺ record was also compared with drought indices in potential dust sources in the Middle East (32-37° N; 38-45° E) and in North Africa (20-35° N; 0-35° E) (Fig. 10).. $-Ca^{2+}$ significantly correlates with drought indices for both regions. The highest correlations were found for the SPEI 3 index which is determined by aridity over the three previous months. The correlation coefficient for the Middle East is statistically significant (p<0.001) and reaches 0.71. The correlation remains

- 5 significant after trend removal (r=0.48, p<0.001 for 19741904-2012 and 0.63 (p<0.001) for 1970-2012) (Fig. 1011b, d). Periods of dryer conditions in the Middle East region coincide with the increased Ca²⁺ concentrations and vice versa. During the period of droughts in both regions, a greater amount of mineral particles are emitted into the atmosphere, and transported to Caucasus glaciers during the spring and summer. The general increasing trend in dust content corresponds to negative trends in precipitation and increased dryness of the soil.
- Ca²⁺ correlates significantly with the SPEI 3 drought index for the North African region (r=0.69 p<0.001) but most of the correlation is defined by similar tendencies in two series. Drier conditions at the dust source region correspond to a positive trend in dust concentrations in the Elbrus ice core. The correlation weakens for the full detrended series (r=0.30, p<0.05) and increases in the later period, reaching 0.51 (p<0.001) in 1970-2012 (Fig. 11a, c). Large portions of the annual dust flux in Elbrus can be deposited during a single strong deposition event. Such events are more often associated with dust transport
- from the Sahara (Kutuzov et al., 2013). Dust emitted from North Africa can also mix with with dust from Middle Eastern sources during transport (Shahgedanova et al., 2013).
 As evident from the Elbrus ice core record, the frequency of dust events and total Ca²⁺ concentrations has increased, where

the most notable trend occurs after the 1950s. This increase corresponds with the recent analysis of centennial climate change trends in Africa which depict a significant northward expansion of the Saharan desert in the winter (Thomas & Nigam, 2018). The desert conditions are observed across larger territories in recent years and are associated with a slightly negative trend in winter precipitation and an increase in surface air temperatures. In coastal North Africa, dry conditions

dominated and rainfall decreased since the 1980s (Nicholson et al., 2018). –The number of <u>dust</u> days in the eastern Mediterranean increased over the period 1958–2006 (Ganor et al., 2010).

Two periods of maximum Ca²⁺ concentrations in the ice core correspond to the two most severe drought episodes in the

25 Middle East since the1940s, and occur during 1998–2000 and 2007-2009 (Barlow et al., 2016). The significant precipitation decline (up to 70%) during these years is explained by the dominance of the high pressure systems over the eastern Mediterranean during the winter and spring months (Trigo et al., 2010).

Our findings are supported by an analysis of the frequency of droughts in Syria. For the period from 1961 to 2009, 25 years with droughts were observed, resulting in ~40% of the years classified as drought years. On average, droughts lasted 4.5

30 years, although in the 1970s a single drought lasted 10 years (Breisinger et al., 2011). A number of droughts lasting 2 or more years significantly impacted agricultural production and livestock in the north-east of the country, where a single drought in 1961 resulted in the loss of 80% of camels and 50 % of sheep. During the drought of 1998–2001, 329,000 people of which 47,000 were nomadic families were forced to eliminate livestock numbers and experienced an acute shortage of food (De Châtel, 2014). Dust emission is influenced by anthropogenic <u>Anthropogenic</u> land use and changes in land cover <u>impact the soil erodibility</u> and <u>dust emission</u>. which impact wind driven soil erosion. The magnitude of such impacts is highly uncertain as both climatic and anthropogenic processes occur simultaneously (Webb & Pierre, 2018). Unsustainable agricultural practices, overgrazing, and deforestation may significantly increase the area of the dust sources. It should be noted however that only

5 around 5% of the land in North Africa and the Middle East is suitable for agriculture; the rest consists of pastures, forests, shrubs, urban zones, badlands, rocky areas, and deserts (Sivakumar & Stefanski, 2007).

4.4.1. Atmospheric circulation patterns

10

The general increase in dust concentrations in the Elbrus ice core are accompanied by a quasi-decadal variability. To identify the possible factors which influence the dryness in the dust sources and dust transport to the Caucasus Mountains, we investigated the Ca^{2+} record in relation with various circulation indices.

- The relationship of precipitation in the Middle East region to the different large-scale circulation patterns are summarized in a recent review of the droughts in the Middle East and Southwest Asia (Barlow et al., 2016 and references therein). The precipitation and occurrence of droughts in the Middle East region may be influenced by several major climatic features such as the NAO, southwest Asian and Indian monsoon, global-scale variability associated with ENSO and state of the western
- 15 Pacific which together determine the strength of the general atmospheric circulation (Barlow et al., 2016). Correlations were calculated for periods of 30 years, using 5 years sliding windows. All series were detrended prior to analysis. The following indices were analyzed: PDO, SOI, AMO, NAO, TPI (IPO) Niño 3, Niño 3.4, Niño 4, Niño 1-2 (https://www.csrl.noaa.gov/psd/geos_wgsp/Timeseries/).

For the 33 year time period (1979-2012), significant correlations exist between Ca²⁺ and PDO, SOI and Niño 4 in the

- preceding winter period (Table 3). A significant correlation (r=0.47 p<0.01) for the Niño 4 index also exists for Dec-June in 1948-2012. The relationship between dust concentrations and circulation indices of the Pacific Ocean is illustrated by the spatial correlation of Ca²⁺ with sea surface temperatures (Supplementary Fig. 1). 23 out of 47 years of high dust deposition recorded in Elbrus ice core coincide with negative Niño 4 phases in 1900-2012. After the 1950s 19 out of 23 years of increased dust concentration correspond to the negative Niño 4 and negative PDO (Fig. 112). No statistically significant correlation was found between the dust in the Elbrus ice core and the NAO index. Ca²⁺ also correlates with the SOI index of the preceding autumn and winter (Nov-Jan) over the last 50 years. It is still unclear whether circulation in the Pacific played a significant role in the precipitation patterns in these dust source regions before the mid-20th century or not. The uncertainties in reanalysis data in this period are large and therefore do not allow us to draw any conclusions about the stability of the revealed correlations in time.
- 30 The relationship of precipitation in the Middle East region to the different large scale circulation patterns are summarized in a recent review of the droughts in the Middle East and Southwest Asia (Barlow et al., 2016 and references). The precipitation and occurrence of droughts in the Middle East region may be influenced by several major climatic features such

as the NAO, southwest Asian and Indian monsoon, global-scale variability associated with ENSO and state of the western Pacific which together determine the strength of the general atmospheric circulation (Barlow et al., 2016).

A significant negative correlation was found between atmospheric dust concentrations in Syria and the PDO in springtime during 2003–2015. It was shown that a positive geopotential height anomaly is formed over the Arabian Peninsula and North

- 5 Africa during the positive phases of PDO (Pu & Ginoux, 2016). A positive PDO is characterised by an increase of cyclonic activity over the northern Pacific and northern Atlantic which occurs together with the intensification of subtropical antyciclones. Despite an increase in geopotential height, the positive PDO increases the probability of moisture transport to North Africa and Middle Eastern regions in the spring (Dai, 2013; Pu & Ginoux, 2016). This moisture is due to increased advection from the Mediterranean Sea which causes deep convection due to unstable stratification in the lower troposphere.
- 10 The negative PDO phases on the other hand are associated with a negative geopotential height anomaly in the middle troposphere over the Mediterranean and the Middle East and cyclonic activity in the region. Low pressure anomalies over Europe, the Southern Arabian Peninsula and north-eastern to eastern Africa create favourable conditions for westerly winds from North Africa and increase the probability of dust transport to the Caucasus.
- Studies of interannual decadal variability of dust activity in the Arabian peninsula and Fertile Crescent suggest that the occurrence of severe droughts and increased dust emission were influenced by the La Niña phase amplified by the negative PDO (Notaro et al., 2015). There are still large uncertainties in such connections due the lack of long-term observations (Pu & Ginoux, 2016). The correlation between Pacific circulation indices and the amount of dust in the Elbrus ice core supports these previous findings and may indicate an increase in the frequency of extreme El Niño and La Niña events with climate warming (Cai et al., 2015) which can then influence circulation over the Pacific Ocean and extend to the Middle East.
- Our results are also in line with previous conclusions about the possible influence of large circulation patterns on the aridity of the tropics. Dai₇ (2011) suggests that the El Niño-Southern Oscillation, tropical Atlantic <u>sea surface temperature</u> (<u>SST)SSTs</u>, and Asian monsoons played a significant role in the increase in global aridity since the 1970s over Africa, southern Europe, East and South Asia, and eastern Australia while recent warming has increased atmospheric moisture demand contributing to the drying-. It is expected that due to anthropogenic climate change the tropical belt may expand
- 25 toward the poles, shift precipitation patterns, and lead to increasing the territory affected by droughts (Seidel et al., 2008). Model studies show that under global warming the large-scale circulation systems (jet streams and storm tracks) may shift poleward (Mbengue & Schneider, 2017). The size and intensity of the Hadley cell and the associated shift of the subtropical anticyclone zone are likely to occur over the 21st century, which should primarily affect the precipitation regime in subtropical latitudes (Lu et al., 2007). An expansion of the global tropics since 1979 by 1° to 3° latitude was identified in
 - 30 both hemispheres although the mechanisms behind this tropical expansion are still unclear (Lucas et al., 2014).

4.5. Comparison to other paleo records

The first record of dust deposition history in the Caucasus was obtained by F.F. Davitaya in 1962 (Davitaya, 1969). This work was based on sampling of the firn layers from a crevasse located at the Kazbek plateau (4600 m asl). Based on dust

layer counting Davitaya estimated that sampled firn layers covered the period between 1793 and 1962. Dust concentrations clearly increased by a factor of 3 was observed since the late 1920's. This increase was attributed to various reasons including local dust influence due to glacier retreat, industry development, fires, World War II and volcanic activity. Despite our-different methodology and location, we qualitatively determined a similar long-term dust trend in the Elbrus ice core

5 record. The average Ca²⁺ concentration over the same periods increases by a factor of 2.5 from 88 ppb in 1793-1925 to 224 ppb in 1925-1962.

An overall increasing trend in dust content and Ca^{2+} exists in Colle Gnifetti (Alps) ice cores after 1870 (Thevenon et al., 2009). Unlike in the Elbrus Ca^{2+} records, no significant long-term trend over the past 300 years exists in the Alps (Bohleber et al., 2018). In Central Asia, negative trends in dust concentrations were found in ice cores in Tien Shan (Grigholm et al.,

10 2017) and in Tibetan Plateau (Grigholm et al., 2015). After the 1950s, calcium concentrations and multiannual and decadal variability significantly decrease in both Central Asian regions due to changes in dust storm frequency in arid areas related to regional trends in reduced zonal wind strengths (Grigholm et al., 2015).

A review of the dust paleo records demonstrates that 16 of the 25 compiled sedimentary archives from across the globe depict a doubling in dust emissions over the past ~250 years which was attributed to the impact of anthropogenic activity in

- 15 source regions (Hooper and Max, 2018). Previous estimates for the atmospheric dust variability also suggests that desert dust doubled during the 20th century with some exceptions (Mahowald et al., 2010). All of the existing regional and global estimates of atmospheric dust variations in the past have large uncertainties due to limited and unevenly distributed paleodata records, which is especially true for the important dust sources of the North Africa, East Asia and Middle East/Central Asia (Mahowald et al., 2010). Tree ring reconstructions of the June-August drought variability over the past 900
- 20 years in the Mediterranean revealed that 1998-2012 was the driest period in the Levant since the 12th century (Cook et al., 2016). Similarly, the latter half of the 20th century was found to be one of the driest in the last nine centuries in north-western Africa (Touchan et al., 2011).

Understanding variations in dust concentrations over the Caucasus leads to a better understanding of the climate change consequences and atmospheric circulation patterns in the dust source regions. Arid and semi-arid regions in North Africa and

25 the Middle East are unstable under the recent climatic changes. Temperature and hydrological anomalies during the last millennium have led to large variations in human occupation and agricultural production, with precipitation variability as a key factor in the productivity in the Middle East region (Kaniewski, 2012). The Elbrus ice core record confirms previous findings (Cook et al., 2016; Kelley et al., 2015) that the recent droughts in 1998-2012 were the most severe over at least the past three centuries.

30 5. Conclusions

This paper presents the Ca^{2+} record from the Elbrus ice core and provides a valuable archive of regional atmospheric dust variability in the past. Ca^{2+} concentrations in the Elbrus ice core are a proxy for atmospheric dust

transported over long distances from primary dust sources in North Africa and the Middle East. The Ca²⁺ record shows that recent decades were characterized by the highest dust emission activity since 1774. From the 1970s onward, Saharan dust deposition events significantly impact the dust deposition in the Caucasus<u>where this The increase in frequency and intensity</u> of dust deposition<u>coincides with events is related to general trend towards</u> the dryer conditions<u>and therefore higher</u>

5

15

probability of dust emission in the dust sources. Ca^{2+} also-significantly spatially correlates spatially with precipitation and soil moisture content in the Middle East. The correlation between circulation indices and amount of dust in the Elbrus ice core indicate that the influence of circulation over the Pacific Ocean extended to the Middle East.

Data availability

Calcium data can be made available for scientific purposes upon request to the authors (contact: kutuzov@igras.ru; 10 suzanne.Preunkert@univ-grenoble-alpes.fr or michel.legrand@univ-grenoble-alpes).

Author contributions

S. Kutuzov performed research, analyzed data and wrote the original manuscript. M. Legrand analyzed ice samples and data, and contributed to the original manuscript. S. Preunkert analyzed ice samples and commented original manuscript. P. Ginot and V.Mikhalenko performed analysis and commented original manuscript. K. Shukurov calculated backward trajectories and commented original manuscript. A. Poliukhov and P. Toropov analyzed climatic data and atmospheric circulation

patterns and commented original manuscript.

Acknowledgments

Research was supported by RSF grant 17-17-01270. The Les Enveloppes Fluides et l'Environnement- Chimie Atmosphérique (CNRS) program entitled "Evolution séculaire de la charge et composition de l'aérosol organique au dessus

20 de l'Europe" provided funding for analysis in France, with the support of Agence de l'Environnement et de la Maîtrise de l'Energie. Backward trajectory analysis was supported by Presidium Program of the Russian Academy of Sciences No. 20. Interpretation of the shallow core was supported by the President Grants for Government Support of Young Russian Scientists and the Leading Scientific Schools of the Russian Federation grant № MK-2508.2017.5.

25 References

De Angelis, M. and Gaudichet, A.: Saharan dust deposition over Mont Blane (French Alps) during the last 30 years, Tellus B, 43(1), 61–75, doi:10.1034/j.1600-0889.1991.00005.x, 1991.

De Angelis, M., Steffensen, J. P., Legrand, M., Clausen, H. and Hammer, C.: Primary acrosol (sea salt and soil dust) deposited in Greenland ice during the last climatic cycle: Comparison with east Antarctic records, J. Geophys. Res. Ocean., 102(C12), 26681–26698, doi:10.1029/97JC01298, 1997.

Achakulwisut, P., Mickley, L. J. and Anenberg, S. C.: Drought-sensitivity of fine dust in the US Southwest: Implications for

5 air quality and public health under future climate change, Environ. Res. Lett., 13(5), 054025, doi:10.1088/1748-9326/aabf20, 2018.

Barlow, M., Zaitchik, B., Paz, S., Black, E., Evans, J. and Hoell, A.: A review of drought in the Middle East and southwest Asia, J. Clim., 29(23), 8547–8574, doi:10.1175/JCLI-D-13-00692.1, 2016.

Bohleber, P., Erhardt, T., Spaulding, N., Hoffmann, H., Fischer, H. and Mayewski, P.: Temperature and mineral dust
variability recorded in two low-accumulation Alpine ice cores over the last millennium, Clim. Past, 14(1), 21–37, doi:10.5194/cp-14-21-2018, 2018.

Breisinger, C., Zhu, T., Al Riffai, P., Nelson, G. C., Robertson, R., Funes, J. and Verner, D.: Global and Local Economic Impacts of Climate Change in Syria and Options for Adaptation, Food Policy, (June), 57 [online] Available from: http://www.ifpri.org/publication/global-and-local-economic-impacts-climate-change-syria-and-options-adaptation, 2011.

15 Cai, W., Santoso, A., Wang, G., Yeh, S. W., An, S. Il, Cobb, K. M., Collins, M., Guilyardi, E., Jin, F. F., Kug, J. S., Lengaigne, M., Mcphaden, M. J., Takahashi, K., Timmermann, A., Vecchi, G., Watanabe, M. and Wu, L.: ENSO and greenhouse warming, Nat. Clim. Chang., 5(9), 849–859, doi:10.1038/nclimate2743, 2015.

De Châtel, F.: The Role of Drought and Climate Change in the Syrian Uprising: Untangling the Triggers of the Revolution, Middle East. Stud., 50(4), 521–535, doi:10.1080/00263206.2013.850076, 2014.

- Chin, M., Diehl, T., Tan, Q., Prospero, J. M., Kahn, R. A., Remer, L. A., Yu, H., Sayer, A. M., Bian, H., Geogdzhayev, I. V., Holben, B. N., Howell, S. G., Huebert, B. J., Hsu, N. C., Kim, D., Kucsera, T. L., Levy, R. C., Mishchenko, M. I., Pan, X., Quinn, P. K., Schuster, G. L., Streets, D. G., Strode, S. A. and Torres, O.: Multi-decadal aerosol variations from 1980 to 2009: A perspective from observations and a global model, Atmos. Chem. Phys., 14(7), 3657–3690, doi:10.5194/acp-14-3657-2014, 2014.
- 25 <u>Chudnovsky, A. A., Koutrakis, P., Kostinski, A., Proctor, S. P. and Garshick, E.: Spatial and temporal variability in desert dust and anthropogenic pollution in Iraq, 1997–2010, J. Air Waste Manag. Assoc., 67(1), 17–26, doi:10.1080/10962247.2016.1153528, 2017.</u>

Cook, B. I., Anchukaitis, K. J., Touchan, R., Meko, D. M. and Cook, E. R.: Spatiotemporal drought variability in the mediterranean over the last 900 years, J. Geophys. Res., 121(5), 2060–2074, doi:10.1002/2015JD023929, 2016.

30 Dai, A.: Drought under global warming: A review, Wiley Interdiscip. Rev. Clim. Chang., 2(1), 45–65, doi:10.1002/wcc.81, 2011.

Davitaya, F. F.: Atmospheric dust content as a factor affecting glaciation and climatic change, Ann. Assoc. Am. Geogr., 59(3), 552–560, doi:10.1111/j.1467-8306.1969.tb00690.x, 1969.

35 De Angelis, M. and Gaudichet, A.: Saharan dust deposition over Mont Blanc (French Alps) during the last 30 years, Tellus
 35 B, 43(1), 61–75, doi:10.1034/j.1600-0889.1991.00005.x, 1991.

De Angelis, M., Steffensen, J. P., Legrand, M., Clausen, H. and Hammer, C.: Primary aerosol (sea salt and soil dust) deposited in Greenland ice during the last climatic cycle: Comparison with east Antarctic records, J. Geophys. Res. Ocean., 102(C12), 26681–26698, doi:10.1029/97JC01298, 1997.

De Châtel, F.: The Role of Drought and Climate Change in the Syrian Uprising: Untangling the Triggers of the Revolution,
 Middle East. Stud., 50(4), 521–535, doi:10.1080/00263206.2013.850076, 2014.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M.,

Mcnally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N. and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, Q. J. R. Meteorol. Soc., 137(656), 553–597, doi:10.1002/qj.828, 2011.

Delmonte, B., Petit, J.-R. and Maggi, V.: Glacial to Holocene implications of the new 27000-year dust record from the EPICA Dome C (East Antarctica) ice core, Clim. Dyn., 18(8), 647–660, doi:10.1007/s00382-001-0193-9, 2002.

Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT_4 modeling system of trajectories, dispersion, and deposition., Aust. Meteor. Mag., 47, 295–308, 1998.

Gabbi, J., Huss, M., Bauder, A., Cao, F. and Schwikowski, M.: The impact of Saharan dust and black carbon on albedo and long-term mass balance of an Alpine glacier, Cryosphere, 9(4), 1385–1400, doi:10.5194/tc-9-1385-2015, 2015.

 Ganor, E., Osetinsky, I., Stupp, A. and Alpert, P.: Increasing trend of African dust, over 49 years, in the eastern Mediterranean, J. Geophys. Res. Atmos., 115(7), 1–7, doi:10.1029/2009JD012500, 2010.

5

25

30

35

Gautam, R., Liu, Z., Singh, R. P. and Hsu, N. C.: Two contrasting dust-dominant periods over India observed from MODIS and CALIPSO data, Geophys. Res. Lett., 36(6), L06813, doi:10.1029/2008GL036967, 2009.

 Ginot, P., Dumont, M., Lim, S., Patris, N., Taupin, J. D., Wagnon, P., Gilbert, A., Arnaud, Y., Marinoni, A., Bonasoni, P.
 and Laj, P.: A 10 year record of black carbon and dust from a Mera Peak ice core (Nepal): Variability and potential impact on melting of Himalayan glaciers, Cryosphere, 8(4), 1479–1496, doi:10.5194/tc-8-1479-2014, 2014.

Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C. and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, Rev. Geophys., 50(3), 1–36, doi:10.1029/2012RG000388, 2012.

20 Grigholm, B., Mayewski, P. A., Kang, S., Zhang, Y., Morgenstern, U., Schwikowski, M., Kaspari, S., Aizen, V., Aizen, E., Takeuchi, N., Maasch, K. A., Birkel, S., Handley, M. and Sneed, S.: Twentieth century dust lows and the weakening of the westerly winds over the Tibetan Plateau, Geophys. Res. Lett., 42(7), 2434–2441, doi:10.1002/2015GL063217, 2015.

Grigholm, B., Mayewski, P. A., Aizen, V., Kreutz, K., Aizen, E., Kang, S., Maasch, K. A. and Sneed, S. B.: A twentieth century major soluble ion record of dust and anthropogenic pollutants from Inilchek Glacier, Tien Shan, J. Geophys. Res., 122(3), 1884–1900, doi:10.1002/2016JD025407, 2017.

Hooper, J. and Marx, S.: A global doubling of dust emissions during the Anthropocene?, Glob. Planet. Change, 169(June), 70–91, doi:10.1016/j.gloplacha.2018.07.003, 2018.

Kalderon-Asael, B., Erel, Y., Sandler, A. and Dayan, U.: Mineralogical and chemical characterization of suspended atmospheric particles over the east Mediterranean based on synoptic-scale circulation patterns, Atmos. Environ., 43(25), 3963–3970, doi:10.1016/j.atmosenv.2009.03.057, 2009.

Kaniewski, D.: Drought is a recurring challenge in the Middle East, Proc., doi:10.1073/pnas.1116304109/-/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1116304109, 2012.

Kaspari, S., Mayewski, P. a., Handley, M., Kang, S., Hou, S., Sneed, S., Maasch, K. and Qin, D.: A High-Resolution Record of Atmospheric Dust Composition and Variability since a.d. 1650 from a Mount Everest Ice Core, J. Clim., 22(14), 3910–3925, doi:10.1175/2009JCLI2518.1, 2009.

Kelley, C. P., Mohtadi, S., Cane, M. A., Seager, R. and Kushnir, Y.: Climate change in the Fertile Crescent and implications of the recent Syrian drought, Proc. Natl. Acad. Sci., 112(11), 3241–3246, doi:10.1073/pnas.1421533112, 2015.

Kistler, R., Collins, W., Saha, S., White, G., Woollen, J., Kalnay, E., Chelliah, M., Ebisuzaki, W., Kanamitsu, M., Kousky, V., van den Dool, H., Jenne, R. and Fiorino, M.: The NCEP–NCAR 50–Year Reanalysis: Monthly Means CD–ROM and

40 Documentation, Bull. Am. Meteorol. Soc., 82(2), 247–267, doi:10.1175/1520-0477(2001)082<0247:TNNYRM>2.3.CO;2, 2001.

Knippertz, P. and Stuut, J. B. W.: Mineral dust: A key player in the earth system, Miner. Dust A Key Play. Earth Syst., 1–509, doi:10.1007/978-94-017-8978-3, 2014.

Koçak, M., Theodosi, C., Zarmpas, P., Séguret, M. J. M., Herut, B., Kallos, G., Mihalopoulos, N., Kubilay, N. and Nimmo, M.: Influence of mineral dust transport on the chemical composition and physical properties of the Eastern Mediterranean aerosol, Atmos. Environ., 57, 266–277, doi:10.1016/j.atmosenv.2012.04.006, 2012.

5

30

Kutuzov, S., Shahgedanova, M., Mikhalenko, V., Lavrentiev, I. and Kemp, S.: Desert dust deposition on Mt. Elbrus, Caucasus Mountains, Russia in 2009–2012 as recorded in snow and shallow ice core: high resolution "provenancing", transport patterns, physical properties and soluble ionic composition, Cryosph. Discuss., 7(2), 1621–1672, doi:10.5194/tcd-7-1621-2013, 2013a.

10 Kutuzov, S., Shahgedanova, M., Mikhalenko, V., Ginot, P., Lavrentiev, I. and Kemp, S.: High-resolution provenance of desert dust deposited on Mt. Elbrus, Caucasus in 2009-2012 using snow pit and firn core records, Cryosph., 7(5), 1481– 1498, doi:10.5194/tc-7-1481-2013, 2013b.

Kutuzov, S. S., Lavrentiev, I. I., Vasilenko, E. V., Macheret, Y. Y., Petrakov, D. A. and Popov, G. V.: Estimation of the greater caucasus glaciers volume, using radio-echo sounding data and modelling, Earth's Cryosph., 19(1), 78–88, 2015a.

15 Kutuzov, S. S., Mikhalenko, V. N., Shahgedanova, M. V., Ginot, P., Kozachek, A. V., Kuderina, T. M., Lavrentiev, I. I. and Popov, G. V.: Ways of far distance dust transport onto Caucasian glaciers and chemical composition of snow on the Western plateau of Elbrus, Ice Snow, 127(3), 5, doi:10.15356/2076-6734-2014-3-5-15, 2015b.

Lee, D. S. and Pacyna, J. M.: An industrial emissions inventory of calcium for Europe, Atmos. Environ., 33(11), 1687–1697, doi:10.1016/S1352-2310(98)00286-6, 1999.

20 Legrand, M.: Chemistry of Antarctic snow and ice, Le J. Phys. Colloq., 48(C1), C1-77-C1-86, doi:10.1051/jphyscol:1987111, 1987.

Legrand, M.: Seasonally resolved Alpine and Greenland ice core records of anthropogenic HCl emissions over the 20th century, J. Geophys. Res., 107(D12), 4139, doi:10.1029/2001JD001165, 2002.

Legrand, M. and Mayewski, P.: Glaciochemistry of polar ice cores: A review, Rev. Geophys., 35(3), 219–243, doi:10.1029/96RG03527, 1997.

Legrand, M., Preunkert, S., May, B., Guilhermet, J., Hoffman, H. and Wagenbach, D.: Major 20th century changes of the content and chemical speciation of organic carbon archived in Alpine ice cores: Implications for the long-term change of organic aerosol over Europe, J. Geophys. Res. Atmos., 118(9), 3879–3890, doi:10.1002/jgrd.50202, 2013.

Li, L. and Sokolik, I. N.: Analysis of dust aerosol retrievals using satellite data in Central Asia, Atmosphere (Basel)., 9(8), doi:10.3390/atmos9080288, 2018.

Lim, S., Faïn, X., Ginot, P., Mikhalenko, V., Kutuzov, S., Paris, J. D., Kozachek, A. and Laj, P.: Black carbon variability since preindustrial times in the eastern part of Europe reconstructed from Mt. Elbrus, Caucasus, ice cores, Atmos. Chem. Phys., 17(5), doi:10.5194/acp-17-3489-2017, 2017.

Lu, J., Vecchi, G. A. and Reichler, T.: Expansion of the Hadley cell under global warming, Geophys. Res. Lett., 34(6), L06805, doi:10.1029/2006GL028443, 2007.

Lucas, C., Timbal, B. and Nguyen, H.: The expanding tropics: a critical assessment of the observational and modeling studies, Wiley Interdiscip. Rev. Clim. Chang., 5(1), 89–112, doi:10.1002/wcc.251, 2014.

Mahowald, N. M., Ballantine, J. a., Feddema, J. and Ramankutty, N.: Global trends in visibility: implications for dust sources, Atmos. Chem. Phys. Discuss., 7(1), 3013–3071, doi:10.5194/acpd-7-3013-2007, 2007.

40 Mahowald, N. M., Kloster, S., Engelstaedter, S., Moore, J. K., Mukhopadhyay, S., McConnell, J. R., Albani, S., Doney, S. C., Bhattacharya, A., Curran, M. A. J., Flanner, M. G., Hoffman, F. M., Lawrence, D. M., Lindsay, K., Mayewski, P. A.,

Neff, J., Rothenberg, D., Thomas, E., Thornton, P. E. and Zender, C. S.: Observed 20th century desert dust variability: Impact on climateand biogeochemistry, Atmos. Chem. Phys., 10(22), 10875–10893, doi:10.5194/acp-10-10875-2010, 2010.

Di Mauro, B., Baccolo, G., Garzonio, R., Giardino, C. and Massabò, D.: Impact of impurities and cryoconite on the optical properties of the Morteratsch Glacier (Swiss Alps), 2393–2409, 2017.

5 Mbengue, C. and Schneider, T.: Storm-Track Shifts under Climate Change: Toward a Mechanistic Understanding Using Baroclinic Mean Available Potential Energy, J. Atmos. Sci., 74(1), 93–110, doi:10.1175/JAS-D-15-0267.1, 2017.

Middleton, N. J.: Desert dust hazards: A global review, Aeolian Res., 24, 53-63, doi:10.1016/j.aeolia.2016.12.001, 2017.

Mikhalenko, V., Sokratov, S., Kutuzov, S., Ginot, P., Legrand, M., Preunkert, S., Lavrentiev, I., Kozachek, A., Ekaykin, A., Fain, X., Lim, S., Schotterer, U., Lipenkov, V. and Toropov, P.: Investigation of a deep ice core from the Elbrus western plateau, the Caucasus, Russia, Cryosphere, 9(6), doi:10.5194/tc-9-2253-2015, 2015.

Moridnejad, A., Karimi, N. and Ariya, P. A.: A new inventory for Middle East dust source points, Environ. Monit. Assess., 187(9), 582, doi:10.1007/s10661-015-4806-x, 2015.

Nicholson, S. E., Funk, C. and Fink, A. H.: Rainfall over the African continent from the 19th through the 21st century, Glob. Planet. Change, 165(May 2017), 114–127, doi:10.1016/j.gloplacha.2017.12.014, 2018.

15 Notaro, M., Yu, Y. and Kalashnikova, O. V: Journal of Geophysical Research : Atmospheres, 229–249, doi:10.1002/2015JD023855.Received, 2015.

10

Osterberg, E., Mayewski, P., Kreutz, K., Fisher, D., Handley, M., Sneed, S., Zdanowicz, C., Zheng, J., Demuth, M., Waskiewicz, M. and Bourgeois, J.: Ice core record of rising lead pollution in the North Pacific atmosphere, Geophys. Res. Lett., 35(5), L05810, doi:10.1029/2007GL032680, 2008.

- 20 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E. and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, Nature, 399, 429–436, doi:10.1038/20859, 1999.
- Preunkert, S. and Legrand, M.: Towards a quasi-complete reconstruction of past atmospheric aerosol load and composition
 (organic and inorganic) over Europe since 1920 inferred from Alpine ice cores, Clim. Past, 9(4), 1403–1416, doi:10.5194/cp-9-1403-2013, 2013.

Preunkert, S., Wagenbach, D., Legrand, M. and Vincent, C.: Col du Dôme (Mt Blanc Massif, French Alps) suitability for ice-core studies in relation with past atmospheric chemistry over Europe, Tellus B Chem. Phys. Meteorol., 52(3), 993–1012, doi:10.3402/tellusb.v52i3.17081, 2000.

30 Pu, B. and Ginoux, P.: The impact of the Pacific Decadal Oscillation on springtime dust activity in Syria, Atmos. Chem. Phys., 16(21), 13431–13448, doi:10.5194/acp-16-13431-2016, 2016.

Ruth, U., Wagenbach, D., Steffensen, J. P. and Bigler, M.: Continuous record of microparticle concentration and size distribution in the central Greenland NGRIP ice core during the last glacial period, J. Geophys. Res., 108(D3), 4098, doi:10.1029/2002JD002376, 2003.

35 Seidel, D. J., Fu, Q., Randel, W. J. and Reiohler, T. J.: Widening of the tropical belt in a changing climate, Nat. Methods, 1(1), 21–24, doi:10.1038/ngeo.2007.38, 2008.

Shahgedanova, M., Kutuzov, S., White, K. H. and Nosenko, G.: Using the significant dust deposition event on the glaciers of Mt.~Elbrus, Caucasus Mountains, Russia on 5 May 2009 to develop a method for dating and "provenancing" of desert dust events recorded in snow pack, Atmos. Chem. Phys., 13(4), 1797–1808, doi:10.5194/acp-13-1797-2013, 2013.

40 Sivakumar, M. V. K. and Stefanski, R.: Climate and Land Degradation --- an Overview, in Climate and Land Degradation, edited by M. V. K. Sivakumar and N. Ndiang'ui, pp. 105–135, Springer Berlin Heidelberg, Berlin, Heidelberg., 2007. Skiles, S. M., Flanner, M., Cook, J. M., Dumont, M. and Painter, T. H.: Radiative forcing by light-absorbing particles in snow, Nat. Clim. Chang., 8(11), 964–971, doi:10.1038/s41558-018-0296-5, 2018.

Sodemann, H., Palmer, A. S., Schwierz, C., Schwikowski, M. and Wernli, H.: The transport history of two Saharan dust events archived in an Alpine ice core, Atmos. Chem. Phys., 6(3), 667–688, doi:10.5194/acp-6-667-2006, 2006.

5 Solomos, S., Kalivitis, N., Mihalopoulos, N., Amiridis, V., Kouvarakis, G., Gkikas, A., Binietoglou, I., Tsekeri, A., Kazadzis, S., Kottas, M., Pradhan, Y., Proestakis, E., Nastos, P. and Marenco, F.: From Tropospheric Folding to Khamsin and Foehn Winds: How Atmospheric Dynamics Advanced a Record-Breaking Dust Episode in Crete, Atmosphere (Basel)., 9(7), 240, doi:10.3390/atmos9070240, 2018.

Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: Noaa's Hysplit atmospheric transport and dispersion modeling system, Bull. Am. Meteorol. Soc., 96(12), 2059–2077, doi:10.1175/BAMS-D-14-00110.1, 2015.

Thevenon, F., Anselmetti, F. S., Bernasconi, S. M. and Schwikowski, M.: Mineral dust and elemental black carbon records

15 from an Alpine ice core (Colle Gnifetti glacier) over the last millennium, J. Geophys. Res., 114(D17), D17102, doi:10.1029/2008JD011490, 2009.

Thomas, N. and Nigam, S.: Twentieth-century climate change over Africa: Seasonal hydroclimate trends and sahara desert expansion, J. Clim., 31(9), 3349–3370, doi:10.1175/JCLI-D-17-0187.1, 2018.

Thompson, L. G.: A High-Resolution Millennial Record of the South Asian Monsoon from Himalayan Ice Cores, Science (80-.)., 289(5486), 1916–1919, doi:10.1126/science.289.5486.1916, 2000.

Tielidze, L. G. and Wheate, R. D.: The Greater Caucasus Glacier Inventory (Russia, Georgia and Azerbaijan), Cryosph., 12(1), 81–94, doi:10.5194/tc-12-81-2018, 2018.

Touchan, R., Anchukaitis, K. J., Meko, D. M., Sabir, M., Attalah, S. and Aloui, A.: Spatiotemporal drought variability in northwestern Africa over the last nine centuries, Clim. Dyn., 37(1–2), 237–252, doi:10.1007/s00382-010-0804-4, 2011.

25 Trigo, R. M., Gouveia, C. M. and Barriopedro, D.: The intense 2007-2009 drought in the Fertile Crescent: Impacts and associated atmospheric circulation, Agric. For. Meteorol., 150(9), 1245–1257, doi:10.1016/j.agrformet.2010.05.006, 2010.

Usher, C. R., Michel, A. E. and Grassian, V. H.: Reactions on Mineral Dust, Chem. Rev., 103(12), 4883–4940, doi:10.1021/cr020657y, 2003.

Vicente-Serrano, S. M., Beguería, S. and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming:
 The Standardized Precipitation Evapotranspiration Index, J. Clim., 23(7), 1696–1718, doi:10.1175/2009JCLI2909.1, 2010.

Wagenbach, D., Preunkert, S., Schäfer, J., Jung, W. and Tomadin, L.: Northward Transport of Saharan Dust Recorded in a Deep Alpine Ice Core, in The Impact of Desert Dust Across the Mediterranean, edited by S. Guerzoni and R. Chester, pp. 291–300, Springer Netherlands, Dordrecht., 1996.

Webb, N. P. and Pierre, C.: Quantifying Anthropogenic Dust Emissions, Earth's Futur., 6(2), 286–295, doi:10.1002/2017EF000766, 2018.

Zhang, Y., Mahowald, N., Scanza, R. A., Journet, E., Desboeufs, K., Albani, S., Kok, J. F., Zhuang, G., Chen, Y., Cohen, D. D., Paytan, A., Patey, M. D., Achterberg, E. P., Engelbrecht, J. P. and Fomba, K. W.: Modeling the global emission, transport and deposition of trace elements associated with mineral dust, Biogeosciences, 12(19), 5771–5792, doi:10.5194/bg-12-5771-2015, 2015.

Stanelle, T., Bey, I., Raddatz, T., Reick, C. and Tegen, I.: Journal of Geophysical Research : Atmospheres, 526–546, doi:10.1002/2014JD022062.Received, 2014.

Table 1. Ca^{2+} concentrations in Elbrus ice cores in different periods including total concentrations and excluding background samples from the large dust deposition events.

	Ca ²⁺ concentration (<i>ppb</i>)				
	summer (total)	summer (background)	winter (total)	winter (background)	
1774-1800	65	64	42	42	
1800-1850	78	69	52	46	
1850-1900	100	69	43	36	
1900-1950	156	83	37	33	
1950-2000	344	181	75	58	
2000-2012	439	212	99	64	
1774-2012	172	103	54	44	

5

Table 2. Composition of aerosol collected at Erdemli during dust events from Middle East and at Heraklion during dust events from Sahara (from Koçak et al., 2012). The magnesium sea-salt contributions were calculated via the Na+ content, after having subtracted its dust contribution calculated as 0.08 times the Ca2+ content (see text).

<u>Species</u>	Erdemli	Heraklion
	(October 2007)	<u>(April 2008)</u>
<u>[Na⁺]</u>	<u>1148 ng m⁻³</u>	<u>1106 ng m⁻³</u>
	<u>(sea-salt: 581 ng m⁻³)</u>	<u>(sea-salt: 445 ng m⁻³)</u>
[Mg ²⁺]	423 ng m^{-3}	407 ng m^{-3}
	<u>(sea-salt: 70 ng m⁻³)</u>	$(\text{sea-salt: 53 ng m}^{-3})$
$\underline{[Ca^{2+}]}$	7085 ng m^{-3}	<u>8264 ng m⁻³</u>
$[Mg^{2+}]/[Ca^{2+}]$ in dust	0.050	<u>0.043</u>

10

Table 23. Correlation coefficient of the Elbrus Ca^{2+} data and large-scale atmospheric circulation indices.

	1948-2012	1979-2012
PDO (Dec-June)	-0.35	-0.42

SOI (Oct-Jan)	0.40	0.49
Niño 4 (Dec-June)	-0.47	-0.57



Figure 1: Location of Elbrus (red star) and dust sources (orange polygons) based on (Ginoux et al., 2012). Annual NOAA HYSPLIT_4 10-day backward trajectory density plots for the period 1948-2013 using NCEP/NCAR Reanalysis. Trajectories were run every 6 hrs.-Colours are the relative contributions of the source areas determined from 10-day backward trajectories in 0.5x0.5 degree grids.



Figure <u>2</u>: (a): Mean summer half-year thickness along the Elbrus deep ice core. (b): Numbers of samples (N) spanning summer half-years. See details in Preunkert et al., this issue.



5

Figure 3: NOAA HYSPLIT_4 10-day backward trajectories density plots for the period 1948-2013 for December – February (a), March-May (b) June-August (c) and September – November (d) using NCEP/NCAR Reanalysis. Trajectories were run every 6 hrs. Only backward trajectories within the boundary layer were considered. Contribution of 10-day backward trajectories in 0.5x0.5 degree grids.



Figure 4: (a): Total (red) and background (after the removal of samples influenced by large dust deposition events, black) Ca²⁺ concentrations (ppb) in Elbrus ice core samples. The full record (a) and the upper section of the ice core (b) are shown.



Figure 5: (a): Averaged concentrations of calcium and ammonium (used for determining seasonality as described in the text), over 1900-1950 (left) and 1950-2010 (right). Total calcium concentrations are the grey bars in the top panels while calcium concentrations calculated after the removal of samples are in red. The middle panels refer to calcium concentrations corresponding to the dust fraction (i.e., $[Ca^{2+}] - [Ca^{2+red}]$, orange bars). Note that the equidistant binning of the summer and winter layers applied to derive these 12 values per year does not necessary correspond to monthly means (only true if there is an absence of seasonality in precipitation). Nevertheless the figure clearly illustrates the occurrence of a dust related calcium peak at the beginning of the summer season.

10



Figure 6: Total and background (grey lines) Ca^{2+} concentrations in Elbrus ice core in summer and winter. 5-year moving averages are shown as black lines.





31



Figure 78: [Mg²⁺_{red.}] versus [Ca²⁺_{red.}] in individual Elbrus ice summer samples (not impacted by dust events) over the recent decades (left) and prior to the 1950 increase in calcium.



Figure 89: Individual summer means of background levels of Ca^{2+} , Mg^{2+} , and of the $[Mg^{2+}_{red.}]/[Ca^{2+}_{red.}]$ mass ratio along the Elbrus ice cores.



Figure 910: Spatial correlation of Elbrus Ca²⁺ concentrations with soil moisture (a), and precipitation (b) ERA-Interim from 1979 to 2013. North African and Middle Eastern domains are shown by orange boxes.



Figure $\frac{1011}{12}$: Ca²⁺ and Standardised Precipitation-Evapotranspiration Index (SPEI3) averaged over the regions of (a), (c) North Africa (20-35°N; 0-35°E; and (b), (d) the Middle East (32-37°N; 38-45°E). Outlines of the regions are presented on Figure $\frac{910}{10}$. Detrended records are shown at the bottom graphs.



Figure 1112: PDO (a) and Niño 4 (b) indices (December-June average), and normalized Ca2+ record (c). Grey shading indicate years when high dust concentrations coincided with negative Niño 4 phases.