

1 *Supplement of*

2 **Sources and formation of carbonaceous aerosols in Xi'an,**
3 **China: primary emissions and secondary formation**
4 **constrained by radiocarbon**

5 Haiyan Ni^{1,2,3}, Ru-Jin Huang^{1*}, Junji Cao¹, Jie Guo¹, Haoyue Deng², Ulrike Dusek²

6 ¹State Key Laboratory of Loess and Quaternary Geology, Key Laboratory of Aerosol Chemistry and Physics,
7 Center for Excellence in Quaternary Science and Global Change, Institute of Earth Environment, Chinese
8 Academy of Sciences, Xi'an, 710061, China

9 ²Centre for Isotope Research (CIO), Energy and Sustainability Research Institute Groningen (ESRIG),
10 University of Groningen, Groningen, 9747 AG, the Netherlands

11 ³University of Chinese Academy of Sciences, Beijing, 100049, China

12 *Correspondence to:* rujin.huang@ieecas.cn

13 **S1. Estimation of the probability density functions (PDFs) of p values**

14 The p values used in Eq. (11) in the main text is the fraction of EC from coal combustion (EC_{coal})
15 in EC from fossil sources (EC_{fossil}). That is,

16
$$p = \frac{EC_{coal}}{EC_{fossil}} = \frac{EC_{coal}}{EC_{coal} + EC_{liq.fossil}} \quad (S1)$$


17 where EC_{fossil} is the sum of EC_{coal} and EC from liquid fossil fuel combustion (i.e., vehicle emissions;
18 $EC_{liq.fossil}$).

19 Eq. (S1) can be formulated as:


20
$$p = \frac{f_{coal}}{f_{fossil}} = \frac{f_{coal}}{f_{coal} + f_{liq.fossil}} \quad (S2)$$

21 where f_{coal} and $f_{liq.fossil}$ is the relative contribution of coal combustion emission and liquid fossil fuel
22 combustion to EC. The sum of f_{coal} and $f_{liq.fossil}$ is f_{fossil} of EC, which is well constrained by $F^{14}C$ of
23 EC.

24 The PDFs of f_{coal} and $f_{liq.fossil}$ (eg., Fig. S4), derived from the Bayesian calculations detailed in Sect.
25 2.6 in the main text, are used to calculated the PDFs of p .




26
27 **Figure S1.** Selected samples for ^{14}C analysis. Three composite samples that represent high (H),
28 medium (M) and low (L) TC concentrations are combined from several individual filter samples
29 per season. Each composite sample is consisting of 2 to 4 24-hr filter pieces with similar TC
30 loadings and air mass backward trajectories (Table S1).




31

32 **Figure S2.** Fraction modern ($F^{14}C$) of elemental carbon (EC), organic carbon (OC), water-insoluble
 33 OC (WIOC) and water-soluble OC (WSOC) ($F^{14}C_{(EC)}$, $F^{14}C_{(OC)}$, $F^{14}C_{(WIOC)}$ and $F^{14}C_{(WSOC)}$
 34 respectively). $F^{14}C_{(WSOC)}$ is calculated from the measured $F^{14}C_{(OC)}$ and $F^{14}C_{(WIOC)}$ following the
 35 isotope mass balance. The blue dashed area for best estimate of $F^{14}C_{(WSOC)}$ (blue filled circle)
 36 indicates ranges of $F^{14}C_{(WSOC)}$ (Sect. 2.5).



37

38 **Figure S3.** **(a)** An example probability density functions (PDFs) of concentrations of POC_{fossil} (red), SOC_{fossil} (light blue) for sample Autumn-L. **(b)**
39 PDFs of concentrations of and OC_{o.nf} (light blue) and POC_{bb} (red) for the same sample. Their concentrations are estimated by ^{14}C -apportioned OC
40 and EC using the EC tracer method (Sect. 2.5). The mean and median are indicated by the dashed and solid vertical lines.



45 **Table S1.** Sample information as well as the fraction modern ($F^{14}C$) of elemental carbon (EC),
 46 organic carbon (OC), water-insoluble OC (WIOC) and water-soluble OC (WSOC) ($F^{14}C_{(EC)}$,
 47 $F^{14}C_{(OC)}$, $F^{14}C_{(WIOC)}$ and $F^{14}C_{(WSOC)}$ respectively), and stable carbon isotopic compositions ($\delta^{13}C$, ‰)
 48 of EC ($\delta^{13}C_{EC}$).

Sample name	Sampling Date	$F^{14}C_{(EC)}^a$	$F^{14}C_{(OC)}^a$	$F^{14}C_{(WIOC)}^a$	$F^{14}C_{(WSOC)}^b$	$\delta^{13}C_{EC}$
Winter-H	2015.12.20	0.340 ± 0.005	0.640 ± 0.009	0.565 ± 0.006	0.704 (0.682–0.717)	-24.6
	2015.12.21					
Winter-M	2015.11.30	0.258 ± 0.005	0.609 ± 0.007	0.558 ± 0.007	0.649 (0.635–0.657)	-25
	2015.12.8					
	2015.12.9					
Winter-L	2015.12.14	0.320 ± 0.005	0.626 ± 0.007	0.553 ± 0.006	0.69 (0.675–0.699)	-24.7
	2015.12.16					
	2015.12.17					
Spring-H	2016.5.5	0.123 ± 0.004	0.534 ± 0.006	0.514 ± 0.006	0.543 (0.541–0.543)	-24.7
	2016.5.10					
Spring-M	2016.4.19	0.145 ± 0.006	0.531 ± 0.007	0.450 ± 0.006	0.577 (0.567–0.583)	-24.8
	2016.4.20					
Spring-L	2016.4.23	0.184 ± 0.004	0.557 ± 0.007	0.445 ± 0.006	0.637 (0.610–0.654)	-24.2
	2016.4.24					
	2016.4.27					
Summer-H	2016.7.21	0.159 ± 0.004	0.549 ± 0.006	0.438 ± 0.006	0.605 (0.587–0.616)	-24.7
	2016.7.23					
Summer-M	2016.7.11	0.191 ± 0.004	0.593 ± 0.007	0.497 ± 0.006	0.651 (0.631–0.663)	-25.2
	2016.7.16					
	2016.7.27					
Summer-L	2016.7.5	0.181 ± 0.006	0.637 ± 0.007	0.394 ± 0.006	0.795 (0.750–0.822)	-25
	2016.7.6					
	2016.7.12					
	2016.7.13					
Autumn-H	11/3/2016	0.169 ± 0.004	0.562 ± 0.007	0.516 ± 0.007	0.599 (0.591–0.603)	-25.2
	11/4/2016					
	11/13/2016					
Autumn-M	10/17/2016	0.154 ± 0.004	0.547 ± 0.007	0.492 ± 0.006	0.587 (0.575–0.595)	-25.5
	10/18/2016					
	11/1/2016					
Autumn-L	10/15/2016	0.194 ± 0.004	0.593 ± 0.006	0.518 ± 0.006	0.635 (0.623–0.643)	-25.1
	10/16/2016					
	10/20/2016					

49 ^a $F^{14}C$ values are given in average \pm measurement uncertainty.

50 ^b $F^{14}C_{(WSOC)}$ is calculated from the measured $F^{14}C_{(OC)}$ and $F^{14}C_{(WIOC)}$ following the isotope mass balance (Eq.
 51 4 in the main text). The range of $F^{14}C_{(WSOC)}$ is presented in the parentheses, calculated following the method
 52 detailed in Sect 2.5.

53 **Table S2.** Consensus value of F¹⁴C secondary standards IAEA- C7 and -C8 along with measured
54 F¹⁴C values. Data corrections for the measured F14C of secondary standards are the same as those
55 for samples.

Standards	Consensus value of F ¹⁴ C	measured F ¹⁴ C	measured mass (μgC)
IAEA-C7	0.4953 ± 0.0012	0.4884 ± 0.0059	76
		0.5017 ± 0.0064	80
IAEA-C8	0.1503 ± 0.0017	0.1511 ± 0.0039	63
		0.1540 ± 0.0038	100

56

57 **Table S3.** Relative contributions of non-fossil sources to EC, OC, WIOC and WSOC ($f_{\text{bb}}(\text{EC})$, $f_{\text{nf}}(\text{OC})$, $f_{\text{nf}}(\text{WIOC})$, $f_{\text{nf}}(\text{WSOC})$), and relative fossil
 58 sources contribution to EC, OC, WIOC and WSOC ($f_{\text{fossil}}(\text{EC})$, $f_{\text{fossil}}(\text{OC})$, $f_{\text{fossil}}(\text{WIOC})$, $f_{\text{fossil}}(\text{WSOC})$) for each sample.

Sample name	$f_{\text{bb}}(\text{EC})$	$f_{\text{fossil}}(\text{EC})$	$f_{\text{nf}}(\text{OC})$	$f_{\text{fossil}}(\text{OC})$	$f_{\text{nf}}(\text{WIOC})$	$f_{\text{fossil}}(\text{WIOC})$	$f_{\text{nf}}(\text{WSOC})$	$f_{\text{fossil}}(\text{WSOC})$
Winter-H	0.310 ± 0.008	0.690 ± 0.008	0.587 ± 0.014	0.413 ± 0.014	0.516 ± 0.012	0.484 ± 0.012	0.639 ± 0.014	0.361 ± 0.014
Winter-M	0.235 ± 0.006	0.765 ± 0.006	0.559 ± 0.012	0.441 ± 0.012	0.509 ± 0.012	0.491 ± 0.012	0.590 ± 0.012	0.410 ± 0.012
Winter-L	0.291 ± 0.007	0.709 ± 0.007	0.574 ± 0.012	0.426 ± 0.012	0.504 ± 0.011	0.496 ± 0.011	0.627 ± 0.013	0.373 ± 0.013
Spring-H	0.112 ± 0.004	0.888 ± 0.004	0.490 ± 0.011	0.510 ± 0.011	0.468 ± 0.011	0.532 ± 0.011	0.495 ± 0.010	0.505 ± 0.010
Spring-M	0.132 ± 0.006	0.868 ± 0.006	0.487 ± 0.011	0.513 ± 0.011	0.410 ± 0.010	0.590 ± 0.010	0.525 ± 0.011	0.475 ± 0.011
Spring-L	0.167 ± 0.005	0.833 ± 0.005	0.511 ± 0.011	0.489 ± 0.011	0.406 ± 0.010	0.594 ± 0.010	0.578 ± 0.014	0.422 ± 0.014
Summer-H	0.144 ± 0.005	0.856 ± 0.005	0.504 ± 0.011	0.496 ± 0.011	0.399 ± 0.009	0.601 ± 0.009	0.550 ± 0.012	0.450 ± 0.012
Summer-M	0.173 ± 0.005	0.827 ± 0.005	0.544 ± 0.012	0.456 ± 0.012	0.454 ± 0.010	0.546 ± 0.010	0.591 ± 0.013	0.409 ± 0.013
Summer-L	0.165 ± 0.006	0.835 ± 0.006	0.585 ± 0.012	0.415 ± 0.012	0.359 ± 0.009	0.641 ± 0.009	0.720 ± 0.019	0.280 ± 0.019
Autumn-H	0.153 ± 0.005	0.847 ± 0.005	0.516 ± 0.011	0.484 ± 0.011	0.470 ± 0.011	0.530 ± 0.011	0.545 ± 0.011	0.455 ± 0.011
Autumn-M	0.140 ± 0.004	0.860 ± 0.004	0.502 ± 0.011	0.498 ± 0.011	0.448 ± 0.010	0.552 ± 0.010	0.534 ± 0.011	0.466 ± 0.011
Autumn-L	0.177 ± 0.005	0.823 ± 0.005	0.544 ± 0.012	0.456 ± 0.012	0.472 ± 0.011	0.528 ± 0.011	0.578 ± 0.012	0.422 ± 0.012

59

60
61**Table S4.** Concentrations of EC, OC, WIOC and WSOC from non-fossil sources (EC_{bb} , OC_{nf} , $WIOC_{nf}$ and $WSOC_{nf}$) and fossil sources (EC_{fossil} , OC_{fossil} , $WIOC_{fossil}$ and $WSOC_{fossil}$) in units of $\mu\text{g m}^{-3}$ for each sample.

Sample name	EC_{bb}	EC_{fossil}	OC_{nf}	OC_{fossil}	$WIOC_{nf}$	$WIOC_{fossil}$	$WSOC_{nf}$	$WSOC_{fossil}$
Winter-H	3.08 ± 0.18	6.86 ± 0.39	27.66 ± 1.56	19.43 ± 1.20	10.78 ± 0.78	10.12 ± 0.74	16.72 ± 1.82	9.43 ± 1.08
Winter-M	1.44 ± 0.09	4.70 ± 0.28	21.17 ± 1.17	16.73 ± 0.97	8.25 ± 0.62	7.95 ± 0.59	12.80 ± 1.36	8.89 ± 0.96
Winter-L	0.82 ± 0.06	1.99 ± 0.14	8.31 ± 0.48	6.16 ± 0.37	3.33 ± 0.17	3.27 ± 0.17	4.95 ± 0.53	2.94 ± 0.32
Spring-H	0.36 ± 0.03	2.86 ± 0.19	5.62 ± 0.33	5.85 ± 0.34	1.56 ± 0.08	1.77 ± 0.09	4.03 ± 0.33	4.12 ± 0.34
Spring-M	0.30 ± 0.03	2.00 ± 0.15	3.68 ± 0.22	3.87 ± 0.23	1.08 ± 0.06	1.56 ± 0.08	2.58 ± 0.24	2.34 ± 0.22
Spring-L	0.22 ± 0.02	1.09 ± 0.10	2.48 ± 0.16	2.37 ± 0.15	0.79 ± 0.06	1.15 ± 0.09	1.68 ± 0.19	1.23 ± 0.14
Summer-H	0.32 ± 0.03	1.88 ± 0.14	3.71 ± 0.23	3.65 ± 0.22	0.94 ± 0.08	1.41 ± 0.11	2.75 ± 0.26	2.25 ± 0.21
Summer-M	0.17 ± 0.02	0.83 ± 0.08	2.25 ± 0.15	1.89 ± 0.13	0.68 ± 0.06	0.82 ± 0.07	1.55 ± 0.17	1.07 ± 0.12
Summer-L	0.12 ± 0.02	0.60 ± 0.07	1.96 ± 0.14	1.39 ± 0.10	0.46 ± 0.03	0.82 ± 0.05	1.49 ± 0.17	0.58 ± 0.08
Autumn-H	1.05 ± 0.07	5.79 ± 0.33	12.05 ± 0.68	11.32 ± 0.64	4.77 ± 0.22	5.37 ± 0.24	7.22 ± 0.72	6.03 ± 0.61
Autumn-M	0.54 ± 0.04	3.29 ± 0.21	5.88 ± 0.35	5.83 ± 0.35	2.13 ± 0.15	2.62 ± 0.18	3.71 ± 0.38	3.24 ± 0.34
Autumn-L	0.28 ± 0.02	1.29 ± 0.11	3.29 ± 0.21	2.76 ± 0.18	0.99 ± 0.07	1.11 ± 0.08	2.29 ± 0.23	1.67 ± 0.17

62

63 **Table S5.** Concentrations ($\mu\text{g m}^{-3}$) of primary OC from biomass burning (POC_{bb}), OC from non-
 64 fossil sources excluding primary biomass burning ($\text{OC}_{\text{o,nf}}$), primary OC from fossil sources
 65 ($\text{POC}_{\text{fossil}}$), secondary OC from fossil sources ($\text{SOC}_{\text{fossil}}$) (median and interquartile range). The
 66 median values for POC_{bb} and $\text{OC}_{\text{o,nf}}$ are very close to their mean values due to their symmetric
 67 PDFs (Fig. S3b).

Sample Name	POC_{bb}	$\text{OC}_{\text{o,nf}}$	$\text{POC}_{\text{fossil}}$	$\text{SOC}_{\text{fossil}}$
Winter-H	12.27 (11.26–13.37)	15.34 (13.87–16.78)	9.24 (7.52–11.64)	10.10 (7.64–11.97)
Winter-M	5.77 (5.26–6.27)	15.37 (14.45–16.29)	5.99 (4.95–7.70)	10.55 (8.92–11.84)
Winter-L	3.26 (2.98–3.55)	5.03 (4.61–5.46)	2.69 (2.19–3.39)	3.42 (2.73–3.99)
Spring-H	1.44 (1.31–1.58)	4.17 (3.92–4.42)	3.87 (3.05–5.05)	1.97 (0.81–2.77)
Spring-M	1.22 (1.11–1.33)	2.46 (2.27–2.64)	2.58 (2.10–3.34)	1.28 (0.52–1.77)
Spring-L	0.87 (0.79–0.96)	1.60 (1.46–1.74)	1.58 (1.25–1.98)	0.77 (0.38–1.12)
Summer-H	1.26 (1.15–1.38)	2.45 (2.26–2.64)	2.49 (2.00–3.22)	1.15 (0.42–1.66)
Summer-M	0.69 (0.62–0.77)	1.55 (1.43–1.67)	1.00 (0.84–1.25)	0.87 (0.60–1.06)
Summer-L	0.47 (0.42–0.53)	1.48 (1.38–1.59)	0.76 (0.62–0.98)	0.62 (0.40–0.78)
Autumn-H	4.20 (3.84–4.56)	7.88 (7.30–8.45)	7.07 (5.93–9.06)	4.21 (2.21–5.43)
Autumn-M	2.14 (1.96–2.34)	3.73 (3.43–4.03)	3.75 (3.23–4.78)	2.02 (0.99–2.61)
Autumn-L	1.11 (1.00–1.22)	2.18 (2.01–2.35)	1.61 (1.34–2.05)	1.13 (0.68–1.43)

68

69 **Table S6.** Relative non-fossil sources contribution to EC, OC, WIOC and WSOC ($f_{\text{bb}}(\text{EC})$, $f_{\text{nf}}(\text{OC})$, $f_{\text{nf}}(\text{WIOC})$, $f_{\text{nf}}(\text{WSOC})$), and relative fossil
 70 sources contribution to EC, OC, WIOC and WSOC ($f_{\text{fossil}}(\text{EC})$, $f_{\text{fossil}}(\text{OC})$, $f_{\text{fossil}}(\text{WIOC})$, $f_{\text{fossil}}(\text{WSOC})$) in different seasons and throughout the year.

Season	$f_{\text{bb}}(\text{EC})$	$f_{\text{fossil}}(\text{EC})$	$f_{\text{nf}}(\text{OC})$	$f_{\text{fossil}}(\text{OC})$	$f_{\text{nf}}(\text{WIOC})$	$f_{\text{fossil}}(\text{WIOC})$	$f_{\text{nf}}(\text{WSOC})$	$f_{\text{fossil}}(\text{WSOC})$
Winter	0.279 ± 0.039	0.721 ± 0.039	0.573 ± 0.014	0.427 ± 0.014	0.510 ± 0.006	0.490 ± 0.006	0.619 ± 0.026	0.381 ± 0.026
Spring	0.137 ± 0.028	0.863 ± 0.028	0.496 ± 0.013	0.504 ± 0.013	0.428 ± 0.035	0.572 ± 0.035	0.533 ± 0.042	0.467 ± 0.042
Summer	0.161 ± 0.015	0.839 ± 0.015	0.544 ± 0.040	0.456 ± 0.040	0.404 ± 0.047	0.596 ± 0.047	0.620 ± 0.089	0.380 ± 0.089
Autumn	0.157 ± 0.019	0.843 ± 0.019	0.521 ± 0.021	0.479 ± 0.021	0.464 ± 0.013	0.536 ± 0.013	0.552 ± 0.023	0.448 ± 0.023
Annual	0.183 ± 0.062	0.817 ± 0.062	0.534 ± 0.037	0.466 ± 0.037	0.451 ± 0.049	0.549 ± 0.049	0.581 ± 0.060	0.419 ± 0.060

71

72 **Table S7.** Concentrations of EC, OC, WIOC and WSOC from non-fossil sources (EC_{bb} , OC_{nf} , $WIOC_{nf}$ and $WSOC_{nf}$) and fossil sources (EC_{fossil} ,
 73 OC_{fossil} , $WIOC_{fossil}$ and $WSOC_{fossil}$) in units of $\mu\text{g m}^{-3}$ in different seasons and throughout the year.

Season	EC_{bb}	EC_{fossil}	OC_{nf}	OC_{fossil}	$WIOC_{nf}$	$WIOC_{fossil}$	$WSOC_{nf}$	$WSOC_{fossil}$
Winter	1.78 ± 1.17	4.52 ± 2.44	19.05 ± 9.85	14.11 ± 7.01	7.45 ± 3.79	7.11 ± 3.50	11.49 ± 5.99	7.09 ± 3.60
Spring	0.29 ± 0.07	1.98 ± 0.89	3.93 ± 1.58	4.03 ± 1.75	1.14 ± 0.39	1.49 ± 0.31	2.76 ± 1.18	2.56 ± 1.46
Summer	0.20 ± 0.10	1.10 ± 0.68	2.64 ± 0.94	2.31 ± 1.19	0.69 ± 0.24	1.02 ± 0.34	1.93 ± 0.71	1.30 ± 0.86
Autumn	0.62 ± 0.39	3.46 ± 2.25	7.07 ± 4.50	6.64 ± 4.34	2.63 ± 1.94	3.03 ± 2.16	4.41 ± 2.54	3.65 ± 2.21
Annual	0.72 ± 0.84	2.76 ± 2.03	8.17 ± 8.23	6.77 ± 5.94	2.98 ± 3.34	3.16 ± 3.06	5.15 ± 4.85	3.65 ± 2.97

74

75 **Table S8.** Fractional contribution of different incomplete combustion sources to EC in different
76 seasons (median, interquartile range (25th-75th percentile)).

Sources		Winter	Spring	Summer	Autumn
Biomass burning	median	0.28	0.146	0.163	0.159
	25th-75th percentile	(0.26–0.31)	(0.13–0.17)	(0.15–0.18)	(0.15–0.18)
Coal combustion	median	0.246	0.296	0.227	0.19
	25th-75th percentile	(0.13–0.41)	(0.15–0.50)	(0.11–0.41)	(0.09–0.36)
Liquid fossil fuel combustion	median	0.459	0.534	0.598	0.638
	25th-75th percentile	(0.29–0.59)	(0.33–0.69)	(0.41–0.72)	(0.45–0.74)

77

78 **Table S9.** EC concentrations (in unit of $\mu\text{g m}^{-3}$) from biomass burning (EC_{bb}), coal combustion
 79 (EC_{coal}) and liquid fossil fuel combustion ($\text{EC}_{\text{liq,fossil}}$) for each sample (median and interquartile
 80 range in unit of $\mu\text{g m}^{-3}$), and the seasonal averaged concentrations ($\mu\text{g m}^{-3}$) calculated by averaging
 81 the median values for each sample in each season^a.

	EC_{bb}		EC_{coal}	$\text{EC}_{\text{liq,fossil}}$
	median	(interquartile range)	median	(interquartile range)
Winter-H	3.07	(2.94–3.22)	2.79	(1.43–4.51)
Winter-M	1.44	(1.38–1.52)	1.42	(0.67–2.60)
Winter-L	0.82	(0.77–0.86)	0.69	(0.36–1.18)
Spring-H	0.36	(0.34–0.38)	1.02	(0.44–1.90)
Spring-M	0.30	(0.29–0.32)	0.70	(0.31–1.30)
Spring-L	0.22	(0.21–0.23)	0.50	(0.24–0.79)
Summer-H	0.32	(0.30–0.34)	0.66	(0.30–1.20)
Summer-M	0.17	(0.16–0.19)	0.20	(0.10–0.39)
Summer-L	0.12	(0.11–0.13)	0.16	(0.08–0.32)
Autumn-H	1.05	(1.00–1.10)	1.46	(0.68–2.99)
Autumn-M	0.54	(0.51–0.56)	0.68	(0.33–1.33)
Autumn-L	0.28	(0.26–0.29)	0.37	(0.18–0.68)
Winter ^a	1.78 ± 1.16		1.63 ± 1.06	
Spring ^a	0.30 ± 0.07		0.74 ± 0.26	
Summer ^a	0.20 ± 0.10		0.34 ± 0.28	
Autumn ^a	0.62 ± 0.39		0.84 ± 0.57	

82 ^aThe seasonal averaged concentrations calculated by averaging the median values for each sample
 83 in each season.