



# Gaseous, PM<sub>2.5</sub> Mass, and Speciated Emission Factors from Laboratory Chamber Peat Combustion

John G. Watson<sup>1,2\*</sup>, Junji Cao<sup>2,3</sup>, L.-W. Antony Chen<sup>4</sup>, Qiyuan Wang<sup>2</sup>, Jie Tian<sup>2,3</sup>, Xiaoliang Wang<sup>1</sup>, Steven Gronstal<sup>1</sup>, Steven Sai Hang Ho<sup>5</sup>, Adam C. Watts<sup>1</sup>, Judith C. Chow<sup>1,2</sup>

<sup>1</sup>Division of Atmospheric Sciences, Desert Research Institute, Reno, Nevada, USA

<sup>2</sup>Key Laboratory of Aerosol Chemistry and Physics, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, China.

<sup>3</sup>CAS Center for Excellence in Quaternary Science and Global Change, Xi'an, China

<sup>4</sup>Department of Environmental and Occupational Health, University of Nevada, Las Vegas, Nevada, USA

<sup>5</sup>Hong Kong Premium Services and Research Laboratory, Hong Kong, China

\*Corresponding Author: [john.watson@dri.edu](mailto:john.watson@dri.edu)



## 23 Abstract

24 Peat fuels representing four biomes of boreal (western Russia and Siberia), temperate  
25 (northern Alaska, U.S.A.), subtropical (northern and southern Florida, U.S.A), and tropical  
26 (Borneo, Malaysia) regions were burned in a laboratory chamber to determine gas and particle  
27 emission factors (EFs). Tests with 25 % fuel moisture were conducted with predominant  
28 smoldering combustion conditions (average modified combustion efficiency [MCE] =  $0.82 \pm$   
29  $0.08$ ). Average fuel-based  $\text{EF}_{\text{CO}_2}$  (carbon dioxide) are highest ( $1400 \pm 38 \text{ g kg}^{-1}$ ) and lowest ( $1073$   
30  $\pm 63 \text{ g kg}^{-1}$ ) for the Alaskan and Russian peats, respectively.  $\text{EF}_{\text{CO}}$  (carbon monoxide) and  $\text{EF}_{\text{CH}_4}$   
31 (methane) are  $\sim 12\text{--}15\%$  and  $\sim 0.3\text{--}0.9\%$  of  $\text{EF}_{\text{CO}_2}$ , in the range of  $157\text{--}171 \text{ g kg}^{-1}$  and  $3\text{--}10 \text{ g kg}^{-1}$ ,  
32 respectively. EFs for nitrogen species are at the same magnitude of  $\text{EF}_{\text{CH}_4}$ , with an average of  
33  $5.6 \pm 4.8$  and  $4.7 \pm 3.1 \text{ g kg}^{-1}$  for  $\text{EF}_{\text{NH}_3}$  (ammonia) and  $\text{EF}_{\text{HCN}}$  (hydrogen cyanide);  $1.9 \pm 1.1 \text{ g kg}^{-1}$   
34 for  $\text{EF}_{\text{NO}_x}$  (nitrogen oxides); as well as  $2.4 \pm 1.4$  and  $2.0 \pm 0.7 \text{ g kg}^{-1}$  for  $\text{EF}_{\text{NO}_y}$  (reactive nitrogen)  
35 and  $\text{EF}_{\text{N}_2\text{O}}$  (nitrous oxide).

36 An oxidation flow reactor (OFR) was used to simulate atmospheric aging times of  $\sim 2$  and  
37  $\sim 7$  days to compare fresh (upstream) and aged (downstream) emissions. Filter-based  $\text{EF}_{\text{PM}_{2.5}}$   
38 varied by  $>4$ -fold ( $14\text{--}61 \text{ g kg}^{-1}$ ) without appreciable changes between fresh and aged emissions.  
39 The majority of  $\text{EF}_{\text{PM}_{2.5}}$  consists of  $\text{EF}_{\text{OC}}$  (organic carbon), with  $\text{EF}_{\text{OC}}/\text{EF}_{\text{PM}_{2.5}}$  ratios in the range  
40 of  $52\text{--}98\%$  for fresh emissions, and  $\sim 15\%$  degradation after aging. Reductions of  $\text{EF}_{\text{OC}}$  ( $\sim 7\text{--}9 \text{ g}$   
41  $\text{kg}^{-1}$ ) after aging are most apparent for boreal peats with the largest degradation in organic carbon  
42 that evolves at  $<140^\circ\text{C}$ , indicating the loss of high vapor pressure semi-volatile organic  
43 compounds upon aging. The highest  $\text{EF}_{\text{Levoglucosan}}$  is found for Russian peat ( $\sim 16 \text{ g kg}^{-1}$ ), with  $\sim 35\text{--}$   
44  $50\%$  degradation after aging. EFs for water-soluble OC ( $\text{EF}_{\text{WSOC}}$ ) accounts for  $\sim 20\text{--}62\%$  of fresh  
45  $\text{EF}_{\text{OC}}$ .

46 The majority ( $>95\%$ ) of the total emitted carbon is in the gas phase with  $54\text{--}75\%$   $\text{CO}_2$ ,  
47 followed by  $8\text{--}30\%$  CO. Nitrogen in the measured species explains  $24\text{--}52\%$  of the consumed  
48 fuel nitrogen with an average of  $35 \pm 11\%$ , consistent with past studies that report  $\sim$ one- to two-  
49 thirds of the fuel nitrogen measured in biomass smoke. The majority ( $>99\%$ ) of the total emitted  
50 nitrogen is in the gas phase, with an average of  $16.7\%$  fuel N emitted as  $\text{NH}_3$  and  $9.5\%$  of fuel N  
51 emitted as HCN.  $\text{N}_2\text{O}$  and  $\text{NO}_y$  constituted  $5.7\%$  and  $2.9\%$  of consumed fuel N. EFs from this  
52 study can be used to refine current emissions inventories.



53 Keywords: Peat combustion, modified combustion efficiency, emission factors, oxidation flow  
54 reactor, carbon balance, nitrogen budget.



## 55 1 Introduction

56 Globally, peatlands occupy ~3 % of the Earth's land surface, but they store as much as 610  
57 gigatonnes (i.e.,  $610 \times 10^{15}$  grams) of carbon, representing 20–30 % of the planet's terrestrial carbon  
58 (Page et al., 2011; Rein et al., 2009). Peatland fires can persist for weeks to months and are  
59 dominated by the smoldering-phase as opposed to the flaming-phase of biomass burning (Hu et  
60 al., 2018; Stockwell et al., 2016). This results in lower combustion efficiencies, increased  
61 particulate matter (PM) emissions, and larger fractions of brown carbon (BrC) compared to black  
62 carbon (BC) or soot (Pokhrel et al., 2016). Peat fires emit reduced nitrogen compounds (e.g.,  
63 ammonia [ $\text{NH}_3$ ] and hydrogen cyanide [ $\text{HCN}$ ]), volatile and semi-volatile organic compounds  
64 (VOCs and SVOCs), and  $\text{PM}_{2.5}$  (PM with aerodynamic diameters  $< 2.5 \mu\text{m}$ ) (Akagi et al.,  
65 2011; Yokelson et al., 2013). Peat smoke and ash affect ecosystem productivity, soil acidity,  
66 biogeochemical cycling, atmospheric chemistry, Earth's radiation balance, and human health.  
67 Warmer climates lower the water table in peatlands and change the pattern, frequency, and  
68 intensity of the peatland fires causing local- and regional-scale air pollution and visibility  
69 impairment (Page et al., 2002; Turetsky et al., 2010; 2015a; 2015b).

70 Nitrogen, one of the most important plant nutrients, affects global carbon and  
71 biogeochemical cycles (Crutzen and Andreae, 1990; Gruber and Galloway, 2008). Deposition of  
72 oxidized and reduced nitrogen species from biomass burning, such as gaseous nitric oxide (NO),  
73 nitrogen dioxide ( $\text{NO}_2$ ), and  $\text{NH}_3$  as well as particulate nitrate ( $\text{NO}_3^-$ ) and ammonium ( $\text{NH}_4^+$ ), alters  
74 terrestrial ecosystems (Chen et al., 2010), while nitric acid ( $\text{HNO}_3$ ) contributes to soil acidification  
75 and excessive nitrification that reduce plant resistance to environmental stresses (Goulding et al.,  
76 1998). Gaseous nitrogen oxides ( $\text{NO}_x$ ) affect atmospheric chemistry through: 1) reactions with  
77 hydroxyl (OH) and peroxy ( $\text{HO}_2 + \text{RO}_2$ ) radicals; 2) conversion to nitrate radical ( $\text{NO}_3$ ), dinitrogen  
78 pentoxide ( $\text{N}_2\text{O}_5$ ), and acyl peroxy nitrates (particularly peroxyacetyl nitrate [PAN]), which are  
79 important  $\text{NO}_x$  reservoirs; and 3) formation of ozone ( $\text{O}_3$ ) and secondary organic aerosols (SOA)  
80 (Alvarado et al., 2010; Cubison et al., 2011; Ng et al., 2007). While  $\text{NH}_3$  neutralizes  $\text{HNO}_3$  to form  
81 particulate ammonium nitrate ( $\text{NH}_4\text{NO}_3$ ), it may also react with alkanolic acids to form alkyl  
82 amides, nitriles, and ammonium salts that can also contribute to SOA formation (Na et al.,  
83 2007; Simoneit et al., 2003; Zhao et al., 2013). In addition,  $\text{NH}_3$  interacts with SOA to form “BrC”  
84 that further influence the aerosol radiative forcing (Updyke et al., 2012).



85 This study quantifies peat burning emission factors (EFs) for fresh and aged multipollutant  
86 mixtures through controlled burns in a laboratory combustion chamber with atmospheric aging  
87 simulated by an oxidation flow reactor (OFR). These tests are applied to peat samples from diverse  
88 parts of the world.

## 89 **2 Experiment**

### 90 **2.1 Fuel types**

91 Peatlands are found all over the world, as illustrated in Fig. 1 (Yu et al., 2010) with large  
92 deposits found in the northern USA and Canada, northern Europe, Russia/Siberia, and southeast  
93 Asia. Eight types of peat fuels from different regions and climates were collected for testing,  
94 including: boreal (i.e., Odintsovo, Russia and Pskov, Siberia); temperate (i.e., black spruce forest,  
95 northern Alaska, USA); subtropical (i.e., northern [Putnam County Lakebed] and southern  
96 [Everglades National Park] Florida, USA and Caohai and Gaopo, Guizhou, southwest China); and  
97 tropical (i.e., Borneo, Malaysia) peats.

98 Representative peat samples of 250–1150 g from the upper 20 cm of the peatland surface  
99 were excavated for each region indicated in Fig. 1. As peat is a heterogeneous mixture of  
100 decomposed plant material, it can be formed in different wetlands under changing climates and  
101 nutrient contents (Turetsky et al., 2015a). Supplemental Fig. S1 shows that the appearance of peat  
102 fuels varies by region.

103 Approximately 2–3 g of each peat fuel were dried in a vacuum oven ( $\sim 105^{\circ}\text{C}$ ) for two  
104 hours prior to elemental analysis of carbon (C), hydrogen (H), nitrogen (N), sulfur (S), and oxygen  
105 (O) (Thermo Flash-EA 1112 CHNS/O Analyzer, Waltham, MA, USA).

### 106 **2.2 Experimental setup with preliminary testing**

107 The laboratory setup shown in Fig. 2 used a biomass combustion chamber with a volume  
108 of  $\sim 8\text{ m}^3$  ( $1.8\text{ m}[\text{W}] \times 1.8\text{ m}[\text{L}] \times 2.2\text{ m}[\text{H}]$ ) (Tian et al., 2015). Instrument specifications and  
109 operating principles are shown in Table S1. The chamber is made of 3 mm-thick aluminum to  
110 withstand high temperature heating. Smoke from the chamber was drawn through a dilution  
111 sampling manifold where the exhaust was diluted with clean air to achieve cooling that allowed  
112 for condensation of SVOCs. A portion of the exhaust was directed through potential aerosol mass  
113 (PAM)-OFR (Aerodyne Research Inc., Billerica, MA, USA) to simulate atmospheric aging prior  
114 to quantification by the sampling instruments shown in Fig. 2. The 185 and 254 nm (OFR185)  
115 ultraviolet (UV) lamps in the OFR were operated at 2 and 3.5 volts with  $10\text{ L min}^{-1}$  flow rate to



116 simulate ~2- and 7-days of aging assuming an average daily OH concentration of  $1.5 \times 10^6$   
117 molecules  $\text{cm}^{-3}$ . The estimated OH exposures ( $\text{OH}_{\text{exp}}$ ) at 2 and 3.5 volts were  $2.6 \times 10^{11}$  and  $8.8 \times$   
118  $10^{11}$  molecules-sec  $\text{cm}^{-3}$  based on the measured decay of sulfur dioxide ( $\text{SO}_2$ ) (Cao et al., 2019).

119 Sampling ports for stack concentrations of carbon dioxide ( $\text{CO}_2$ ) and multiple gases by  
120 Fourier transform infrared (FTIR) spectroscopy were located ~1 m above the top of the chamber  
121 roof in the exhaust duct. The FTIR spectrometer measured gaseous emissions prior to dilution to  
122 obtain enhanced signal-to-noise ratios for trace gases (Jaakkola et al., 1998).

123 Oxides of nitrogen were measured as  $\text{NO}_x$  (the sum of NO and  $\text{NO}_2$ ) and total reactive  
124 nitrogen ( $\text{NO}_y$ , including NO,  $\text{NO}_2$ ,  $\text{N}_2\text{O}_5$ ,  $\text{HNO}_3$ ,  $\text{HNO}_4$ ,  $\text{ClONO}_2$ , HONO, alkyl nitrates, and PAN)  
125 by chemiluminescence  $\text{NO}_x$  and  $\text{NO}_y$  analyzers (Allen et al., 2018; Ballenthin et al., 2003). The  
126  $\text{NO}_x$  analyzers placed upstream and downstream of the OFR determined  $\text{NO}_x$  changes with  $\text{OH}_{\text{exp}}$   
127 in the OFR. There are known interferences for the non-selective catalytic converter in the  
128 chemiluminescent  $\text{NO}_x$  analyzer and for the spectroscopic absorption in FTIR (Allen et al.,  
129 2018; Prenni et al., 2014; Villena et al., 2012). The chemiluminescence monitor converts most  
130 nitrogenous compounds to NO, with  $\text{HNO}_3$  and PAN being the most important potential  
131 interferents (Winer et al., 1974). However, much of the available  $\text{HNO}_3$  and PAN is removed by  
132 the tubing leading to the molybdenum converter in the standard  $\text{NO}_x$  analyzer, which is why the  
133  $\text{NO}_y$  analyzer locates the converter at the inlet. Allen et al. (2018) found no significant differences  
134 between  $\text{NO}_x$  measurements of biomass burning plumes when comparing a chemiluminescent  
135 analyzer with more specific UV absorption measurements. The following analyses are based on:  
136 1) the commercial  $\text{NO}_x$  analyzers for NO,  $\text{NO}_2$ , and  $\text{NO}_x$  (NO +  $\text{NO}_2$  as equivalent  $\text{NO}_2$ ); 2) the  
137  $\text{NO}_y$  analyzer for total reactive nitrogen; and 3) the FTIR spectrometer for trace gas measurements  
138 of methane ( $\text{CH}_4$ ),  $\text{NH}_3$ , hydrogen cyanide (HCN), nitrous oxide ( $\text{N}_2\text{O}$ ), and 14 low molecular-  
139 weight VOCs (C2–C6).

140  $\text{PM}_{2.5}$  filter packs were sampled upstream and downstream of the OFR to characterize fresh  
141 and aged emissions, respectively, with Minivol  $\text{PM}_{10}/\text{PM}_{2.5}$  samplers (Airmetrics, Springfield, OR,  
142 USA) operated at  $5 \text{ L Min}^{-1}$  flow rate per channel.  $\text{PM}_{2.5}$  mass, elements, carbon, water-soluble  
143 organic carbon (WSOC), ions, carbohydrates, organic acids, as well as gaseous  $\text{NH}_3$  and  $\text{HNO}_3$   
144 were obtained from the paired upstream and downstream filter samples to examine changes in  
145 speciated EFs and source profiles with photochemical aging. Average filter-based EFs are



146 examined by peat types and aging times (i.e., denoted as Fresh 2 vs Aged 2 and Fresh 7 vs. Aged  
147 7) (Cao et al., 2019; Chow et al., 2019).

148 Peat fuels were heated to 90°C and weighed every 24 hours to achieve a stable dry mass  
149 with ~0.16 % moisture by weight content (after ~96 hours of heating). The low heating  
150 temperature (i.e., below the water boiling point) minimized VOC losses, although some  
151 compounds with high volatilities could have been removed at 90°C. To better simulate the field  
152 conditions during peat fires, distilled-deionized water (DDW) was added to rehydrate the dry peat  
153 and achieve a fuel moisture of ~25 % (by weight) before each experiment (Yatavelli et al., 2017).  
154 To examine the effects of fuel moisture on emissions, additional experiments (n=3) were  
155 conducted at 60 % moisture content (by weight) for the Putnam (FL) peat.

156 For each test, ~10–30 g of dried peat was placed in an asbestos insulated circular container  
157 on top of an induction heater that provided heating during the first ~5–10 minutes of combustion  
158 (see Fig. S2). This method replaced a propane torch used in initial test burns to minimize non-  
159 peat burning emissions. The smoldering process is usually self-propagating and sustained by heat  
160 conduction and radiation with fuel mass continuously monitored by a scale underneath the  
161 induction heater (Ohlemiller et al., 1979). Continuous PM<sub>2.5</sub> mass concentrations were monitored  
162 with a DustTrak (Wang et al., 2009) (Table S1). When PM<sub>2.5</sub> concentrations reached their  
163 maximum and started to decline, the induction heater was turned off. The fuel was consumed with  
164 diminished smoke emissions after ~20 minutes. Preliminary tests were conducted using ~10–20  
165 g of fuel and a dilution ratio of ~3 to 5, yielding sufficient particle loadings on the filters (~150–  
166 290 µg per 47 mm filter disc). To achieve higher filter deposits of 300–600 µg per filter that  
167 accommodate comprehensive organic speciation, additional fuels (~15–20 g) were added with the  
168 induction heater turned on for another ~10 minutes. Sampling continued until the concentrations  
169 returned to background level.

### 170 **2.3 PM<sub>2.5</sub> mass and carbon**

171 PM<sub>2.5</sub> mass and major chemical species concentrations were obtained from the parallel  
172 Teflon-membrane and quartz-fiber filters (Teflo<sup>®</sup>, 2 µm pore size, R2PJ047 and Tissuquartz 2500  
173 QAFUP, Pall Life Sciences, Port Washington, NY, USA). Teflon-membrane filters were  
174 equilibrated in a temperature (20–23 °C) and relative humidity (30–40 %) controlled environment  
175 for a minimum of 48 hours prior to gravimetric analysis by a microbalance with ± 1 µg sensitivity  
176 (Watson et al., 2017). This was followed by elemental analysis by x-ray fluorescence (Watson et



al., 1999). Quartz-fiber filters were pre-fired at 900° C for four hours to minimize organic artifacts. An aliquots (0.5 cm<sup>2</sup>) of the quartz-fiber filter were submitted for organic, elemental, and brown carbon (OC, EC, and BrC) analyses following the IMPROVE\_A thermal/optical reflectance (TOR) protocol (Chow et al., 2007;2015). Half of the quartz-fiber filters was extracted in DDW for ionic speciation (i.e., chloride [Cl<sup>-</sup>], nitrate [NO<sub>3</sub><sup>-</sup>], nitrite [NO<sub>2</sub><sup>-</sup>], sulfate [SO<sub>4</sub><sup>=</sup>], water-soluble sodium [Na<sup>+</sup>] and potassium [K<sup>+</sup>], ammonium [NH<sub>4</sub><sup>+</sup>], 17 carbohydrates, and 10 organic acids) by ion chromatography (Chow and Watson, 2017) and for WSOC by combustion and non-dispersive infrared detection. Citric acid and sodium chloride impregnated cellulose-fiber filters placed behind the Teflon-membrane and quartz-fiber filters, respectively, acquired NH<sub>3</sub> as NH<sub>4</sub><sup>+</sup> and nitric acid (HNO<sub>3</sub>) as volatilized nitrate, respectively, with analysis by ion chromatography. Details on chemical analyses can be found in Chow et al. (2019).

The open face sampling manifold allows homogenous particle deposits on 47-mm filters (Watson et al., 2017). To test the `uniformity of particle deposits, five individual punches were removed from the center and each quadrant of the 47-mm quartz-fiber filter disc for carbon analyses. Table S2 shows total carbon (TC = OC + EC) concentration variations of 1.7% to 5% across the filters for the five test burns, within the overall uncertainty of the emission estimates. Standard deviations from the five filter punches for each experiment are low with coefficients of variation of 1.7–5.0 %.

## 2.4 Modified combustion efficiency and fuel-based emission factors

The modified combustion efficiency (MCE) is defined as the ratio of background-subtracted CO<sub>2</sub> to the sum of CO<sub>2</sub> and CO (Ward and Radke, 1993):

$$\text{MCE} = \frac{\Delta\text{CO}_2}{\Delta\text{CO}_2 + \Delta\text{CO}} \quad (1)$$

where ΔCO<sub>2</sub> and ΔCO are CO<sub>2</sub> and CO concentrations above background. MCE provides a real-time indicator of the combustion status (e.g., MCE > ~0.9 for flaming and MCE < ~0.85 for smoldering).

Each burn was completed when concentrations of pollutants measured on-line (i.e., CO, NO<sub>x</sub>, NO<sub>y</sub>, and PM<sub>2.5</sub>) returned to the baseline/background levels. Dilution ratios ranging from 2.7 to 5 were taken into account when calculating EFs. As shown in Fig. 2, air passed through a charcoal bed to remove VOCs and a high-efficiency particulate air (HEPA) filter provided particle-





free clean air for dilution. Fuel-based EFs are calculated based on carbon mass balance, expressed as grams of emission per kilogram of dry fuel ( $\text{g kg}^{-1}$ ) (Wang et al., 2012). For gaseous and particle species  $i$ , the time-integrated EF <sub>$i$</sub>  is:

$$\text{EF}_i = \text{CMF}_{\text{fuel}} \frac{C_i}{C_{\text{CO}_2} \left( \frac{M_c}{M_{\text{CO}_2}} \right) + C_{\text{CO}} \left( \frac{M_c}{M_{\text{CO}}} \right) + C_{\text{CH}_4} \left( \frac{M_c}{M_{\text{CH}_4}} \right) + \sum_j C_{\text{VOC}_j} \left( \frac{n_j \times M_c}{M_{\text{VOC}_j}} \right) + \text{PM}_c} \times 1000 \quad (2)$$

where  $\text{CMF}_{\text{fuel}}$  is the carbon mass fraction of the fuel in kg carbon per kg of fuel;  $C_i$ ,  $C_{\text{CO}_2}$ ,  $C_{\text{CO}}$ ,  $C_{\text{CH}_4}$ , and  $C_{\text{VOC}_j}$  are the background-subtracted concentrations for species  $i$  (e.g., nitrogen or  $\text{PM}_{2.5}$  species),  $\text{CO}_2$ ,  $\text{CO}$ ,  $\text{CH}_4$ , and  $\text{VOC}$  ( $\text{C}_2$ – $\text{C}_6$ ) species  $j$  in  $\text{mg m}^{-3}$  under standard conditions (temperature = 293 °K and pressure = 1 atm), respectively;  $\text{PM}_c$  is the total carbon concentration of  $\text{PM}_{2.5}$  in  $\text{mg m}^{-3}$ ;  $M_c$ ,  $M_{\text{CO}_2}$ ,  $M_{\text{CO}}$ ,  $M_{\text{CH}_4}$ , and  $M_{\text{VOC}_j}$  are the atomic or molecular weights of carbon,  $\text{CO}_2$ ,  $\text{CO}$ ,  $\text{CH}_4$ , and  $\text{VOC}$  species  $j$  in mg per mole, respectively;  $n_j$  is the number of carbon atom in  $\text{VOC}$  species  $j$ ; and the factor 1000 converts kg to g. All concentrations are converted to stack concentration, i.e., species measured after dilution are adjusted with dilution ratio. This equation assumes that the carbon mass in unmeasured VOCs and other emissions not listed above is negligible compared to that in  $\text{CO}$ ,  $\text{CO}_2$ ,  $\text{CH}_4$ , measured VOCs ( $\text{C}_2$ – $\text{C}_6$ ), and  $\text{PM}_{2.5}$  carbon.

### 3 Results and discussion

#### 3.1 Fuel composition

Table 1 shows that peat contains 44–57 % C and 31–39 % O with the exception of the two Guizhou, China peats (20–30 % C and 21–24 % O). The carbon content ( $50.6 \pm 2.5$  % C) in Borneo, Malaysian peat is within the range of carbon fractions reported for Kalimantan and Sumatra, Indonesia peat (44–60 % C) (Christian et al., 2003; Hatch et al., 2015; Iinuma et al., 2007; May et al., 2014; Setyawati et al., 2017). The low carbon content (20–30 % C) of Guizhou peats is similar to the 28–30 % C reported for two eastern North Carolina, USA peats (Black et al., 2016).

Hydrogen contents of 2–7 % H in Table 1 are consistent with abundances found elsewhere, including: 1) ~6 % H for northern Minnesota, USA peat (Yokelson et al., 1997); 2) ~2–3 % H for the eastern North Carolina peat (Black et al., 2016); and 3) ~5–7 % H for Indonesian peats (Christian et al., 2003; Hatch et al., 2015; Iinuma et al., 2007). Sulfur (S) contents are below



235 detection limits ( $<0.01\%$ ), and the nitrogen contents are  $1\text{--}4\%$  N. Ratios of N/C are  $0.02\text{--}0.08$ ,  
236 consistent with the reported N/C ratios of: 1)  $0.036$  for Neustädter Moor, northern Germany  
237 (Iinuma et al., 2007); 2)  $0.017\text{--}0.04$  for Ireland and United Kingdom (Wilson et al., 2015); 3)  $0.02\text{--}$   
238  $0.03$  for Alberta and Ontario, Canada (Stockwell et al., 2014); 4)  $0.062$  for Minnesota, U.S.A.  
239 (Yokelson et al., 1997); 5)  $0.022\text{--}0.03$  for the eastern coast of North Carolina, U.S.A. (Black et  
240 al., 2016); and 6)  $0.036\text{--}0.039$  for Kalimantan and Sumatra, Indonesia (Christian et al., 2003; Hatch  
241 et al., 2015).

242 The sum of elements (i.e., C, H, N, S, and O) accounts for  $91\text{--}98\%$  of total mass except  
243 for the Guizhou peats ( $47\text{--}56\%$ ). As Guizhou peats appear to be a mixture of peat and soil, these  
244 samples may represent degraded peats (Miettinen et al., 2017) or contain additional minerals or  
245 high ash contents, similar to North Carolina peats ( $44\text{--}62\%$  ash, Black et al., 2016). Therefore,  
246 these peats were only used for preliminary testing of sample ignition and heating to optimize  
247 burning conditions. Overall, the six other peats in Table 1 represent biomes from different regions  
248 of the world.

### 249 **3.2 Emission factors (EFs)**

250 Table S3 summarizes the experimental setup for the 40 peat combustion tests and contains  
251 the peat mass measurements before and after each burn. The after burn residue may have contained  
252 unburned peat as well as non-combustible ash, and dry-mass based emission factors are normalized  
253 to the difference between before and after mass. The residues were not analyzed for C and N  
254 contents. A few samples were voided due to sampling abnormalities. The following analyses are  
255 based on the 32 paired (Fresh vs Aged) samples at  $25\%$  fuel moisture and 3 paired samples at  $60\%$   
256 fuel moisture. The amount of fuel consumed per test ranged from  $21\text{--}48\text{ g}$  for all but Russian  
257 peat ( $14\text{--}15\text{ g}$ ) due to limited supply.

258  $\text{PM}_{2.5}$  mass concentrations were high, in the range of  $328\text{--}2277\text{ }\mu\text{g}/\text{m}^3$ . Typical sample  
259 durations from ignition to completion were  $\sim 40\text{--}60$  minutes, except for the Everglades (FL) peats  
260 that took longer (up to 135 minutes). Similar particle loadings (mostly within  $\pm 20\%$ ) were found  
261 between downstream (aged) and upstream (fresh) samples. The exception is Everglades (FL) peat,  
262 where prolonged sample durations and 7-days aging times resulted in higher downstream particle  
263 loadings with ratios of aged/fresh ranging from  $1.6\text{--}2.0$ .



### 3.2.1 Gaseous carbon emission factors

Individual and average carbonaceous gas EFs are summarized in Table S4. Apparent variations by biome are found among the different peats with relative standard deviations ranging from 2% to 27%. The largest EFs are found for CO<sub>2</sub> (EF<sub>CO<sub>2</sub></sub>), ranging from 994–1455 g kg<sup>-1</sup>, which are 1–2 orders of magnitude higher than the corresponding EF<sub>CO</sub> and EF<sub>CH<sub>4</sub></sub>. Average EF<sub>CO<sub>2</sub></sub> varied by >30 % among biomes, ranging from 1073 ± 61 to 1400 ± 38 g kg<sup>-1</sup> for the Russian and Alaskan peats, respectively.

Muraleedharan et al. (2000) reported the first laboratory-combustion EFs of 150–185 g kg<sup>-1</sup> for EF<sub>CO<sub>2</sub></sub>, 15–37 g kg<sup>-1</sup> for EF<sub>CO</sub>, and 6–11 g kg<sup>-1</sup> for EF<sub>CH<sub>4</sub></sub> on a wet mass basis for Brunei peat with a 51.4% moisture content. Table 2 shows studies conducted over the past decade, with more field monitoring during the 2015 ENSO period in Indonesia. Open path (OP)-FTIR was commonly used to acquire gaseous emissions with MCEs ranging 0.77–0.86, consistent with smoldering combustion. A limited number of burns (n of 1 to 6) were conducted in the laboratory using combustion chambers, whereas a larger number of in situ field-burn samples (n of 17 to 35) were acquired in southeast Asia (Setyawati et al., 2017; Stockwell et al., 2016; Wooster et al., 2018).

Table 2 shows over 2-fold variations in EF<sub>CO<sub>2</sub></sub> among studies. The highest EF<sub>CO<sub>2</sub></sub> with the lowest variability was found for tropical peats (ranges 1331–1831 g kg<sup>-1</sup> for smoldering). The average EF<sub>CO<sub>2</sub></sub> (1331 ± 78 g kg<sup>-1</sup>) for Malaysian peat (n=6) from this study is ~16 % and ~18 % lower than the 1579 ± 58 and 1615 ± 184 g kg<sup>-1</sup> for Peninsula, Malaysia (Smith et al., 2018) and average boreal/temperate peats (Hu et al., 2018), respectively. Malaysian peat EF<sub>CO<sub>2</sub></sub> measured in this study is 20 % lower than the 1681 ± 96 g kg<sup>-1</sup>, averaged from seven studies of Kalimantan and Sumatra, Indonesia peats (Christian et al., 2003; Huijnen et al., 2016; Nara et al., 2017; Stockwell et al., 2014).

Overall average EF<sub>CO<sub>2</sub></sub> (1269 ± 139 g kg<sup>-1</sup>, n=32) from this study (Table S4) are ~19–25 % lower than the 1563 ± 65 g kg<sup>-1</sup> for peatland fires used in atmospheric models (Akagi et al., 2011); 1550 ± 130 g kg<sup>-1</sup> in a recent review (Andreae, 2019); and 1703 g kg<sup>-1</sup> (Christian et al., 2003) adopted by the 2014 Intergovernmental Panel on Climate Change (IPCC) for organic soil fire inventories (IPCC, 2014). EFs derived from this study cover four biomes and may improve global emission estimates.



293 Average  $EF_{CO}$  is typically  $\sim 12\text{--}15\%$  of  $EF_{CO_2}$  in the range of  $157\text{--}171\text{ g kg}^{-1}$  for all but the  
 294 two Florida peats with  $394 \pm 46\text{ g kg}^{-1}$  ( $MCE = 0.65 \pm 0.04$ ) and  $93 \pm 21\text{ g kg}^{-1}$  ( $MCE = 0.90 \pm$   
 295  $0.03$ ) for the Putnam and Everglades (FL) peats, respectively (Table S4 and Table 2). This is  
 296 consistent with a higher  $EF_{CO}$  under lower MCEs reported by Setyawati et al. (2017) –a 45-fold  
 297 increase from  $3.1 \pm 7.2\text{ g kg}^{-1}$  for flaming ( $MCE = 0.998 \pm 0.005$ ) to  $138 \pm 72\text{ g kg}^{-1}$  for smoldering  
 298 ( $MCE = 0.894 \pm 0.055$ ) combustion.

299 Average  $EF_{CO}$  of  $157\text{--}161\text{ g kg}^{-1}$  for boreal and temperate peats are  $\sim 10\%$  lower than the  
 300  $179 \pm 61\text{ g kg}^{-1}$  from Hu et al. (2018). The overall average  $EF_{CO}$  of  $175 \pm 92\text{ g kg}^{-1}$  from this study  
 301 is  $\sim 4\%$  lower than the  $182 \pm 60\text{ g kg}^{-1}$  in Akagi et al. (2011),  $\sim 30\%$  lower than the  $250 \pm 23\text{ g kg}^{-1}$   
 302 <sup>1</sup> in Andreae (2019), and  $\sim 15\%$  lower than the  $207\text{--}210\text{ g kg}^{-1}$  used in IPCC (2014).

303 Average  $EF_{CH_4}$  is  $\sim 0.3\text{--}0.9\%$  of  $EF_{CO_2}$ , lowest for cold climates with  $3.2\text{--}6.9\text{ g kg}^{-1}$  for the  
 304 boreal and temperate peats and  $6.7\text{--}10.4\text{ g kg}^{-1}$  for the subtropical and tropical peats (Table S4).  
 305 Table 2 shows that  $EF_{CH_4}$  for Malaysian and Indonesian peats exceeding  $\sim 10\text{ g kg}^{-1}$  in five of the  
 306 eight past studies. These EFs are more in line with the  $11.8 \pm 7.8\text{ g kg}^{-1}$  in Akagi et al. (2011),  $9.3$   
 307  $\pm 1.5\text{ g kg}^{-1}$  in Andreae (2019), and  $9\text{--}21\text{ g kg}^{-1}$  in IPCC (2014), but are higher than the average  
 308 ( $6.6 \pm 2.4\text{ g kg}^{-1}$ ) found in this study.

### 309 3.2.2 Gaseous nitrogen emission factors

310 Individual and average gaseous nitrogen species EFs are summarized in Table S5.  $EF_{NO}$   
 311 and  $EF_{NO_2}$  are low in the range of  $0.2\text{--}2.1\text{ g kg}^{-1}$ . For fresh emissions, most of the  $NO_x$  ( $NO + NO_2$ )  
 312 are present as NO. After the OFR, NO decreased while  $NO_2$  increased, as shown in Fig. S3, but  
 313 depleted after the OFR as shown in Fig. S3. A low correlation coefficient ( $r = 0.67$ ) between the  
 314 downstream and upstream  $EF_{NO_x}$  indicates large variabilities among tests.

315 Table 3 shows that most studies do not report  $EF_{NO}$  or  $EF_{NO_2}$ , partially due to the low  
 316 concentrations and large variabilities under atmospheric aging. Stockwell et al. (2014;2016)  
 317 reported  $0.31\text{--}1.85\text{ g kg}^{-1}$   $EF_{NO}$  and  $2.31\text{--}2.36\text{ g kg}^{-1}$   $EF_{NO_2}$  for Indonesia peats. These levels are  
 318 much higher than the  $EF_{NO_x}$  (as  $NO_2$ ) of  $0.75 \pm 0.10\text{ g kg}^{-1}$  for Malaysian peat in this study.

319 Emissions for reactive nitrogen,  $EF_{NO_y}$  (as  $NO_2$ ), ranged  $0.61\text{--}6.3\text{ g kg}^{-1}$  with an average of  
 320  $2.4 \pm 1.4\text{ g kg}^{-1}$  (Table S5).  $EF_{NO_y} > 2.5\text{ g kg}^{-1}$  are found for the two Florida peats with an average  
 321 of  $4.3 \pm 1.1\text{ g kg}^{-1}$  for Everglades (FL), which reports higher N content ( $3.93 \pm 0.08\%$ ) among



322 peats (Table 1). Figure S4 shows that ~74 % of the  $\text{NO}_y$  is  $\text{NO}_x$  with high correlation coefficient  
323 ( $r = 0.93$ ). Nitrogen oxides are typically converted to other oxidized nitrogen species within 24  
324 hours after emission (Prenni et al., 2014; Seinfeld and Pandis, 1998). The ratio of  $\text{NO}_x/\text{NO}_y$  has  
325 been used to infer photochemical aging (Kleinman et al., 2003; Kleinman et al., 2007; Olszyna et  
326 al., 1994; Parrish et al., 1992). The high  $\text{NO}_x/\text{NO}_y$  ratios suggest that  $\text{NO}_x$  had not converted to  
327 other reactive nitrogen species in the diluted peat plume.

328  $\text{N}_2\text{O}$ , an inert form of oxide from nitrogen with an atmospheric lifetime of ~110 years,  
329 commonly emitted from fossil fuel, solid waste fertilizers, and biomass combustion, is a  
330 greenhouse gas defined by U.S. EPA (2016). Table S5 shows that  $\text{EF}_{\text{N}_2\text{O}}$  are similar to  $\text{EF}_{\text{NO}_y}$ , in  
331 the range of 1.1–4.4  $\text{g kg}^{-1}$  and average of  $2.0 \pm 0.7 \text{ g kg}^{-1}$ . The highest average  $\text{EF}_{\text{N}_2\text{O}}$  ( $3.6 \pm 0.6 \text{ g}$   
332  $\text{kg}^{-1}$ ) is found for Putnam (FL) peat.

333  $\text{HCN}$ , a known emission from biomass burning (Li et al., 2000; Stockwell et al., 2014),  
334 exhibits >7-fold differences (1.8–14  $\text{g kg}^{-1}$ ) in  $\text{EF}_{\text{HCN}}$  (Table S5). The average  $\text{EF}_{\text{HCN}}$  ( $11.5 \pm 2.3$   
335  $\text{g kg}^{-1}$ ) for Putnam (FL) peat is 2- to 5-fold higher than for the other biomes. Table 3 shows large  
336  $\text{EF}_{\text{HCN}}$  variations among studies, from  $0.73 \pm 0.50 \text{ g kg}^{-1}$  (Ireland, Wilson et al., 2015) to  $5.75 \pm$   
337  $1.60 \text{ g kg}^{-1}$  (Indonesia, Stockwell et al., 2016). More consistent  $\text{EF}_{\text{HCN}}$  are found for tropical peats  
338 in the range of 3–6  $\text{g kg}^{-1}$ . Average  $\text{EF}_{\text{HCN}}$  of this study,  $4.7 \pm 3.1 \text{ g kg}^{-1}$ , are in-line with the  $5.0 \pm$   
339  $4.9$  and  $4.4 \pm 1.2 \text{ g kg}^{-1}$  reported by Akagi et al. (2011) and Andreae (2019).

340  $\text{EF}_{\text{NH}_3}$  (0.4–8.3  $\text{g kg}^{-1}$ ) are of the same magnitude as  $\text{EF}_{\text{HCN}}$  except for the Everglades (FL)  
341 peat (9–18  $\text{g kg}^{-1}$ ). Total reduced nitrogen emissions,  $\text{EF}_{\text{NH}_3} + \text{EF}_{\text{HCN}}$ , for the two Florida peats  
342 (12–25  $\text{g kg}^{-1}$ ) are ~2- to 3-fold higher than those for other regions. Table 3 also shows high  
343 variabilities in  $\text{EF}_{\text{NH}_3}$  among studies (1–11  $\text{g kg}^{-1}$ ). The overall average of  $5.6 \pm 4.8 \text{ g kg}^{-1}$  in this  
344 study is consistent with the  $4.2 \pm 3.2 \text{ g kg}^{-1}$  in Andreae (2019), but ~50 % of the  $10.8 \pm 12.4 \text{ g kg}^{-1}$   
345 in Akagi et al. (2011). The high standard deviations associated with these averages signify large  
346 variabilities among experiments.

347 Figure S5a shows some scatter in  $\text{EF}_{\text{NH}_3}$  determined by FTIR and the impregnated filter,  
348 especially at high concentrations. The regression slope shows that  $\text{EF}_{\text{NH}_3}$  by the FTIR was ~22 %  
349 lower than that by filters with a correlation coefficient of 0.76. Variable baselines in the FTIR  
350 measurements along with some nitrogen content in the diluted air and breath  $\text{NH}_3$  (Hibbard and  
351 Killard, 2011) in the testing laboratory may have contributed to these variations. The impregnated



filter collects all of the  $\text{NH}_3$  over the sampling period, including amounts that are below the FTIR detection limits, so it is probably better representing the time-integrated  $\text{EF}_{\text{NH}_3}$ . Reduction of  $\text{EF}_{\text{NH}_3}$  is most apparent after atmospheric aging in Fig. S5b, with  $2\text{--}14 \text{ g kg}^{-1}$  in fresh emissions and reduced to  $\sim 0.5\text{--}3 \text{ g kg}^{-1}$  after aging with a regression slope of 0.11.

### 3.2.3 $\text{PM}_{2.5}$ and carbon emission factors

Continuous  $\text{PM}_{2.5}$  from the DustTrak with the factory calibration factor yielded  $\text{PM}_{2.5}$  EFs 3- to 5-fold of those derived from gravimetric analyses, higher than the 2-fold mass differences by Wooster et al. (2018). This discrepancy is well known as the factory calibration uses Arizona road dust with a size distribution that is much coarser than that of biomass burning. Therefore,  $\text{EF}_{\text{PM}_{2.5}}$  is calculated from the filter samples. Chow et al. (2019) present the species abundances from this study in  $\text{PM}_{2.5}$  mass based on the average fresh and aged profiles, separated by 2- and 7-day photochemical aging times simulated with the oxidation flow reactor (Aerodyne, 2019). The same approach is used in Table S6 to compare fresh and aged particle EFs. Comparisons between combined fresh vs. aged EFs for  $\text{PM}_{2.5}$  mass, carbon (OC, EC, and TC), and levoglucosan for individual tests are shown in Table S7.

$\text{EF}_{\text{PM}_{2.5}}$  varies >4-fold ( $14\text{--}61 \text{ g kg}^{-1}$ ) for different peats without large differences between fresh and aged emissions.  $\text{EF}_{\text{OC}}$  varied from  $9\text{--}44 \text{ g kg}^{-1}$  while  $\text{EF}_{\text{EC}}$  ( $0.00\text{--}2.2 \text{ g kg}^{-1}$ ) were low (Table S7). The majority of  $\text{EF}_{\text{PM}_{2.5}}$  consist of  $\text{EF}_{\text{OC}}$ , with average  $\text{EF}_{\text{OC}}/\text{EF}_{\text{PM}_{2.5}}$  ratios by peat type of 52–98 % in fresh emissions, followed by  $\sim 14\text{--}23$  % reductions after aging, with the exception of Putnam (FL) peats (remained at 69–70 %).

Reductions of  $\text{EF}_{\text{OC}}$  after  $\sim 7$  days of photochemical aging are most apparent ( $\sim 7\text{--}9 \text{ g kg}^{-1}$ ) for the boreal peats, with the largest degradation for low temperature OC1 (evolved at  $140^\circ\text{C}$ ), indicating losses of high vapor pressure SVOCs upon aging (Table S6). The two Florida peats exhibit an initial  $\text{EF}_{\text{OC}}$  decrease of  $\sim 2 \text{ g kg}^{-1}$  after 2-days aging, but with an increase of  $1.8\text{--}4.0 \text{ g kg}^{-1}$  after 7 days. However, these changes are less than the standard deviations associated with the averages.

$\text{EF}_{\text{WSOC}}$  varies by 5-fold ( $3\text{--}16 \text{ g kg}^{-1}$ ) with over a  $\sim 50$  % increase for the Putnam (FL) and Malaysian peats after 7 days. Average  $\text{EF}_{\text{WSOC}}$  by peat type accounts for  $\sim 16\text{--}36$  % and  $\sim 20\text{--}62$  % of fresh  $\text{EF}_{\text{PM}_{2.5}}$  and  $\text{EF}_{\text{OC}}$ , respectively, with  $\sim 6\text{--}16$  % increases in  $\text{EF}_{\text{WSOC}}/\text{EF}_{\text{OC}}$  ratios after aging, partially due to the reduction in  $\text{EF}_{\text{OC}}$  (Table S7). Figure S6 shows reduced correlation coefficients



( $r$  from 0.82 to 0.77 for  $\text{PM}_{2.5}$ , from 0.88 to 0.84 for OC, and 0.94 to 0.68 for WSOC) from 2- to 7-day aging.

Table 4 compares filter-based PM mass and carbon from different studies. As different carbon protocols yield different fractions of OC and EC (Watson et al 2005), so the analytical protocols are listed. Most studies follow either IMPROVE\_A thermal/optical reflectance (TOR) (Chow et al., 2007) or NIOSH thermal/optical transmittance (TOT) protocols (NIOSH, 1999). As the transmittance pyrolysis correction (i.e., TOT) accounts for charred OC both on the filter surface and organic vapor within the filter substrate, lower  $\text{EF}_{\text{EC}}$  are expected as compared to TOR (Chow et al., 2004). To remove the OC and EC split uncertainty, TC to PM mass ratios are listed for comparison. Two studies reported black carbon (BC) from a micro-aethalometer (Wooster et al., 2018) or a single particle soot photometer (SP2; May et al., 2014). As BC levels are very low, not much differences can be distinguished between EC and BC.

Most studies report  $\text{EF}_{\text{PM}_{2.5}}$  with a few exceptions for  $\text{EF}_{\text{PM}_{10}}$  (Iinuma et al., 2007; Kuwata et al., 2018) and  $\text{EF}_{\text{PM}_1}$  (May et al., 2014). As most of the  $\text{PM}_{10}$  is in the  $\text{PM}_{2.5}$  fraction for biomass combustion, particle size fractions have a minor effect on PM EFs (Geron and Hays, 2013; Hu et al., 2018).

Table 4 shows that the majority of  $\text{EF}_{\text{PM}_{2.5}}$  lies in the range of  $\sim 20\text{--}50 \text{ g kg}^{-1}$  with the exception of very low  $\text{EF}_{\text{PM}_{2.5}}$  of 4–8 and 6–7  $\text{g kg}^{-1}$  reported by Bhattarai et al. (2018) and Black et al. (2016). These are probably due to low filter mass loadings and limited testing ( $n$  of 1 to 3), which may result in large uncertainties in gravimetric mass.

Despite different carbon analysis methods, most  $\text{EF}_{\text{OC}}$  lies in the range of  $\sim 5\text{--}30 \text{ g kg}^{-1}$  with the exception of  $\text{EF}_{\text{OC}}$  ( $37 \text{ g kg}^{-1}$ ) for Putnam (FL) and  $\text{EF}_{\text{OA}}$  ( $34.5 \text{ g kg}^{-1}$ ) for Indonesian peat measured by a time-of-flight/mass spectrometer (May et al., 2014). Most studies show that  $\text{EF}_{\text{TC}}$  accounts for  $\sim 60\text{--}85 \%$  of the  $\text{EF}_{\text{PM}_{2.5}}$ , with low  $\text{EF}_{\text{EC}}$  ( $0.02\text{--}1.3 \text{ g kg}^{-1}$ ).

$\text{EF}_{\text{WSOC}}$  of 6–7 and 4–6  $\text{g kg}^{-1}$  for the Alaskan and Malaysian peats from this study are consistent with the 6.7 and 3.1  $\text{g kg}^{-1}$  from German and Indonesian peats in Iinuma et al. (2007), respectively.  $\text{EF}_{\text{Levoglucosan}}$  exhibits  $>2$  orders of magnitude variabilities among the biomes with 0.24–16  $\text{g kg}^{-1}$  and 0.24–9.6  $\text{g kg}^{-1}$  in fresh and aged emissions, respectively.

The highest  $\text{EF}_{\text{Levoglucosan}}$  is found for the average fresh Russian peats ( $15.8 \pm 2.9 \text{ g kg}^{-1}$ ), and this is diminished by 45 % after 7-day aging ( $8.8 \pm 2.1 \text{ g kg}^{-1}$ ). Few studies report  $\text{EF}_{\text{Levoglucosan}}$





and results are highly variable. The  $EF_{Levoglucosan}$  of  $0.57 \text{ g kg}^{-1}$  in  $PM_{2.5}$  (converted from  $46 \text{ mg/g}$  OC) by Jayarathne et al. (2018) is  $\sim 23 \%$  of the  $2.5 \text{ g kg}^{-1}$  by Iinuma et al. (2007), both for Indonesia peats. The  $EF_{Levoglucosan}$  of  $0.5\text{--}1.0 \text{ g kg}^{-1}$  from fresh Malaysian peat in this study is comparable to  $0.57 \text{ g kg}^{-1}$  by Jayarathne et al. (2018). The  $4.6 \text{ g kg}^{-1}$  of  $EF_{Levoglucosan}$  for the northern German peat (Iinuma et al., 2007) is higher than the  $1.2\text{--}4.7 \text{ g kg}^{-1}$  found for the average Siberian and Alaskan peats in this study.

EFs for ionic nitrogen species are low ( $<0.1 \text{ g kg}^{-1}$ ) in fresh emissions. Both  $EF_{NH_4^+}$  and  $EF_{NO_3^-}$  increase with 7-day aging –  $>0.5 \text{ g kg}^{-1}$   $EF_{NH_4^+}$  for all peat and  $>1 \text{ g kg}^{-1}$   $EF_{NO_3^-}$  for all but Russian ( $0.79 \pm 0.08 \text{ g kg}^{-1}$ ) and Putnam (FL) peats ( $0.66 \pm 0.08 \text{ g kg}^{-1}$ ), consistent with the formation of secondary inorganic aerosol.

### 3.3 Effect of fuel moisture content on emission factors

Only a few studies examine the effects of fuel moisture on peat emissions with inconsistent results. An early study by McMahon et al. (1980) reported high emissions for total suspended particle (TSP,  $\sim 30\text{--}60 \mu\text{m}$ ) of  $30 \pm 20 \text{ g kg}^{-1}$  for dry ( $<11 \%$  moisture) as compared to  $4.1 \pm 3.8 \text{ g kg}^{-1}$  (after the first 24 hours) for wet ( $53\text{--}97 \%$  moisture) organic soil. Rein et al. (2009) found higher  $CO_2$  (but not CO) yields while increasing fuel moisture to 600% for tests of boreal Scotland peats in a cone calorimeter which continuously supplies heat to the fuel. Smoldering combustion is possible at high measured *in situ* fuel-moisture contents when surrounding peat provides insulation and heat from combustion is available for drying just in advance of the advancing front, but such samples will not burn in the laboratory. Watts (2013) sustained lab-based peat smoldering from a cypress swamp (FL) at  $\sim 250\%$  moisture content, which appears to be a maximum.

Table 2 shows that increasing moisture content from  $\sim 25 \%$  to  $\sim 60 \%$  for the three Putnam (FL) peats resulted in a 11 % increase in  $EF_{CO_2}$  but 20 % and 12 % reductions for  $EF_{CO}$  and  $EF_{CH_4}$ , respectively. No consistent changes are found for nitrogen species (Table 3), with negligible changes in  $EF_{NH_3}$  and  $EF_{HCN}$ ; 13–30 % reduction in  $EF_{NO}$ ,  $EF_{NO_x}$ , and  $EF_{NO_y}$ , as well as 45 % and 9 % increase in  $EF_{NO_2}$  and  $EF_{N_2O}$  respectively. On the other hand, a reduction of  $\sim 30 \%$  is found for  $EF_{PM_{2.5}}$  (Table 4) from 25 % to 60 % fuel moisture. Higher fuel moisture contents typically result in less efficient burning conditions, thereby increasing CO and reducing MCE (Chen et al., 2010). However, an opposite trend is found with  $EF_{CO}$  reduced from  $394 \pm 46$  to  $315 \pm 10 \text{ g kg}^{-1}$





and MCEs increased from  $0.65 \pm 0.04$  to  $0.72 \pm 0.01$ . Overall, the EFs for ~60 % moisture contents are comparable to EFs for the six other peats with ~25 % moisture content.

Increased (~25 to 60 %) fuel moisture yields a ~20 % reduction in fresh  $EF_{OC}$ , much lower than the 35–43 % reduction (~25 to 50 % moisture) reported by Chakrabarty et al. (2016) for the Siberian and Alaskan peats. By increasing fuel moisture, Chakrabarty et al (2016) also reported an increase in  $EF_{CO_2}$  by 20 % but a ~75% reduction and 35% increase in  $EF_{CO}$  for Siberian and Alaskan peats, respectively, based on a single sample.

### 3.4 Distribution of carbon and nitrogen species

Figure 3 shows the distribution of carbonaceous species. Because the EFs are calculated based on the carbon mass balance method (Eq. (2)), the total emitted carbon is assumed to be the same as total consumed carbon. The majority (>90 %) of total emitted carbon are present in the gas phase with 54–75 %  $CO_2$ , followed by 8–30 %  $CO$ . On average, emitted carbon includes  $69.8 \pm 7.5$  %  $CO_2$ ,  $14.8 \pm 6.5$  %  $CO$ ,  $1.0 \pm 0.3$  %  $CH_4$ ,  $9.4 \pm 2.4$  % carbon compounds, and  $4.8 \pm 1.3$  %  $PM_{2.5}$  TC. The highest ( $30 \pm 4$  %) and lowest ( $8.4 \pm 1.9$  %)  $CO$  abundances for the Putnam and Everglades (FL) peats are consistent with the lowest and highest average MCEs of 0.65 and 0.90, respectively.

The nitrogen budget in Fig. 4 accounts for 24–52% of N in the consumed fuel. Since burn temperatures are below those at which  $NO_x$  forms from oxygen reactions with  $N_2$  in the air, most of the N in emissions derives from the N content of the fuels. Kuhlbusch et al. (1991) found  $N_2$  emissions constituted an average of  $31 \pm 20$  % of N in consumed grass, hay, pine needle, clover, and wood fuels. Since  $N_2$  measurements require combustion in  $N_2$ -free atmosphere (e.g., a He- $O_2$  mixture),  $N_2$  was not quantified here, but it was probably emitted in similar quantities. Isocyanic acid (HNCO) is another important N-containing compound found in biomass burning emissions (Roberts et al., 2011). Koss et al. (2018) report a 0.16 g/kg nitrogen-equivalent EF (0.5 g/kg for HNCO) for a peat sample, comparable to EFs for several of the measured nitrogen compounds summarized in Table 3. Other nitrogen-containing gases reported by Koss et al. (2018) with EFs >0.1 g/kg include acetonitrile ( $CH_3CN$ ), acetamide ( $CH_3CONH_2$ ), benzonitrile ( $C_6H_5CN$ ), and pyridine + pentadienenitriles ( $C_5H_5N$ ), which could account for part of the unmeasured N in emissions. Neff et al. (2002) found that organic nitrogen formed from photochemical reactions of hydrocarbon with  $NO_x$  plays an important role in the global nitrogen cycle. Approximately  $30 \pm 16$  % of Neff et al.'s total nitrogen was from organic nitrogen, similar to the 25 % of total nitrogen



deposition flux reported by Jickells et al. (2013). Alkaloids, dissolved organic nitrogen, along with nitroaromatic compounds have been reported (e.g., Benitez et al., 2009; Kopacek and Posch, 2011; Koppmann et al., 2005; Kuhlbusch et al., 1991; Laskin et al., 2009; Stockwell et al., 2015).

The majority (>99 %) of the measured nitrogen in emissions is in the gas phase. On average, 16.7 % of the fuel nitrogen was emitted as  $\text{NH}_3$  and 9.5 % was emitted as  $\text{HCN}$ .  $\text{N}_2\text{O}$  and  $\text{NO}_y$  constituted 5.7 % and 2.9 % of N in the consumed fuel.  $\text{NH}_3$  emissions accounted for 26–28 % of consumed N for Everglades (FL) and Malaysian peats while  $\text{HCN}$  emissions dominated fuel N (13–17 %) for the Putnam (FL) and Malaysian peats. The fraction of  $\text{N}_2\text{O}$  emissions in Malaysian peat N ( $10.3 \pm 1.1$  %) was more than twice the fractions found for the other regions with reactive nitrogen ( $\text{NO}_y$ ) only accounting for 2–4 % of the fuel N. The sum of  $\text{NH}_3$  and  $\text{HCN}$  nitrogen ranged 35–39 % of consumed N for the Malaysian and Everglades (FL) peats, which is about 3 times the fraction for Russian peat.

Lobert et al. (1990) point out the importance of nitrogen-containing gases in biomass burning for the atmospheric nitrogen balance. On average, the emitted nitrogen includes  $17 \pm 10$  %  $\text{NH}_3$ ,  $9.5 \pm 3.8$  %  $\text{HCN}$ ,  $5.7 \pm 2.5$  %  $\text{N}_2\text{O}$ ,  $2.8 \pm 1.0$  %  $\text{NO}_y$  (including  $\text{NO}_x$ ), and  $0.14 \pm 0.18$  % of PM nitrogen (sum of  $\text{NO}_2^-$ ,  $\text{NO}_3^-$ , and  $\text{NH}_4^+$ ). The average nitrogen budget accounts for  $35 \pm 11$  % of the total consumed nitrogen, consistent with past studies showing that ~one- to two-thirds of the fuel nitrogen is accounted for during biomass combustion.

#### 4 Summary and conclusions

This paper reports fuel composition and emission factors (EFs) from laboratory chamber combustion of six types of peat fuels representing boreal (Russia and Siberia), temperate (northern Alaska, USA), subtropical (northern and southern Florida, USA), and tropical (Borneo, Malaysia) climate regions. Dried peat contains 44–57 % carbon (C), 31–39 % oxygen (O), 5–6 % hydrogen (H), 1–4 % nitrogen (N), and <0.01 % Sulfur (S). The N content only accounts for 2–8 % of the C, consistent with peat compositions reported in other studies.

Thirty-two tests with 25 % fuel moisture were reported with predominant smoldering combustion conditions ( $\text{MCE} = 0.82 \pm 0.08$ ). Average fuel-based EFs for  $\text{CO}_2$  ( $\text{EF}_{\text{CO}_2}$ ) are highest ( $1400 \pm 38 \text{ g kg}^{-1}$ ) and lowest ( $1073 \pm 63 \text{ g kg}^{-1}$ ) for the Alaskan and Russian peats, respectively.  $\text{EF}_{\text{CO}}$  and  $\text{EF}_{\text{CH}_4}$  are ~12–15 % and ~0.3–0.9 % of  $\text{EF}_{\text{CO}_2}$  in the range of ~157–171 g  $\text{kg}^{-1}$  and 3–



10 g kg<sup>-1</sup>, respectively. The exception is the two Florida peats, reporting the highest (394 ± 46 g kg<sup>-1</sup>) and lowest (93 ± 21 g kg<sup>-1</sup>) EF<sub>CO</sub> for the Putnam and Everglades (FL) peats, respectively.

Filter-based EF<sub>PM<sub>2.5</sub></sub> varied by over 4-fold (14–61 g kg<sup>-1</sup>) without appreciable changes between fresh and aged emissions. The majority of EF<sub>PM<sub>2.5</sub></sub> consists of EF<sub>OC</sub>, with average EF<sub>OC</sub>/EF<sub>PM<sub>2.5</sub></sub> ratios by peat type in the range of 52–98 % in fresh emissions, followed by ~14–23 % reduction after aging with the exception of Putnam (FL) peats (retained 69–70 %). Reduction of EF<sub>OC</sub> (~7–9 g kg<sup>-1</sup>) are most apparent for boreal peats with the largest decrease in low temperature OC1 (evolved at 140 °C), suggesting the loss of high vapor pressure semi-volatile organic compounds during aging. EFs for water-soluble OC (EF<sub>WSOC</sub>) accounts for ~20–62 % of EF<sub>OC</sub> with ~6–16 % increase in EF<sub>WSOC</sub>/EF<sub>OC</sub> ratios after aging. The highest EF<sub>Levoglucon</sub> is found for average Russian peat (15.8 ± 2.9 g kg<sup>-1</sup>) with a degradation of 45 % after aging.

The majority (>90 %) of the total emitted carbon is in the gas phases with 54–75 % CO<sub>2</sub>, followed by 8–30 % CO. Nitrogen budget only explains 24–52 % of the consumed nitrogen with an average of 35 ± 11 %, consistent with past studies that ~one- to two-thirds of the total nitrogen are lost upon biomass combustion. The majority (>99 %) of the total emitted nitrogen is in the gas phase, dominated by the two reduced nitrogen species with 16.7 % for NH<sub>3</sub> and 9.5 % for HCN. N<sub>2</sub>O and NO<sub>y</sub> are detectable at 5.7 % and 2.9 % abundance. EFs from this study can be used to refine current emissions inventories.

## 5 Author contributions

JGW, JCC, JC, L-WAC, and XW jointly designed the study, performed the data analyses, and prepared the manuscript. QW, JT, and SSHH carried out the peat combustion experiments. SG conducted emission factor calculations. ACW provided peat fuels.

## 6 Competing interests

The authors declare that there are no conflicts of interest.

## 7 Acknowledgements

This research was primarily supported by the National Science Foundation (NSF, AGS-1464501 and CHE-1214163) as well as internal funding from both the Desert Research Institute, Reno, NV, USA, and Institute of Earth Environment, Chinese Academy of Sciences, Xian, China. The Caohai and Gaopo peat samples were provided by Dr. Pinhua Xia of Guizhou Normal University, Guizhou, China and Dr. Chunmao Zhu of the Japan Agency for Marine-Earth Science and Technology, Yokosuka, Japan.



## 8 References

- Aerodyne: Potential Aerosol Mass (PAM) oxidation flow reactor, Aerodyne Research Inc., Billerica, MA, 2019.  
<http://www.aerodyne.com/sites/default/files/u17/PAM%20Potential%20Aerosol%20Mass%20Reactor.pdf>
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, 2011.
- Allen, C., Carrico, C. M., Gomez, S. L., Andersen, P. C., Turnipseed, A. A., Williford, C., Birks, J. W., Salisbury, D., Carrion, R., Gates, D., Macias, F., Rahn, T., Aiken, A. C., and Dubey, M. K.: NO<sub>x</sub> instrument intercomparison for laboratory biomass burning source studies and urban ambient measurements in Albuquerque, New Mexico, *J. Air Waste Manage. Assoc.*, 68, 1175–1189, 10.1080/10962247.2018.1487347, 2018.
- Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K. E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., St Clair, J. M., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C., and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmos. Chem. Phys.*, 10, 9739–9760, 2010.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning- An updated assessment, *Atmospheric Chemistry and Physics Discussion*, 1–27, 10.5194/acp-2019-303, 2019.
- Ballenthin, J. O., Thorn, W. F., Miller, T. M., Viggiano, A. A., Hunton, D. E., Koike, M., Kondo, Y., Takegawa, N., Irie, H., and Ikeda, H.: In situ HNO<sub>3</sub> to NO<sub>y</sub> instrument comparison during SOLVE, *Journal of Geophysical Research*, 108, ACH 7-1–ACH 7-11, 10.1029/2002JD002136, 2003.
- Benitez, J. M. G., Cape, J. N., Heal, M. R., van Dijk, N., and Diez, A. V.: Atmospheric nitrogen deposition in south-east Scotland: Quantification of the organic nitrogen fraction in wet, dry and bulk deposition, *Atmos. Environ.*, 43, 4087–4094, 2009.
- Bhattacharai, C., Samburova, V., Sengupta, D., Iaukea-Lum, M., Watts, A. C., Moosmuller, H., and Khlystov, A. Y.: Physical and chemical characterization of aerosol in fresh and aged emissions from open combustion of biomass fuels, *Aerosol Sci. Technol.*, 52, 1266–1282, 10.1080/02786826.2018.1498585, 2018.
- Black, R. R., Aurell, J., Holder, A., George, I. J., Gullett, B. K., Hays, M. D., Geron, C. D., and Tabor, D.: Characterization of gas and particle emissions from laboratory burns of peat, *Atmos. Environ.*, 132, 49–57, 10.1016/j.atmosenv.2016.02.024, 2016.
- Cao, J. J., Wang, Q. Y., Tan, J., Zhang, Y. G., Wang, W. J., Zhong, B. L., Ho, S. S. H., Chen, L.-W. A., Wang, X. L., Watson, J. G., and Chow, J. C.: Evaluation of the oxidation flow reactor for particulate matter emission limit certification, *Atmos. Environ.*, submitted, 2019.
- Chakrabarty, R. K., Gyawali, M., Yatavelli, R. L. N., Pandey, A., Watts, A. C., Knue, J., Chen, L. W. A., Pattison, R. R., Tsibert, A., Samburova, V., and Moosmuller, H.: Brown carbon aerosols from burning of boreal peatlands: microphysical properties, emission factors, and implications for direct radiative forcing, *Atmos. Chem. Phys.*, 16, 3033–3040, 10.5194/acp-16-3033-2016, 2016.
- Chen, Y., Randerson, J. T., van der Werf, G. R., Morton, D. C., Mu, M., and Kasibhatla, P. S.: Nitrogen deposition in tropical forests from savanna and deforestation fires, *Global Change Biology*, 16, 2024–2038, 2010.
- Cheng, Y. H., Shiu, B. T., Lin, M. H., and Yan, J. W.: Levels of black carbon and their relationship with particle number levels-observation at an urban roadside in Taipei City  
 CHENG2013A, *Environmental Science and Pollution Research*, 20, 1537–1545, 2013.



- 575 Chow, J. C., Watson, J. G., Chen, L.-W. A., Chang, M.-C. O., Robinson, N. F., Trimble, D. L., and Kohl, S. D.: The  
 576 IMPROVE A temperature protocol for thermal/optical carbon analysis: Maintaining consistency with a long-term  
 577 database, *J. Air Waste Manage. Assoc.*, 57, 1014-1023, 2007.
- 578 Chow, J. C., Wang, X. L., Sumlin, B. J., Gronstal, S. B., Chen, L.-W. A., Trimble, D. L., Kohl, S. D., Mayorga, S.  
 579 R., Riggio, G. M., Hurbain, P. R., Johnson, M., Zimmermann, R., and Watson, J. G.: Optical calibration and  
 580 equivalence of a multiwavelength thermal/optical carbon analyzer, *Aerosol Air Qual. Res.*, 15, 1145-1159,  
 581 doi:10.4209/aaqr.2015.02.0106, 2015.
- 582 Chow, J. C., and Watson, J. G.: Enhanced ion chromatographic speciation of water-soluble PM<sub>2.5</sub> to improve  
 583 aerosol source apportionment, *Aerosol Science and Engineering*, 1, 7-24, doi:10.1007/s41810-017-0002-4, 2017.
- 584 Chow, J. C., Cao, J., Chen, L.-W. A., Wang, X. L., Wang, Q. Y., Tian, J., Ho, S. S. H., Carlson, T. N., Kohl, S. D.,  
 585 and Watson, J. G.: Evaluating changes in PM<sub>2.5</sub> peat combustion source profiles with atmospheric aging in an  
 586 oxidation flow reactor, *Atmospheric Measurement Techniques Discussions*, 1-44, 10.5194/amt-2019-198, 2019.
- 587 Christian, T. J., Kleiss, B., Yokelson, R. J., Holzinger, R., Crutzen, P. J., Hao, W. M., Saharjo, B. H., and Ward, D.  
 588 E.: Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from Indonesian, African,  
 589 and other fuels, *Journal of Geophysical Research*, 108, 2003.
- 590 Crutzen, P. J., and Andreae, M. O.: Biomass burning in the tropics: Impact on atmospheric chemistry and  
 591 biogeochemical cycles, *Science*, 250, 1669-1678, 1990.
- 592 Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., Diskin,  
 593 G. S., Fisher, J. A., Fuelberg, H. E., Hecobian, A., Knapp, D. J., Mikoviny, T., Riemer, D., Sachse, G. W., Sessions,  
 594 W., Weber, R. J., Weinheimer, A. J., Wisthaler, A., and Jimenez, J. L.: Effects of aging on organic aerosol from  
 595 open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049-12064, 2011.
- 596 Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian,  
 597 K. L., Borrmann, S., and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) - Instrument  
 598 description and first field deployment, *Aerosol Sci. Technol.*, 39, 637-658, 2005.
- 599 Geron, C., and Hays, M.: Air emissions from organic soil burning on the coastal plain of North Carolina, *Atmos.*  
 600 *Environ.*, 64, 192-199, 2013.
- 601 Goulding, K. W. T., Bailey, N., Bradbury, N. J., Hargreaves, P., Howe, M., Murphy, D. V., Poulton, P. R., and  
 602 Willison, T. W.: Nitrogen deposition and its contribution to nitrogen cycling and associated soil processes, *New*  
 603 *Phytologist*, 139, 49-58, 1998.
- 604 Gruber, N., and Galloway, J. N.: An Earth-system perspective of the global nitrogen cycle, *Nature*, 451, 293-296,  
 605 2008.
- 606 Hatch, L. E., Luo, W., Pankow, J. F., Yokelson, R. J., Stockwell, C. E., and Barsanti, K. C.: Identification and  
 607 quantification of gaseous organic compounds emitted from biomass burning using two-dimensional gas  
 608 chromatography-time-of-flight mass spectrometry, *Atmos. Chem. Phys.*, 15, 1865-1899, 2015.
- 609 Hibbard, T., and Killard, J.: Breath ammonia levels in a normal human population study as determined by  
 610 photoacoustic laser spectroscopy, *Journal of Breath Research*, 5, 1-8, 2011.
- 611 Hu, Y. Q., Fernandez-Anez, N., Smith, T. E. L., and Rein, G.: Review of emissions from smouldering peat fires and  
 612 their contribution to regional haze episodes, *International Journal of Wildland Fire*, 27, 293-312, 10.1071/wf17084,  
 613 2018.



- 614 Huijnen, V., Wooster, M. J., Kaiser, J. W., Gaveau, D. L. A., Flemming, J., Parrington, M., Inness, A., Murdiyarso,  
 615 D., Main, B., and van Weele, M.: Fire carbon emissions over maritime southeast Asia in 2015 largest since 1997,  
 616 Scientific Reports, 6, 10.1038/srep26886, 2016.
- 617 Iinuma, Y., Bruggemann, E., Gnauk, T., Muller, K., Andreae, M. O., Helas, G., Parmar, R., and Herrmann, H.:  
 618 Source characterization of biomass burning particles: The combustion of selected European conifers, African  
 619 hardwood, savanna grass, and German and Indonesian peat, J. Geophys. Res. Atmos., 112, 2007.
- 620 IPCC: 2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands,  
 621 Switzerland, 2014. [https://www.ipcc.ch/site/assets/uploads/2018/03/Wetlands\\_Supplement\\_Entire\\_Report.pdf](https://www.ipcc.ch/site/assets/uploads/2018/03/Wetlands_Supplement_Entire_Report.pdf)
- 622 Jaakkola, P. T., Vahlman, T. A., Roos, A. A., Saarinen, P. E., and Kauppinen, J. K.: On-line analysis of stack gas  
 623 composition by a low resolution FT-IR gas analyzer, Water Air and Soil Pollution, 101, 79-92, 1998.
- 624 Jayarathne, T., Stockwell, C. E., Gilbert, A. A., Daugherty, K., Cochrane, M. A., Ryan, K. C., Putra, E. I., Saharjo,  
 625 B. H., Nurhayati, A. D., Albar, I., Yokelson, R. J., and Stone, E. A.: Chemical characterization of fine particulate  
 626 matter emitted by peat fires in Central Kalimantan, Indonesia, during the 2015 El Nino, Atmos. Chem. Phys., 18,  
 627 2585-2600, 10.5194/acp-18-2585-2018, 2018.
- 628 Jickells, T., Baker, A. R., Cape, J. N., Cornell, S. E., and Nemitz, E.: The cycling of organic nitrogen through the  
 629 atmosphere, Philosophical Transactions: Biological Sciences, 368, 1-7, 2013.
- 630 Kleinman, L. I., Daum, P. H., Lee, Y.-N., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J., Hyde, P.,  
 631 Doskey, P. V., Rudolph, J., Fast, J., and Berkowitz, C.: Photochemical age determinations in the Phoenix  
 632 metropolitan area, Journal of Geophysical Research, 108, 2003.
- 633 Kleinman, L. I., Daum, P. H., Lee, Y. N., Senum, G. I., Springston, S. R., Wang, J., Berkowitz, C., Hubbe, J.,  
 634 Zaveri, R. A., Brechtel, F. J., Jayne, J., Onasch, T. B., and Worsnop, D. R.: Aircraft observations of aerosol  
 635 composition and ageing in New England and Mid-Atlantic States during the summer 2002 New England Air Quality  
 636 Study field campaign, J. Geophys. Res. Atmos., 112, 2007.
- 637 Kopacek, J., and Posch, M.: Anthropogenic nitrogen emissions during the Holocene and their possible effects on  
 638 remote ecosystems, Global Biogeochemical Cycles, 25, 1-17, 2011.
- 639 Koppmann, R., von Czapiewski, K., and Reid, J. S.: A review of biomass burning emissions, part 1: gaseous  
 640 emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen containing compounds,  
 641 Atmospheric Chemistry and Physics Discussion, 5, 10455-10516, 2005.
- 642 Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M.,  
 643 Brown, S. S., Jimenez, J. L., Krechmer, J., Roberts, J. M., Warneke, C., Yokelson, R. J., and de Gouw, J.: Non-  
 644 methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-  
 645 ToF during the FIREX 2016 laboratory experiment, Atmos. Chem. Phys., 18, 3299-3319, 10.5194/acp-18-3299-  
 646 2018, 2018.
- 647 Kuhlbusch, T. A., Lobert, J. M., Crutzen, P. J., and Warneck, P.: Molecular nitrogen emissions from denitrification  
 648 during biomass burning, Nature, 351, 135-137, 1991.
- 649 Kuwata, M., Neelam-Naganathan, G. G., Miyakawa, T., Khan, M. F., Kozan, O., Kawasaki, M., Sumin, S., and  
 650 Latif, M. T.: Constraining the emission of particulate matter from Indonesian peatland burning using continuous  
 651 observation data, J. Geophys. Res. Atmos., 123, 9828-9842, 10.1029/2018jd028564, 2018.
- 652 Laskin, A., Smith, J. S., and Laskin, J.: Molecular Characterization of Nitrogen-Containing Organic Compounds in  
 653 Biomass Burning Aerosols Using High-Resolution Mass Spectrometry, Environ. Sci. Technol., 43, 3764-3771,  
 654 2009.





- 655 Li, Q., Jacob, B. D. J., Bey, I., Yantosca, R. M., Zhao, Y. J., Kondo, Y., and Notholt, J.: Atmospheric hydrogen  
 656 cyanide (HCN): Biomass burning source, ocean sink?, *Geophysical Research Letters*, 27, 357-360, 2000.
- 657 Lobert, J. M., Scharffe, D. H., Hao, W. M., and Crutzen, P. J.: Importance of biomass burning in the atmospheric  
 658 budgets of nitrogen-containing gases, *Nature*, 346, 552-554, 1990.
- 659 May, A. A., McMeeking, G. R., Lee, T., Taylor, J. W., Craven, J. S., Burling, I., Sullivan, A. P., Akagi, S., Collett,  
 660 J. L., Flynn, M., Coe, H., Urbanski, S. P., Seinfeld, J. H., Yokelson, R. J., and Kreidenweis, S. M.: Aerosol  
 661 emissions from prescribed fires in the United States: A synthesis of laboratory and aircraft measurements, *J.*  
 662 *Geophys. Res. Atmos.*, 119, 11826-11849, 2014.
- 663 McMahon, C. K., Wade, D. D., and Tsoukalas, S. N.: Combustion characteristics and emissions from burning  
 664 organic soils, in: *Proceedings, 73rd Annual Meeting of the Air Pollution Control Association, Air & Waste*  
 665 *Management Association*, Pittsburgh, PA, 1980.
- 666 Miettinen, J., Hooijer, A., Vernimmen, R., Liew, S. C., and Page, S. E.: From carbon sink to carbon source:  
 667 Extensive peat oxidation in insular Southeast Asia since 1990, *Environmental Research Letters*, 12, 2017.
- 668 Muraleedharan, T. R., Radojevic, M., Waugh, A., and Caruana, A.: Emissions from the combustion of peat: An  
 669 experimental study, *Atmos. Environ.*, 34, 3033-3035, 2000.
- 670 Na, K., Song, C., Switzer, C., and Cocker, D. R.: Effect of ammonia on secondary organic aerosol formation from  
 671 alpha-Pinene ozonolysis in dry and humid conditions, *Environ. Sci. Technol.*, 41, 6096-6102, 2007.
- 672 Nara, H., Tanimoto, H., Tohjima, Y., Mukai, H., Nojiri, Y., and Machida, T.: Emission factors of CO<sub>2</sub>, CO and CH<sub>4</sub>  
 673 from Sumatran peatland fires in 2013 based on shipboard measurements, *Tellus Series B-Chemical and Physical*  
 674 *Meteorology*, 69, 10.1080/16000889.2017.1399047, 2017.
- 675 Neff, J. C., Holland, E. A., Dentener, F. J., McDowell, W. H., and Russell, K. M.: The origin, composition and rates  
 676 of organic nitrogen deposition: a missing piece of the nitrogen cycle?, *Biogeochemistry*, 57/58, 99-136, 2002.
- 677 Ng, N. L., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Kroll, J. H., Kwan, A. J., McCabe, D. C., Wennberg, P. O.,  
 678 Sorooshian, A., Murphy, S. M., Dalleska, N. F., Flagan, R. C., and Seinfeld, J. H.: Effect of NO<sub>x</sub> level on secondary  
 679 organic aerosol (SOA) formation from the photooxidation of terpenes, *Atmos. Chem. Phys.*, 7, 5159-5174, 2007.
- 680 NIOSH: Method 5050, Elemental carbon (diesel particulate), in: *NIOSH Manual of Analytical Methods*, 4th ed. ed.,  
 681 *National Institute of Occupational Safety and Health*, Cincinnati, OH, 1999.
- 682 Ohlemiller, T. J., Bellan, J., and Rogers, F.: A model of smoldering combustion applied to flexible polyurethane  
 683 foams, *Combustion and Flame*, 36, 197-215, 1979.
- 684 Olszyna, K. J., Bailey, E. M., Simonaitis, R., and Meagher, J. F.: O<sub>3</sub> and NO<sub>y</sub> relationships at a rural site, *Journal of*  
 685 *Geophysical Research*, 99, 14557-14563, 1994.
- 686 Page, S. E., Siegert, F., Rieley, J. O., Boehm, H. D. V., Jaya, A., and Limin, S.: The amount of carbon released from  
 687 peat and forest fires in Indonesia during 1997, *Nature*, 420, 61-65, 10.1038/nature01131, 2002.
- 688 Page, S. E., Rieley, J. O., and Banks, C. J.: Global and regional importance of the tropical peatland carbon pool,  
 689 *Global Change Biology*, 17, 798-818, 2011.
- 690 Parrish, D. D., Hahn, C. J., Williams, E. J., Norton, E. B., and Fehsenfeld, F. C.: Indications of photochemical  
 691 histories of Pacific air masses from measurements of atmospheric trace species at Point Arena, California, *Journal of*  
 692 *Geophysical Research Letters*, 97, 15833-15901, 1992.



- 693 Pokhrel, R. P., Wagner, N. L., Langridge, J. M., Lack, D. A., Jayarathne, T., Stone, E. A., Stockwell, C. E.,  
 694 Yokelson, R. J., and Murphy, S. M.: Parameterization of single-scattering albedo (SSA) and absorption Angstrom  
 695 exponent (AAE) with EC /OC for aerosol emissions from biomass burning, *Atmos. Chem. Phys.*, 16, 9549-9561,  
 696 10.5194/acp-16-9549-2016, 2016.
- 697 Prenni, A. J., Levin, E. J. T., Benedict, K. B., Sullivan, A. P., Schurman, M. I., Gebhart, K. A., Day, D. E., Carrico,  
 698 C. M., Malm, W. C., Schichtel, B. A., Collett, J. L., and Kreidenweis, S. M.: Gas-phase reactive nitrogen near Grand  
 699 Teton National Park: Impacts of transport, anthropogenic emissions, and biomass burning, *Atmos. Environ.*, 89,  
 700 749-756, 2014.
- 701 Rein, G., Cohen, S., and Simeoni, A.: Carbon emissions from smouldering peat in shallow and strong fronts,  
 702 *Proceedings of the Combustion Institute*, 32, 2489-2496, 0.1016/j.proci.2008.07.008, 2009.
- 703 Roberts, J. M., Veres, P. R., Cochran, A. K., Warneke, C., Burling, I. R., Yokelson, R. J., Lerner, B., Gilman, J. B.,  
 704 Kuster, W. C., Fall, R., and de Gouw, J.: Isocyanic acid in the atmosphere and its possible link to smoke-related  
 705 health effects, *Proc. Natl. Acad. Sci. USA*, 108, 8966-8971, 2011.
- 706 Roulston, C., Paton-Walsh, C., Smith, T. E. L., Guerette, E. A., Evers, S., Yule, C. M., Rein, G., and Van der Werf,  
 707 G. R.: Fine particle emissions from tropical peat fires decrease rapidly with time since ignition, *J. Geophys. Res.*  
 708 *Atmos.*, 123, 5607-5617, 10.1029/2017jd027827, 2018.
- 709 Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John  
 710 Wiley & Sons, New York, NY, 1998.
- 711 Setyawati, W., Damanhuri, E., Lestari, P., and Dewi, K.: Emission factor from small scale tropical peat combustion,  
 712 in: *1st Annual Applied Science and Engineering Conference*, edited by: Abdullah, A. G., Nandiyanto, A. B. D., and  
 713 Danuwijaya, A. A., IOP Conference Series-Materials Science and Engineering, 2017.
- 714 Simoneit, B. R. T., Rushdi, A. I., Bin Abas, M. R., and Didyk, B. M.: Alkyl amides and nitriles as novel tracers for  
 715 biomass burning, *Environ. Sci. Technol.*, 37, 16-21, 2003.
- 716 Smith, T. E. L., Evers, S., Yule, C. M., and Gan, J. Y.: In situ tropical peatland fire emission factors and their  
 717 variability, as determined by field measurements in peninsula Malaysia, *Global Biogeochemical Cycles*, 32, 18-31,  
 718 10.1002/2017gb005709, 2018.
- 719 Stephens, M., Turner, N., and Sandberg, J.: Particle identification by laser-induced incandescence in a solid-state  
 720 laser cavity, *Appl. Opt.*, 42, 3726-3736, 10.1364/ao.42.003726, 2003.
- 721 Stockwell, C. E., Yokelson, R. J., Kreidenweis, S. M., Robinson, A. L., Demott, P. J., Sullivan, R. C., Reardon, J.,  
 722 Ryan, K. C., Griffith, D. W. T., and Stevens, L.: Trace gas emissions from combustion of peat, crop residue,  
 723 domestic biofuels, grasses, and other fuels: configuration and Fourier transform infrared (FTIR) component of the  
 724 fourth Fire Lab at Missoula Experiment (FLAME-4), *Atmos. Chem. Phys.*, 14, 9727-9754, 2014.
- 725 Stockwell, C. E., Veres, P. R., Williams, J., and Yokelson, R. J.: Characterization of biomass burning emissions  
 726 from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight  
 727 mass spectrometry, *Atmos. Chem. Phys.*, 15, 845-865, 2015.
- 728 Stockwell, C. E., Jayarathne, T., Cochrane, M. A., Ryan, K. C., Putra, E. I., Saharjo, B. H., Nurhayati, A. D., Albar,  
 729 I., Blake, D. R., Simpson, I. J., Stone, E. A., and Yokelson, R. J.: Field measurements of trace gases and aerosols  
 730 emitted by peat fires in Central Kalimantan, Indonesia, during the 2015 El Nino, *Atmos. Chem. Phys.*, 16, 11711-  
 731 11732, 10.5194/acp-16-11711-2016, 2016.
- 732 Tian, J., Chow, J. C., Cao, J. J., Han, Y. M., Ni, H. Y., Chen, L.-W. A., Wang, X. L., Huang, R. J., Moosmüller, H.,  
 733 and Watson, J. G.: A biomass combustion chamber: Design, evaluation, and a case study of wheat straw combustion  
 734 emission tests, *Aerosol Air Qual. Res.*, 15, 2104-2114, 2015.





- 735 Turetsky, M. R., Kane, E. S., Harden, J. W., Ottmar, R. D., Manies, K. L., Hoy, E., and Kasischke, E. S.: Recent  
 736 acceleration of biomass burning and carbon losses in Alaskan forests and peatlands, *Nature Geoscience*, 4, 27-31,  
 737 2010.
- 738 Turetsky, M. R., Benscoter, B., Page, S., Rein, G., van der Werf, G. R., and Watts, A.: Global vulnerability of  
 739 peatlands to fire and carbon loss, *Nature Geoscience*, 8, 11-14, 10.1038/ngeo2325, 2015a.
- 740 Turetsky, M. R., Benscoter, B., Page, S. E., Rein, G., van der Werf, G. R., and Watts, A. C.: Controls on global peat  
 741 fires and consequences for the carbon cycle, *Nature*, 2015b.
- 742 Updyke, K. M., Nguyen, T. B., and Nizkorodov, S. A.: Formation of brown carbon via reactions of ammonia with  
 743 secondary organic aerosols from biogenic and anthropogenic precursors, *Atmos. Environ.*, 63, 22-31, 2012.
- 744 VDI: Measurement of soot (ambient air) - Thermographic determination of elemental carbon after thermal  
 745 desorption of organic carbon, Verein Deutscher Ingenieure, Dusseldorf, Germany, 1999.
- 746 Villena, G., Bejan, I., Kurtenbach, R., Wiesen, P., and Kleffmann, J.: Interferences of commercial NO<sub>2</sub> instruments  
 747 in the urban atmosphere and in a smog chamber, *Atmos. Meas. Tech.*, 5, 149-159, 2012.
- 748 Wang, X. L., Chancellor, G., Evenstad, J., Farnsworth, J. E., Hase, A., Olson, G. M., Sreenath, A., and Agarwal, J.  
 749 K.: A novel optical instrument for estimating size segregated aerosol mass concentration in real time, *Aerosol Sci.*  
 750 *Technol.*, 43, 939-950, 2009.
- 751 Wang, X. L., Watson, J. G., Chow, J. C., Gronstal, S., and Kohl, S. D.: An efficient multipollutant system for  
 752 measuring real-world emissions from stationary and mobile sources, *Aerosol Air Qual. Res.*, 12, 145-160, 2012.
- 753 Ward, D. E., and Hardy, C. C.: Advances in the characterization and control of emissions from prescribed fires, 77th  
 754 Annual Meeting of the Air pollution Control Association, San Francisco, CA, 1984.
- 755 Ward, D. E., and Radke, L. F.: Emissions measurements from vegetation fires: A comparative evaluation of methods  
 756 and results, *Fire in the Environment: The Ecological, Atmospheric and Climatic Importance of Vegetation Fires*, 13,  
 757 53-76, 1993.
- 758 Watson, J. G., Chow, J. C., and Frazier, C. A.: X-ray fluorescence analysis of ambient air samples, in: *Elemental*  
 759 *Analysis of Airborne Particles*, Vol. 1, edited by: Landsberger, S., and Creatchman, M., *Advances in Environmental,*  
 760 *Industrial and Process Control Technologies*, Gordon and Breach Science, Amsterdam, The Netherlands, 67-96,  
 761 1999.
- 762 Watson, J. G., Tropp, R. J., Kohl, S. D., Wang, X. L., and Chow, J. C.: Filter processing and gravimetric analysis for  
 763 suspended particulate matter samples, *Aerosol Science and Engineering*, 1, 193-205, 2017.
- 764 Watts, A. C.: Organic soil combustion in cypress swamps: Moisture effects and landscape implications for carbon  
 765 release, *Forest Ecology and Management*, 294, 178-187, 10.1016/j.foreco.2012.07.032, 2013.
- 766 Wilson, D., Dixon, S. D., Artz, R. R. E., Smith, T. E. L., Evans, C. D., Owen, H. J. F., Archer, E., and Renou-  
 767 Wilson, F.: Derivation of greenhouse gas emission factors for peatlands managed for extraction in the Republic of  
 768 Ireland and the United Kingdom, *Biogeosciences*, 12, 5291-5308, 2015.
- 769 Winer, A. M., Peters, J. W., Smith, J. P., and Pitts, J. N., Jr.: Response of commercial chemiluminescence NO-NO<sub>2</sub>  
 770 analyzers to other nitrogen-containing compounds, *Environ. Sci. Technol.*, 8, 1118-1121, 1974.
- 771 Wooster, M. J., Gaveau, D. L. A., Salim, M. A., Zhang, T. R., Xu, W. D., Green, D. C., Huijnen, V., Murdiyarso,  
 772 D., Gunawan, D., Borchard, N., Schirrmann, M., Main, B., and Sepriando, A.: New tropical peatland gas and  
 773 particulate emissions factors indicate 2015 Indonesian fires released far more particulate matter (but less methane)  
 774 than current inventories imply, *Remote Sensing*, 10, 10.3390/rs10040495, 2018.



- 775 Yatawelli, R. L. N., Chen, L.-W. A., Knue, J., Samburova, V., Gyawali, M., Watts, A. C., Chakrabarty, R. K.,  
776 Moosmuller, H., Hodzic, A., Wang, X. L., Zielinska, B., Chow, J. C., and Watson, J. G.: Emissions and partitioning  
777 of intermediate-volatility and semi-volatile polar organic compounds (I/SV-POCs) during laboratory combustion of  
778 boreal and sub-tropical peat, *Aerosol Science and Engineering*, 1, 25-32, 2017.
- 779 Yokelson, R. J., Susott, R., Ward, D. E., Reardon, J., and Griffith, D. W. T.: Emissions from smoldering combustion  
780 of biomass measured by open- path Fourier transform infrared spectroscopy, *Journal of Geophysical Research*, 102,  
781 18865-18877, 1997.
- 782 Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., de Gouw, J., Akagi, S. K., Urbanski, S.  
783 P., Veres, P., Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, T. J., Hosseini, S., Miller, J.  
784 W., Cocker, D. R., Jung, H., and Weise, D. R.: Coupling field and laboratory measurements to estimate the emission  
785 factors of identified and unidentified trace gases for prescribed fires, *Atmos. Chem. Phys*, 13, 89-116, 2013.
- 786 Yu, Z. C., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last  
787 Glacial Maximum, *Geophysical Research Letters*, 37, 10.1029/2010gl043584, 2010.
- 788 Zhao, Y. L., Kreisberg, N. M., Worton, D. R., Isaacman, G., Weber, R. J., Liu, S., Day, D. A., Russell, L. M.,  
789 Markovic, M. Z., VandenBoer, T. C., Murphy, J. G., Hering, S. V., and Goldstein, A. H.: Insights into secondary  
790 organic aerosol formation mechanisms from measured gas/particle partitioning of specific organic tracer  
791 compounds, *Environ. Sci. Technol.*, 47, 3781-3787, 2013.
- 792



Table 1. Average peat composition<sup>a</sup> (dry weight percentage) for total carbon (C), hydrogen (H), nitrogen (N), sulfur (S), and oxygen (O).

Peat Location	C (%)	H (%)	N (%)	S (%)	O (%)	N/C mass ratio	Sum (CHNSO; %)
Odintsovo, Russia	44.20 ± 1.01	6.43 ± 0.16	1.50 ± 0.52	<0.01	38.64 ± 0.78	0.034	90.8
Pskov, Siberia	52.03 ± 0.23	6.30 ± 0.05	2.92 ± 0.12	<0.01	36.83 ± 0.39	0.056	98.1
Northern Alaska, USA	50.94 ± 0.81	6.05 ± 0.07	1.79 ± 0.09	<0.01	36.62 ± 0.30	0.035	95.4
Putnam County Lakebed, Florida, USA	56.64 ± 0.37	6.25 ± 0.40	3.53 ± 0.05	<0.01	31.43 ± 0.36	0.062	97.8
Everglades, Florida, USA	47.22 ± 0.57	5.15 ± 0.16	3.93 ± 0.08	<0.01	34.18 ± 0.87	0.083	90.5
Caohai, Guizhou, Southeast China	19.74 ± 2.01	2.09 ± 1.26	1.35 ± 0.16	<0.01	23.95 ± 1.15	0.068	47.1
Gaopo, Guizhou, Southeast China	29.70 ± 2.09	3.13 ± 0.16	2.08 ± 0.22	<0.01	21.46 ± 1.27	0.070	56.4
Borneo, Malaysia	50.55 ± 2.53	6.46 ± 0.99	1.16 ± 0.08	<0.01	33.72 ± 0.30	0.023	91.9

<sup>a</sup>Elemental analyses were performed using Elemental Analyzer (Flash EA1112 CHNS/O Analyzer, Thermo Fisher Scientific, Waltham, MA, USA). Each dried peat sample (~2–3 g) was submitted for combustion analysis at 900°C for C, H, N, and S in a helium/oxygen atmosphere and at 1060°C for O in a helium atmosphere. Three to four replicate sample analyses were conducted for each type of peat to obtain the average and standard deviations.



Table 2. Peat combustion emission factors (EFs) for CO<sub>2</sub>, CO, and CH<sub>4</sub>.<sup>a</sup>

Sampling Location or Review (Reference)	Sampling Method (No. of samples) <sup>b</sup>	Modified Combustion Efficiency (MCE)	Measurement Method	Average Emission Factors (g/kg)			Ratio (EF <sub>CO</sub> /EF <sub>CO2</sub> )
				EF <sub>CO2</sub>	EF <sub>CO</sub>	EF <sub>CH4</sub>	
<b>Boreal</b>							
Odintsovo, Russia (This study)	Lab (n=6, 25 % FM <sup>c</sup> )	0.81 ± 0.03	CO/CO <sub>2</sub> monitors and FTIR <sup>d</sup>	1073 ± 63	157 ± 24	3.20 ± 0.69	0.15
Pskov, Siberia (This study)	Lab (n=7, 25 % FM <sup>c</sup> )	0.85 ± 0.01	CO/CO <sub>2</sub> monitors and FTIR <sup>d</sup>	1380 ± 27	159 ± 14	6.94 ± 1.48	0.12
Western Siberia (Chakrabarty et al., 2016)	Lab (n=1, 25 % FM <sup>c</sup> ) (n=1, 50 % FM <sup>c</sup> )	Smoldering	CO/CO <sub>2</sub> monitors	1432 1698	204 49	NA	0.14 0.029
<b>Temperate</b>							
Northern Alaska, USA (This study)	Lab (n=6, 25 % FM <sup>c</sup> )	0.86 ± 0.03	CO/CO <sub>2</sub> monitors and FTIR <sup>d</sup>	1400 ± 38	161 ± 19	5.69 ± 1.07	0.12
Northern Alaska, USA (Chakrabarty et al., 2016)	Lab (n=1, 25 % FM <sup>c</sup> ) (n=1, 50 % FM <sup>c</sup> )	Smoldering	CO/CO <sub>2</sub> monitors	1238 1598	83 128	NA	0.067 0.08
Hudson Bay lowland, Ontario, Canada	Lab	0.81 ± 0.009	FTIR	1274 ± 19	197 ± 9	6.25 ± 2.17	0.15
(Stockwell et al., 2014)	Lab	0.81 ± 0.327	FTIR	1395 ± 52 <sup>e</sup>	209 ± 68 <sup>e</sup>	6.85 ± 5.66 <sup>e</sup>	0.15
Alaska and Minnesota, USA (Yokelson et al., 1997)	Lab	Smoldering	Infrared system	420 ± 134	170 ± 33	NA	0.40
Edinburg, Scotland, UK (Rein et al., 2009)	Lab	0.84 ± 0.019	FTIR	1346 ± 31	218 ± 22	8.35 ± 1.3	0.16
Sphagnum moss peat, Ireland (Wilson et al., 2015)	Lab (n=5)						
<b>Subtropical</b>							
Putnam County Lakebed, FL, USA (This study)	Lab (n=6, 25 % FM <sup>c</sup> ) (n=3, 60 % FM <sup>c</sup> )	0.65 ± 0.04 0.72 ± 0.01	CO/CO <sub>2</sub> monitors and FTIR <sup>d</sup>	1126 ± 89 1262 ± 27	394 ± 46 315 ± 10	10.42 ± 1.81 9.18 ± 0.26	0.35 0.25
Everglades National Park, FL, USA	Lab	0.90 ± 0.03	CO/CO <sub>2</sub> monitors	1292 ± 80	93 ± 21	7.65 ± 1.36	0.07



(This study)	(n=3, 25 % FM <sup>c</sup> )	(mix of flaming and smoldering)	and FTIR <sup>d</sup>		
Pocosin Lake NWR <sup>f</sup> , NC, USA (Geron and Hays, 2013)	Field (Feb & Aug 2008) (n=3)	0.77–0.83	CO and Infrared gas monitoring	230–300	NA
Green Swamp Preserve, NC, USA (Geron and Hays, 2013))	Field (Feb 2009) (n=8)	0.80–0.81	CO and Infrared gas monitoring	10–280	NA
Alligator River (AR) NWR <sup>f</sup> , NC, USA (Geron and Hays, 2013)	Field (May 2011) (n=8)	0.79–0.86	CO and Infrared gas monitoring	125–290	NA
Pocosin Lake NWR <sup>f</sup> , NC, USA (Black et al., 2016)	Lab (n=2)	0.83 ± 0.02	CO/CO <sub>2</sub> monitors	122 ± 14	0.13
Alligator River NWR <sup>f</sup> , NC, USA (Black et al., 2016)	Lab (n=2)	0.86 ± 0.02	CO/CO <sub>2</sub> monitors	108 ± 20	0.13
<b>Tropical</b>					
Borneo, Malaysia (This study)	Lab (n=6, 25 % FM <sup>c</sup> )	0.85 ± 0.02	CO/CO <sub>2</sub> monitors and FTIR <sup>d</sup>	171 ± 22	0.13
Peninsula, Malaysia (Smith et al., 2018)	Field (Aug 2015) (n=10)	0.80 ± 0.03	FTIR	251 ± 39	0.16
Central Kalimantan, Indonesia (Wooster et al., 2018)	Field (Sep/Oct 2015) (n=23)	0.81 ± 0.032	Cavity-enhanced laser absorption spectrometer and FTIR	279 ± 44	0.16
Central Kalimantan, Indonesia <sup>j</sup> (Stockwell et al., 2016)	Field (Oct/Nov 2015) (n=35)	0.77 ± 0.053	FTIR	291 ± 49	0.19
Central Kalimantan, Indonesia (Huijnen et al., 2016)	Field (Oct 2015)	0.8	Cavity-ring down spectrometer	255 ± 39	0.16
West Kalimantan, Indonesia	Lab	Flaming (0.998 ± 0.005)	CO/CO <sub>2</sub> monitors	3.10 ± 7.17	0.0015



(Setyawati et al., 2017)	(n=17 each)	Smoldering (0.89 ± 0.06)	and gas chromatography	1831 ± 131	138 ± 72	17 ± 1.2	0.075
South Kalimantan, Indonesia (Stockwell et al., 2014)	Lab (n=3)	0.82 ± 0.065	FTIR	1637 ± 204	233 ± 72	12.8 ± 6.61	0.14
South Sumatra, Indonesia (Christian et al., 2003)	Lab (n=1)	0.84	FTIR	1703	210	20.8	0.12
North-Central Sumatra, Indonesia (Nara et al., 2017)	Shipboard (June-Aug 2013) (n=5)	0.84	Infrared and cavity ring-down spectrometer	1663 ± 54	205 ± 23	7.6 ± 1.6	0.12
<b>Reviews<sup>a</sup></b>							
Atmospheric Modeling (Akagi et al., 2011)	NA	NA	NA	1563 ± 56	182 ± 60	11.8 ± 7.8	0.12
Boreal/Temperate Tropical (IPCC, 2014)	NA	NA	NA	362 ± 41 <sup>h</sup>	207 ± 70 <sup>h</sup>	9 ± 4 <sup>h</sup>	NA
Boreal/Temperate Tropical (Hu et al., 2018)	NA	NA	NA	1703 <sup>i</sup>	210 <sup>i</sup>	21 <sup>i</sup>	NA
Peat Fire (Andreae, 2019)	NA	NA	NA	1134 ± 139	179 ± 61	8.1 ± 4.1	0.16
		Smoldering		1615 ± 184	248 ± 50	12.3 ± 5.0	0.40
				1550 ± 130	250 ± 23	9.3 ± 1.5	0.45

<sup>a</sup>Data acquired from this study are highlighted in green  
<sup>b</sup>Only included number of samples reported  
<sup>c</sup>FM: Fuel moisture content  
<sup>d</sup>FTIR: Fourier transform infrared spectroscopy. CH<sub>4</sub> was acquired by FTIR in this study  
<sup>e</sup>Obtained from Stockwell et al. (2014) as only the ratios of moles compound/total moles carbon detected was reported in Yokelson et al. (1997)  
<sup>f</sup>NWR: National Wildlife Reserve  
<sup>g</sup>Reviews for atmospheric modeling and emission inventory development  
<sup>h</sup>From Ward and Hardy (1984); Yokelson et al. (1997;2013)  
<sup>i</sup>From Christian et al. (2003) for tropical peats  
<sup>j</sup>Detailed volatile organic gas emission factors for one of these samples are reported by Koss et al. (2018)

Table 3. Peat combustion emission factors (EFs) for gaseous nitrogen species<sup>a</sup>.

Sampling Location (Reference)	No. of samples	Average Emission Factors (g/kg)							Percent NO <sub>x</sub> /NO <sub>y</sub>
		EF <sub>NH<sub>3</sub></sub> <sup>b</sup>	EF <sub>HCN</sub> <sup>b</sup>	EF <sub>NO</sub> <sup>c</sup>	EF <sub>NO<sub>2</sub></sub> <sup>c</sup>	EF <sub>NO<sub>2</sub></sub> <sup>d</sup> (as NO <sub>x</sub> )	EF <sub>NO</sub> <sup>b</sup>	EF <sub>HONO</sub>	
<b>Boreal</b>									
Odintsovo, Russia (This study)	6	0.99 ± 0.47	2.45 ± 0.43	0.34 ± 0.04	0.48 ± 0.11	1.01 ± 0.14	1.06 ± 0.11	1.64 ± 0.32	NA
Pskov, Siberia (This study)	7	4.65 ± 1.38	5.00 ± 0.74	0.84 ± 0.12	0.42 ± 0.03	1.70 ± 0.20	2.22 ± 0.27	2.29 ± 0.29	NA
Pskov, Siberia (Bhattarai et al., 2018)	3	NA	NA	NA	NA	0.08 ± 0.04 <sup>e</sup>	NA	NA	NA
<b>Temperate</b>									
Northern Alaska, USA (This study)	6	2.7 ± 0.62	2.33 ± 0.22	0.84 ± 0.44	0.37 ± 0.13	1.67 ± 0.76	2.10 ± 0.85	1.57 ± 0.16	NA
Hudson Bay lowland, Ontario, Canada (Stockwell et al., 2014)	NA	2.21 ± 0.24	1.77 ± 0.55	NA	NA	NA	NA	0.18	NA
Alaska and Minnesota, USA (Yokelson et al., 1997)	NA	8.76 ± 13.76	5.09 ± 5.64	NA	NA	NA	NA	NA	NA
Sphagnum moss peat, Ireland (Wilson et al., 2015)	5	2.20 ± 0.35	0.73 ± 0.50	NA	NA	NA	NA	NA	NA
Coastal Swamp land, NC, USA (Stockwell et al., 2014)	NA	1.87 ± 0.37	4.45 ± 3.02	NA	NA	NA	NA	8.48 ± 0.05	NA
<b>Subtropical</b>									
Putnam County Lakebed, FL, USA (This study)	6 (25 % FM) 3 (60 % FM)	3.2 ± 0.26 3.3 ± 0.05 11.9 ± 2.01	11.5 ± 2.3 11.7 ± 0.3 5.12 ± 1.60	1.01 ± 0.33 0.71 ± 0.07 1.78 ± 0.31	0.35 ± 0.28 0.65 ± 0.05 0.83 ± 0.16	2.01 ± 0.68 1.74 ± 0.15 3.56 ± 0.58	2.91 ± 0.34 2.39 ± 0.19 4.33 ± 1.10	3.57 ± 0.63 3.89 ± 0.01 1.46 ± 0.28	NA NA NA
Putnam County Lakebed, FL, USA (Bhattarai et al., 2018)	6	NA	NA	NA	NA	0.11 ± 0.05 <sup>e</sup>	NA	NA	73 ± 5 %
<b>Tropical</b>									
Borneo, Malaysia (This study)	6	3.66 ± 0.27	2.84 ± 0.44	0.26 ± 0.04	0.35 ± 0.05	0.75 ± 0.10	1.07 ± 0.56	1.88 ± 0.19	NA
Peninsula, Malaysia (Smith et al., 2018)	35	7.82 ± 4.37	3.79 ± 1.97	NA	NA	NA	NA	NA	NA
Central Kalimantan, Indonesia (Stockwell et al., 2016)	3	2.86 ± 1.00	5.75 ± 1.60	0.31 ± 0.36	NA	NA	NA	0.208 ± 0.059	NA
South Kalimantan, Indonesia (Stockwell et al., 2014)	NA	3.38 ± 3.02	3.66 ± 2.43	0.51 ± 0.12	2.31 ± 1.46	NA	NA	NA	NA
<b>Reviews<sup>g</sup></b>									
Atmospheric Modeling (Akagi et al., 2011)	NA	10.8 ± 12.4	5.0 ± 4.93	NA	NA	1.23 ± 0.87 <sup>f</sup>	NA	NA	NA
Smoldering Boreal/Temperate Smoldering Tropical (Hu et al., 2018)	3	3.39 ± 6.89 8.0 ± 3.04	3.38 ± 3.21 5.24 ± 1.55	NA	2.31 ± 1.46 2.36 ± 0.03	NA	NA	NA	NA
Peat Fire (Andreae, 2019)	3	4.2 ± 3.2	4.4 ± 1.2	NA	1.84 <sup>f</sup> (± 0.48 to 3.4)	NA	NA	NA	NA

<sup>a</sup>Data acquired from this study is highlighted in green  
<sup>b</sup>Data acquired from Fourier Transform Infrared (FTIR) spectroscopy for this study



<sup>c</sup>Data acquired from the NO<sub>x</sub> instrument upstream of the oxidation flow reactor for this study

<sup>d</sup>Data acquired from the NO<sub>y</sub> instrument for this study

<sup>e</sup>Reported as NO<sub>x</sub>

<sup>f</sup>The reported NO<sub>x</sub> as NO was converted to NO<sub>x</sub> as NO<sub>2</sub> for comparison

<sup>g</sup>Reviews for atmospheric modeling and emission inventory development



Table 4. Peat combustion emission factors (EFs) for PM<sub>2.5</sub> mass and carbon<sup>a</sup>.

Sampling Location (Reference)	Sampling Method (No. of samples)	Modified Combustion Efficiency (MCE)	Carbon Analysis Method <sup>b</sup>	Average Emission Factor (g/kg)			Ratio (EF <sub>TC</sub> /EF <sub>PM</sub> )
				EF <sub>PM<sub>2.5</sub></sub> <sup>c</sup> (PM size)	EF <sub>OC</sub>	EF <sub>EC</sub>	
<b>Boreal</b>							
Odintsovo, Russia (This study) <sup>a</sup>	Lab (n=6, 25% FM) <sup>d</sup>	0.81 ± 0.03	IMPROVE_A	42.6 ± 5.2 (Fresh) <sup>e</sup>	25.1 ± 3.3 (Fresh) <sup>e</sup>	0.77 ± 0.38 (Fresh) <sup>e</sup>	0.61 ± 0.05
				40.5 ± 7.2 (Aged) <sup>e</sup>	17.2 ± 2.7 (Aged) <sup>e</sup>	0.69 ± 0.19 (Aged) <sup>e</sup>	0.45 ± 0.07
Siberia (This study) <sup>a</sup>	Lab (n=7, 25% FM) <sup>d</sup>	0.85 ± 0.01	IMPROVE_A	33.9 ± 6.3 (Fresh) <sup>e</sup>	26.0 ± 3.4 (Fresh) <sup>e</sup>	0.69 ± 0.58 (Fresh) <sup>e</sup>	0.80 ± 0.08
				30.7 ± 10.2 (Aged) <sup>e</sup>	18.1 ± 4.5 (Aged) <sup>e</sup>	0.78 ± 0.31 (Aged) <sup>e</sup>	0.64 ± 0.13
Pskov, Siberia (Bhattarai et al., 2018)	Lab (n=3)	NA	IMPROVE_A	7.98 ± 1.58	6.52 ± 1.4	0.02 ± 0.01	0.82
Western Siberia (Chakrabarty et al., 2016)	Lab (n=1, 25% FM) <sup>d</sup> (n=1, 50% FM) <sup>d</sup>	<0.7	IMPROVE_A	NA	17	0.2	NA
					11	0.1	
Neustädter Moor, Northern Germany (Iinuma et al., 2007)	Lab	0.84	VDI	44 (PM <sub>10</sub> ) <sup>g</sup>	12.8	0.96	0.31
<b>Temperate</b>							
Northern Alaska, USA (This study) <sup>a</sup>	Lab (n=6, 25% FM) <sup>d</sup>	0.85 ± 0.02	IMPROVE_A	24.0 ± 7.6 (Fresh) <sup>e</sup>	17.4 ± 4.1 (Fresh) <sup>e</sup>	0.60 ± 0.24 (Fresh) <sup>e</sup>	0.77 ± 0.12
				24.8 ± 5.3 (Aged) <sup>e</sup>	14.9 ± 3.9 (Aged) <sup>e</sup>	0.55 ± 0.42 (Aged) <sup>e</sup>	0.63 ± 0.16
Interior Alaska, USA (Chakrabarty et al., 2016)	Lab (n=1, 25% FM) <sup>d</sup> (n=1, 50% FM) <sup>d</sup>	0.7 0.7	IMPROVE_A	NA	7 4	0.1 0.2	NA
<b>Subtropical</b>							
Putnam County Lakebed, FL, USA (This study) <sup>a</sup>	Lab (n=4, 25% FM) <sup>d</sup> Lab (n=2, 25% FM) <sup>d</sup> Lab (n=3, 60% FM) <sup>d</sup>	0.65 ± 0.04 0.67 ± 0.02 0.72 ± 0.01	IMPROVE_A	53.1 ± 6.8 (Fresh) <sup>e</sup>	36.6 ± 1.9 (Fresh) <sup>e</sup>	1.33 ± 0.60 (Fresh) <sup>e</sup>	0.72 ± 0.08
				53.9 ± 8.3 (Aged) <sup>e</sup>	37.3 ± 6.7 (Aged) <sup>e</sup>	0.95 ± 0.07 (Aged) <sup>e</sup>	0.71 ± 0.04
				51.6 ± 7.9 (Fresh 2) <sup>f</sup>	36.6 ± 1.8 (Fresh 2) <sup>f</sup>	1.8 ± 0.61 (Fresh 2) <sup>f</sup>	0.85 ± 0.04
				48.2 ± 8.4 (Aged 2) <sup>f</sup>	34.0 ± 8.3 (Aged 2) <sup>f</sup>	0.99 ± 0.15 (Aged 2) <sup>f</sup>	0.66 ± 0.10
Everglades National Park, FL, USA (This study) <sup>a</sup>	Lab (n=3, 60% FM) <sup>d</sup> Lab (n=7, 25% FM) <sup>d</sup>	0.72 ± 0.01 0.90 ± 0.03	IMPROVE_A	35.9 ± 4.3 (Fresh 2) <sup>f</sup>	29.3 ± 2.2 (Fresh 2) <sup>f</sup>	1.00 ± 0.07 (Fresh 2) <sup>f</sup>	0.75 ± 0.11
				34.7 ± 2.6 (Aged 2) <sup>f</sup>	22.1 ± 2.3 (Aged 2) <sup>f</sup>	0.85 ± 0.85 (Aged 2) <sup>f</sup>	0.72 ± 0.05
				23.6 ± 5.1 (Fresh) <sup>e</sup>	19.0 ± 4.4 (Fresh) <sup>e</sup>	0.78 ± 0.45 (Fresh) <sup>e</sup>	0.85 ± 0.15
				33.5 ± 11.4 (Aged) <sup>e</sup>	18.8 ± 5.2 (Aged) <sup>e</sup>	0.67 ± 0.30 (Aged) <sup>e</sup>	0.60 ± 0.12



Pocosin Lakes NWR <sup>a</sup> , NC, USA (Geron and Hays, 2013)	Field (n=3) (Feb & Aug 2008)	0.77-0.83	NA	34-55	NA	NA	NA
Green Swamp Preserve, NC, USA (Geron and Hays, 2013)	Field (n=8) (Feb 2009)	0.80-0.81	NA	44-53	NA	NA	NA
Alligator River NWR <sup>b</sup> , NC, USA (Geron and Hays, 2013)	Field (n=8) (May 2011)	0.79-0.86	NA	48-79	NA	NA	NA
Pocosin Lakes NWR <sup>b</sup> , NC, USA (Black et al., 2016)	Lab (n=2)	0.83 ± 1.02	NIOSH	5.9 ± 6.7	4.3 ± 4.1	0.082 ± 0.091	0.74
Alligator River NWR <sup>b</sup> , NC, USA (Black et al., 2016)	Lab (n=2)	0.86 ± 0.02	NIOSH	7.1 ± 5.6	6.3 ± 4.1	0.052 ± 0.057	0.89
Putnam County Lakebed, FL, USA (Bhattarai et al., 2018)	Lab (n=3)	NA	IMPROVE_A	6.89 ± 1.28	6.56 ± 1.10	0.04 ± 0.02	0.96
<b>Tropical</b>							
Borneo, Malaysia (This study) <sup>a</sup>	Lab (n=4, 25% FM) <sup>d</sup>	0.83 ± 0.03	IMPROVE_A	22.6 ± 3.1 (Fresh) <sup>e</sup> 22.6 ± 5.0 (Aged) <sup>e</sup>	18.0 ± 2.0 (Fresh) <sup>e</sup> 14.4 ± 1.7 (Aged) <sup>e</sup>	0.28 ± 0.11 (Fresh) <sup>e</sup> 0.29 ± 0.20 (Aged) <sup>e</sup>	0.81 ± 0.02 0.68 ± 0.16
Borneo, Malaysia (Bhattarai et al., 2018)	Lab (n=1)	NA	IMPROVE_A	3.9	9.62	0.1	2.4
Selangor, Malaysia (Roulston et al., 2018)	Field (n=6) (Jul/Aug 2016)	0.8-0.85	NA	28.0 ± 18.0	NA	NA	NA
Sumatra, Indonesia (Christian et al., 2003)	Lab (n=1)	Smoldering	Unspecified	NA	6.02	0.04	NA
Southern Sumatra, Indonesia (Iinuma et al., 2007)	Lab	Smoldering	VDI	33.0 (PM <sub>10</sub> ) <sup>g</sup>	8	0.57	0.26
Riau, Indonesia (Kuwata et al., 2018)	Field (June 2013) Field (Feb-Mar 2014)	NA	NA	13.0 ± 2.0 (PM <sub>10</sub> ) 19.0 ± 2.0 (PM <sub>10</sub> )	NA	NA	NA
Central Kalimantan, Indonesia (Wooster et al., 2018)	Field (n=23) (Sep/Oct 2015)	0.81 ± 0.032	NA	17.82 ± 6.86	NA	0.106 ± 0.043 (BC) <sup>y</sup>	NA
Central Kalimantan, Indonesia (Jayarathne et al., 2018)	Field (n=21) (Oct/Nov 2015)	0.78 ± 0.04	NIOSH	17.3 ± 6.0	12.4 ± 5.4	0.24 ± 0.1	0.73
Indonesia (location not specified) (May et al., 2014)	Lab	0.89	TOF-AMS and SP2	34.9 (PM <sub>1</sub> ) <sup>k</sup>	34.5 (OA) <sup>k</sup>	0.01 (BC) <sup>k</sup>	0.99



Reviews <sup>1</sup>						
Peatlands from tropical forest (Akagi et al., 2011)	NA	NA	NA	6.23 ± 3.6	0.2 ± 0.11	NA
Smoldering Boreal/Temperate Smoldering Tropical (Hu et al., 2018)	NA	NA	NA	8.38 ± 4.14 8.8 ± 4.24	0.36 ± 0.28 0.28 ± 0.18	0.46 0.52
Peat fires (Andreae, 2019)	NA	NA	NA	12.4	0.19	0.73

<sup>a</sup>Data acquired from this study are highlighted in green  
<sup>b</sup>The IMPROVE\_A protocol reports OC and EC by Thermal/Optical reflectance (TOR, Chow et al., 2007); The NIOSH and NIOSH5040 reports OC and EC by Thermal/Optical transmittance (NIOSH, 1999); VDI is German Industrial Standard (VDI, 1999); TOF-MS: time-of-flight mass spectrometer (Drewnack et al., 2005); and Single Particle Soot Photometer (SP2, DMT Inc., Boulder, CO, USA) measures black carbon (BC) by laser-induced incandescence technique (Stephens et al., 2003).  
<sup>c</sup>Size fraction is PM<sub>2.5</sub> except where otherwise noted.  
<sup>d</sup>FM; Fuel Moisture  
<sup>e</sup>Includes averages of all fresh and all aged emission factors (EFs) for the 25% fuel moisture (i.e., grouped Fresh 2 and Fresh 7 vs Aged 2 and Aged 7 shown in Table S7)  
<sup>f</sup>Comparison between 25% and 60% fuel moisture content are only made with Fresh 2 vs. Aged 2 of Putnam (FL) peats.  
<sup>g</sup>Sum of five stages of Berner Impactor with 0.05-0.14, 0.14-0.42, 0.42-1.2, 1.2-3.5, and 3.5-10 µm size ranges.  
<sup>h</sup>National Wildlife Refuge, eastern NC  
<sup>i</sup>From Jayathne et al. (2018)  
<sup>j</sup>BC by MicroAethalometer (AE 51) (Cheng et al., 2013; Wooster et al., 2018)  
<sup>k</sup>PM<sub>1</sub> and organic aerosol (OA) acquired from Time-of-Flight Mass Spectrometry (TOF-MS) measurements (Drewnack et al., 2005)  
<sup>l</sup>Reviews for atmospheric modeling and emission inventory development.




Figure 1. Global distribution of peatlands (Yu et al., 2010). Samples were obtained from Odintsovo, Russia; Pskov, Siberia; black spruce forest, Northern Alaska, USA; Putnam County Lakebed and Everglades National Park, Florida, USA; Caohai and Gaopo, Guizhou, China; and Borneo, Malaysia.




Figure 2. Configuration for peat combustion experimental set up. (FTIR: Fourier Transform Infrared Spectrometer; OFR: oxidation flow reactor; OFR lamps were operated at 2 and 3.5 volts to simulate aging of ~2 and 6.79 days, respectively).




Figure 3. Average carbonaceous species abundances in total emitted carbon (the sum of carbon in CO<sub>2</sub>, CO, CH<sub>4</sub>, VOCs, and PM<sub>2.5</sub> total carbon (TC = OC + EC). Numbers on top of the bars are average Modified Combustion Efficiencies (MCE) and the number of samples in each average. The Carbon Compounds include hydrogen cyanide (HCN), formaldehyde (CH<sub>2</sub>O), methanol (CH<sub>3</sub>OH), formic acid (HCOOH), carbonyl sulfide (COS), ethylene (C<sub>2</sub>H<sub>4</sub>), ethane (C<sub>2</sub>H<sub>6</sub>), acetaldehyde (C<sub>2</sub>H<sub>4</sub>O), ethanol (C<sub>2</sub>H<sub>5</sub>OH), acetic acid (CH<sub>3</sub>COOH), propane (C<sub>3</sub>H<sub>8</sub>), acrolein (C<sub>3</sub>H<sub>4</sub>O), acetone (C<sub>3</sub>H<sub>6</sub>O), 3-butadiene (C<sub>4</sub>H<sub>6</sub>), benzene (C<sub>6</sub>H<sub>6</sub>), hexane (C<sub>6</sub>H<sub>14</sub>), phenol (C<sub>6</sub>H<sub>5</sub>OH), and chlorobenzene (C<sub>6</sub>H<sub>5</sub>Cl) acquired by Fourier Transfer Infrared Spectrometry.




Figure 4. Ratio of emitted over consumed nitrogen for each type of peat (emitted nitrogen is the sum of nitrogen in HCN, NH<sub>3</sub>, NO, NO<sub>2</sub>, and NO<sub>x</sub> [NO<sub>y</sub>-NO<sub>x</sub>], N<sub>2</sub>O, HNO<sub>3</sub>, and PM<sub>2.5</sub> ions [NO<sub>2</sub><sup>-</sup> + NO<sub>3</sub><sup>-</sup> + NH<sub>4</sub><sup>+</sup>]; and the consumed nitrogen is the product of percent fuel nitrogen content and mass of fuel burned).