

Is Positive Correlation between Cloud Droplet Effective Radius and Aerosol Optical Depth over Land Due to Retrieval Artifacts or Real Physical Processes?

Hailing Jia^{1,2}, Xiaoyan Ma^{1,2}, Johannes Quaas³, Yan Yin^{1,2}, and Tom Qiu⁴

5 ¹Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science & Technology, Nanjing 210044, China

²Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, School of Atmospheric Physics, Nanjing University of Information Science & Technology, Nanjing 210044, China

³Institute for Meteorology, Universität Leipzig, Leipzig, Germany

10 ⁴University of Victoria, Victoria, Canada

Correspondence to: Xiaoyan Ma (xma@nuist.edu.cn)

Abstract. The Moderate Resolution Imaging Spectroradiometer (MODIS) C6 L3, Clouds and the Earth's Radiant Energy System (CERES) Edition-4 L3 products, and the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA-Interim reanalysis data are employed to systematically study aerosol-cloud correlations over three anthropogenic aerosol
15 regions and their adjacent oceans, as well as explore the effect of retrieval artifacts and underlying physical mechanisms. This study is confined to warm phase and single layer clouds without precipitation during the summertime (June, July, and August). Our analysis suggest that cloud effective radius (CER) is positively correlated with aerosol optical depth (AOD) over land (positive slopes), but negatively correlated with aerosol index (AI) over oceans (negative slopes) even with small ranges of liquid water path (quasi-constant). The changes in albedo at the top of atmosphere (TOA) corresponding to aerosol-induced
20 changes in CER also lends credence to the authenticity of this opposite aerosol-cloud correlation between land and ocean. It is noted that potential artifacts, such as the retrieval biases of both cloud (partially cloudy and 3-D shaped clouds) and aerosol, can result in a serious overestimation of the slope of CER-AOD/AI. Our results show that collision-coalescence seems not to be the dominant cause for positive slope over land, but the increased CER caused by increased aerosol might further increase CER by initializing collision-coalescence, generating a positive feedback. By stratifying data according to the lower
25 tropospheric stability and relative humidity near cloud top, it is found that the positive correlations more likely occur in case of drier cloud top and stronger turbulence in clouds, while negative correlations occur in case of moister cloud top and weaker turbulence in clouds, which implies entrainment mixing might be a possible physical interpretation for such a positive CER-AOD slope.

1 Introduction

30 Atmospheric particles influence the Earth's radiation budget and hence climate change by directly scattering and absorbing solar radiation and indirectly by acting as cloud condensation nuclei (CCN), altering cloud properties and precipitation (Twomey, 1977; Albrecht, 1989). The latter historically has been referred to as the aerosol indirect effect, and more recently as the effective radiative forcing due to aerosol-cloud interactions, which remain to have the largest uncertainty in assessing the anthropogenic contribution to present climate change (IPCC, 2013). An increase in CCN number concentration will
35 generate a cloud that consists of more but smaller drops—under constant cloud liquid water path (LWP). The consequence is scattering of more solar radiation back to space. The decrease in cloud effective radius (CER) with increasing aerosol concentrations was historically termed aerosol first indirect effect (AIE), cloud albedo effect, or Twomey effect (Twomey, 1977; Feingold et al., 2003), and in the more recent literature, as radiative forcing due to aerosol-cloud interactions (IPCC, 2013).

40 There have been many observational evidences from in-situ aircraft measurements (Pawlowska & Brenguier, 2000; Wilcox et al., 2006; Roberts et al., 2008; Kleinman et al., 2012; Werner et al., 2014), ground- (Feingold et al., 2003; Kim et al., 2003; Garrett et al., 2004; Qiu et al., 2017) and satellite-based (Nakajima et al., 2001; Bréon et al., 2002; Kaufman et al., 2005; Koren et al., 2005, 2010; Quaas et al., 2008; Costantino & Bréon, 2010, 2013; Lihavainen et al., 2010; Chen et al., 2014; Christensen et al., 2016) remote sensing in support of this negative correlation between aerosol concentrations and CER. Moreover, solid
45 evidences for Twomey effect were also found from aerosol-induced changes in top of atmosphere (TOA) albedo. Su et al. (2010) found a significant increase in TOA albedo associated with continental aerosols relative to those associated with oceanic aerosols under all LWP ranges when cloud fraction is constrained; similar results have been reported by Chen et al. (2014) and Christensen et al. (2016). However, relationships of aerosol optical depth and aerosol index with cloud albedo, cloud fraction, and cloud liquid water path have been shown to be difficult to interpret due to the confounding influence of relative humidity
50 fluctuations that impact both quantities (Quaas et al., 2009, 2010; Gryspeerd et al. 2014; 2016).

In addition to the explorations on aerosol-cloud interactions at the cloud top (from satellite-based remote sensing instruments) and cloud base (from ground-based remote sensing instruments), the role of vertical observations in the cloud-aerosol-precipitation interaction has been studied by using satellite-based radar/lidar. For example, the measurements provided by CALIOP sensors were widely used to obtain the respective position of aerosol and cloud layers (Costantino and Bréon,
55 2010, 2013; Zuidema, et al., 2016; Liu et al., 2017), which can improve the estimation of the amount of aerosols that actually enter the cloud. The CPR (cloud profiling radar) onboard CloudSat is able to penetrate optically thick clouds layers (Wang et al., 2013), and thus has been used to differentiate cloud regimes (Peng et al., 2015; Christensen et al., 2016) and investigate the response of cloud vertical structure to aerosols (Chen et al., 2016). Additionally, the aerosol-induced changes in the vertical structure of precipitation were also explored by using the Tropical Rainfall Measuring Mission (TRMM) products (Wall et al.,
60 2014; Li et al., 2016; Guo et al., 2018).

In addition to the widely observed Twomey effect, positive correlations between aerosol concentrations and CER were also found in some regions, such as southeastern US and southeastern China (Yuan et al., 2008), eastern China (Tang et al., 2014; Wang et al., 2014, 2015; Liu et al., 2017), and Indian (Panicker et al., 2010; Manoj et al., 2012). Overall, positive correlations occur over land while negative correlations dominate over ocean (Grandey and Stier, 2010; Ma et al., 2018). The
65 lack of consensus on these relationships motivates further exploration of underlying physical reasons for these opposite correlations.

It is acknowledged that quantification of aerosol first indirect effect is highly uncertain due to various influencing factors, including (1) aerosol microphysics such as size (Feingold et al., 2001; Dusek et al., 2006; Zhang et al., 2011), chemical composition (Nenes et al., 2002; Lance et al., 2004; Ervens et al., 2005; McFiggans et al., 2006; Almeida et al., 2014) and
70 mixing state (Wang et al., 2008, 2010), (2) meteorological conditions such as vertical velocity (Koren et al., 2010; Lu et al., 2012), lower tropospheric stability (Wang et al., 2014; Saponaro et al., 2017; Ma et al., 2018), wind shear (Fan et al., 2009), and precipitable water vapor (Yuan et al., 2008; Qiu et al., 2017), (3) cloud types (Gryspeerd and Stier, 2012; Chen et al., 2016), and (4) vertical overlapping status of aerosol and cloud layers (Costantino and Bréon, 2010, 2013; Huang et al., 2015; Liu et al., 2017). It is extremely difficult to completely isolate the response of CER to aerosol perturbations from the above-
75 mentioned influencing factors. Yuan et al. (2008) examined the positive correlation between CER and aerosol optical depth (AOD) by using Moderate Resolution Imaging Spectroradiometer (MODIS) satellite products, and speculated that slightly soluble organics particles, which induced a decrease of aerosol activation, might be a possible explanation for the positive correlation. Wang et al. (2014) explored AIE over Eastern China by using both MODIS satellite and National Center for Environmental Prediction (NCEP) reanalysis data and found that positive correlations are more likely to be found under
80 unstable atmosphere conditions. Ma et al. (2018) employed MODIS satellite products and ERA-Interim reanalysis data to systematically explore the impact of meteorological conditions on the correlations over the major industrial regions and over relatively clean oceans, and concluded that positive correlations are more likely corresponding to relatively high cloud top height (CTH) and low lower tropospheric stability (LTS), while negative relationships were predominantly found for low CTH and high LTS. Tang et al. (2014) pointed out that covariation of wind field and relative humidity in North China Plain may
85 contribute to such positive correlations, that is, relatively wet and polluted southerly wind leads to simultaneous increases in both AOD and CER, while dry and clean northerly wind results in coincident decreases in AOD and CER. Gryspeerd and Stier (2012) reported that the strongest positive correlation occurs in the shallow cumulus cloud regimes, while the negative one is analyzed for the stratiform cloud regimes. By using the profiles from the CALIPSO lidar that identifies the respective position of aerosol and cloud layers, Costantino and Bréon (2010, 2013) found that the aerosol indirect effect is stronger for
90 well-mixed aerosol and cloud layers than for separated one.

In addition to real physical-chemical processes stated above, the positive correlations between AOD (or Aerosol Index; AI) and CER may also result from artificial correlations due to the biases of the retrievals. The MODIS aerosol retrieval algorithm is conducted only for clear pixels determined by a cloud mask. AOD could be overestimated due to either cloud contaminations where spurious clouds might be present in pixels that are erroneously identified as completely clear pixels (Kaufman et al.,

95 2005; Remer et al., 2005; Zhang et al., 2005), or cloud adjacency (or 3-D) effect where cloud-free pixels are brightened by reflected light from surrounding clouds (Cahalan et al., 2001; Wen et al., 2006, 2007; Várnai and Marshak, 2009). Moreover, cloud retrievals applied to partially cloudy and 3-D shaped clouds pixels are expected to deviate from the retrieval assumptions of overcast homogenous cloud and 1D plane-parallel radiative transfer, and tend to result in overestimation of CER (Han et al., 1994; Coakley et al., 2005; Matheson et al., 2006; Zhang and Platnick, 2011; Zhang et al., 2012; Grosvenor et al., 2018).
100 Therefore, covariation of biases in CER and AOD (AI) may incur a false correlation between the two variables.

As stated above, aerosol-cloud correlations derived from satellite remote sensing are potentially veiled by large variations in both retrieval biases and meteorological conditions. By employing MODIS and CERES satellite data as well as ERA-Interim reanalysis data, this study aims to: (a) examine whether the positive relationship between AI and CER over land is true, (b) assess how and to what extent the satellite retrieval biases may affect the satellite-diagnosed aerosol-cloud correlations, and
105 (c) explore the underlying physical mechanisms. This paper is organized as follows: the data descriptions of both satellite and reanalysis and data processing are presented in Sect. 2, and major findings are discussed in Sect. 3. A summary and discussion is given in Sect. 4.

2 Data

In this study, three regions with strong anthropogenic emissions over land, namely, East China (EC), East U.S. (EU), and
110 West Europe (WE), as well as their neighboring oceans (ECO, EUO, and WEO, respectively), are chosen to systematically examine aerosol-cloud correlations under different anthropogenic emissions and dynamic and thermodynamic conditions (Fig. 1). The data used include aerosol and cloud properties gathered from the MODIS/Aqua Collection 6 Level-3 (L3) daily product (Levy et al., 2013; Platnick et al., 2017), albedo at the top of the atmosphere (TOA) obtained from the Clouds and the Earth's Radiant Energy System (CERES) Aqua Edition-4 L3 products (Wielicki et al., 1996), as well as meteorological variables
115 extracted from the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA-Interim reanalysis data (Dee et al., 2011). The 14 consecutive years (2003–2016) of daily $1^\circ \times 1^\circ$ gridded data are used for statistical analysis in order to obtain statistically significant results from enough samples.

2.1 Satellite Data

The MODIS C6 L3 product provides AOD retrieved at several wavelengths globally (Levy et al., 2013; Sayer et al., 2014),
120 which has been validated extensively (Remer et al., 2005; Tripathi et al., 2005). In this study, the AOD products retrieved from the Dark Target (DT) algorithm are used. Aerosol index (AI), in comparison to AOD, is believed to be a better proxy for CCN since it contains the information of aerosol size (Nakajima et al., 2001; Stier, 2016). The AI can be derived on the basis of AOD and Ångström exponent (AE), with the former provided directly by the MODIS product and the latter calculated from AOD at wavelength of 460 and 660 nm. However, the MODIS retrieval of AE over land may be problematic (Levy et al.,
125 2013; Sayer et al., 2013). For collection 6 products, several previous studies have found little quantitative skill in MODIS-

retrieved aerosol size parameters over land (Levy et al., 2010; Mielonen et al., 2011). For this reason, we use AOD and AI as proxies for CCN over land and ocean, respectively.

130 The MODIS C6 L3 product provides cloud macrophysical parameters, including cloud fraction (CF), cloud top temperature, cloud top pressure, and cloud top height, as well as cloud microphysical parameters (CER, LWP, and cloud optical depth) with statistics (mean, minimum, maximum, and standard deviation) at three wavelengths (1.6, 2.1, and 3.7 μm) for individual cloud phases (liquid, ice, and undetermined) separately. We filtered the MODIS cloud data according to the criteria employed by Saponaro et al. (2017) to ensure the data used for analysis are only limited to warm liquid phase and single layer clouds, and non-precipitating cases. In previous versions, before MODIS C6, all pixels identified as partly cloudy (either partially cloud-covered or at cloud edge) were restored to clear sky, and the corresponding cloud retrievals were missing, that is, cloud
135 retrievals were only performed in overcast cloudy pixels. In C6, however, the retrievals of cloud microphysical properties are now attempted on these pixels, and successful retrievals are reported in the Level-2 product and aggregated to the Level-3 product, which are reported separately in partly cloudy (PCL) science data sets (SDSs) and are segregated from the normal “overcast” SDSs. Therefore, the simultaneous availability of PCL and normal “overcast” SDSs provides a great opportunity to investigate the effect of partly cloudy retrievals on aerosol-cloud correlations.

140 The CERES SSF1deg Edition-4 product provides radiative fluxes and albedo at TOA for all-sky and clear conditions in the longwave, shortwave, and window regions (Loeb et al. 2005). The albedo at TOA (plenary albedo) is the ratio of broadband (0.2-5 μm) shortwave reflected and the incoming solar flux at the top of the atmosphere. In comparison with the previous version, the CERES instrument calibration and CERES instrument spectral response function corrections have been improved in Edition-4 (Doelling et al., 2016a, 2016b).

145 The satellite-based measurements can provide long-term data sets of aerosols and clouds in high spatial and temporal resolution on a global scale, and have been believed to be a promising tool for the study of aerosol-cloud interactions. However, several inherent limitations prevent a definite quantitative interpretation of aerosol indirect effects. The relationship between aerosol and cloud droplet number, N_d , at cloud base fundamentally is controlled by both the critical supersaturation of aerosol particles, which described by Köhler theory, and the maximum supersaturation at cloud base, which determined by the updraft
150 speed and the competition of the existing CCN for the available humidity. N_d at cloud base thus is a function of cloud-scale, cloud-base vertical wind, aerosol size distribution, and solubility. N_d at cloud base thus necessarily is positively correlated to CCN, with an approximately logarithmic relationship. In this study, we use the satellite-retrieved cloud-top (at approximately 1~5 optical depth into the cloud, depending on the wavelength employed to retrieve it (Platnick, 2000)) CER stratified by classes of LWP as cloud quantity. It deviates from cloud-base N_d since

- 155
- the stratification by LWP bins might not be sufficient to disentangle LWP and N_d impacts on CER
 - cloud-top N_d might be related only loosely to cloud-base N_d in case the clouds are non-adiabatic (entrain environmental air from the sides or top), and/or coagulation of cloud droplets occurs
 - the retrieved CER might be contaminated by various retrieval problems and thus is possibly only loosely related to the real cloud-top CER.

160 As aerosol quantity, we use AOD/AI retrieved in the pixels determined as cloud-free in the retrieval algorithm. This may be only loosely related to the cloud-base CCN. Stier (2016) stated that 52 % of the area of the globe shows correlation coefficients between $CCN_{0.2\%}$ at cloud base and AI below 0.5. This is largely due to:

- AOD/AI is a vertical integral and so might be dominated by aerosol layers that do not affect cloud-base CCN concentrations (Stier, 2016)
- 165 - the clear-sky column retrievals might not be representative of the cloud-base CCN in the neighbouring cloudy pixels
- AOD/AI might be loosely related to cloud-base CCN if affected by deliquescence in high relative humidity environments, or if dust or other insoluble aerosol has a substantial share of the total aerosol.

2.2 Reanalysis Data

To explore the extent to which meteorological conditions affect the CER correlation with AOD/AI, ERA-Interim reanalysis data was employed to derive the lower tropospheric stability (LTS) and relative humidity near cloud top (RH_{CT}). LTS is computed as the difference between the potential temperature at 700 hPa and at the surface (Klein and Hartmann, 1993), representing the magnitude of the inversion strength in the lower troposphere. Large LTS is associated with stable conditions in which vertical mixing is suppressed. RH_{CT} is the relative humidity over the pressure levels closest to the cloud top pressure. The daily $1^\circ \times 1^\circ$ gridded reanalysis data at 14:00 local solar time are used to match the cloud parameters obtained from MODIS L3 data.

3 Results

3.1 Correlations between CER and AOD/AI

According to Twomey (1977), the presence of anthropogenic aerosols increases the cloud droplet number concentration but decreases CER for a constant LWP. Since CER is a function of both AOD/AI and LWP, under conditions where LWP changes with aerosol concentration, the variation in LWP could act to modulate the relationship between AOD/AI and CER. Therefore, the constant LWP assumption should be highlighted in assessing aerosol first indirect effect. However, many previous studies did not constrain the LWP (Breon et al., 2002; Tang et al., 2014; Wang et al., 2014, 2015; Liu et al., 2017), or constrained the LWP into coarse intervals (Nakajima et al., 2001; Sekiguchi et al., 2003; Yuan et al., 2008), which would induce the uncertainty in correlating CER and AOD/AI. In our previous study (Ma et al., 2018), the CER and AOD/AI are grouped over LWP bins with an interval of 40 g m^{-2} , and then a linear regression analysis with the logarithms of the CER and AOD/AI in each LWP bin is performed (as shown in Fig. 2a). It is found that CER is positively correlated with AOD over land (positive slopes), but negatively correlated with AI over their adjacent oceans (negative slopes), and the positive slopes over land become weaker while negative slopes over ocean change slightly as LWP increases. To examine whether such intervals of LWP bin can result in the uncertainty in quantifying CER-AOD/AI correlation, especially the positive ones over land shown in Fig. 2a, we conducted a similar statistical analysis for the LWP versus AOD/AI within each LWP bin. As shown in Fig. 2b,

in the smallest LWP bin (20-60 g m⁻², thin cloud), LWP is more sensitive to AOD/AI, with a significant positive correlation over land but negative correlation over ocean, which will amplify the positive CER-AOD correlation over land and the negative CER-AI correlation over ocean shown in Fig. 2a. For a larger LWP bin (>100 g m⁻²), LWP is insensitive to both AOD and AI, indicating that the increase of cloud water is governed by meteorological conditions instead of aerosol in case of thick clouds.

195 Therefore, coarse intervals of LWP bin might be a possible cause for the appearance that the positive slopes of CER versus AI over land become weaker as LWP increases (Ma et al., 2018).

It is noted that the positive CER-AOD correlation over land can still be found even within bins where LWP is insensitive to AI, implying that the variation of LWP within bin may not be the dominant cause for the observed positive relationship over land for large LWP. However, for the smaller LWP bin where LWP is susceptible to AI, it is still not clear whether the sign of CER-AI correlation is interfered by LWP changes. To this end, we constrain the LWP to a smaller range (5 g m⁻²) to ensure more constant LWP. As data samples in such small intervals are insufficient to calculate the slope of CER versus AOD/AI, the difference of CER between polluted and clean conditions was thus employed as an alternative to the slopes. In this study, two-dimensional (“joint”) histograms of CER and LWP are used to show the probability distribution function of CER at each LWP bin. Meanwhile, the 25th and 75th percentiles of the AOD/AI are computed for each bin with certain CER and LWP, and then the samples with AOD/AI lower than 25th percentiles and with AOD/AI greater than 75th percentiles are classified as ‘clean case’ and ‘polluted case’, respectively. The difference between the joint histograms in polluted and clean cases demonstrates how CER varies with AOD over land and AI over ocean in a quasi-constant LWP bin. Figure 3 indicates that CER increases significantly with increasing LWP over all regions, and the dependence of CER on LWP in polluted case is weaker than that in clean case. By looking at the difference between the joint histograms in polluted and clean cases (the third column in Fig. 3), it is clearly shown that, over land, the samples of polluted case tend to concentrate in the larger CER bin than clean case, while the opposite is found over ocean. That is, as AOD/AI increases, CER increases over land but decreases over ocean even with the tighter constraint on LWP. It can thus be concluded that the variation of LWP within a bin do not change the sign of CER-AOD/AI slope, although it would overestimate the correlation between CER and AOD/AI to some extent.

200
205
210

215 Additionally, previous studies have noted a nonlinear effect of aerosols on cloud microphysics (Breon et al., 2002; Lohmann and Lesins, 2002; Jin et al., 2008; Gryspeerdt et al., 2017). Using the polarization and directionality of the earth reflectances (POLDER) satellite data, Breon et al. (2002) found that the slope of CER versus AI on a global scale is highly significant for AI less than 0.15, whereas a saturation effect occurs as the aerosol keeps rising. The similar saturation effect was also reproduced by the simulations in global climate models (Lohmann and Lesins, 2002; Gryspeerdt et al., 2017). In this study, such saturation effect was found in three ocean regions (ECO, EUO, and WEO) adjacent to the anthropogenic regions (Fig. S1), but with a higher threshold (AI = 0.3) than that derived from the analysis on global oceans (Breon et al., 2002). However, it is noted that no saturation effect was found in the regions with strong anthropogenic emissions over land (EC, EU, and WE), i.e., CER shows a continuous increase as AOD increases (Fig. S1), suggesting that the saturation effects does not appear to interfere with the observed positive slopes over land.

220

225 3.2 Evidences for the Positive Correlation between CER and AOD over Land

Although the positive correlation between CER and AOD/AI over land has been reported by several satellite-based studies (Tang et al., 2014; Wang et al., 2014, 2015; Liu et al., 2017; Ma et al., 2018;), its reliability is still controversial due to the limitations of satellite retrievals. The main concerns include: (a) satellite-retrieved aerosol and cloud properties remain to have biases; (b) the cloud properties observed from satellites are confined to cloud tops rather than entire cloud; and (c) the AI or
230 AOD may not actually represent the magnitude of aerosols entering the clouds. Therefore, it is necessary to confirm whether the positive correlation is real before discussing possible physical causes.

Under cloudy sky, the response of TOA in-cloud albedo at a given LWP on aerosol changes is considered to be a more direct indicator for the Twomey effect than CER (Su et al., 2010), given that the albedo is calculated from CERES TOA upwelling fluxes while CER is retrieved from MODIS reflectance based on multiple assumptions and thus tends to have larger
235 retrieval biases (Matheson et al., 2006; Vant-Hull et al., 2007; Zhang and Platnick, 2011; Zhang et al., 2012). In addition, aerosol and cloud retrievals can also interfere with each other, and thus have potential to result in an artifact correlation. These retrieval biases will be discussed in more detail in section 3.3.1. Therefore, using TOA albedo (α) as a proxy to examine Twomey effect can better exclude the influence of artifacts in the retrievals. In general, the variations of all-sky TOA albedo are controlled by the following five parts: (a) cloud fraction (CF, representing the horizontal extent of cloud); (b) cloud liquid
240 water path (representing the vertical development of a cloud); (c) cloud droplet size (that is, the Twomey effect when considering aerosol perturbations at a certain LWP); (d) clear-sky aerosol loading and above-cloud absorbing aerosol, and (e) surface albedo. In the presence of clouds, the aerosol-induced TOA albedo changes are minimal compared to cloud-induced changes, and therefore (d) are not considered in our analysis. Additionally, for a specific season and region, surface albedo is also assumed to be uniform (Robock, 1980; Wang et al., 2004), so the effect of (e) is ruled out. In this study, CERES TOA
245 albedo and AI are stratified according to CF (with an interval of 0.04) and LWP (with an interval of 10 g m⁻²) in order to isolate the Twomey effect from the interference by them, and ‘clean case’ and ‘polluted case’ are distinguished according to the method used in section 3.1. Thereby, for a specific bin determined by both CF and LWP, the difference of TOA albedo between polluted and clean conditions ($\Delta\alpha$) can be used to represent the changes in TOA albedo caused only by the Twomey effect (Fig. 4). The number of samples for each LWP and CF bin is shown in Fig. S2. A positive $\Delta\alpha$ means that the observed TOA
250 albedo has a larger value in the polluted case than in clean one, and the reverse is true when $\Delta\alpha$ has a negative value. Figure 4 shows that $\Delta\alpha$ is normally negative over land (top), but positive over ocean (bottom). The ratio of positive to total samples over EC, EU, and WE is only 37%, 11%, and 13%, respectively, which is evidently less than that over ECO (85%), EUO (61%), and WEO (84%). This result implies that the reflected solar shortwave radiation at TOA generally tends to reduce with AOD over land while to increase with AI over ocean. As mentioned above, under the constant CF and LWP, the response
255 of TOA albedo to elevated aerosol loading can be interpreted as the result of changes in CER. That is, under polluted conditions, larger droplets over land scatter less solar radiation back to space, inducing lower TOA albedo, whereas smaller droplets over

ocean do the opposite. This finding hence lends credibility to the positive CER-AOD correlations over land and negative CER-AI correlations over ocean, as derived in section 3.1.

3.3 Factors Influencing the Slope of CER versus AOD/AI

260 3.3.1 Artifact Correlations

Although the positive correlations between CER and AOD over land is believed to be a real relationship as discussed in section 3.2, the magnitude of this slope may still be subject to artificial correlations due to the biases of both aerosol retrievals (cloud contamination and cloud adjacency effect) and cloud retrievals. In other words, if biases in CER are correlated with biases in AOD, a false correlation may occur between the two variables.

265 The MODIS aerosol retrieval algorithm is applied only for clear pixels determined by cloud mask (Remer et al., 2005). To screen out cloudy pixels, a 3×3 standard deviation test was used to detect low clouds (Martins et al., 2002) and IR channels to identify high clouds in addition to the standard cloud mask product. The 20% darkest and 50% brightest pixels were also removed for possible cloud contaminations. However, a part of pixels associated with cloud contaminations might still be erroneously identified as clear (Kaufman et al., 2005; Remer et al., 2005; Zhang et al., 2005), inducing an overestimation of

270 AOD. Meanwhile, cloud-free pixels may be brightened by reflected light from surrounding clouds, i.e., cloud adjacency effect (Cahalan et al., 2001; Wen et al., 2006, 2007; Varnai and Marshak, 2009), which could also overestimate AOD. The overestimation of AOD caused by both cloud contaminations and cloud adjacency effect was found to increase as cloud fraction (CF) increases (Zhang et al., 2005). Additionally, cloud retrievals applied to the partly cloudy (PCL) pixels are expected to deviate from the retrieval assumptions of overcast homogenous cloud, and tend to overestimate CER (Han et al.,

275 1994; Coakley et al., 2005; Matheson et al., 2006). Therefore, a spurious positive correlation could occur in case of co-existence of overestimation of AOD and CER. To look into the contribution of retrieval biases on artifact correlations, the correlations of CER and AOD/AI derived from PCL retrievals and normal “overcast” retrievals are compared, where data are stratified according to CF in order to account for the impact of biases in aerosol retrievals (Fig. 5). Considering the limited data samples for the LWP bins of 140-180 and 180-220 g m^{-2} , the associated statistical results are not included into our analysis. As seen in

280 Fig. 5, the CER-AOD/AI slopes in PCL retrieval (circles) are significantly larger than that in overcast retrieval (dots). It is interesting to see that, over ocean, where the Twomey effect dominates actually, the slope even becomes positive in the PCL case, which suggests that CER biases in PCL retrievals can result in a serious overestimation of the positive slope. In addition to PCL retrievals, the biases in aerosol retrievals could also affect this positive slope. The differences between the slope in PCL and in overcast retrievals in case of $\text{CF} > 60\%$ ($\Delta S_{>60}$) and $\text{CF} < 60\%$ ($\Delta S_{<60}$) over three anthropogenic regions and their

285 adjacent oceans are summarized in Table 1, which clearly shows that $\Delta S_{>60}$ is overall larger than $\Delta S_{<60}$ under all LWP bins, implying that the overestimation of AOD due to retrieval biases under larger CF can amplify the positive slope caused by PCL retrievals. Therefore, in the actual retrieval process, if MODIS algorithm erroneously identifies PCL pixels as overcast pixels, the artifact positive correlation will be introduced, especially for condition of larger CF.

In addition to PCL retrievals, the cloud retrievals for 3-D shaped clouds are also problematic due to deviating from the
290 retrieval assumptions of plane-parallel clouds (Nakajima and King, 1990). The variations in the sun-satellite scattering
geometries can interfere satellite-measured signals and hence induce biases in retrievals for 3-D shaped clouds, which are
interpreted as shadowing and illumination effects (Vant-Hull et al., 2007). Such retrieval biases in CER were found to be larger
in 2.1 μm channel than in 3.7 μm channel (Zhang and Platnick, 2011; Zhang et al., 2012). Our previous study (Ma et al., 2018)
295 stated that the results of positive slope over land and negative slope over ocean changed little when using CER in 3.7 μm
(CER_{3.7 μm}) rather than CER in 2.1 μm (CER_{2.1 μm}) for the regression analysis. In order to examine the effect of biases in 3-D
shaped cloud retrievals on artifact correlations, the differences between CER_{2.1 μm} -AOD slope and CER_{3.7 μm} -AOD slope
over land and the differences between CER_{2.1 μm} -AI slope and CER_{3.7 μm} -AI slope over ocean under all LWP bins are calculated,
respectively (Fig. 6). The number of samples for each bin for regression analysis is shown in Fig. S3. It is clearly shown that
300 that retrieval biases for 3-D shaped cloud tend to result in a higher slope than its physically correct value.

3.3.2 Possible Physical Explanations

3.3.2.1 Collision and Coalescence

In warm clouds, collision-coalescence is a much more effective process to increase droplet size compared with diffusion
of water vapor (Langmuir, 1948; Kogan, 1993; Beard and Ochs, 1993; Pruppacher and Klett, 1997). The impact of collision-
305 coalescence on CER is believed to be minimal when CER is smaller than $\sim 14 \mu\text{m}$, while the coalescence rate increases very
rapidly when CER is greater than this value (Gerber, 1996; Freud and Rosenfeld, 2012), i.e., collision-coalescence becomes
dominant. To explore the possible contribution of collision-coalescence processes to the correlation between CER and AOD/AI,
the data are thus stratified into two subsets with CER greater and less than $14 \mu\text{m}$, respectively. As shown in Fig. 7, the
dependence of CER on AOD/AI becomes fairly weak with the slope close to zero over both land and ocean when collision-
310 coalescence process is dominant (blue; CER $> 14 \mu\text{m}$), while CER-AOD/AI slopes are still positive over land and negative
over ocean when collision-coalescence is negligible, which suggests that collision-coalescence is not likely the major cause
for the opposite CER-AOD/AI correlations over land and ocean. Figure 8 shows the probability density functions of AOD/AI
for the above two subsets over land and ocean, respectively. It is clearly demonstrated that the occurrence frequency of AOD/AI
presents a significant difference between the case of CER $> 14 \mu\text{m}$ and CER $< 14 \mu\text{m}$, with larger AOD for the former and
315 smaller AOD for the later over land, while the reverse is true over ocean. This result implies that collision-coalescence are
more likely to occur in case of larger AOD over land while smaller AI over ocean, which may further enhance the positive
CER-AOD slope over the land and the negative CER-AI slope over ocean, as demonstrated in Fig.7. From the above analyses,
we assume that although collision-coalescence is not the dominant cause for positive slope over land, the increased CER caused
by increased aerosol might further increase CER by initializing collision-coalescence, generating a positive feedback.

320 3.3.2.2 Entrainment mixing

In addition to the aforementioned collision-coalescence process, entrainment mixing is also believed to be an effective way to reduce cloud droplet number concentration and increase CER, which of course depends on the specific details of entrainment-mixing mechanisms, as discussed in depth by Kim et al. (2006). The possible impacts of the entrainment mixing process on the estimated aerosol first indirect effect (AIE) have been reported by previous studies. For instance, Kim et al. (2006) found that the AIE is relatively weak in sub-adiabatic clouds compared with adiabatic clouds based on ground-based remote sensing at the Southern Great Plains, and pointed out that this may be due to interference from heterogeneous entrainment mixing that change the droplet number concentrations in a manner that attenuates the AIE. Additionally, Shao and Liu (2006) suggested the AIE is about half of those estimated by many previous studies, and attributed this difference to evaporation resulting from entrainment mixing processes. By using a mixed-layer model, Dal Gesso et al. (2014) found that the entrainment rate in stratocumulus clouds tends to be larger for the smaller lower tropospheric stability (LTS). Although LTS is an indicator of temperature inversion proposed as relevant for marine stratocumulus clouds (Klein and Hartmann, 1993), it has also been widely used to characterize atmospheric stability over both land and ocean, and thereby indicating the vertical mixing (Chen et al., 2014; Chen et al., 2016; Liu et al., 2017; Ma et al., 2018). As known, the degree of entrainment mixing is subject to not only the intensity of turbulence in clouds but also the relative humidity of air outside clouds (De Rooy et al., 2013). In this study, the former is represented by lower tropospheric stability (LTS), and the latter is described by relative humidity near cloud top (RH_{CT}).

To investigate the extent to which entrainment mixing affect the slopes of CER versus AOD/AI, the dataset is thus stratified according to LTS and RH_{CT} to differentiate environmental regimes. In order to satisfy the constant LWP assumption for aerosol first indirect effect, LWP bin range from 60 to 100 $g\ m^{-2}$ is employed, among which the LWP does not change significantly with AOD/AI (as demonstrated in Fig. 1b) and statistical samples are sufficient for regression analysis. The slopes of CER versus AOD over land and the slopes of CER versus AI over ocean on log-log scale under low, medium, and high LTS and RH_{CT} conditions are shown in Fig. 9, in which nine categories are classified to make sure that each category has roughly the same number of samples (Table 2). It is clearly shown that slopes under low LTS (weak stable condition) are more positive than that under medium (medium stable) and high LTS (stable condition) over both land and ocean. Also, the slopes under low RH_{CT} (dry condition) are overall more positive than that under high RH_{CT} (moist condition), although the responses of slopes to RH_{CT} are not as significant as that with LTS. That is, under drier cloud top and stronger turbulence in clouds, positive correlations more likely occur, which are associated with stronger entrainment.

Figure 10 shows the distributions of sample number for LTS and RH_{CT} . It is clearly demonstrated that there exists a systemic difference in both LTS and RH_{CT} between land and ocean; the frequencies of clouds under lower LTS (Fig. 10a) and RH_{CT} (Fig. 10b) are higher over land than over ocean. The average LTS and RH_{CT} over EC (10.3 ± 2.3 and 67.8 ± 14.8), EU (13.2 ± 2.5 and 68.4 ± 15.6), and WE (12.5 ± 3.2 and 65.1 ± 16.0) is generally lower than that over ECO (14.2 ± 1.4 and 73.8 ± 13.3), EUO (14.1 ± 1.7 and 71.9 ± 16.8), and WEO (17.9 ± 3.1 and 75.8 ± 23.5). Therefore, the opposite correlations over major industrial

regions and their adjacent oceans discussed in section 3.1 is probably associated with this systemic difference in the indicators of entrainment mixing, i.e., LTS and RH_{CT} . As shown in Fig. 9, positive correlations are more likely occur under the condition of drier cloud top and stronger turbulence in clouds, which is more prevalent over land, while negative correlations normally occur under the condition of moister cloud top and weaker turbulence in clouds, which is common over ocean. The opposite CER-AOD/AI correlations over land and the negative one over ocean have also been generally found on a global scale (Grandey and Stier, 2010), but this is not always the case for some regions (Nakajima et al., 2001; Bréon et al., 2002), indicating that aerosol-cloud interactions remain challenging due to the interference of other influencing factors, such as different aerosol microphysical properties and coincidentally changing meteorological conditions.

4 Summary

Using fourteen years of co-located aerosol and cloud observations from MODIS C6 L3 products, together with ERA-Interim reanalysis data, we systematically investigated correlations between cloud effective radius (CER) and aerosol optical depth (AOD) over three anthropogenic regions and correlations between CER and aerosol index (AI) over their adjacent oceans, assessed the effects of satellite retrieval biases on the aerosol-cloud correlations, and then explored the underlying physical mechanisms. We also verified the reliability of the CER-AOD/AI correlation derived from satellite data by employing CERES Edition-4 L3 TOA albedo product.

Our analysis indicated that cloud effective radius is overall positively correlated with aerosol optical depth over land (positive slopes), but negatively correlated with aerosol index over oceans (negative slopes). Since CER is a function of both AOD/AI and LWP, the assumption of constant LWP must be taken into consideration when estimating aerosol first indirect effect. The results shown that variations of LWP with AOD/AI in coarse LWP bins (40 g m^{-2}) can amplify the positive slope over land and the negative one over ocean. Therefore, we reanalyze the correlation between CER and AOD/AI for narrow intervals of LWP (5 g m^{-2}) in order to exclude the interference of the covariation of LWP and AOD/AI, and still get positive relationships over land and negative one over ocean.

The reliability of positive correlation between CER and AOD based on satellite remote sensing is still controversial (Zhao et al., 2018) due to the limitations of retrieval biases. Therefore, it is necessary to examine whether the positive correlation over land is real before exploring possible physical causes. We stratified data according to CF (with an interval of 0.04) and LWP (with an interval of 10 g m^{-2}) to isolate the Twomey effect. The results indicated that the changes in albedo at TOA corresponding to aerosol-induced changes in CER also lends credence to the reliability of positive correlation over land and negative one over ocean.

Although the positive correlations between CER and AOD over land is believed to be a real relationship as discussed above, the magnitude of slope is still subject to artificial correlations due to the retrieval biases in both aerosol (cloud contamination and cloud adjacency effect) and cloud (partially cloudy and 3-D shaped clouds). To evaluate the contribution of retrieval biases on artificial correlations, we compared the correlations between CER and AOD/AI derived from PCL(partly cloudy) retrievals

385 and normal “overcast” retrievals, and took the impact of retrieval biases in aerosol into account by stratifying data according to CF. It is suggested that CER biases in PCL retrievals can induce an overestimation of the CER-AOD/AI slope, and the retrieval biases of aerosol can amplify the overestimation of slope caused by PCL retrievals. Additionally, our analysis shown that retrieval biases for 3-D shaped cloud also tend to result in a more positive slope of CER versus AOD/AI than its physically correct value. Therefore, the artificial positive correlation will be introduced if the following occurs: (a) PCL pixels are
390 erroneously identified as overcast pixels; (b) cloudy pixels are erroneously identified as clear; and (c) 3-D shaped clouds affect both cloud retrievals in their own cloud pixels and aerosol retrievals in surrounding cloud-free pixels.

We also explored potential physical mechanisms that can help to explain observed positive correlation between CER and AOD over land, including collision-coalescence and entrainment mixing. It is suggested that collision-coalescence seems not to be the dominant cause for positive slope of CER versus AOD over land, but there might exist a positive feedback that the
395 increased CER caused by increased aerosol might further increase CER by initializing collision-coalescence. Additionally, we stratified data according to the lower tropospheric stability (LTS) and relative humidity near cloud top (RH_{CT}) to differentiate environmental regimes, and found that the positive correlations more likely occur in case of drier cloud top and stronger turbulence in clouds, corresponding to stronger entrainment mixing. Furthermore, there exists systematic differences in LTS and RH_{CT} between land dominated by the positive correlation and ocean dominated by the negative one, i.e., LTS and RH_{CT}
400 are lower over land than over ocean. Therefore, it is inferred that entrainment mixing might be a possible physical interpretation for such positive CER-AI slope.

It is acknowledged that although we have explored some uncertainties in clarifying the correlation between aerosols and clouds, the complexity of this issue and inherent limitations of polar-orbiting satellite measurements make it difficult to reach a definitive conclusion regarding causal relationships between CER and AOD/AI. Additional potential sources of the
405 uncertainties include:

- Satellite-derived cloud-top CER might deviate from cloud-base or whole cloud CER in case the clouds are non-adiabatic, and/or coagulation of cloud droplets occurs.
- As a vertical integral quantity, AOD/AI is not always a good proxy for cloud-base CCN concentrations in some cases (Stier, 2016).
- 410 - AOD/AI might be loosely related to cloud-base CCN if affected by deliquescence in high relative humidity environments, or if dust or other insoluble aerosol has a substantial share of the total aerosol.
- AOD/AI of clear-sky column retrievals might not be representative of the cloud-base CCN in the neighbouring cloudy pixels.

Unfortunately, it is quite difficult to evaluate these uncertainties in this study due to the limitations of the data. However,
415 our study still present sufficient evidence to warrant further investigations of the physical mechanisms using detailed in-situ field measurements of aerosol and cloud properties, especially vertical profiles.

Data availability. All data used in this study are publicly available. The MODIS/Aqua Level 3 Collection 6 datasets were acquired from the Level-1 and Atmosphere Archive & Distribution System (LAADS) Distributed Active Archive Center (DAAC), located in the Goddard Space Flight Center in Greenbelt, Maryland (<https://ladsweb.nascom.nasa.gov/>). CERES SSF data were obtained from the NASA Langley Research Center Atmospheric Science Data Center (https://ceres.larc.nasa.gov/order_data.php). The ECMWF ERA-Interim data were collected from the ECMWF data server <http://www.ecmwf.int/en/research/climate-reanalysis/>.

Author contributions. HJ and XM designed the study and the statistical analysis. HJ processed the satellite data and drafted the manuscript. XM, JQ, YY, and TQ validated and debugged the results. All authors contributed to revising the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study is supported by the National Natural Science Foundation of China grants (41475005 and 41675004) and the National Key R&D Program of China grants (2016YFA0600404). Yan Yin acknowledges funding support by the National Natural Science Foundation of China grants (41590873). We are grateful to the ease access to MODIS and CERES, provided by NASA. We also thank ECMWF for providing daily ERA-Interim reanalysis data in our work.

References

- Albrecht, B.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245(4923), 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Almeida, G. P., Brito, J., Morales, C. A., Andrade, M. F., and Artaxo, P.: Measured and modelled cloud condensation nuclei (CCN) concentration in São Paulo, Brazil: the importance of aerosol size-resolved chemical composition on CCN concentration prediction, *Atmos. Chem. Phys.*, 14, 7559–7572, doi:10.5194/acp-14-7559-2014, 2014.
- Beard, K. V. and Ochs, H. T.: Warm-Rain Initiation: An Overview of Microphysical Mechanisms, *J. Appl. Meteorol.*, 32, 608–625, 1993.
- Bréon, F.-M., Tanré, D., and Generoso, S.: Aerosol effect on cloud droplet size monitored from satellite, *Science*, 295, 834–838, 2002.
- Cahalan, R. F., Oreopoulos L., Wen G., Marshak A., Tsay S.-C., and DeFelice T.: Cloud characterization and clear sky correction from Landsat 7, *Remote Sens. Environ.*, 78, 83–98, doi:10.1016/S0034-4257(01)00251-6, 2001.
- Chen, T. M., Guo, J. P., Li, Z. Q., Zhao, C., Liu, H., Cribb, M., Wang, F., and He, J.: A CloudSat perspective on the cloud climatology and its association with aerosol perturbation in the vertical over East China, *J. Atmos. Sci.*, 73, 3599–3616, <https://doi.org/10.1175/JAS-D-15-0309.1>, 2016.

- Chen, Y.-C., Christensen, M. W., Stephens, G. L., and Seinfeld, J. H.: Satellite-based estimate of global aerosol-cloud radiative forcing by marine warm clouds, *Nat. Geosci.*, 7, 643–646, <https://doi.org/10.1038/ngeo2214>, 2014.
- Christensen, M. W., Chen, Y.-C., and Stephens, G. L.: Aerosol indirect effect dictated by liquid clouds, *J. Geophys. Res.*, 121, 14636–14650, <https://doi.org/10.1002/2016JD025245>, 2016.
- Coakley, J. A., Friedman, M. A., and Tahnk, W. R.: Retrieval of cloud properties for partly cloudy imager pixels, *J. Atmos. Ocean. Techn.*, 22(1), 3–17, 2005.
- Costantino, L. and Breon, F.-M.: Analysis of aerosol-cloud interaction from multi-sensor satellite observations, *Geophys. Res. Lett.*, 37, L11801, doi:10.1029/2009GL041828, 2010.
- Costantino, L. and Bréon, F.-M.: Aerosol indirect effect on warm clouds over South-East Atlantic, from co-located MODIS and CALIPSO observations, *Atmos. Chem. Phys.*, 13, 69–88, doi:10.5194/acp-13-69-2013, 2013.
- Dee, D., Uppala, S., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- de Rooy, W. C., Bechtold, P., Frohlich, K., Hohenegger, C., Jonker, H., Mironov, D., Siebesma, A. P., Teixeira, J., Yano, J.-I.: Entrainment and detrainment in cumulus convection: an overview, *Q. J. Roy. Meteorol. Soc.*, 139, 1–19, doi:10.1002/qj.1959, 2013.
- Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S., and Andreae, M. O.: Size matters more than chemistry for cloud-nucleating ability of aerosol particles, *Science*, 312, 1375–1378, doi:10.1126/science.1125261, 2006.
- Doelling, D. R., Haney, C. O., Scarino, B. R., Gopalan, A., and Bhatt, R.: Improvements to the Geostationary Visible Imager Ray-Matching Calibration Algorithm for CERES Edition 4, *J. Atmos. Ocean. Technol.*, 33, 2679–2698, <https://doi.org/10.1175/jtech-d-16-0113.1>, 2016.
- Doelling, D. R., Sun, M., Nguyen, L. T., Nordeen, M. L., Haney, C. O., Keyes, D. F., Mlynchak, P. E.: Advances in Geostationary-Derived Longwave Fluxes for the CERES Synoptic (SYN1deg) Product, *J. Atmos. Ocean. Techn.*, 33, 503–521, <https://doi.org/10.1175/JTECH-D-15-0147.1>, 2016.
- Ervens, B., Feingold, G., and Kreidenweis, S. M.: Influence of water-soluble organic carbon on cloud drop number concentration, *J. Geophys. Res.*, 110(D18), 18 211, doi:10.1029/2004JD005634, 2005.
- Fan, J., Yuan, T., Comstock, J. M., Ghan, S., Khain, A., Leung, L. R., Li, Z., Martins, J. V., and Ovchinnikov, M.: Dominant role by vertical wind shear in regulating aerosol effects on deep convective clouds, *J. Geophys. Res.*, 114, D22206, doi:10.1029/2009JD012352, 2009.

- Feingold, G., Eberhard, W., Veron, D., and Previdi, M.: First measurements of the Twomey indirect effect using ground-based remote sensors, *Geophys. Res. Lett.*, 30, 1287, doi:10.1029/2002GL016633, 2003.
- Feingold, G., Remer, L. A., Ramaprasad, J., and Kaufman, Y. J.: Analysis of smoke impact on clouds in Brazilian biomass burning regions: An extension of Twomey's approach, *J. Geophys. Res.*, 106, 22 907–22 922, 2001.
- 485 Freud, E. and Rosenfeld, D.: Linear relation between convective cloud drop number concentration and depth for rain initiation, *J. Geophys. Res.*, doi:10.1029/2011JD016457, in preparation, 2011.
- Garrett, T. J., Zhao, C., Dong, X., Mace, G. G., and Hobbs, P. V.: Effects of varying aerosol regimes on low-level Arctic stratus, *Geophys. Res. Lett.*, 31, L17105, doi:10.1029/2004GL019928, 2004.
- 490 Gerber, H.: Microphysics of marine stratocumulus clouds with two drizzle modes, *J. Atmos. Sci.*, 53, 1649–1662, 1996.
- Grandey, B. S. and Stier, P.: A critical look at spatial scale choices in satellite-based aerosol indirect effect studies, *Atmos. Chem. Phys.*, 10, 11459–11470, doi:10.5194/acp-10-11459-2010, 2010.
- Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., Boers, R., Cairns, B., Chiu, J. C., Christensen, M., Deneke, H., Diamond, M., Feingold, G., Fridlind, A., Hünerbein, A., Knist, C., Kollias, P., Marshak, A., McCoy, D., Merk, D., Painemal, D., Rausch, J., Rosenfeld, D., Russchenberg, H., Seifert, P., Sinclair, K., Stier, P., van Diedenhoven, B., Wendisch, M., Werner, F., Wood, R., Zhang, Z., and Quaas, J.: Remote Sensing of Droplet Number Concentration in Warm Clouds: A review of the current state of knowledge and perspectives, *Rev. Geophys.*, <https://doi.org/10.1029/2017RG000593>, 2018.
- 495 Gryspeerdt, E. and Stier, P.: Regime-based analysis of aerosol-cloud interactions, *Geophys. Res. Lett.*, 39, L21802. <https://doi.org/10.1029/2012GL053221>, 2012.
- 500 Gryspeerdt, E., Stier, P., and Grandey, B.: Cloud fraction mediates the aerosol optical depth - cloud top height relationship, *Geophys. Res. Lett.*, 41, <https://doi.org/10.1002/2014GL059524>, 2014.
- Gryspeerdt, E., Quaas, J., and Bellouin, N.: Constraining the aerosol influence on cloud fraction, *J. Geophys. Res.*, 121, 3566–3583, <https://doi.org/10.1002/2015JD023744>, 2016.
- 505 Gryspeerdt, E., Quaas, J., Ferrachat, S., Gettelman, A., Ghan, S., Lohmann, U., Morrison, H., Neubauer, D., Partridge, D. G., Stier, P., Takemura, T., Wang, H., Wang, M., and Zhang, K.: Constraining the instantaneous aerosol influence on cloud albedo, *Proc. Natl. Acad. Sci. USA*, 114, 4899–4904, <https://doi.org/10.1073/pnas.1617765114>, 2017.
- Guo, J., Liu, H., Li, Z., Rosenfeld, D., Jiang, M., Xu, W., Jiang, J. H., He, J., Chen, D., Min, M., and Zhai, P.: Aerosol-induced changes in the vertical structure of precipitation: a perspective of TRMM precipitation radar, *Atmos. Chem. Phys.*, 18, 13329–13343, <https://doi.org/10.5194/acp-18-13329-2018>, 2018.
- 510 Han, Q. Y., Rossow, W. B., and Lacis, A. A.: Near-global survey of effective droplet radii in liquid water clouds using ISCCP data, *J. Climate*, 7(4), 465–497, 1994.
- Huang, J., Guo, J., Wang, F., Liu, Z., Jeong, M. J., Yu, H., and Zhang, Z.: CALIPSO inferred most probable heights of global dust and smoke layers, *J. Geophys. Res.-Atmos.*, 120, 5085–5100, 2015.

- 515 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., doi:10.1017/CBO9781107415324, 2013.
- Jin, M. and Shepherd, J. M.: Aerosol relationships to warm season clouds and rainfall at monthly scales over east China: Urban
520 land versus ocean, *J. Geophys. Res.*, 113, D24S90, doi:10.1029/2008JD010276, 2008
- Kaufman, Y. J., Koren, I., Remer, L. A., Rosenfeld, D. and Rudich, Y.: The effect of smoke, dust, and pollution aerosol on shallow cloud development over the Atlantic Ocean. *Proc. Natl. Acad. Sci. USA*, 102(32), 11 207–11 212. <https://doi.org/10.1073/pnas.0505191102>, 2005.
- Kim, B.-G., Schwartz, S. E., Miller, M. A., and Min, Q.: Effective radius of cloud droplets by ground-based remote sensing:
525 Relationship to aerosol, *J. Geophys. Res.*, 108, 4740, doi:10.1029/2003JD003721, 2003.
- Kim, B.-G., Miller, M. A., Schwartz, S. E., Liu, Y., and Min, Q.: The role of adiabaticity in the aerosol first indirect effect, *J. Geophys. Res.*, 113, D05210, doi:10.1029/2007JD008961, 2008.
- Klein, S. A. and Hartmann, D. L.: The seasonal cycle of low stratiform clouds, *J. Climate*, 6, 1588–1606, 1993.
- Kleinman, L. I., Daum, P. H., Lee, Y.-N., Lewis, E. R., Sedlacek III, A. J., Senum, G. I., Springston, S. R., Wang, J., Hubbe,
530 J., Jayne, J., Min, Q., Yum, S. S., and Allen, G.: Aerosol concentration and size distribution measured below, in, and above cloud from the DOE G-1 during VOCALS-REx, *Atmos. Chem. Phys.*, 12, 207– 223, doi:10.5194/acp-12-207-2012, 2012.
- Koren, I., Feingold, G., and Remer, L. A.: The invigoration of deep convective clouds over the Atlantic: Aerosol effect, meteorology or retrieval artifact? *Atmos. Chem. Phys.*, 10(18), 8855–8872, <https://doi.org/10.5194/acp-10-8855-2010>,
535 2010.
- Koren, I., Kaufman, Y. J., Rosenfeld, D., Remer, L. A., and Rudich, Y.: Aerosol invigoration and restructuring of Atlantic convective clouds, *Geophys. Res. Lett.*, 32, L14828, <https://doi.org/10.1029/2005GL023187>, 2005.
- Kogan, Y. L.: Drop Size Separation in Numerically Simulated Convective Clouds and Its Effect on Warm Rain Formation, *J. Atmos. Sci.*, 50, 1238–1253, 1993.
- 540 Lance, S., Nenes, A., and Rissman, T. A.: Chemical and dynamical effects on cloud droplet number: Implications for estimates of the aerosol indirect effect, *J. Geophys. Res.*, 109, D22208. <https://doi.org/10.1029/2004JD004596>, 2004.
- Langmuir, I.: The production of rain by a chain reaction in cumulus clouds at temperatures above freezing, *J. Atmos. Sci.*, 5, 175–192, 1948
- Levy, R. C., Remer, L. A., Kleidman, R. G., Mattoo, S., Ichoku, C., Kahn, R., and Eck, T. F.: Global evaluation of the
545 Collection 5 MODIS dark-target aerosol products over land, *Atmos. Chem. Phys.*, 10, 10399–10420, doi:10.5194/acp-10-10399-2010, 2010.
- Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmos. Meas. Tech.*, 6, 2989–3034, doi:10.5194/amt-6-2989-2013, 2013.

- Li, Z., Lau, W.M., Ramanathan, V., Wu, G., Ding, Y., Manoj, M. G., Liu, J., Qian, Y., Li, J., Zhou, T., Fan, J., Rosenfeld, D.,
550 Ming, Y., Wang, Y., Huang, J., Wang, B., Xu, X., Lee, S.-S., Cribb, M., Zhang, F., Yang, X., Zhao, C., Takemura, T.,
Wang, K., Xia, X., Yin, Y., Zhang, H., Guo, J., Zhai, P. M., Sugimoto, N., Babu, S. S., and Brasseur, G. P.: Aerosol and
monsoon climate interactions over Asia, *Rev. Geophys.*, 54, 866–929, <https://doi.org/10.1002/2015RG000500>, 2016.
- Lihavainen, H., Kerminen, V. M. and Remer, L. A.: Aerosol-cloud interaction determined by both in situ and satellite data
over a northern high-latitude site, *Atmos. Chem. Phys.*, 10(22), 10987–10995, [https://doi.org/10.5194/acp-10-10987-](https://doi.org/10.5194/acp-10-10987-2010)
555 [2010, 2010](https://doi.org/10.5194/acp-10-10987-2010).
- Liu, Y., de Leeuw, G., Kerminen, V.-M., Zhang, J., Zhou, P., Nie, W., Qi, X., Hong, J., Wang, Y., Ding, A., Guo, H., Krüger,
O., Kulmala, M., and Petäjä, T.: Analysis of aerosol effects on warm clouds over the Yangtze River Delta from multi-
sensor satellite observations, *Atmos. Chem. Phys.*, 17, 5623–5641, <https://doi.org/10.5194/acp-17-5623-2017>, 2017.
- Loeb, N. G., Kato, S., Loukachine, K., and Manalo-Smith, N.: Angular distribution models for top-of-atmosphere radiative
560 flux estimation from the Clouds and Earth’s Radiant Energy System instrument on the Terra satellite. Part 1 methodology.,
J. Atmos. Ocean. Tech., 22, 338–351, 2005.
- Lohmann, U. and Lesins, G.: Stronger constraints on the anthropogenic indirect aerosol effect, *Science*, 298, 1012–1016, 2002.
- Lu, C., Liu Y., Niu S., and Vogelmann A. M.: Observed impacts of vertical velocity on cloud microphysics and implications
for aerosol indirect effects, *Geophys. Res. Lett.*, 39, L21808, doi:10.1029/2012GL053599, 2012.
- 565 Ma, X., Jia, H., Yu, F., and Quaas, J.: Opposite aerosol index-cloud droplet effective radius correlations over major industrial
regions and their adjacent oceans, *Geophys. Res. Lett.*, 45, 5771–5778, 2018.
- Matheson, M. A., Coakley Jr., J. A., and Tahnk, W. R.: Multiyear advanced very high resolution radiometer observations of
summertime stratocumulus collocated with aerosols in the northeastern Atlantic, *J. Geophys. Res.*, 111, D15206,
doi:10.1029/2005JD006890, 2006.
- 570 Manoj, M. G., Devara, P. C. S., Joseph, S., and Sahai, A. K.: Aerosol indirect effect during the aberrant Indian Summer
Monsoon breaks of 2009, *Atmos. Environ.*, 60, 153–163, 2012.
- Martins, J. V., Tanre, D., Remer, L. A., Kaufman, Y. J., Mattoo, S., Levy, R.: MODIS cloud screening for remote sensing of
aerosol over oceans using spatial variability, *Geophys. Res. Lett.*, 29(12), doi:10.1029/2001GL013252, 2002.
- McFiggans, G., Artaxo, P., Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S., Gysel, M., Laaksonen, A.,
575 Lohmann, U., Mentel, T. F., Murphy, D. M., O’Dowd, C. D., Snider, J. R., and Weingartner, E.: The effect of physical
and chemical aerosol properties on warm cloud droplet activation, *Atmos. Chem. Phys.*, 6, 2593–2649,
<https://doi.org/10.5194/acp-6-2593-2006>, 2006.
- Mielonen, T., Levy, R. C., Aaltonen, V., Komppula, M., de Leeuw, G., Huttunen, J., Lihavainen, H., Kolmonen, P., Lehtinen,
K. E. J., and Arola, A.: Evaluating the assumptions of surface reflectance and aerosol type selection within the MODIS
580 aerosol retrieval over land: the problem of dust type selection, *Atmos. Meas. Tech.*, 4, 201–214, doi:10.5194/amt-4-201-
2011, 2011.

- Nakajima, T. and King, M. D.: Determination of optical thickness and effective particle radius of clouds from reflected solar radiation measurements. Part I: Theory, *J. Atmos. Sci.*, 47, 1878–1893, 1990.
- 585 Nakajima, T., Higurashi, A., Kawamoto, K., and Penner, J. E.: A possible correlation between satellite-derived cloud and aerosol microphysical parameters, *Geophys. Res. Lett.*, 28, 1171–1174, 2001.
- Nenes, A., Charlson, R. J., Facchini, M. C., Kulmala, M., Laaksonen, A., and Seinfeld, J. H.: Can chemical effects on cloud droplet number rival the first indirect effect? *Geophys. Res. Lett.*, 29(17), 1848, <https://doi.org/10.1029/2002GL015295>, 2002.
- 590 Panicker, A. S., Pandithurai, G., and Dipu, S.: Aerosol indirect effect during successive contrasting monsoon seasons over Indian subcontinent using MODIS data, *Atmos. Environ.*, 44, 1937–1943, 2010.
- Platnick, S.: Vertical photon transport in cloud remote sensing problems, *J. Geophys. Res.*, 105, 22 919–22 935, 2000.
- Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS cloud optical and microphysical products: Collection 6 updates and examples from Terra and Aqua, *IEEE T. Geosci. Remote*, 55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.
- 595 Pawlowska, H., and Brenguier, J. L.: Microphysical properties of stratocumulus clouds during ACE-2, *Tellus B*, 52(2), 868–887, <https://doi.org/10.1034/j.1600-0889.2000.00076.x>, 2000.
- Pruppacher, H. R., and J. D. Klett, 1997: *Microphysics of Clouds and Precipitation*. 2nd ed. Kluwer Academic, 954 pp.
- 600 Qiu, Y., Zhao, Z., Guo, J., and Li, J.: 8-Year ground-based observational analysis about the seasonal variation of the aerosol-cloud droplet effective radius relationship at SGP site, *Atmospheric Environment*, 164, 139–146. <https://doi.org/10.1016/j.atmosenv.2017.06.002>, 2017.
- Quaas, J., Boucher, O., Bellouin, N., and Kinne, S.: Satellite-based estimate of the direct and indirect aerosol climate forcing, *J. Geophys. Res.*, 113, D05204, <https://doi.org/10.1029/2007JD008962>, 2008.
- 605 Quaas, J., Ming, Y., Menon, S., Takemura, T., Wang, M., Penner, J. E., Gettelman, A., Lohmann, U., Bellouin, N., Boucher, O., Sayer, A. M., Thomas, G. E., McComiskey, A., Feingold, G., Hoose, C., Kristjánsson, J. E., Liu, X., Balkanski, Y., Donner, L. J., Ginoux, P. A., Stier, P., Grandey, B., Feichter, J., Sednev, I., Bauer, S. E., Koch, D., Grainger, R. G., Kirkevåg, A., Iversen, T., Seland, Ø., Easter, R., Ghan, S. J., Rasch, P. J., Morrison, H., Lamarque, J.-F., Iacono, M. J., Kinne, S., and Schulz, M.: Aerosol indirect effects – general circulation model intercomparison and evaluation with satellite data, *Atmos. Chem. Phys.*, 9, 8697–8717, <https://doi.org/10.5194/acp-9-8697-2009>, 2009.
- 610 Quaas, J., Stevens, B., Stier, P., and Lohmann, U.: Interpreting the cloud cover – aerosol optical depth relationship found in satellite data using a general circulation model, *Atmos. Chem. Phys.*, 10, 6129–6135, doi:10.5194/acp-10-6129-2010, 2010.
- Remer, L. A., Kaufman, Y. J., Tanre, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R. R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS aerosol algorithm, products, and validation, *J. Atmos. Sci.*, 62, 947–973, 2005.
- 615

- Roberts, G. C., Ramana, M. V., Corrigan, C., Kim, D., and Ramanathan, V.: Simultaneous observations of aerosol-cloudalbedo interactions with three stacked unmanned aerial vehicles, *P. Natl. Acad. Sci.*, 105, 7370–7375, <https://doi.org/10.1073/pnas.0710308105>, 2008.
- Robock, A.: The seasonal cycle of snow cover, sea ice and surface albedo. *Mon. Wea. Rev.*, 108, 267–285, 1980.
- 620 Saponaro, G., Kolmonen, P., Sogacheva, L., Rodriguez, E., Virtanen, T., and Leeuw, G. D.: Estimates of the aerosol indirect effect over the Baltic Sea region derived from 12 years of MODIS observations, *Atmos. Chem. Phys.*, 17(4), 3133–3143, <https://doi.org/10.5194/acp-17-3133-2017>, 2017.
- Sayer, A. M., Hsu, N. C., Bettenhausen, C., and Jeong, M.-J.: Validation and uncertainty estimates for MODIS Collection 6 “Deep Blue” aerosol data, *J. Geophys. Res.*, 118, 7864–7872, <https://doi.org/10.1002/jgrd.50600>, 2013.
- 625 Sekiguchi, M., Nakajima, T., Suzuki, K., Kawamoto, K., Higurashi, A., Rosenfeld, D., Sano, I., and Mukai, S.: A study of the direct and indirect effects of aerosols using global satellite data sets of aerosol and cloud parameters, *J. Geophys. Res.*, 108(D22), 4699, doi:10.1029/2002JD003359, 2003.
- Shao, H., and Liu, G.: Influence of mixing on evaluation of the aerosol first indirect effect, *Geophys. Res. Lett.*, 33, L14809, doi:10.1029/2006GL026021, 2006.
- 630 Stier, P.: Limitations of passive remote sensing to constrain global cloud condensation nuclei. *Atmos. Chem. Phys.*, 16(10), 6595–6607, <https://doi.org/10.5194/acp-16-6595-2016>, 2016.
- Su, W., Loeb, N. G., Xu, K. M., Schuster, G. L., and Eitzen, Z. A.: An estimate of aerosol indirect effect from satellite measurements with concurrent meteorological analysis, *J. Geophys. Res.*, 115, D18219, doi:10.1029/2010JD013948, 2010.
- 635 Tang, J., Wang, P., Mickley, L. J., Xia, X., Liao, H., Yue, X., Sun, L., and Xia, J.: Positive relationship between liquid cloud droplet effective radius and aerosol optical depth over Eastern China from satellite data, *Atmos. Environ.*, 84, 244–253, doi:10.1016/j.atmosenv.2013.08.024, 2014.
- Tripathi, S. N., Dey, S., Chandel, A., Srivastava, S., Singh, R. P., and Holben, B. N.: Comparison of MODIS and AERONET derived aerosol optical depth over the Ganga Basin, India, *Ann. Geophys.*, 23, 1093–1101, doi:10.5194/angeo-23-1093-2005, 2005.
- 640 Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, 1977.
- Vant-Hull, B., Marshak, A., Remer, L. A., and Li, Z. Q.: The effects of scattering angle and cumulus cloud geometry on satellite retrievals of cloud droplet effective radius, *IEEE T. Geosci. Remote*, 45, 1039–1045, doi:10.1109/TGRS.2006.890416, 2007.
- 645 Várnai, T. and Marshak, A.: MODIS observations of enhanced clear-sky reflectance near clouds, *Geophys. Res. Lett.*, 36, L06807, doi:10.1029/2008GL037089, 2009.
- Wall, C., Zipser, E., and Liu, C. T.: An investigation of the aerosol indirect effect on convective intensity using satellite observations, *J. Atmos. Sci.*, 71(1), 430–447, doi:10.1175/jas-d-13-0158.1., 2014.

- 650 Wang, F., Guo, J., Wu, Y., Zhang, X., Deng, M., Li, X., Zhang, J., and Zhao, J.: Satellite observed aerosol-induced variability in warm cloud properties under different meteorological conditions over eastern China, *Atmos. Environ.*, 84, 122–132, doi:10.1016/j.atmosenv.2013.11.018, 2014.
- Wang, F., Guo, J., Zhang, J., Huang, J., Min, M., Chen, T., Liu, H., Deng, M., and Li, X.: Multi-sensor quantification of aerosol-induced variability in warm clouds over eastern China, *Atmos. Environ.*, 113, 1–9, doi:10.1016/j.atmosenv.2015.04.063, 2015.
- 655 Wang, J., Lee, Y. N., Daum, P. H., Jayne, J., and Alexander, M. L.: Effects of aerosol organics on cloud condensation nucleus (CCN) concentration and first indirect aerosol effect, *Atmos. Chem. Physics*, 8(3), 9783–9818. <https://doi.org/10.5194/acpd-8-9783-2008>, 2008.
- Wang, J., Cubison, M. J., Aiken, A. C., Jimenez, J. L., and Collins, D. R.: The importance of aerosol mixing state and size-resolved composition on CCN concentration and the variation of the importance with atmospheric aging of aerosols, *Atmos. Chem. Phys.*, 10, 7267–7283, doi:10.5194/acp-10-7267-2010, 2010.
- 660 Wang, K., Liu, J., Zhou, X., Sparrow, M., Ma, M., Sun, Z., and Jiang, W.: Validation of the MODIS global land surface albedo product using ground measurements in a semidesert region on the Tibetan Plateau, *J. Geophys. Res.*, 109, D05107, <https://doi.org/10.1029/2003JD004229>, 2004.
- Wang, Z., Vane, D., Stephens, G., and Reinke, D.: CloudSat Project: Level 2 Combined Radar and Lidar Cloud Scenario Classification Product Process Description and Interface Control Document, 61 pp., California Institute of Technology, Calif, 2013.
- 665 Wen, G., Marshak, A., and Cahalan, R. F.: Impact of 3-D clouds on clear sky reflectance and aerosol retrieval in a biomass burning region of Brazil, *IEEE Geosci. Remote*, 3(1), 169–172, 2006.
- Wen, G., Marshak, A., Cahalan, R., Remer, L., and Kleidman, R.: 3-D aerosol-cloud radiative interaction observed in collocated MODIS and ASTER images of cumulus cloud fields, *J. Geophys. Res.-Atmos.*, 112(D13), D13204, doi:10.1029/2006JD008267, 2007.
- 670 Werner, F., Ditas, F., Siebert, H., Simmel, M., Wehner, B., Pilewskie, P., Schmeissner, T., Shaw, R. A., Hartmann, S., Wex, H., Roberts, G. C., and Wendisch, M.: Twomey effect observed from collocated microphysical and remote sensing measurements over shallow cumulus, *J. Geophys. Res.-Atmos.*, 119, 1534–1545, doi:10.1002/2013JD020131, 2014.
- 675 West, R. E. L., Stier, P., Jones, A., Johnson, C. E., Mann, G. W., Bellouin, N., Partridge, D. G., and Kipling, Z.: The importance of vertical velocity variability for estimates of the indirect aerosol effects, *Atmos. Chem. Phys.*, 14, 6369–6393, doi:10.5194/acp14-6369-2014, 2014.
- Wielicki, B. A., Barkstrom, B. R., Harrison, E. F., Lee, R. B. I., Smith, G. L., and Cooper, J. E.: Clouds and the Earth's radiant energy system (CERES): An Earth observing system experiment, *Bull. Amer. Meteorol. Soc.*, 77, 853–868, 1996.
- 680 Wilcox, E. M., Roberts, G., and Ramanathan, V.: Influence of aerosols on the shortwave cloud radiative forcing from north Pacific Oceanic Clouds: Results from the Cloud Indirect Forcing Experiment (CIFEX), *Geophys. Res. Lett.*, 33, L21804, doi:10.1029/2006GL027150, 2006.

- Yuan, T., Li, Z., Zhang, R., and Fan, J.: Increase of cloud droplet size with aerosol optical depth: An observation and modeling study, *J. Geophys. Res.*, 113, D04201, <https://doi.org/10.1029/2007JD008632>, 2008.
- 685 Zhang, Q., Quan, J., Tie, X., Huang, M., and Ma, X.: Impact of aerosol particles on cloud formation: Aircraft measurements in China, *Atmos. Environ.*, 45(3), 665–672, <https://doi.org/10.1016/j.atmosenv.2010.10.025>, 2011.
- Zhang, J., Reid, J., and Holben, B.: An analysis of potential cloud artifacts in MODIS over ocean aerosol optical thickness products, *Geophys. Res. Lett.*, 32, L15803, doi:10.1029/2005GL023254, 2005.
- Zhang, Z., Ackerman, A. S., Feingold, G., Platnick, S., Pincus, R., and Xue, H.: Effects of cloud horizontal inhomogeneity and
690 drizzle on remote sensing of cloud droplet effective radius: Case studies based on large-eddy simulations, *J. Geophys. Res.*, 117, D19208, doi:10.1029/2012JD017655, 2012.
- Zhang, Z., and Platnick S.: An assessment of differences between cloud effective particle radius retrievals for marine water clouds from three MODIS spectral bands, *J. Geophys. Res.*, 116, D20215, doi:10.1029/2011JD016216, 2011.
- Zuidema, P., Redemann, J., Haywood, J., Wood, R., Piketh, S., Hipondoka, M., and Formenti, P.: Smoke and Clouds above
695 the Southeast Atlantic: Upcoming Field Campaigns Probe Absorbing Aerosol's Impact on Climate, *B. Am. Meteorol. Soc.*, 97, 1131–1135, <https://doi.org/10.1175/BAMS-D-15-00082.1>, 2016.

700 **Table 1. The differences between the slope of CER versus aerosol loading (AOD for land, AI for ocean) in PCL and in overcast retrievals in case of CF > 60% ($\Delta S_{>60}$) and CF < 60% ($\Delta S_{<60}$) under three LWP bins (g m^{-2}).**

Region	20 < LWP ≤ 60		60 < LWP ≤ 100		100 < LWP ≤ 140	
	$\Delta S_{<60}$	$\Delta S_{>60}$	$\Delta S_{<60}$	$\Delta S_{>60}$	$\Delta S_{<60}$	$\Delta S_{>60}$
EC	0.03	0.036	0.039	0.059	0.068	0.089
EU	0.05	0.069	0.023	0.046	0.014	0.047
WE	0.07	0.09	0.048	0.078	0.02	0.85
ECO	0.089	0.150	0.069	0.140	0.055	0.080
EUO	0.049	0.176	0.052	0.060	0.061	0.110
WEO	0.159	0.160	0.092	0.190	0.055	0.185

Table 2. The range of LTS and RH_{CT} for low, medium and high conditions, respectively.

Region	LTS (K)			RH _{CT} (%)		
	Low	Medium	High	Low	Medium	High
EC	1.9 ~ 9.4	9.4 ~ 11.3	11.3 ~ 22.1	5.1 ~ 65.8	65.8 ~ 76.9	76.9 ~ 99.8
EU	5.8 ~ 12.1	12.1 ~ 14.1	14.1 ~ 26.3	4.7 ~ 64.4	64.4 ~ 76.4	76.4 ~ 99.8
WE	5.0 ~ 11.2	11.2 ~ 13.9	13.9 ~ 26.8	6.5 ~ 60.0	60.0 ~ 73.4	73.4 ~ 99.8
ECO	11.6 ~ 13.6	13.6 ~ 14.5	14.5 ~ 20.1	7.6 ~ 71.3	71.3 ~ 81.3	81.3 ~ 99.8
EUO	9.7 ~ 13.4	13.4 ~ 14.6	14.6 ~ 21.9	5.2 ~ 68.2	68.2 ~ 80.8	80.8 ~ 99.8
WEO	9.8 ~ 16.9	16.9 ~ 19.4	19.4 ~ 26.8	3.4 ~ 66.6	66.6 ~ 87.3	87.3 ~ 99.8






Figure 1. Regions analyzed in this study: three anthropogenic regions (EC, EU, and WE) and their adjacent oceans (ECO, EOU, and WEO).




715 Figure 2. The computed slopes of (a) CER versus aerosol loading (AOD for land, AI for ocean) (Ma et al., 2018) and (b) LWP versus aerosol loading on log-log scale over six regions, in which data are stratified according to LWP.




720 **Figure 3. Joint histograms between LWP (x axis) and CER (y axis) for EC, EU, WE, ECO, EUO, and WEO. The first and second columns show the LWP-CER joint histograms for the low and high aerosol loading (AOD for land, AI for ocean) cases, respectively. The histograms are normalized so each column sums to 1, such that the histograms show the probability of observing a specific CER, given a certain LWP. The black line indicates the mean CER at each LWP. The third column shows the difference between the polluted and clean cases.**

725



730

Figure 4. The difference of TOA albedo for polluted (AOD/AI > 75th percentile) and clean case (AOD/AI < 25th percentile) ($\Delta\alpha$) over (a) EC, (b) EU, (c) WE, (d) ECO, (e) EUO, and (f) WEO, in which data are stratified according to both LWP (x axis) and CF (y axis).



735 **Figure 5.** Cloud effective radius (CER) varies with aerosol loading (AOD for land, AI for ocean) under three liquid water path (LWP) categories (20~60, 60~100, 100~140 g m⁻²). The figures from top to bottom show the CER-AOD/AI correlation over different regions (EC, EU, WE, ECO, EUO, WEO). The circles (dots) show the mean CER from PCL (overcast) retrieval at each AOD (AI) bin in case of CF > 60% (red) and CF < 60% (blue), respectively. The slopes on log-log scale along with the total number of samples

740 (bracketed) for these four categories are also provided in each panel, in which the numbers in *italic (bold)* represent PCL (overcast)
retrieval.

745




Figure 6. The differences between CER_{2.1µm}-AOD (AI) slope and CER_{3.7µm}-AOD (AI) slope under all LWP bins over three anthropogenic regions (EC, EU, and WE) and their adjacent oceans (ECO, EUO, and WEO).






Figure 7. Cloud effective radius (CER) varies with aerosol loading (AOD for land, AI for ocean) under three liquid water path (LWP) categories (20~60, 60~100, 100~140 g m⁻²). The figures from top to bottom represent different regions (EC, EU, WE, ECO, EUO, and WEO). The blue (red) dots show the mean CER at each category of AOD (AI) in case of CER > 14 μm (< 14 μm), and the black circles indicate the results that do not differentiate CER. The slopes on log-log scale along with the total number of samples (bracketed) for these three categories are also provided in each panel.




760 **Figure 8. Probability density functions of AOD (AI) for CER > 14 μm (blue) and CER < 14 μm (red) over three anthropogenic regions (EC, EU, and WE) and their adjacent oceans (ECO, EUO, and WEO).**



765

Figure 9. The slopes of CER versus AOD (AI) on log-log scale under low, medium, and high LTS and RH_{CT} conditions, respectively, over (a) EC, (b) EU, (c) WE, (d) ECO, (e) EUO, and (f) WEO.

770



775

Figure 10. The distributions of sample number for (a) LTS and (b) RH_{CT} over land (upper; EC, EU, and WE) and ocean (lower; ECO, EUO, and WEO). Vertical lines show the mean values. Texts with the corresponding color represent the mean and standard deviation for each region.