

1

SUPPORTING INFORMATION AVAILABLE

2

3

4

5

**Inferreding the anthropogenic NO_x emission trend
over the United States during 2003 - 2017 from
satellite observations: Was there a flattening of
the emission tend after the Great Recession?**

6

Jianfeng Li¹, Yuhang Wang^{1*}

7

8

¹School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia,
USA

9

* Correspondence to Yuhang Wang (yuhang.wang@eas.gatech.edu)

10

11

12 **Table Captions**

13 Table S1. Summary of major satellite instruments for remote sensing of atmospheric NO₂ VCD in
14 the past decade

15 Table S2. Summary of satellite NO₂ TVCD products and their retrieval information

16 Table S3. Selection criteria for satellite NO₂ TVCD pixel data

17 Table S4. Summary of annual trends of AQS NO₂ surface concentrations and satellite NO₂ TVCD
18 products in each region during different periods

19 **Table S1. Summary of major satellite instruments for remote sensing of atmospheric NO₂ VCD in the past decade**

Instrument	Satellite	Launch date	End date	Operator	Equator crossing time (local time)	UV/Vis Spectral range (nm)	Spectral resolution (nm)	Swath length (km)	Nadir pixel resolution (km × km)	Global coverage (days)
SCIAMACHY	ENVISAT ¹	03/01/2002 ²	04/08/2012 ²	ESA ³	10:00 ¹	240 – 805 ⁴	0.24 – 0.48 ⁴	960 ⁵	60 × 30 ⁵	6 ⁵
GOME-2A	MetOp-A ⁶	10/19/2006 ⁶	in operation	EUMETSAT ⁷	9:30 ⁸	240 – 790 ⁸	0.26 – 0.51 ⁸	1920 before Jul. 15 th , 2013; 960 after Jul. 15 th , 2013 ⁸	80 × 40 before Jul. 15 th , 2013; 40 × 40 after Jul. 15 th , 2013 ⁸	1.5 ⁹
GOME-2B	MetOp-B ⁶	09/17/2012 ⁶	In operation	EUMETSAT	9:30 ⁸	240 – 790 ⁸	0.26 – 0.51 ⁸	1920 ⁸	80 × 40 ⁸	1.5 ⁹
OMI	EOS-Aura ¹⁰	07/15/2004 ¹⁰	In operation	NASA	13:45 ¹⁰	270 – 500 ¹¹	0.45 – 1.0 ¹¹	2600 ¹¹	24 × 13 ¹¹	1 ¹¹

20 ¹ Refer to <https://earth.esa.int/web/guest/missions/esa-operational-eo-missions/envisat>21 ² Refer to <https://en.wikipedia.org/wiki/Envisat>22 ³ The European Space Agency23 ⁴ Refer to <http://www.iup.uni-bremen.de/sciamachy/instrument/performance/index.html>24 ⁵ Refer to Boersma et al. (2008), Boersma et al. (2009), and (Lee et al., 2009)25 ⁶ Refer to <https://www.eumetsat.int/website/home/Satellites/CurrentSatellites/Metop/index.html>26 ⁷ The European Organization for the Exploitation of Meteorological Satellites27 ⁸ Refer to EUMETSAT (2015)28 ⁹ Refer to Lee et al. (2009) and Wang et al. (2017)29 ¹⁰ Refer to <https://aura.gsfc.nasa.gov/>30 ¹¹ Refer to <https://aura.gsfc.nasa.gov/omi.html>

31 **Table S2. Summary of satellite NO₂ TVCD products and their retrieval information**

NO ₂ TVCD products	Version	Available period	DOAS fitting method	Stratosphere–troposphere separation	Fitting window (nm)	Albedo / reflectance	A priori profiles	Radiative transfer model	Cloud	Uncertainty
GOME-2B	TM4NO2A (2.3)	12/20/2012 – current	Intensity fit ¹	Assimilation of satellite total slant columns in the TM4 model ^{2, 3}	405 – 465 ¹	Climatology albedo from 3 years of OMI data ⁴	TM4 (2° × 3°) ²	DAK ²	FRESCO+ (Oxygen A-band around 760 nm) ⁵	1.0 × 10 ¹⁵ molecules/cm ² + 25% ²
SCIAMACHY	QA4ECV (v1.1)	08/02/2002 – 04/08/2012	Optical Density ^{1, 6}	Assimilation of OMI total slant columns in the TM5 - MP model ^{6, 7}	425 – 465 ⁶	Climatology albedo based on SCIAMACHY ⁸	TM5-MP (1° × 1°) ⁶	DAK	FRESCO+	35% - 45% over polluted scenes; > 100% over background regions (Pacific Ocean) ⁶
GOME-2A	QA4ECV (v1.1)	02/01/2007 – 12/31/2016			405 – 465 ^{1, 6}	Climatology albedo based on GOME-2A ⁸			FRESCO+	
OMI-QA4ECV	QA4ECV (v1.1)	10/01/2004 – Current			405 – 465 ^{1, 6}	Climatology albedo from 5 years of OMI data ⁶			Improved O ₂ -O ₂ (477 nm) ⁹	
OMI-NASA	SPv3	01/01/2005 – 07/31/2017	Stepwise intensity fit with monthly averaged solar irradiance spectrum ^{1, 10}	Based on OMI total slant columns over regions with low estimated TVCD contributions (TVCD contributions less than 0.3 × 10 ¹⁵ molecules/cm ²) ¹⁰	402 – 465 ^{1, 10}	OMI climatology albedo ¹⁰	GMI (1° × 1.25°) ¹⁰	TMORAD ¹⁰	O ₂ -O ₂ (477 nm) ^{10, 11}	SPv2.1 TVCD has uncertainties of about 30% under clear-sky conditions to about 60% under cloudy conditions ¹² , and the relative difference between SPv3 and SPv2.1 is less than ~20% ¹⁰ .
OMI-BEHR ¹³	v3.0B	01/01/2005 – 07/31/2017				Based on MCD43D BRDF product (for land) and model parameterization (for ocean)	WRF-Chem (12 km)			~ 45% on average ¹⁴

32 ¹ Refer to Zara et al. (2018)33 ² Refer to Boersma et al. (2011). “TM4” is the Tracer Model, version 4. “DAK” is the Doubling-Adding KNMI (DAK) radiative transfer model.34 ³ Refer to Williams et al. (2009)35 ⁴ Refer to Kleipool et al. (2008)36 ⁵ Refer to Wang et al. (2017) and Wang et al. (2008)37 ⁶ Refer to Boersma et al. (2018)38 ⁷ Refer to Williams et al. (2017)39 ⁸ Refer to Tilstra et al. (2017)40 ⁹ Refer to Veefkind et al. (2016)41 ¹⁰ Refer to Bucsela et al. (2013), Bucsela et al. (2016), Krotkov et al. (2017), and Marchenko et al. (2015). “TMORAD” is the TMOS radiative transfer model.42 ¹¹ Refer to Acarreta et al. (2004)43 ¹² Refer to Lamsal et al. (2014), Oetjen et al. (2013), and Tong et al. (2015)44 ¹³ Refer to Laughner et al. (2018). OMI-BEHR uses the SCD from OMI-NASA SPv3 but updates inputs for the AMF calculation, such as a prior NO₂ vertical profiles and surface reflectance. Besides, OMI-BEHR only provides NO₂ TVCD over the contiguous United States (CONUS). As in this study, we used the OMI-NASA datasets archived in the OMI-BEHR product, so we only obtained OMI-NASA datasets extended to July 31, 2017.45 ¹⁴ Average uncertainty over the CONUS is calculated based on the file from <http://behr.cchem.berkeley.edu/behr/BEHR-us-uncertainty.hdf>

47

Table S3. Selection criteria for satellite NO₂ TVCD pixel data

NO ₂ TVCD products	Period	Solar zenith angle	albedo	Cloud radiance fraction	Snow or ice covered	AMFtrop/AMFgeo	Flag for retrieval success	Retrieval quality flag	Rows in swath
GOME-2B	01/01/2013 – 12/31/2017	< 80°	<= 0.3	<= 50%	No	> 0.2	Yes		All
SCIAMACHY	01/01/2003 – 12/31/2011	< 80°	<= 0.3	<= 50%	No	> 0.2	Yes		All
GOME-2A	01/01/2008 – 12/31/2016	< 80°	<= 0.3	<= 50%	No	> 0.2	Yes		All
OMI-QA4ECV ¹	01/01/2005 – 12/31/2017	< 80°	<= 0.3	<= 50%	No	> 0.2	Yes		6 - 21
OMI-NASA ¹	01/01/2005 – 12/31/2016	< 80°	<= 0.3	<= 50%			Yes	Yes	6 – 21
OMI-BEHR ¹	01/01/2005 – 12/31/2016	< 80°	<= 0.3	<= 50%			Yes	Yes	6 - 21

48

¹ Rows 6-21 are selected to remove the anomalies developed in the OMI sensor (Boersma et al., 2018; Zhang et al., 2018).

49 **Table S4. Summary of annual trends of AQS NO₂ surface concentrations and satellite NO₂ TVCD products in each region during different periods¹**

		Northeast		Midwest		South		West	
		AQS site	CONUS	AQS site	CONUS	AQS site	CONUS	AQS site	CONUS
AQS NO ₂ VMR at 13:00–14:00	2003–2011	-6.8 ± 0.7%		-6.1 ± 1.2%		-6.6 ± 0.7%		-7.6 ± 1.2%	
	2011–2017	-8.0 ± 1.2%		-6.4 ± 0.8%		-5.8 ± 0.6%		-7.2 ± 1.6%	
AQS NO ₂ VMR at 10:00–11:00	2003–2011	-6.6 ± 0.5%		-5.8 ± 1.5%		-6.5 ± 1.3%		-7.1 ± 1.6%	
	2011–2017	-7.6 ± 1.0%		-6.8 ± 0.5%		-5.7 ± 0.1%		-6.1 ± 1.1%	
SCIAMACHY	2003–2011	-17.1 ± 2.7%	-11.0 ± 3.3%	-12.9 ± 6.8%	-6.5 ± 0.8%	-9.1 ± 1.0%	-6.2 ± 1.5%	-9.1 ± 1.8%	-7.0 ± 1.4%
	2011–2017								
GOME2B	2003–2011								
	2013–2017	-11.4 ± 3.7%	-10.8 ± 3.9%	-9.9 ± 13.1%	-4.4 ± 27.2%	-8.9 ± 3.0%	-7.5 ± 3.6%	-11.8 ± 3.0%	-10.6 ± 2.3%
OMI-QA4ECV	2005–2011	-14.2 ± 6.3%	-10.6 ± 3.8%	-9.2 ± 4.2%	-8.4 ± 2.8%	-9.2 ± 2.7%	-8.2 ± 1.5%	-10.5 ± 1.6%	-8.7 ± 0.9%
	2011–2017	-18.0 ± 16.2%	-7.6 ± 4.2%	-7.6 ± 3.3%	-7.0 ± 1.7%	-4.8 ± 1.4%	-4.6 ± 1.0%	-6.4 ± 1.4%	-4.8 ± 1.2%
OMI-NASA	2005–2011	-11.8 ± 1.3%	-11.0 ± 1.8%	-10.9 ± 4.8%	-10.0 ± 4.1%	-10.0 ± 3.5%	-9.5 ± 1.9%	-10.2 ± 1.8%	-8.5 ± 0.9%
	2011–2016	-10.0 ± 4.9%	-8.5 ± 3.8%	-13.2 ± 3.2%	-9.2 ± 2.7%	0.3 ± 19.2%	-8.0 ± 5.5%	-9.0 ± 5.7%	-6.6 ± 3.9%
OMI-BEHR	2005–2011	-11.8 ± 1.8%	-10.9 ± 1.9%	-12.2 ± 7.3%	-9.8 ± 4.4%	-9.5 ± 3.1%	-8.8 ± 2.0%	-9.9 ± 1.1%	-8.2 ± 0.4%
	2011–2016	-8.2 ± 3.4%	-6.6 ± 1.7%	-27.4 ± 24.3%	-8.1 ± 3.0%	-7.2 ± 2.3%	-5.0 ± 1.3%	-13.2 ± 14.5%	-7.0 ± 4.8%

50 ¹ Annual trends are the averages of regional seasonal trends (e.g., Figure 7).

52 **References**

- 53 Acarreta, J. R., de Haan, J. F., and Stammes, P.: Cloud pressure retrieval using the O₂ - O₂
54 absorption band at 477 nm, *Journal of Geophysical Research: Atmospheres*, 109,
55 10.1029/2003JD003915, 2004.
- 56 Boersma, K. F., Jacob, D. J., Eskes, H. J., Pinder, R. W., Wang, J., and Van Der A, R. J.:
57 Intercomparison of SCIAMACHY and OMI tropospheric NO₂ columns: Observing the diurnal
58 evolution of chemistry and emissions from space, *Journal of Geophysical Research:*
59 *Atmospheres*, 113, 10.1029/2007JD008816, 2008.
- 60 Boersma, K. F., Jacob, D. J., Trainic, M., Rudich, Y., De Smedt, I., Dirksen, R., and Eskes, H. J.:
61 Validation of urban NO₂ concentrations and their diurnal and seasonal variations observed from
62 the SCIAMACHY and OMI sensors using in situ surface measurements in Israeli cities,
63 *Atmospheric Chemistry and Physics*, 9, 3867-3879, 10.5194/acp-9-3867-2009, 2009.
- 64 Boersma, K. F., Eskes, H. J., Dirksen, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool,
65 Q. L., Sneep, M., Claas, J., and Leitão, J.: An improved tropospheric NO₂ column retrieval
66 algorithm for the Ozone Monitoring Instrument, *Atmospheric Measurement Techniques*, 4, 1905-
67 1928, 10.5194/amt-4-1905-2011, 2011.
- 68 Boersma, K. F., Eskes, H. J., Richter, A., De Smedt, I., Lorente, A., Beirle, S., van Geffen, J. H.,
69 Zara, M., Peters, E., and Roozendael, M. V.: Improving algorithms and uncertainty estimates for
70 satellite NO₂ retrievals: results from the quality assurance for the essential climate variables
71 (QA4ECV) project, *Atmospheric Measurement Techniques*, 11, 6651-6678, 10.5194/amt-11-
72 6651-2018, 2018.
- 73 Bucsela, E. J., Krotkov, N. A., Celarier, E. A., Lamsal, L. N., Swartz, W. H., Bhartia, P. K.,
74 Boersma, K. F., Veefkind, J. P., Gleason, J. F., and Pickering, K. E.: A new stratospheric and
75 tropospheric NO₂ retrieval algorithm for nadir-viewing satellite instruments: applications to OMI,
76 *Atmospheric Measurement Techniques*, 6, 2607-2626, 10.5194/amt-6-2607-2013, 2013.
- 77 Bucsela, E. J., Celarier, E. A., Gleason, J. L., Krotkov, N. A., Lamsal, L. N., Marchenko, S. V.,
78 and Swartz, W. H.: OMNO2 README Document Data Product Version 3.0, NASA, 38, 2016.
- 79 EUMETSAT: GOME_FACTSHEET, Germany, 33, 2015.
- 80 Kleipool, Q. L., Dobber, M. R., de Haan, J. F., and Levelt, P. F.: Earth surface reflectance
81 climatology from 3 years of OMI data, *Journal of Geophysical Research: Atmospheres*, 113,
82 10.1029/2008JD010290, 2008.
- 83 Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J.,
84 Chan, K. L., Wenig, M., and Zara, M.: The version 3 OMI NO₂ standard product, *Atmospheric*
85 *Measurement Techniques*, 10, 3133-3149, 10.5194/amt-10-3133-2017, 2017.
- 86 Lamsal, L. N., Krotkov, N. A., Celarier, E. A., Swartz, W. H., Pickering, K. E., Bucsela, E. J.,
87 Gleason, J. F., Martin, R. V., Philip, S., and Irie, H.: Evaluation of OMI operational standard NO₂

- 88 column retrievals using in situ and surface-based NO₂ observations, Atmospheric Chemistry and
89 Physics, 14, 11587-11609, 10.5194/acp-14-11587-2014, 2014.
- 90 Laughner, J. L., Zhu, Q., and Cohen, R. C.: The Berkeley High Resolution Tropospheric NO₂
91 product, Earth System Science Data, 10, 2069-2095, 10.5194/essd-10-2069-2018, 2018.
- 92 Lee, C., Martin, R. V., van Donkelaar, A., Richter, A., Burrows, J. P., and Kim, Y. J.: Remote
93 Sensing of Tropospheric Trace Gases (NO₂ and SO₂) from SCIAMACHY, in: Atmospheric and
94 Biological Environmental Monitoring, Springer, 63-72, 2009.
- 95 Marchenko, S., Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., and Bucsela, E. J.:
96 Revising the slant column density retrieval of nitrogen dioxide observed by the Ozone
97 Monitoring Instrument, Journal of Geophysical Research: Atmospheres, 120, 5670-5692,
98 10.1002/2014JD022913, 2015.
- 99 Oetjen, H., Baidar, S., Krotkov, N. A., Lamsal, L. N., Lechner, M., and Volkamer, R.: Airborne
100 MAX - DOAS measurements over California: Testing the NASA OMI tropospheric NO₂
101 product, Journal of Geophysical Research: Atmospheres, 118, 7400-7413, 10.1002/jgrd.50550,
102 2013.
- 103 Tilstra, L. G., Tuinder, O. N. E., Wang, P., and Stammes, P.: Surface reflectivity climatologies
104 from UV to NIR determined from Earth observations by GOME - 2 and SCIAMACHY, Journal
105 of Geophysical Research: Atmospheres, 122, 4084-4111, 10.1002/2016JD025940, 2017.
- 106 Tong, D., Lamsal, L., Pan, L., Ding, C., Kim, H., Lee, P., Chai, T., Pickering, K. E., and Stajner,
107 I.: Long-term NO_x trends over large cities in the United States during the great recession:
108 Comparison of satellite retrievals, ground observations, and emission inventories, Atmospheric
109 Environment, 107, 70-84, 10.1016/j.atmosenv.2015.01.035, 2015.
- 110 Veefkind, J. P., de Haan, J. F., Sneep, M., and Levelt, P. F.: Improvements to the OMI O₂-O₂
111 operational cloud algorithm and comparisons with ground-based radar-lidar observations, Atmos.
112 Meas. Tech., 9, 6035-6049, 10.5194/amt-9-6035-2016, 2016.
- 113 Wang, P., Stammes, P., van der A, R., Pinardi, G., and van Roozendael, M.: FRESCO+: an
114 improved O₂ A-band cloud retrieval algorithm for tropospheric trace gas retrievals, Atmos.
115 Chem. Phys., 8, 6565-6576, 10.5194/acp-8-6565-2008, 2008.
- 116 Wang, Y., Beirle, S., Lampel, J., Koukouli, M., De Smedt, I., Theys, N., Ang, L., Wu, D., Xie, P.,
117 and Liu, C.: Validation of OMI, GOME-2A and GOME-2B tropospheric NO₂, SO₂ and HCHO
118 products using MAX-DOAS observations from 2011 to 2014 in Wuxi, China: investigation of the
119 effects of priori profiles and aerosols on the satellite products, Atmospheric Chemistry and
120 Physics, 17, 5007, 10.5194/acp-17-5007-2017, 2017.
- 121 Williams, J. E., Scheele, M. P., van Velthoven, P. F. J., Cammas, J.-P., Thouret, V., Galy-Lacaux,
122 C., and Volz-Thomas, A.: The influence of biogenic emissions from Africa on tropical
123 tropospheric ozone during 2006: a global modeling study, Atmospheric Chemistry and Physics, 9,
124 5729-5749, 10.5194/acp-9-5729-2009, 2009.

125 Williams, J. E., Boersma, K. F., Sager, P. L., and Verstraeten, W. W.: The high-resolution version
126 of TM5-MP for optimized satellite retrievals: description and validation, Geoscientific Model
127 Development, 10, 721-750, 10.5194/gmd-10-721-2017, 2017.

128 Zara, M., Boersma, K. F., De Smedt, I., Richter, A., Peters, E., Van Geffen, J. H. G. M., Beirle,
129 S., Wagner, T., Van Roozendael, M., and Marchenko, S.: Improved slant column density retrieval
130 of nitrogen dioxide and formaldehyde for OMI and GOME-2A from QA4ECV: intercomparison,
131 uncertainty characterization, and trends, Meas. Tech. Discuss, 1-47, 10.5194/amt-11-4033-2018,
132 2018.

133 Zhang, R., Wang, Y., Smeltzer, C., Qu, H., Koshak, W., and Boersma, K. F.: Comparing OMI-
134 based and EPA AQS in situ NO₂ trends: towards understanding surface NO_x emission changes,
135 Atmospheric Measurement Techniques, 11, 3955-3967, 10.5194/amt-11-3955-2018, 2018.

136