

Interactive comment on “Investigation of the global methane budget over 1980–2017 using GFDL-AM4.1” by Jian He et al.

Anonymous Referee #1

Received and published: 5 August 2019

The work describes CH₄ budget using an atmospheric chemistry-transport model that is developed at the GFDL. The authors have taken in to account all possible causes of variabilities in CH₄ budget, such as the emissions and loss due to tropospheric hydroxyl (OH). As shown in the manuscript, OH variability is of as much importance as the emissions in explaining the CH₄ growth rate variabilities in different decades in the period of 1980s to 2010s. The manuscript is generally well written. However, I felt toward the end of the manuscript is a bit of stretch and could be reduced (I have made some suggestions in my specific comments). The manuscript can be accepted after a major revision.

Specific comments: Line 49 & 62ff: can the growth rate discussions in the introduction be made concise and put together at one place.

C1

Line 80ff: I think there are other prominent inverse modelling results trying to explain the recent regrowth of CH₄ concentrations.

Line 135-137: This is a quite strange statement. After reading the whole manuscript I do not believe you have tried to address these couple of issues to a great extent. May be remove?

Line 156: Not from wetchart? I mean does wetchart not have IAV?

Line 206: Not clear if this is after LNOx scaling? please make this statement precise (e.g., Control).

Line 249ff: "The meridional curve" needed some clarifications here, e.g., selected sites within a latitude band to get the mean CH₄ at 5 different latitude bands or something like that.

Line 296: Sometimes the sites like Key Biscayne are sampled by moving the model grids to the ocean side. You might check that out.

Line 315ff: The tropical bias in all HIPPO is a bit strange! Not OH but transport (or emissions)? I am suspecting this because the bias due to OH would appear at all altitudes (timescale ~1yr), because the bias is in the lower troposphere, if the vertical transport is slow, you would find more CH₄ is accumulated in the lower troposphere (timescale~week)

Line 326: do you run CH₃CCl₃ & SF₆?, say within the CCM framework?

Line 346ff: suggesting too much emissions in the NH, where most of Anthro emissions are...May be you can test this better by site-level comparisons.

Line 353ff: I cannot find this 1 year mismatch (please be clear), instead I find a persistent offset during 1984-1991 (how the major and minor ticks marked in Fig. 5; the labeled ticks only should be major?)

Line 369: How can you say that? I thought your optimization was not good for this

C2

period, because the number of observation sites may not have covered the global reasonably well. I mean biased high toward the NH. Could you check how many SH sites you have data before 1988.

Line 374: Most likely due to an overestimation of China emissions (e.g., Saeki and Patra, GOSL, 2017, and references therein) (regional inversion is needed for adjusting such regional emission biases)

Line 378: "...which is also a remote site" and remote from China emissions

Lines 394ff: I am not very sure if the comparison with GOSAT/SCIA are adding any values to this work. Better be kept aside for a full paper, unless the reasons for the mean offsets are figured out and discussed. For instance you could compare your results with the ACE-FTS data to find out if there is any bias in the stratospheric CH₄ as there is no significant offsets in the tropospheric CH₄ is seen in comparison with surface data and HIPPO.

Line 426ff: The emission increase in the 1990s is apparently linked to OH increase in AM; which sector can provide this extra emissions. I think this result is very different from what I have seen in the literature, and thus needing some explanation. Surprisingly, the emission increase rate in the 1990s is greater than the recent regrowth period.

Lines 484ff: The discussions using Fig 9-11 aren't that interesting as presented. I would recommend the authors to move these plots to the supplement or show 1-2 panels in the main text; for example all the 4 panels in Fig 9 & 10 are essentially showing very similar distributions. The S0Aopt and S0Wopt are also showing similar behaviour. This is mainly because the emission (E)-a priori emissions are the same in both the simulations, and the correction emissions Del-E following Anthropogenic or Wetland emission patterns only play minor role.

I am actually curious if you could use some of the continental sites, e.g., NWR, LEF,

C3

SGP or TAP, and use the model-measurement comparisons to say whether the S0Aopt or S0Wopt are more realistic.

Line 519: Such high correlations are a bit surprising, if I see the lines in Fig. 12. For example AGR show -ve trend, yet show positive correlation. How is that possible?

Line 633: This is similar to the essential conclusion in some other publications as well, where ENE and Animals were made responsible for the post-2006 CH₄ growth rate. I guess it is extremely difficult to separate emissions from Animals and Wetlands by ¹³C signature in CH₄.

Lines 638ff: I am curious if inconsistency between the tropospheric OH and CH₄-loss by OH are arising from the spin-up. Did you spun-up the simulations using different OH from the 1970s?

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-529>, 2019.

C4