

Supplement to “Technical Note: Reanalysis of Aura MLS Chemical Observations”

Quentin Errera¹, Simon Chabrillat¹, Yves Christophe¹, Jonas Debosscher¹, Daan Hubert¹, William Lahoz^{2,†}, Michelle L. Santee³, Masato Shiotani⁴, Sergey Skachko⁵, Thomas von Clarmann⁶, and Kaley Walker⁷

¹Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

²Norsk Institutt for Luftforskning, NILU, Norway

³Jet Propulsion Laboratory, California Institute of Technology, USA

⁴Research Institute for Sustainable Humanosphere, Kyoto University, Japan

⁵Environment and Climate Change Canada

⁶Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research, Germany

⁷Department of Physics, University of Toronto, Canada

[†]Deceased, 1 April 2019

Correspondence: Quentin Errera (quentin.errera@aeronomie.be)




Figure S1. As Fig. 5 but between 60°S-30°S.




Figure S2. As Fig. 5 but between 30°S-30°N.




Figure S3. As Fig. 6 but between 60°S-30°S.




Figure S4. As Fig. 6 but between 30°S-30°N.




Figure S5. As Fig. 8 but in the outer Antarctic polar vortex between (75°S-60°S of equivalent latitude).




Figure S6. As Fig. 8 but in the outer Arctic polar vortex between (60°N-75°N of equivalent latitude).




Figure S7. As Fig. 8 but in the outer Arctic polar vortex between (75°N-90°N of equivalent latitude).




Figure S8. As Fig. 10 but between 90°S-60°S