



Modeling of Gas-Wall Partitioning of Organic Compounds Using a Quantitative Structure-Activity Relationship

Sanghee Han, Myoseon Jang, and Huanhuan Jiang

Department of Environmental Engineering Science, University of Florida, Gainesville, Florida, USA

5 Correspondence to: Myoseon Jang (mjang@ufl.edu)

Abstract. This study streamlines modeling of the gas–wall process (GWP) of semivolatile organic compounds (SVOC) by predicting gas–wall equilibrium partitioning constant ($K_{w,i}$) and accommodation coefficient ($\alpha_{w,i}$) of SVOC(i) using a quantitative structure–activity relationship. PaDEL-Descriptor, software that calculates molecular descriptors, is employed to obtain physicochemical parameters (i.e., hydrogen bond acidity ($H_{a,i}$), hydrogen bond basicity ($H_{b,i}$), dipolarity/polarizability (S_i), and polarizability (α_i)) of SVOC(i). For the prediction of $K_{w,i}$, activity coefficients ($\gamma_{w,i}$) of SVOC(i) to the chamber wall are semiempirically predicted using chamber data in the form of a polynomial equation coupled with the physicochemical parameters. $\gamma_{w,i}$ of various SVOCs differ in functionalities and molecular sizes ranging from 10^0 to 10^4 . We conclude that the estimation of $\gamma_{w,i}$ is essential to improve the prediction of $K_{w,i}$. To predict the impact of relative humidity (RH) on GWP, each coefficient in the polynomial equation for $\ln(K_{w,i})$ was correlated to RH. Increasing RH enhanced GWP significantly for all polar SVOCs. For example, the predicted $K_{w,i}$ of 1-heptanoic acid increased more than three times (from 0.58 to 1.96) by increasing RH from 0.4 to 0.75 due to the reduction in $\gamma_{w,i}$. The characteristic time for GWP are estimated using $K_{w,i}$ and $\alpha_{w,i}$ to evaluate the effect of GWP on secondary organic aerosol (SOA) mass. It might be significant in the absence of inorganic aerosol, but insignificant in the presence of electrolytic salts, where aqueous reactions dominate SOA growth.

1. Introduction

20 Organic Aerosol (OA) can be produced via the atmospheric secondary process of reactive hydrocarbons which emitted from both vegetation and anthropogenic sources as well as the emission from primary sources such as fuel combustion, industries, and vehicles into the ambient air. Secondary organic aerosol (SOA), generated by atmospheric process of reactive hydrocarbons, constitutes a large proportion (up to 40%) of OA in the ambient air (Hallquist et al., 2009; Volkamer et al., 2006) and it can significantly impact on climate (Seinfeld and Pandis, 2016), visibility (Park et al., 2003), and human health (Cohen et al., 2017). Thus, a large effort has given to the prediction of SOA budget in regional and global scales (Carlton et al., 2009; Volkamer et al., 2006; Barsanti et al., 2017). SOA models have been developed based on the mass balance of the hydrocarbon mass consumption during the atmospheric process. The prediction of SOA formation has been approached using a partitioning-based model with semiempirical parameters that are obtained from several semivolatile surrogate products (Odum et al., 1996; Donahue et al., 2006). Historically, the Community Multi-scale Air Quality Model (CMAQ) tends to predict SOA mass lower than that observed because of missing precursors and reaction mechanisms (Barsanti et al., 2013), especially during the summer (Appel et al., 2017). Hence, significant efforts have been made to add in-particle chemistry into SOA models to allow the formation of nonvolatile oligomers in the aerosol phase (Jang et al., 2002; Cao and Jang, 2010; McNeill et al., 2012; Im et al., 2014; Beardsley and Jang, 2016; Zhou et al., 2019).

Recent studies have also reported that the underestimated SOA mass in the model prediction can be attributed to the chamber because oxygenated products can partition on the reactor wall. Consequently, this negative bias in the model parameters can mislead modeling efforts to predict SOA formation at regional scales. McMurry and Grosjean (1985) was the first to address



the loss of gaseous compounds of very volatile organic species (i.e., *n*-butane, *n*-pentane, and toluene) and atmospherically important inorganic tracers (i.e., NO, NO₂, ozone, and SO₂) on the wall. The gas–wall process (GWP) of Semivolatile Organic Compounds (SVOCs) has been shown only recently to constitute a significant potential bias in atmospheric chamber studies and has been noted as a source of substantial underestimation of SOA burdens. For example, Zhang et al. (2014) reported that the GWP can lead to a SOA yield lower by a factor of 1.1 to 4.2 and particularly, that of toluene was found from 2.1 to 4.2. La et al. (2016) indicated that the SOA yield inferred from a chamber study can be underrated by more than 50% in experiments and explicit model simulations for alkane and alkene series. To date, the influence of GWP on the predicted SOA yields remains controversial due to the limitations of the experimental database, which measures the vapor concentration. The accurate measurement of vapor concentration in the initial point is burdensome, without the delay attributable to the time required to finish injecting organic species into a chamber.

The gas–wall partitioning coefficient ($K_{OM,i}$, m³ μg⁻¹) of SVOC(*i*) to the absorbing organic matter (OM_{wall}) on the Teflon film and the mass accommodation coefficient ($\alpha_{w,i}$) are key parameters to predict GWP of various SVOCs. The unitless partitioning coefficient ($K_{w,i}$) is estimated by multiplying $K_{OM,i}$ by the concentration of OM_{wall} (mg m⁻³). $K_{w,i}$ is influenced by OM_{wall} , which can be altered by the chamber history (i.e., the fresh wall and the aged wall). The vapor pressure of SVOC(*i*) ($p_{L,i}^{\circ}$) is frequently employed to predict multiphase partitioning of SVOCs. For example, a simple relationship between the gas-particle partitioning coefficient ($K_{p,i}$) of SVOCs and their saturated vapor pressures (Pankow, 1994; Im et al., 2014) was employed to predict $K_{p,i}$. $p_{L,i}^{\circ}$ is useful in predicting $K_{p,i}$ for a homologous series that is similar in activity coefficient ($\gamma_{p,i}$). However, the variation of $\gamma_{p,i}$ of various SVOCs in different functionalities can be large, ranging from 10⁰ to 10³ and it can significantly influence the predictability of $K_{p,i}$ (Jang et al., 1997; Jang and Kamens, 1998; Beardsley and Jang, 2016). Similar to $K_{p,i}$, several studies have attempted to predict $K_{OM,i}$ using SVOC(*i*)’s saturated concentration (C_i^*) calculated as:

$$C_i^* = \frac{MW_{OM} p_{L,i}^{\circ} \gamma_{w,i}}{RT} \quad (1)$$

where MW_{OM} is the mean molecular weight of the organic matter on the Teflon film, R is the gas constant (8.21 × 10⁻⁵ m³ atm K⁻¹ mol⁻¹), and T (K) is the temperature, and $p_{L,i}^{\circ}$ (atm) assuming that the activity coefficient ($\gamma_{w,i}$) of SVOCs to the OM_{wall} is one (Krechmer et al., 2016). However, the $\gamma_{w,i}$ values of different SVOCs on the chamber wall can vary depending on SVOC’s polarities and molecular size and influence of $K_{OM,i}$. SOA products originating from the oxidation of hydrocarbons are diverse in functionalities. For example, the oxygen to carbon (O:C) ratio of α -pinene products is 0.43 on average (Zhang et al., 2015a; Chen et al., 2011) but that of isoprene products is about 0.8 (Bertram et al., 2011; Chen et al., 2011; Kuwata et al., 2013). The variation of the O:C ratios of products is large even in an individual SOA. For example, the O:C ratios of 1,3,5-trimethylbenzene SOA products range from 0.11 to 3.0 (Jenkin et al., 2012; Im et al., 2014; Zhou et al., 2019). The hygroscopicity of the organic-coated Teflon wall also can influence $\gamma_{w,i}$ (Krechmer et al., 2016; Huang et al., 2018). To improve the accuracy in the prediction of $K_{OM,i}$, the estimation of $\gamma_{w,i}$ is called for.

$\alpha_{w,i}$ is the fraction that reversible uptake of a gas-phase species will occur upon collision with the chamber wall. $\alpha_{w,i}$ can vary from 1 to below 10⁻⁷ depending on volatility of the species and compound of surface (Brune, 2019). The magnitude of $\alpha_{w,i}$ is also disputably estimated semiempirically using chamber data as a range from 10⁻⁵ to 10⁻⁸ (Bian et al., 2015; Zhang et al., 2015b; Krechmer et al., 2016).

A quantitative structure–activity relationship (QSAR) approach was applied for the first time in this study to predict $K_{w,i}$ and $\alpha_{w,i}$. The QSAR theory was traditionally developed in the form of a linear solvation energy relationship (LSER) (Alan, 1992; Puzyn et al., 2010), by predicting the solvent effect on the rate or equilibrium constant of an analyte using the multi regression



75 equation associate with physicochemical parameters. The LSER-based prediction of a solvation property (SP) is typically expressed as:

$$\log SP = c + aH_{d,i} + bH_{a,i} + eE_i + sS_i + \dots \quad (2)$$

where $H_{d,i}$, $H_{a,i}$, E_i , and S_i indicate hydrogen bond acidity, hydrogen bond basicity, excess molar refraction, and dipolarity/polarizability, respectively (Abraham and McGowan, 1987; Abraham et al., 1991; Platts et al., 1999). The
80 physicochemical parameters are not limited to the terms in Eq. (2) and can be extended to additional parameters depending upon the properties of SP, series of solutes, and the medium. QSAR has also been applied widely in environmental studies, such as partitioning of an organic compound in airborne particles (Jang and Kamens, 1999; Arp and Goss, 2009; Endo and Goss, 2014).

In this study, the estimation of $\gamma_{w,i}$ was performed using the QSAR approach, which utilizes physicochemical parameters
85 such as $H_{d,i}$, $H_{a,i}$, E_i , S_i , and polarizability (α_i) to improve the predictability of $K_{w,i}$. Each coefficient in the QSAR-based equation used to predict $K_{w,i}$ was correlated to relative humidity (RH) using chamber data under different RH levels. To assess GWP's significance, the characteristic time of GWP (τ_{GWP}) calculated using $K_{w,i}$ and $\alpha_{w,i}$ was compared with the characteristic times of important atmospheric processes, such as gas-particles partitioning, gas-phase oxidation, and aerosol phase reactions.

90 2. Experiment

2.1. Chamber experiment

The University of Florida Atmospheric PHotochemical Outdoor Reactor (UF-APHOR) chamber is bisected into East and West chambers by an airtight door in the middle. Each chamber's volume is 52 m³ with an 86 m² surface area. Table 1 summarizes the experimental conditions of the UF-APHOR chamber data and the literature data (Matsunaga and Ziemann, 2010; Yeh and
95 Ziemann, 2015) that were used to develop the model to predict $K_{w,i}$ and $\alpha_{w,i}$ under different levels of RH. The experimental procedure is described in Sect. S1. To reduce the delay attributable to inject chemicals into the chamber, a cocktail of the organic mix (Table 2) was introduced to the chamber using the two chambers (Fig. S1). The East chamber was used to inject chemicals with a typical manifold heated under a clean air stream. Then, the organic vapor in the East chamber was transferred to the West chamber through the door between the two chambers for 10 minutes. The large size of UF-APHOR benefits the
100 increase in time to reach equilibrium. The impact of the surface area (A) to volume (V) ratio of the chamber on $K_{w,i}$ and GWP equilibrium time will be discussed in Sect. 4.5 and 4.6. The SVOC vapor was collected from the West chamber using a 40-cm 5-channel annular denuder immediately after closing the door between the two chambers (Sect. S1). The concentrations of SVOC vapor were corrected for the chamber dilution by monitoring the concentration of CCl₄ using gas chromatography with a flame ionization detector (GC-FID, Hewlett-Packard 5890, Palo Alto, CA, USA). To ensure particle formation, the particle
105 size distribution was monitored using a scanning mobility particle sizer (SMPS, TSI 3080, Shoreview, MN, USA) and a condensation particle counter (CPC, TSI 3022A, Shoreview, MN, USA).

2.2. Chemical analysis

The organic vapor was collected using a 40-cm 5-channel annular denuder coated with XAD-4 resin powder (Gundel et al., 1995; Leach et al., 1999). Details of the sampling and workup procedures, chemical purity, and analytical precision are reported
110 in Sect. S1 in SI. In brief, the denuder sample was extracted with dichloromethane. Extracted samples were concentrated with an evaporator (Heidolph Rotary Evaporator Laborota 4001, Schwabach, Germany) at 80°C. The SVOC's concentrations were



analyzed with a gas chromatography/mass spectrometer (GC/MS) (Varian 3800/2000 GC/MS, Palo Alto, CA, USA). Six deuterated polycyclic aromatic hydrocarbons (naphthalene-d₈, acenaphthylene-d₁₀, anthracene-d₁₀, fluoranthene-d₁₀, pyrene-d₁₀, and chrysene-d₁₂) were used as internal standards.

115 2.3. Absorbing organic matter on the Teflon film and its hygroscopicity

The functional-group's distribution of the nonvolatile organic matter adsorbed on the Teflon film ($M_{wall-OM}$) was analyzed using a Fourier Transform Infrared (FTIR) spectrometer (Nicolet Magma 560, Madison, WI, USA). $M_{wall-OM}$ was collected by wiping the Teflon film (surface area: 20 cm × 20 cm) with an acetone-drenched nylon filter. The solution extracted with 10 mL of acetone was concentrated by clean air and placed on a silicon FTIR disc (13 mm × 2 mm, Sigma-Aldrich, St. Louis, MO, USA). The mass of collected $M_{wall-OM}$ was determined using the mass difference between the measure taken before and after placing $M_{wall-OM}$ on the FTIR disk by an analytical balance (Mettler Toledo MX5 microbalance, Columbus, OH, USA). The FTIR spectrum of $M_{wall-OM}$ (Fig. 1) is decoupled into the functional bends (i.e., C–H stretching, O–H stretching (water, alcohol, and carboxylic acid), C=O stretching (i.e., carboxylic acid and carbonyls), C=C stretching, H₂O bending, ester C–O) using a curve-fitting method assuming that each bend has a Gaussian shape (Li et al., 2016). The construction of the functional composition was calculated using the peak area of each decoupled FTIR bend and the database of the relative intensity of various FTIR bends originating from various pure compounds.

3. Model Description

The structure of the predictive model for $K_{w,i}$ and $\alpha_{w,i}$ is illustrated in Fig. 2. $K_{w,i}$ and $\alpha_{w,i}$ are predicted in the form of a polynomial equation coupled with QSAR descriptors using the experimental data that measured gas-phase concentrations of SVOCs in the UF-APHOR chamber. The GWP is assumed as a partitioning of SVOCs to OM_{wall} , which consists of the organic composition of OM_{wall} ($M_{wall-OM}$) and the water content ($M_{wall-water}$) associated with $M_{wall-OM}$. The permeability of SVOCs through the Teflon film is assumed to be negligible.

The Teflon film is highly crystalline with high density (2.15 kg m⁻³) (Leivo et al., 2004). Thus, the Teflon film has low permeability to liquid, organic vapors, and moisture (Kaur et al., 2001). Teflon film is more hydrophobic than aliphatic compounds. For example, the solubility parameter of Teflon film (FEP, Fluorinated Ethylene Propylene) based on Hansen's parameter is only 12.7, that of polyethylene film (or long-chain alkane) is 17.6, and that of ethylene glycol is 24.8 (Barton, 2017). Thus, wax-like long-chain alkanes presumably have a higher affinity to Teflon film than polar compounds. *n*-Alkanes are common species in both primary emissions (combustion of fossil fuels) and biogenic emissions (Simoneit, 1989; Bi et al., 2003; Lyu et al., 2019). These long-chain alkanes can enter the chamber during the ventilation process, and they can also constantly intrude the chamber via chamber dilution. Based on FTIR spectrum (Fig. 1), $M_{wall-OM}$ is hydrophobic, showing a low O:C ratio. It seems that polar products that are created during chamber studies (i.e., SOA formation) and partition to the chamber wall degrade more quickly than wax-like alkanes under ambient sunlight. Oxygenated compounds such as aldehydes and conjugated alkenes are generally more labile for photolysis and atmospheric oxidants than long-chain alkanes. The chemical characteristics of OM_{wall} will be discussed more in Sect. 4.1.

145 3.1 Gas-wall process based on absorption-desorption kinetics

SVOC(*i*)'s gas-wall partitioning is described by absorption-desorption kinetics and can be expressed as:





in which $k_{on,i}$ and $k_{off,i}$ are the SVOC's rate constants for deposition and evaporation (desorption), respectively. Based on a mass balance, SVOC's steady-state concentration in the gas phase ($C_{g,i}$, $\mu\text{g m}^{-3}$) is expressed as:

$$150 \quad \frac{dC_{g,i}}{dt} = -k_{on,i}C_{g,i} + k_{off,i}C_{w,i} \quad (4)$$

where $C_{w,i}$ is SVOC(*i*)'s concentration ($\mu\text{g m}^{-3}$) on the Teflon wall. The sum of $C_{g,i}$ and $C_{w,i}$ is denoted as $C_{T,i}$. From the analytical solution of Eq. (4), $C_{g,i}$ can be expressed as follows:

$$C_{g,i} = \frac{k_{on,i}C_{T,i}}{k_{on,i}+k_{off,i}} e^{-(k_{on,i}+k_{off,i})t} + \frac{k_{off,i}C_{T,i}}{k_{on,i}+k_{off,i}} \quad (5)$$

At equilibrium ($\frac{dC_{g,i}}{dt} = 0$), SVOC(*i*)'s $K_{OM,i}$ onto OM_{wall} is described in the form of traditional gas-particle partitioning as (Pankow, 1994):

$$155 \quad K_{OM,i} = \frac{7.501RT}{10^9 MW_{OM} \gamma_{w,i} p_{L,i}} = \frac{C_{w,i}}{C_{g,i} OM_{wall}} = \frac{k_{on,i}}{k_{off,i} OM_{wall}} \quad (6)$$

R is the ideal gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$) and T is the temperature (K). MW_{OM} is OM_{wall} 's average molecular weight, and $p_{L,i}^\circ$ is liquid vapor pressure (mmHg) of SVOC(*i*). $K_{w,i}$ is described as:

$$K_{w,i} = K_{OM,i} OM_{wall} = \frac{k_{on,i}}{k_{off,i}} \quad (7)$$

160 By applying $K_{w,i}$ to Eq. (5), $C_{g,i}$ is rewritten as:

$$C_{g,i} = \frac{K_{w,i} C_{T,i}}{K_{w,i} + 1} e^{-k_{on,i} \left(1 + \frac{1}{K_{w,i}}\right) t} + \frac{C_{T,i}}{K_{w,i} + 1} \quad (8)$$

SVOC's diffusion from the gas phase to the Teflon wall was assumed to be the only significant mechanism of the vapor loss to the wall in a well-stirred chamber. $k_{on,i}$ is a fractional loss rate (McMurry and Grosjean, 1985) expressed as:

$$k_{on,i} = \left(\frac{A}{V}\right) \frac{\alpha_{w,i} \bar{v}_i / 4}{1 + \frac{\pi \alpha_{w,i} \bar{v}_i}{8(K_e D)^{1/2}}} \quad (9)$$

165 \bar{v}_i is the gas molecules' mean thermal speed. D ($1.0 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$) and K_e (0.12 s^{-1}) are referred to as the diffusion coefficient and coefficient of eddy diffusion applied as a fixed number, respectively. Increasing the A/V ratio can increase $k_{on,i}$ and thus $K_{w,i}$ suggesting that GWP is higher with a smaller chamber.

3.2 Prediction of $K_{w,i}$ and $\alpha_{w,i}$ using QSAR

By taking the logarithm, the theoretical $K_{w,i}$ is described as:

$$170 \quad \ln(K_{w,i}) = -\ln(\gamma_{w,i}) - \ln(p_{L,i}^\circ) + \ln\left(\frac{7.501RT OM_{wall}}{10^9 MW_{OM}}\right) \quad (10)$$

In this study, the term $\ln(\gamma_{w,i})$ was predicted by the QSAR approach using the physicochemical descriptors $H_{d,i}$, $H_{a,i}$, S_i , E_i , and α_i . Then, Eq. (10) can be rewritten as:

$$\ln(K_{w,i}) = -(a_p H_{d,i} + b_p H_{a,i} + s_p S_i + e_p E_i + r_p \alpha_i + c_p) - \ln(p_{L,i}^\circ) + \ln\left(\frac{7.501RT OM_{wall}}{10^9 MW_{OM}}\right) \quad (11)$$

For each compound, descriptors ($H_{d,i}$, $H_{a,i}$, S_i , E_i , and α_i) were obtained using PaDEL-Descriptor, (Yap, 2011) and $p_{L,i}^\circ$ was



175 calculated with group contributions (Stein and Brown, 1994; Zhao et al., 1999).

The prediction of $\alpha_{w,i}$, which is dependent on SVOC's physicochemical properties and the chamber's characteristics (A/V), also can be approached by using QSAR as follows:

$$\ln(\alpha_{w,i}) = a_d H_{d,i} + b_d H_{a,i} + s_d S_i + e_d E_i + r_d \alpha_i + c_d \quad (12)$$

$K_{w,i}$ and $\alpha_{w,i}$ were obtained in the form of a polynomial equation by fitting the calculated $C_{g,i}$ (Eq. (8)) to the observed $C_{g,i}$. The significance of the coefficient associated with each QSAR parameter in Eq. (11) and Eq. (12) was determined based on the statistical values (Sect. S3). The estimation of the characteristic time (τ) helps understand the role of $K_{w,i}$ and $\alpha_{w,i}$ in GWP for various SVOCs because τ_{GWP} indicates the way SVOC can reach equilibrium quickly. As shown in Eq. (5), τ_{GWP} can be calculated from $k_{on,i}$ and $k_{off,i}$ as follows:

$$\tau_{GWP} = \frac{1}{k_{on,i} + k_{off,i}} \quad (13)$$

185 4. Results and Discussion

4.1. Chemical characteristics of OM_{wall}

As seen in Fig. S2 in Sect. S2, the change in the functional-group distributions in OM_{wall} is small year around. Therefore, the composition of $M_{wall-OM}$ is fixed as seen in Fig. 1. Table S1 shows the functional-group distribution of $M_{wall-OM}$. The O:C ratio of $M_{wall-OM}$ is 0.27 supporting that $M_{wall-OM}$ is hydrophobic. The amount of OM_{wall} was set to 18.52 mg m^{-3} by averaging its measurements. $M_{wall-water}$ was determined using FTIR spectra obtained by interfacing the FTIR spectrometer with a specially fabricated, RH-controlling flow tube (Fig. S3 in Sect. S2) (Jang et al., 2010; Beardsley et al., 2013; Zhong and Jang, 2014). To determine MW_{OM} , the number of functional groups within the constrained functional distribution is optimized with the UNIQUAC Functional-group Activity Coefficients (UNIFAC) model by fitting the activity coefficient of water in OM_{wall} to RH. The elemental composition of $M_{wall-OM}$ is determined as $C_{15}O_4H_{24}$ using the functional composition and MW_{OM} . A detailed description of the estimation of chemical characteristics of $M_{wall-OM}$ and $M_{wall-water}$ is found in Sect. S2.

4.2. Impact of humidity on GWP

The hygroscopicity of OM_{wall} can impact on $K_{w,i}$ due to the changing $M_{wall-water}$ as a function of RH. To predict the influence of RH on $K_{w,i}$, each coefficient in the QSAR-based polynomial equation was correlated to RH using the chamber data obtained under the three different RH levels. The two datasets obtained at RH = 0.40 (Oct 8, 2018) and RH = 0.75 (May 15, 2018) from the UF-APHOR chamber (this study) and the data under the dry condition (RH < 0.001) were obtained from the literature (Yeh and Ziemann, 2015; Matsunaga and Ziemann, 2010). The optimal coefficients of a multilinear regression model for $K_{w,i}$ and $\alpha_{w,i}$ were determined by the backward elimination procedure considering the adjusted R^2 and p -value. Table S2 summarizes the resulting linear regression coefficients for each experiment to predict $\ln(K_{w,i})$. Parameters $H_{d,i}$, $H_{a,i}$, S_i , and α_i were chosen as independent variables for $K_{w,i}$ (Table S2). The $K_{p,i}$ was related exponentially to RH, as reported in previous studies (Jang and Kamens, 1998; Pankow et al., 1993). Fig. S4 illustrates the correlation between coefficient of each physicochemical parameter to predict $\ln(\gamma_{w,i})$ and RH. Thus, each coefficient in the polynomial equation for $\ln(K_{w,i})$ was correlated linearly to RH using the three sets of experimental data, as follows:

$$\ln(K_{w,i}) = (3.09RH - 5.17)H_{d,i} + (2.71RH - 2.80)H_{a,i} - (0.01RH + 0.19)\alpha_i + (1.73RH - 1.31)S_i + 9.00 - \ln(p_{L,i}^\circ) + \ln\left(\frac{7.501RTOM_{wall}}{10^9 MW_{OM}}\right). \quad (14)$$



The prediction is that $\ln(K_{w,i})$ was correlated positively with RH for QSAR descriptors $H_{a,i}$, $H_{d,i}$, and S_i but not for α_i . The $\ln\left(\frac{7.501RTOM_{wall}}{10^9MW_{OM}}\right)$ term slightly increased with increasing RH due to the variation in the MW_{OM} , calculated as a function of RH using hygroscopicity data (Fig. S3).

Fig. 3 illustrates the tendency of $C_{g,i}$ at the two different RH levels (0.4 and 0.75) over the course of the chamber experiment (May 15, 2018 and October 08, 2018). Fig. 4 shows $K_{w,i}$ prediction as a function of RH. The increasing RH enhanced $K_{w,i}$ significantly for all polar SVOCs. For example, $K_{w,i}$ of 1-heptanoic acid increased by a factor of three (from 0.58 to 1.96) when RH increased from 0.4 to 0.75. $K_{w,i}$ of the hydrophobic SVOC such as *n*-eicosane was, however insensitive to RH as reported by Huang et al. (2018).

Parameters $H_{d,i}$, $H_{a,i}$, S_i , and α_i were chosen to predict $\alpha_{w,i}$ based on the statistical analysis (Table S3) semiempirically using the three UF-APHOR data as:

$$\ln(\alpha_{w,i}) = -0.92H_{d,i} - 1.3H_{a,i} - 0.039\alpha_i - 0.98S_i - 10.69. \quad (15)$$

$\alpha_{w,i}$ decreased with increasing molecular size, a trend similar to that reported by Ye et al. (2016). In addition, $\alpha_{w,i}$ also decreased when $H_{d,i}$, $H_{a,i}$, and S_i increased. The effect of RH on $\alpha_{w,i}$ was not derived in this study.

4.3. $K_{w,i}$, $\alpha_{w,i}$, and τ_{GWP} of various SVOCs: chamber data vs prediction

Values of SVOCs' $K_{w,i}$, $\alpha_{w,i}$, and τ_{GWP} were predicted and are summarized in Table 2 with the values of all the SVOCs' descriptors. $K_{w,i}$ was calculated using Eq. (14) at a given RH (0.75) and temperature (298 K) with 18.52 mg m^{-3} of $M_{wall-OM}$ and 273.3 g mol^{-1} of MW_{OM} (Sect. 4.1 and S2). Overall, by reducing volatility, the SVOC has a greater $K_{w,i}$ with a large carbon number or strong hydrogen bonding. However, volatility alone cannot fully explain the tendency of $K_{w,i}$. For example, 1-decanoic acid's $K_{w,i}$ is nearly 7 times higher (12.80) than that of *n*-nonadecane (1.93), although *n*-nonadecane's estimated $p_{L,i}^\circ$ ($1.09 \times 10^{-3} \text{ mmHg}$) is close to that of 1-decanoic acid ($1.18 \times 10^{-3} \text{ mmHg}$). In general, a longer τ_{GWP} was found for SVOCs with high $K_{w,i}$ and low $\alpha_{w,i}$. For example, *n*-eicosane's τ_{GWP} (95 min) was significantly higher than that of 2-heptanol (21 min), while *n*-eicosane and 2-heptanol had a similar $\alpha_{w,i}$ (2.9×10^{-6} and 3.3×10^{-6} , respectively). The dissimilar $K_{w,i}$ values of *n*-eicosane and 2-heptanol (1.93 and 0.28, respectively) occurred with large differences in τ_{GWP} . The volatile (small $K_{w,i}$) and hydrophobic SVOCs can reach equilibrium quickly (short τ_{GWP}). However, the predicted $C_{g,i}$ of citral deviated significantly from our observations. Aldehydes generally are more reactive than ketones because aldehydes can be oligomerized in the aerosol's aqueous layer or the wall (Jang and Kamens, 2001; Jang et al., 2002; Tong et al., 2006; Liggio and Li, 2006). Citral, a conjugated aldehyde, was reactive on the wall, and thus was excluded from the derivation of the QSAR model used to predict $K_{w,i}$ and $\alpha_{w,i}$.

The τ_{GWP} values obtained from this study ranged from 21 to 144 minutes (Table 2). Overall, the τ_{GWP} values obtained in this study are greater than those reported previously (Zhang et al., 2015b; Krechmer et al., 2016). For example, the τ_{GWP} of 2-tridecanone of this study was 71 minutes, while that in Yeh and Ziemann (2015) reported as less than 35 minutes. The difference in τ_{GWP} between laboratory studies can be explained with the difference in RH (0.75 at UF-APHOR vs 0.001 at Ziemann's chamber (Yeh and Ziemann, 2015) and the A/V ratio (i.e., 1.65 for the UF-APHOR chamber and 3.0 at Ziemann's chamber). The A/V ratio originating from the chamber dimension affects the calculation of $k_{on,i}$ as seen in Eq. (9) as well as the estimation of OM_{wall} . In addition, increasing RH can increase τ_{GWP} for polar compounds because the lower $\gamma_{w,i}$ (or elevated $K_{w,i}$) at the higher RH increases the time required to reach equilibrium. $k_{on,i}$ associated with SVOC flux to the chamber wall is not (or little) affected by RH.



4.4. Impact of $\gamma_{w,i}$ on the prediction of $K_{w,i}$

The observed $K_{w,i}$ are plotted against predicted $1/C_i^*$ (estimated from $RT/MW_{OM}\gamma_{w,i}P_L^\circ$ with unity $\gamma_{w,i}$) and $K_{w,i}$ in Fig. 5(a) and Fig. 5(b), respectively. Fig. 5 was constructed using the three datasets in this study and the data reported in the literature (Matsunaga and Ziemann, 2010; Yeh and Ziemann, 2015). When observed $K_{w,i}$ is plotted to $1/C_i^*$ (Fig 5(a)), a regression line with some scatter was obtained ($R^2 = 0.12$, except aldehydes). However, the inclusion of $\gamma_{w,i}$ (Fig 5(b)), as calculated from the QSAR method, dramatically improves the predictability of $K_{w,i}$ ($R^2 = 0.64$, except aldehydes). To demonstrate the potential impact of $\gamma_{w,i}$ on $K_{w,i}$, $1/C_i^*$ was plotted vs the predicted $K_{w,i}$ in Fig. 5(c). For various organic compounds that differ in functionalities and molecular sizes, $\gamma_{w,i}$ ranges 10^0 – 10^4 . Overall, the $K_{w,i}$ values of monofunctional oxygenated compounds, such as *n*-alcohols, 1-carboxylic acids, ketones, and *n*-alkanes reasonably accord with the predicted $K_{w,i}$ using Eq. (14). However, the prediction of diols and aldehydes deviated from observation by one order of magnitude (Fig. 5(b)).

The study of multifunctional SVOCs' (e.g., diols) GWP generally would be more difficult than that of monofunctional compounds because of the experimental difficulty of injecting multifunctional SVOCs into the chamber and the detection techniques. In addition, estimating multifunctional alcohols' $p_{L,i}^\circ$ could be problematic because of intramolecular hydrogen bonding's contribution when estimating their boiling point. Estimating multifunctional alcohols' descriptors may also be uncertain because of the lack of a database (Miller, 1990; Yap, 2011).

The vapor fraction ($F_{g,i} = C_{g,i}/C_{T,i}$) of the total concentration of SVOC(*i*) at equilibrium can be calculated from $K_{w,i}$ with the following equation:

$$F_{g,i} = \frac{1}{1+K_{w,i}} \quad (16)$$

Fig. S5 shows the model predicted $F_{g,i}$ and experimentally measured $F_{g,i}$ using the same dataset applied to $K_{w,i}$. The smaller $F_{g,i}$ indicates the significance of GWP on the chemical loss to the chamber wall.

4.5. Characteristic time analyses of major atmospheric processes

To assess the importance of GWP on SOA formation, the τ_{GWP} predicted using $K_{w,i}$ and $\alpha_{w,i}$ was compared with the characteristic time (τ) associated with other important atmospheric processes. Table S4 summarizes equations to estimate τ_{GWP} , τ_{OH} (reaction with an OH radical at $[OH] = 1.33 \times 10^7$ molecules cm^{-3} , at noon during a typical chamber experiment), τ_{or} (reaction in the organic aerosol phase), τ_{in} (reaction in the inorganic aqueous phase), and τ_{GP} (SVOC's gas-particle partitioning). The characteristic times (τ_{GWP} , τ_{OH} , τ_{or} , τ_{in} , and τ_{GP}) are estimated in Fig. 6 for the eight surrogate SVOCs differentiated by two different levels of $K_{w,i}$ (10^{-1} and 10^1), $\alpha_{w,i}$ (1×10^{-6} and 4×10^{-6}), and the reactivity scale (fast and no reaction) in the aerosol phase (Im et al., 2014; Beardsley and Jang, 2016; Zhou et al., 2019).

To evaluate the significance of GWP on chamber-generated SOA mass, Figure 6 also provides $F_{g,i}$ along with the characteristic times of various processes of various SVOCs. Overall, GWP is significant with inert oxygenated products with large $K_{w,i}$ and large $\alpha_{w,i}$. τ_{GP} is as short as $\sim 1 \times 10^1$ s and it is always shorter than τ_{GWP} . τ_{GWP} increases by an order of magnitude by increasing $K_{w,i}$ from 10^{-1} to 10^1 . The large $\alpha_{w,i}$ (4×10^{-6}) shortens τ_{GWP} compared with the small $\alpha_{w,i}$ (1×10^{-6}) at given conditions. τ_{OH} is $\sim 4 \times 10^3$ s under the given conditions. τ_{GWP} is shorter than τ_{OH} when $K_{w,i}$ is small (10^{-1}). Both τ_{or} and τ_{in} change with the reactivity scale in the aerosol phase by six orders of magnitude (fast (A–D) vs. no reactivity (E–H) in Fig. 6). When the reactivity scale is fast, τ_{GWP} is significantly larger than both τ_{or} and τ_{in} . When



oxygenated products in the phase are dominated by heterogeneously reactive organic species or hygroscopic inorganic
285 electrolytes are present in the aerosol, GWP may be less influential on SOA formation.

In addition, τ_{GP} , τ_{OH} , τ_{GWP} , τ_{or} , and τ_{in} of oxygenated products that are found in SOA (toluene, α -pinene, or isoprene)
are also estimated in Table S5. The products in Table S5 are chosen because of their high concentrations obtained in simulations
using the explicit gas mechanisms (i.e., Master Chemical Mechanism (MCM) version 3.3.1 (Jenkin et al., 2012)) under the
low NO_x condition (HC ppbC/NO_x ppb = 6.5 (toluene), 20 (isoprene), and 6.9 (α -pinene)). τ_{or} and τ_{in} of the products in
290 Table S5 were estimated using the rate constants obtained from the UNIPAR model (Im et al., 2014; Beardsley and Jang, 2016;
Zhou et al., 2019). SVOCs are classified by reactivity in the aerosol phase in UNIPAR: very fast (VF, more than two aldehydes
or epoxides), fast (F, at least two aldehydes or epoxides), medium (M, one aldehyde or epoxide), slow (S, ketones), and no
reaction (P). For most toluene products, τ_{in} is much shorter than those of τ_{GWP} suggesting that GWP's influence on the
formation of toluene SOA might be trivial in the presence of the electrolytic aqueous phase. However, its effect on SOA
295 formation becomes significant in the absence of the inorganic aerosol based on τ_{GWP} and τ_{or} (Table S5).

Glyoxal, a major product from the photooxidation of toluene, has a high $p_{L,i}^{\circ}$ but can contribute to SOA formation significantly
because of its high reactivity (Fu et al., 2008). As shown in Table S5, both glyoxal's τ_{or} and τ_{in} are much shorter than τ_{GWP}
and thus, GWP has little effect. Of equal importance is the impact of GWP on toluene SOA, which would be less significant
than the prediction of previous studies (Zhang et al., 2014; La et al., 2016) because of the contribution of reactive SVOCs'
300 heterogeneous reactions in the aerosol phase, particularly in the electrolytic aqueous phase. The importance of the reaction of
an SVOC with an OH radical is comparable to that of GWP. IEPOXB, an isoprene oxidation product, is known to contribute
significantly to isoprene SOA (Hu et al., 2015). Similar to toluene SOA products, the GWP of IEPOXB's influence on isoprene
SOA may not be important in the presence of an aqueous inorganic seed ($\tau_{GWP} \sim 3.3$ h; $\tau_{in} \sim 0.1$ h; $\tau_{or} \sim 5.8$ h). The volatility
of such α -pinene oxidation products as pinic acid, pinonic acid, and pinonaldehyde is relatively low compared with that of
305 toluene SVOCs. Pinic acid, which is not reactive in the aerosol phase, can contribute quickly to SOA growth, but GWP will
influence it in the long term. Pinonic acid's reactivity (slow) and pinonaldehyde's (medium) are comparably weak, but their
low volatility and short τ_{in} (τ_{or}) can lead SOA to form quickly without GWP's significant impact.

4.6. Model sensitivity and uncertainties

To determine the model sensitivity (%), the $F_{g,i}$ prediction for the major variables was performed by changing the values of
310 RH ($0.001 < RH < 0.8$), temperature (288–308 K), OM_{wall} (6.9 to 37.5 mg m⁻³), and the A/V ratio (from 1 to 3) selectively. In
Fig. 7(a) and (b), the modulation of $F_{g,i}$ for the two SVOCs (1-decanoic acid and *n*-nonadecane) was scaled to the standard
condition (RH = 0.50, temperature = 298 K, $OM_{wall} = 18.52$ mg m⁻³, and A/V ratio = 1.65 based on the possible range of the
conditions). At the standard condition, the predicted $F_{g,i}$ of *n*-nonadecane and 1-decanoic acid are 0.23 and 0.17, respectively.
Among the four variables, RH had the greatest influence on $F_{g,i}$, particularly for the polar compound (1-decanoic acid). Both
315 1-decanoic acid and *n*-nonadecane have similar $p_{L,i}^{\circ}$ (Table 2), but their sensitivity to RH differs. RH's effect on 1-decanoic
acid is related to $K_{w,i}$ exponentially, as seen in Eq. (14). As shown in Fig. 7, the chamber dimension (the A/V ratio) and
 OM_{wall} also influenced $F_{g,i}$ significantly. The A/V ratio increases $K_{w,i}$ by increasing $k_{on,i}$ (Eq. (9)) and impacts on the
estimation of OM_{wall} . Thus, the A/V ratio can affect $F_{g,i}$ more than OM_{wall} .

Fig. 7(c) and (d) illustrates the uncertainty in $F_{g,i}$ attributable to the uncertainties of the major model parameters used to
320 predict $K_{w,i}$ and $\alpha_{w,i}$: four descriptors (S_i , α_i , $H_{d,i}$, and $H_{a,i}$) and $p_{L,i}^{\circ}$ in Eq. (14) and Eq. (15). Jover et al. (2004) reported
the uncertainty associated with descriptors $H_{d,i}$, $H_{a,i}$, and S_i as 8.4%, 3.6%, and 3.1%, respectively. The error associated with



α_i was estimated to be 5% based on other descriptors' errors and the database used for the additive method (Miller, 1990). Stein and Brown (1994) and Zhao et al. (1999) reported that the uncertainty associated with the group contribution method used for $p_{L,i}^\circ$ estimation is a factor of 1.45. Among the five model parameters in Fig. 7(c) and (d), $p_{L,i}^\circ$ was most influential on uncertainties of $F_{g,i}$.

The uncertainties of the QSAR model equations to predict $K_{w,i}$ and $\alpha_{w,i}$ can be determined from the experimental data. The time consumed for transferring organic vapors from the East chamber to the West chamber can impact the uncertainty in measurements of $C_{g,i}$. For most of the organic vapors, τ_{GWP} is significantly longer than 10 minutes (600 s). Therefore, the impact of the uncertainties of $C_{g,i}$ on the QSAR equations, which are attributed to various SVOCs, could be small.

4.7. Atmospheric implications

The QSAR-based model platform in this study allows the GWP for a variety of SVOCs' formation during the atmospheric processes of hydrocarbons to be predicted. We concluded that the $\gamma_{w,i}$ expressed with QSAR descriptors is essential to predict $K_{w,i}$, although $p_{L,i}^\circ$ is the most critical. Overall, GWP appears to be more significant with the more polar and larger SVOCs (section 4.3). Among the variables (temperature, RH, A/V ratio, and OM_{wall}), RH was the most influential in predicting $K_{w,i}$ (or $F_{g,i}$) for polar compound (in Sect. 4.6) and the A/V ratio also influenced $F_{g,i}$ significantly due to the impact of the A/V ratio on OM_{wall} . The accuracy of the data and the collected chemical species influence the coefficients' suitability in the QSAR-based model. The suitability of the QSAR-based model to predict simple monofunctional organic species' $K_{w,i}$ and $\alpha_{w,i}$ was examined, but it needs to be extended to accommodate various multifunctional organics.

The relative importance of GWP to form SOA was assessed by comparison of τ_{GWP} with τ_{GP} , τ_{OH} , τ_{or} , and τ_{in} (in Sect. 4.5). τ_{GWP} is comparable to τ_{OH} , although their values depend on SVOC's molecular structure as well as its reactivity toward an OH radical. In the presence of an aqueous inorganic seed ($10 \mu\text{g m}^{-3}$ of effloresced inorganic seed), reactive SVOCs' τ_{in} (Table S5) is significantly shorter than that of τ_{GWP} by six orders of magnitude, suggesting that GWP's role in SOA growth may not be significant. In the absence of an inorganic seed (or in the presence of a dry seed), τ_{or} and τ_{in} are comparable to τ_{GWP} for many SVOCs. However, τ_{or} decreases for highly reactive organics (i.e., glyoxal) in aerosol or volatile SVOCs (i.e., pinonic acid and pinonaldehyde, $p_L^\circ < 10^{-3}$ mmHg). Many studies have found a high fraction of oligomeric matter in SOA mass, (Jang et al., 2002; Tolocka et al., 2004; Gao et al., 2004; Beardsley and Jang, 2016; Lemaire et al., 2016), evidence that heterogeneous chemistry of organic species is an important mechanism by which SOA forms. For example, Lemaire et al. (2016) reported 50% of the oligomeric mass of isoprene SOA. If the inorganic aerosol is acidic, oligomerization is accelerated and τ_{in} becomes even shorter than the value shown in Table S5. For example, Beardsley and Jang (2016) reported that at least 65% of total SOA is oligomer, and an oligomer can constitute 85% of SOA in the presence of an acidic seed. Because of the attribution of aqueous chemistry in aerosol, the bias in SOA mass by GWP could be varied. Inert organic products' wall deposition also can influence SOA mass by modulating the formation of reactive SVOCs and the quantity of oxidants (i.e., OH radicals and ozone), and consequently affect the SOA mass.

As shown in Fig. 3(f), the product with an aldehyde group might be affected by heterogeneous reactions on the chamber wall and cause a bias in the estimation of SOA mass. The heterogeneous reactions of reactive aldehydes can be significant when electrolytic inorganic species build up on the Teflon wall. However, noticing that τ_{GP} is typically much shorter than τ_{GWP} , the heterogeneous reaction of the reactive organic species in the aerosol phase should be much more significant than that on the wall surfaces when the reaction is irreversible.

To date, GWP has been studied by measuring individual SVOCs. However, there is a gap between GWP studies and their



360 application to conventional SOA models that are approached by an apparent gas-particle partitioning with the semiempirical
parameters of only a small number of semivolatile surrogate products (Odum et al., 1996; Donahue et al., 2006). The model
equations used to predict the GWP parameters ($K_{w,i}$, $k_{on,i}$, and $k_{off,i}$) in this study can be coupled with explicit SOA models.
For example, the UNIPAR model predicts SOA mass by counting the volatility and aerosol phase reactions of the lumping
species sourced from the SVOCs that are produced explicitly during hydrocarbons' photooxidation (Im et al., 2014; Beardsley
365 and Jang, 2016; Zhou et al., 2019). The integration of GWP parameters with the explicit SOA model improves the
quantification of the bias in SOA mass attributable to GWP. GWP's predictability also improves the study of the gaseous
SVOCs' degradation because numerous chemical species' degradation kinetics rely on chamber data. For example, Bertrand
et al. (2018) reported that biomass burning markers' decay in particulate matter was not attributable to reactions with OH
radicals, but gas-particle partitioning and GWP dominated it instead.

370

Author contribution. MJ designed the experiments and MJ, SH, and HJ carried them out. SH prepared the manuscript with
contributions from all co-authors.

Acknowledgments. This research was supported by the awards from the NSF (AGS 1923651) and the National Strategic
375 Project-Fine particle of the National Research Foundation of Korea (NRF) funded by the Ministry of Science and ICT (MSIT),
the Ministry of Environment (ME), and the Ministry of Health and Welfare (MOHW) (2017M3D8A1090654).



References

- Abraham, M., and McGowan, J.: The use of characteristic volumes to measure cavity terms in reversed phase liquid-chromatography, *Chromatographia*, 23, 243-246, 10.1007/BF02311772, 1987.
- 380 Abraham, M., Whiting, G., Doherty, R., and Shuely, W.: Hydrogen-bonding .16. A new solute solvation parameter, PI-2(H), from gas-chromatographic data, *Journal of Chromatography*, 587, 213-228, 10.1016/0021-9673(91)85158-C, 1991.
- Alan, R.: Model solvent systems for QSAR. Part 3. An LSER analysis of the 'critical quartet.' New light on hydrogen bond strength and directionality, *Journal of the Chemical Society, Perkin Transactions 2*, 705-722, 1992.
- Appel, K., Napelenok, S., Foley, K., Pye, H., Hogrefe, C., Luecken, D., Bash, J., Roselle, S., Pleim, J., Foroutan, H., Hutzell, W., Pouliot, G., Sarwar, G., Fahey, K., Gantt, B., Gilliam, R., Heath, N., Kang, D., Mathur, R., Schwede, D., Spero, T., Wong, D., and Young, J.: Description and evaluation of the Community Multiscale Air Quality (CMAQ) modeling system version 5.1, *Geoscientific Model Development*, 10, 1703-1732, 10.5194/gmd-10-1703-2017, 2017.
- 385
- Arp, H., and Goss, K.: Ambient Gas/Particle Partitioning. 3. Estimating Partition Coefficients of Apolar, Polar, and Ionizable Organic Compounds by Their Molecular Structure, *Environmental Science & Technology*, 43, 1923-1929, 10.1021/es8025165, 2009.
- 390
- Barsanti, K., Carlton, A., and Chung, S.: Analyzing experimental data and model parameters: implications for predictions of SOA using chemical transport models, *Atmospheric Chemistry and Physics*, 13, 12073-12088, 10.5194/acp-13-12073-2013, 2013.
- Barsanti, K. C., Kroll, J. H., and Thornton, J. A.: Formation of low-volatility organic compounds in the atmosphere: Recent advancements and insights, *The journal of physical chemistry letters*, 8, 1503-1511, 2017.
- 395
- Barton, A. F.: *CRC handbook of solubility parameters and other cohesion parameters*, Routledge, 2017.
- Beardsley, R., Jang, M., Ori, B., Im, Y., Delcomyn, C., and Witherspoon, N.: Role of sea salt aerosols in the formation of aromatic secondary organic aerosol: yields and hygroscopic properties, *Environmental Chemistry*, 10, 167-177, 10.1071/EN13016, 2013.
- 400
- Beardsley, R., and Jang, M.: Simulating the SOA formation of isoprene from partitioning and aerosol phase reactions in the presence of inorganics, *Atmospheric Chemistry and Physics*, 16, 5993-6009, 10.5194/acp-16-5993-2016, 2016.
- Bertram, A., Martin, S., Hanna, S., Smith, M., Bodsworth, A., Chen, Q., Kuwata, M., Liu, A., You, Y., and Zorn, S.: Predicting the relative humidities of liquid-liquid phase separation, efflorescence, and deliquescence of mixed particles of ammonium sulfate, organic material, and water using the organic-to-sulfate mass ratio of the particle and the oxygen-to-carbon elemental ratio of the organic component, *Atmospheric Chemistry and Physics*, 11, 10995-11006, 2011.
- 405
- Bertrand, A., Stefenelli, G., Pieber, S., Bruns, E., Temime-Roussel, B., Slowik, J., Wortham, H., Prevot, A., El Haddad, I., and Marchand, N.: Influence of the vapor wall loss on the degradation rate constants in chamber experiments of levoglucosan and other biomass burning markers, *Atmospheric Chemistry and Physics*, 18, 10915-10930, 10.5194/acp-18-10915-2018, 2018.
- 410
- Bi, X., Sheng, G., Peng, P. a., Chen, Y., Zhang, Z., and Fu, J.: Distribution of particulate-and vapor-phase n-alkanes and polycyclic aromatic hydrocarbons in urban atmosphere of Guangzhou, China, *Atmospheric Environment*, 37, 289-298, 2003.
- Bian, Q., May, A., Kreidenweis, S., and Pierce, J.: Investigation of particle and vapor wall-loss effects on controlled wood-smoke smog-chamber experiments, *Atmospheric Chemistry and Physics*, 15, 11027-11045, 10.5194/acp-15-11027-2015, 415 2015.
- Brune, W. H.: *The Chamber Wall Index for Gas-Wall Interactions in Atmospheric Environmental Enclosures*, Environmental science & technology, 2019.



- Cao, G., and Jang, M.: An SOA Model for Toluene Oxidation in the Presence of Inorganic Aerosols, *Environmental Science & Technology*, 44, 727-733, 10.1021/es901682r, 2010.
- 420 Carlton, A., Wiedinmyer, C., and Kroll, J.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmospheric Chemistry and Physics*, 9, 4987-5005, 10.5194/acp-9-4987-2009, 2009.
- Chen, Q., Liu, Y., Donahue, N. M., Shilling, J. E., and Martin, S. T.: Particle-phase chemistry of secondary organic material: modeled compared to measured O: C and H: C elemental ratios provide constraints, *Environmental science & technology*, 45, 4763-4770, 2011.
- 425 Cohen, A., Brauer, M., Burnett, R., Anderson, H., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope, C., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C., and Forouzanfar, M.: Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015, *Lancet*, 389, 1907-1918, 10.1016/S0140-6736(17)30505-6, 2017.
- 430 Donahue, N., Robinson, A., Stanier, C., and Pandis, S.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environmental Science & Technology*, 40, 2635-2643, 10.1021/es052297c, 2006.
- Endo, S., and Goss, K.: Applications of Polyparameter Linear Free Energy Relationships in Environmental Chemistry, *Environmental Science & Technology*, 48, 12477-12491, 10.1021/es503369t, 2014.
- 435 Fu, T., Jacob, D., Wittrock, F., Burrows, J., Vrekoussis, M., and Henze, D.: Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *Journal of Geophysical Research-Atmospheres*, 113, 10.1029/2007JD009505, 2008.
- Gao, S., Ng, N., Keywood, M., Varutbangkul, V., Bahreini, R., Nenes, A., He, J., Yoo, K., Beauchamp, J., Hodyss, R., Flagan, R., and Seinfeld, J.: Particle phase acidity and oligomer formation in secondary organic aerosol, *Environmental Science & Technology*, 38, 6582-6589, 10.1021/es049125k, 2004.
- 440 Gundal, L., Lee, V., Mahanama, K., Stevens, R., and Daisey, J.: Direct determination of the phase distributions of semivolatile polycyclic aromatic-hydrocarbons using annular denuders, *Atmospheric Environment*, 29, 1719-1733, 10.1016/1352-2310(94)00366-S, 1995.
- Hallquist, M., Wenger, J., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N., George, C., 445 Goldstein, A., Hamilton, J., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M., Jimenez, J., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T., Monod, A., Prevot, A., Seinfeld, J., Surratt, J., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmospheric Chemistry and Physics*, 9, 5155-5236, 10.5194/acp-9-5155-2009, 2009.
- Hu, W., Campuzano-Jost, P., Palm, B., Day, D., Ortega, A., Hayes, P., Krechmer, J., Chen, Q., Kuwata, M., Liu, Y., de Sa, S., 450 McKinney, K., Martin, S., Hu, M., Budisulistiorini, S., Riva, M., Surratt, J., St Clair, J., Isaacman-Van Wertz, G., Yee, L., Goldstein, A., Carbone, S., Brito, J., Artaxo, P., de Gouw, J., Koss, A., Wisthaler, A., Mikoviny, T., Karl, T., Kaser, L., Jud, W., Hansel, A., Docherty, K., Alexander, M., Robinson, N., Coe, H., Allan, J., Canagaratna, M., Paulot, F., and Jimenez, J.: Characterization of a real-time tracer for isoprene epoxydiols-derived secondary organic aerosol (IEPOX-SOA) from aerosol mass spectrometer measurements, *Atmospheric Chemistry and Physics*, 15, 11807-11833, 10.5194/acp-15-11807-455 2015, 2015.
- Huang, Y., Zhao, R., Charan, S., Kenseth, C., Zhang, X., and Seinfeld, J.: Unified Theory of Vapor-Wall Mass Transport in Teflon-Walled Environmental Chambers, *Environmental Science & Technology*, 52, 2134-2142, 10.1021/acs.est.7b05575, 2018.
- Im, Y., Jang, M., and Beardsley, R.: Simulation of aromatic SOA formation using the lumping model integrated with explicit



- 460 gas-phase kinetic mechanisms and aerosol-phase reactions, *Atmospheric Chemistry and Physics*, 14, 4013-4027,
10.5194/acp-14-4013-2014, 2014.
- Jang, J., Jang, M., Mui, W., Delcomyn, C., Henley, M., and Hearn, J.: Formation of Active Chlorine Oxidants in Saline-Oxone
Aerosol, *Aerosol Science and Technology*, 44, 1018-1026, 10.1080/02786826.2010.507612, 2010.
- Jang, M., Kamens, R., Leach, K., and Strommen, M.: A thermodynamic approach using group contribution methods to model
465 the partitioning of semivolatile organic compounds on atmospheric particulate matter, *Environmental Science &
Technology*, 31, 2805-2811, 10.1021/es970014d, 1997.
- Jang, M., and Kamens, R.: A thermodynamic approach for modeling partitioning of semivolatile organic compounds on
atmospheric particulate matter: Humidity effects, *Environmental Science & Technology*, 32, 1237-1243,
10.1021/es970773w, 1998.
- 470 Jang, M., and Kamens, R.: A predictive model for adsorptive gas partitioning of SOCs on fine atmospheric inorganic dust
particles, *Environmental Science & Technology*, 33, 1825-1831, 10.1021/es9805167, 1999.
- Jang, M., and Kamens, R.: Atmospheric secondary aerosol formation by heterogeneous reactions of aldehydes in the presence
of a sulfuric acid aerosol catalyst, *Environmental Science & Technology*, 35, 4758-4766, 10.1021/es010790s, 2001.
- Jang, M., Czoschke, N., Lee, S., and Kamens, R.: Heterogeneous atmospheric aerosol production by acid-catalyzed particle-
475 phase reactions, *Science*, 298, 814-817, 10.1126/science.1075798, 2002.
- Jenkin, M., Wyche, K., Evans, C., Carr, T., Monks, P., Alfarra, M., Barley, M., McFiggans, G., Young, J., and Rickard, A.:
Development and chamber evaluation of the MCM v3.2 degradation scheme for beta-caryophyllene, *Atmospheric
Chemistry and Physics*, 12, 5275-5308, 10.5194/acp-12-5275-2012, 2012.
- Jover, J., Bosque, R., and Sales, J.: Determination of Abraham solute parameters from molecular structure, *Journal of Chemical
480 Information and Computer Sciences*, 44, 1098-1106, 10.1021/ci049943w, 2004.
- Kaur, I., Misra, B., and Kohli, A.: Synthesis of Teflon-FEP grafted membranes for use in water desalination, *Desalination*, 139,
357-365, 10.1016/S0011-9164(01)00331-9, 2001.
- Krechmer, J., Pagonis, D., Ziemann, P., and Jimenez, J.: Quantification of Gas-Wall Partitioning in Teflon Environmental
Chambers Using Rapid Bursts of Low-Volatility Oxidized Species Generated in Situ, *Environmental Science & Technology*,
485 50, 5757-5765, 10.1021/acs.est.6b00606, 2016.
- Kuwata, M., Shao, W., Leboutteiller, R., and Martin, S.: Classifying organic materials by oxygen-to-carbon elemental ratio to
predict the activation regime of Cloud Condensation Nuclei (CCN), *Atmospheric Chemistry and Physics*, 13, 5309-5324,
2013.
- La, Y., Camredon, M., Ziemann, P., Valorso, R., Matsunaga, A., Lannuque, V., Lee-Taylor, J., Hodzic, A., Madronich, S., and
490 Aumont, B.: Impact of chamber wall loss of gaseous organic compounds on secondary organic aerosol formation: explicit
modeling of SOA formation from alkane and alkene oxidation, *Atmospheric Chemistry and Physics*, 16, 1417-1431,
10.5194/acp-16-1417-2016, 2016.
- Leach, K., Kamens, R., Strommen, M., and Jang, M.: Partitioning of semivolatile organic compounds in the presence of a
secondary organic aerosol in a controlled atmosphere, *Journal of Atmospheric Chemistry*, 33, 241-264,
495 10.1023/A:1006108430033, 1999.
- Leivo, E., Wilenius, T., Kinoshita, T., Vuoristo, P., and Mantyla, T.: Properties of thermally sprayed fluoropolymer PVDF, ECTFE,
PFA and FEP coatings, *Progress in Organic Coatings*, 49, 69-73, 10.1016/j.porgcoat.2003.08.011, 2004.
- Lemaire, V., Coll, I., Couvidat, F., Mouchel-Vallon, C., Seigneur, C., and Siour, G.: Oligomer formation in the troposphere:
from experimental knowledge to 3-D modeling, *Geoscientific Model Development*, 9, 1361-1382, 10.5194/gmd-9-1361-
500 2016, 2016.
- Li, J., Jang, M., and Beardsley, R.: Dialkylsulfate formation in sulfuric acid-seeded secondary organic aerosol produced using



- an outdoor chamber under natural sunlight, *Environmental Chemistry*, 13, 590-601, 10.1071/EN15129, 2016.
- Liggio, J., and Li, S.: Reactive uptake of pinonaldehyde on acidic aerosols, *Journal of Geophysical Research-Atmospheres*, 111, 10.1029/2005JD006978, 2006.
- 505 Lyu, R., Shi, Z., Alam, M. S., Wu, X., Liu, D., Vu, T. V., Stark, C., Xu, R., Fu, P., and Feng, Y.: Alkanes and aliphatic carbonyl compounds in wintertime PM_{2.5} in Beijing, China, *Atmospheric Environment*, 202, 244-255, 2019.
- Matsunaga, A., and Ziemann, P.: Gas-Wall Partitioning of Organic Compounds in a Teflon Film Chamber and Potential Effects on Reaction Product and Aerosol Yield Measurements, *Aerosol Science and Technology*, 44, 881-892, 10.1080/02786826.2010.501044, 2010.
- 510 McMurry, P., and Grosjean, D.: Gas and aerosol wall losses in Teflon film smog chambers, *Environmental Science & Technology*, 19, 1176-1182, 10.1021/es00142a006, 1985.
- McNeill, V., Woo, J., Kim, D., Schwier, A., Wannell, N., Sumner, A., and Barakat, J.: Aqueous-Phase Secondary Organic Aerosol and Organosulfate Formation in Atmospheric Aerosols: A Modeling Study, *Environmental Science & Technology*, 46, 8075-8081, 10.1021/es3002986, 2012.
- 515 Miller, K: Additivity methods in molecular polarizability, *Journal of the American Chemical Society*, 112, 8533-8542, 10.1021/ja00179a044, 1990.
- Odum, J., Hoffmann, T., Bowman, F., Collins, D., Flagan, R., and Seinfeld, J.: Gas/particle partitioning and secondary organic aerosol yields, *Environmental Science & Technology*, 30, 2580-2585, 10.1021/es950943+, 1996.
- Pankow, J., Storey, J., and Yamasaki, H: Effects of relative-humidity on gas-particle partitioning of semivolatile organic-compounds to urban particulate matter, *Environmental Science & Technology*, 27, 2220-2226, 10.1021/es00047a032, 1993.
- 520 Pankow, J.: An absorption-model of gas-particle partitioning of organic-compounds in the atmosphere, *Atmospheric Environment*, 28, 185-188, 10.1016/1352-2310(94)90093-0, 1994.
- Pankow, J. F.: An absorption model of the gas/aerosol partitioning involved in the formation of secondary organic aerosol, *Atmospheric Environment*, 28, 189-193, 1994.
- 525 Park, R., Jacob, D., Chin, M., and Martin, R.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, *Journal of Geophysical Research-Atmospheres*, 108, 10.1029/2002JD003190, 2003.
- Platts, J., Butina, D., Abraham, M., and Hersey, A.: Estimation of molecular linear free energy relation descriptors using a group contribution approach, *Journal of Chemical Information and Computer Sciences*, 39, 835-845, 10.1021/ci980339t, 1999.
- 530 Puzyn, T., Leszczynski, J., and Cronin, M. T.: Recent advances in QSAR studies: methods and applications, Springer Science & Business Media, 2010.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.
- Simoneit, B. R.: Organic matter of the troposphere—V: application of molecular marker analysis to biogenic emissions into 535 the troposphere for source reconciliations, *Journal of Atmospheric Chemistry*, 8, 251-275, 1989.
- Stein, S., and Brown, R.: Estimation of normal boiling points from group contributions, *Journal of Chemical Information and Computer Sciences*, 34, 581-587, 10.1021/ci00019a016, 1994.
- Tolocka, M., Jang, M., Ginter, J., Cox, F., Kamens, R., and Johnston, M.: Formation of oligomers in secondary organic aerosol, *Environmental Science & Technology*, 38, 1428-1434, 10.1021/es035030r, 2004.
- 540 Tong, C., Blanco, M., Goddard, W., and Seinfeld, J.: Secondary organic aerosol formation by heterogeneous reactions of aldehydes and ketones: A quantum mechanical study, *Environmental Science & Technology*, 40, 2333-2338, 10.1021/es0519785, 2006.
- Volkamer, R., Jimenez, J., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L., Worsnop, D., and Molina, M.:



- 545 Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophysical Research Letters*, 33, 10.1029/2006GL026899, 2006.
- Yap, C.: PaDEL-Descriptor: An Open Source Software to Calculate Molecular Descriptors and Fingerprints, *Journal of Computational Chemistry*, 32, 1466-1474, 10.1002/jcc.21707, 2011.
- Ye, P., Ding, X., Hakala, J., Hofbauer, V., Robinson, E., and Donahue, N.: Vapor wall loss of semi-volatile organic compounds in a Teflon chamber, *Aerosol Science and Technology*, 50, 822-834, 10.1080/02786826.2016.1195905, 2016.
- 550 Yeh, G., and Ziemann, P.: Alkyl Nitrate Formation from the Reactions of C-8-C-14 n-Alkanes with OH Radicals in the Presence of NO_x: Measured Yields with Essential Corrections for Gas-Wall Partitioning, *Journal of Physical Chemistry a*, 118, 8147-8157, 10.1021/jp500631v, 2014.
- Yeh, G., and Ziemann, P.: Gas-Wall Partitioning of Oxygenated Organic Compounds: Measurements, Structure-Activity Relationships, and Correlation with Gas Chromatographic Retention Factor, *Aerosol Science and Technology*, 49, 726-737, 555 10.1080/02786826.2015.1068427, 2015.
- Zhang, X., Cappa, C., Jathar, S., Mcvay, R., Ensberg, J., Kleeman, M., and Seinfeld, J.: Influence of vapor wall loss in laboratory chambers on yields of secondary organic aerosol, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 5802-5807, 10.1073/pnas.1404727111, 2014.
- Zhang, X., McVay, R. C., Huang, D. D., Dalleska, N. F., Aumont, B., Flagan, R. C., and Seinfeld, J. H.: Formation and evolution 560 of molecular products in alpha-pinene secondary organic aerosol, *Proc Natl Acad Sci U S A*, 112, 14168-14173, 10.1073/pnas.1517742112, 2015a.
- Zhang, X., Schwantes, R., McVay, R., Lignell, H., Coggon, M., Flagan, R., and Seinfeld, J.: Vapor wall deposition in Teflon chambers, *Atmospheric Chemistry and Physics*, 15, 4197-4214, 10.5194/acp-15-4197-2015, 2015b.
- Zhao, L., Li, P., and Yalkowsky, S.: Predicting the entropy of boiling for organic compounds, *Journal of Chemical Information and Computer Sciences*, 39, 1112-1116, 10.1021/ci990054w, 1999.
- 565 Zhong, M., and Jang, M.: Dynamic light absorption of biomass-burning organic carbon photochemically aged under natural sunlight, *Atmospheric Chemistry and Physics*, 14, 1517-1525, 10.5194/acp-14-1517-2014, 2014.
- Zhou, C., Jang, M., and Yu, Z.: Simulation of SOA formation from the photooxidation of monoalkylbenzenes in the presence of aqueous aerosols containing electrolytes under various NO_x levels, *Atmospheric Chemistry and Physics*, 19, 5719-5735, 570 2019.



Table 1. Chamber experiments used to model the gas-wall process of SVOCs.

Index (Fig. 5)	Date	RH	Temp (°C)	Dilution factor ^{a)}
a	October 8, 2018	0.40	297~301	$y = 0.38e^{0.049t}$
b	May 15, 2018	0.75	294~296	$y = 0.60e^{0.076t}$
c	August 6, 2018	0.53	297~301	$y = 0.40e^{0.040t}$
d	(Yeh and Ziemann, 2015)	<0.001	297~299	
e	(Matsunaga and Ziemann, 2010)			

575 ^{a)} The factor for chamber dilution is calculated by measuring CCl₄ with GC-FID




Table 2. Chemicals injected into the chamber and their physicochemical parameters (at 298 K) predicted using PaDEL-Descriptor and estimated $K_{w,i}$, $\alpha_{w,i}$, and τ_{GWP} under a given condition (298K, RH=0.75).

No.	Chemical	MW	Molecular descriptors ^{a)}						α_i	p_L° (b)	$K_{w,i}$	$\alpha_{w,i}$ ($\times 10^{-6}$)	τ_{GWP} (s)
			$H_{d,i}$	$H_{a,i}$	E_f	S_f	$H_{d,i}$	$H_{a,i}$					
1	1-hexanoic acid	116	0.59	0.41	0.15	0.51	14.08	0.078	1.12	2.7	3462		
2	1-heptanoic acid	130	0.59	0.41	0.15	0.51	16.26	0.026	2.18	2.5	4900		
3	1-octanoic acid	144	0.59	0.41	0.15	0.51	18.45	0.009	4.13	2.3	6276		
4	1-decanoic acid	172	0.59	0.41	0.15	0.51	22.82	0.001	12.80	1.9	8630		
5	Benzoic acid	122	0.59	0.46	0.78	0.93	7.52	0.020	15.73	2.2	7171		
6	Pyruvic acid	88	0.43	0.68	0.29	0.88	8.48	0.256	1.33	1.9	4244		
7	2-heptanol	116	0.35	0.40	0.19	0.41	17.49	0.317	0.28	3.3	1281		
8	1-octanol	130	0.35	0.39	0.21	0.45	19.68	0.109	0.53	3.0	2220		
9	1-nonanol	144	0.35	0.39	0.21	0.45	21.86	0.036	1.05	2.7	3566		
10	Benzyl alcohol	108	0.35	0.56	0.83	0.88	8.75	0.093	4.77	2.4	5716		
11	Phenol	94	0.55	0.43	0.83	0.88	6.56	0.988	0.44	2.6	1938		
12	2,5-dimethylphenol	122	0.55	0.43	0.85	0.83	10.93	0.123	1.48	2.3	4390		
13	2,6-dimethoxyphenol	154	0.04	0.284	0.74	1.36	14.76	0.041	9.47	2.2	7335		
14	n-heptadecane	240	0.00	0.08	0.04	0.13	39.36	0.005	0.76	3.8	2875		
15	n-nonadecane	308	0.00	0.08	0.04	0.13	43.73	0.001	1.47	3.2	4771		
16	n-eicosane	324	0.00	0.08	0.04	0.13	45.91	0.001	1.93	2.9	5700		
17	2-Dodecanone	184	0.00	0.42	0.18	0.60	27.19	0.053	0.63	2.5	3076		
18	2-Tridecanone	198	0.00	0.42	0.18	0.60	29.38	0.022	0.97	2.3	4235		
19	Decanal	156	0.00	0.42	0.15	0.51	22.82	0.193	0.41	2.9	2018		
20	Citral	152	0.00	0.46	0.15	0.51	18.45	0.142	1.31	2.8	3928		
21	Benzaldehyde	106	0.00	0.47	0.15	0.51	7.52	0.982	1.66	3.3	3596		

^{a)} Based on the QSAR approach with PaDEL-Descriptor (Yap, 2011).

^{b)} Calculated through group contribution (Zhao et al., 1999; Stein and Brown, 1994).




582

583

584

Figure 1. FTIR spectra of the absorbing matter on the Teflon film. The absorbing matter was extracted with acetone and concentrated on a silicon disc. The FTIR spectra were obtained at the two different humidity (RH=0.1 and 0.8).

585



587 **Figure 2. Structure of the gas-wall partitioning model for various SVOCs.**




Figure 3. The observed concentrations of gaseous SVOCs ($C_{g,i}$, plot) in the chamber air for the two experiments at two different humidity (October 8, 2018 at RH=0.40 and May 15, 2018 at RH= 0.75) and the predicted $C_{g,i}$ (line) as a time series. The error bar associate with $C_{g,i}$ is estimated using standard deviation of the measured concentrations of internal standard (7–11 %).




Figure 4. The predicted $K_{w,i}$ with the QSAR-based GWP model as a function of RH.




Figure 5. The observed $K_{w,i}$ is plotted vs the estimated $1/C_i^*$ (a), activity coefficient (= 1) and the predicted $K_{w,i}$ (b) using activity coefficients estimated with the QSAR model (Eq. (14)) for various organic compound data sets from our study and literatures. The labels in (a) and (b) indicate the experiment summarized in Table 1 under different humidity conditions ($RH_a=0.4$, $RH_b=0.75$, $RH_c=0.53$, $RH_d=0.001$). The experimental data set of alkyl nitrate compounds is additionally obtained from Yeh and Ziemann (2014). The predicted $K_{w,i}$ is also plotted to $1/C_i^*$ for various SVOCs in different functionalities ((c), *n*-alkanes (C_{2n} , $n=6-15$), ketones (C_{2n} , $n=4-13$), 1-alcohols ($C_{12-C_{16}}$), 1-carboxylic acids (C_6-C_{17}), α,ω -diols (C_5-C_{11}), and α,ω -dicarboxylic acids (C_2-C_7)). This Figure shows that the $\gamma_{w,i}$ of SVOCs can influence the prediction of gas-wall partitioning in the order of the 4 (up to 10^4).




Figure 6. The characteristic time of the gas phase reaction of SVOC with an OH radical (τ_{OH}), gas-particle partitioning (τ_{GP}), gas-wall partitioning (τ_{GWP}), the reaction of SVOC in the organic phase (τ_{or}), and the reaction of SVOC in the inorganic phase (τ_{in}) are illustrated at the bottom (see Table S4). $K_{w,i}$ and $\alpha_{w,i}$ influence τ_{GWP} and reactivity scale influences τ_{or} and τ_{in} . The reactivity follows the class of the lumping in UNIPAR model (Fast and No reactivity). (Im et al., 2014; Beardsley and Jang, 2016; Zhou et al., 2019). The value on the bar chart indicates τ_{GWP} . The propagation error is calculated based on the uncertainties associated with parameters (Table S4).




Figure 7. The sensitivity (%) of the predicted $F_{g,i}$ (the vapor fraction of the total concentration of SVOC, Eq. (16)) to A/V, OM_{wall}, temperature, and humidity for *n*-nonadecane and 1-decanoic acid (a, b). The information inside the parentheses is associated with the reference value of $F_{g,i}$ and the variation of the variable. $F_{g,i}$ uncertainties (%) of the major model parameters, such as S_i , α_i , $H_{a,i}$, $H_{d,i}$, and P_L^o for *n*-nonadecane and 1-decanoic acid (c, d). The information inside the parentheses is associated with the range of uncertainties for each model parameter to predict $F_{g,i}$. The $F_{g,i}$ values at the standard condition for *n*-nonadecane and 1-decanoic acid are 0.23 and 0.17, respectively.