

Interactive comment on “How emissions uncertainty influences the distribution and radiative impacts of smoke from fires in North America” by Therese S. Carter et al.

Anonymous Referee #1

Received and published: 21 November 2019

The goal of this study presented in this paper is to quantify the uncertainties in the air quality and radiative impacts associated with BB aerosol emissions due to biomass (fuel) consumption and emission factors. The approach employed drives GEOS-Chem simulations with different BB emission inventories, two bottom-up (GFED4s and FINN) and two top-down (GFAS1.2 and QFED2.4). Results from the GEOS-Chem simulations are compared with observations of carbonaceous aerosols, black carbon (BC) and organic aerosol (OA), and satellite-derived aerosol optical depth (AOD) over North America. The authors find large differences in emissions and resultant air quality impacts across the four inventories explored. The authors conclude that differences between emission inventories in US, Alaska-Canada, and globally are driven by differ-

Printer-friendly version

Discussion paper



ences in fuel consumption, not emission factors.

The topic of this study is of great interest to the atmospheric chemistry and climate communities. The study was well designed and the authors are careful in their interpretation of the results. The manuscript is very well written and the presentation of methods and results is thorough, yet concise. I have only one significant issue with the paper and that is Section 6 (see comments below).

Specific Comments

L41: define AR5

L49-50: It should be noted that deep penetration of lungs and most acute health impacts are generally associated with fine PM (PM_{2.5}) fraction of PM.

L78-79: This insertion of “climate forcing” seems incorrect. Even if one defines climate forcing as a perturbation of the Earth’s energy balance, it need not be anthropogenic. Large volcanic eruptions result in climate forcing. And anthropogenic activities, e.g. land use and fire suppression (Andela et al., 2017), can reduce natural fire activity.

L169-170: Did the authors select the WRAP profile as opposed to Mu et al. (2011) since the focus was North America? Were there any complications/problems with using the WRAP cycle, which is intended to represent western US wildfires, for fires globally?

Mu et al. (2011) Daily and 3-hourly variability in global fire emissions and consequences for atmospheric model predictions of carbon monoxide, JGR Atmospheres, 116, D24303

L247-256: The authors should better describe the challenge of FRP methods using MODIS data associated with the sparse temporal coverage. The most significant weakness in FRP based methods using observations from MODIS is the need to estimate FRP (for time integration to get FRE) between temporally sparse observations. Under cloud free conditions, in mid-latitudes Terra & Aqua provide 4 observations a day, maybe 6 depending on swath overlaps. FRP methods often require estimating FRP

Printer-friendly version

Discussion paper



between the Aqua over-pass ~13:30 LT and Terra ~22:30 LT over-pass, the period of peak fire activity in the western US and western Canada. (At high latitudes swath overlap increases and temporal coverage is much better).

L266-267: 0.16 ug/m³ several orders of magnitude lower than typical field BBOA concentrations? That suggests typical field BBOA around 2000 ug/m³. Is this typical? PM₁ level of 2000 ug/m³ seems like a somewhat concentrated smoke plume. Please clarify.

L336-338: Liu et al. (2017) findings imply that the EFPM values used for western US wildfires may be higher than those used shown Fig 3. This should be clarified.

L343-353: Are the results for Boreal NA and CONUS similar if one uses EFOC to derive DM?

L383-384: It is likely that prescribed understory burning of forests in the southeast US are also a significant contributor to the CONUS springtime peak.

L395-398: “The southeastern US, in particular, 395 is of interest to the public health and policy communities because a prevalence of agricultural burning there, which dominates burned surface area (Nowell et al. 2018), may have a stronger impact on low altitude air quality in a relative sense than large wildfires that inject higher into the air.”

While agricultural fires comprise a large share of fires and area burned in the southeast it does not dominate surface burned area. Nowell et al. (2018) reports that in Florida “silviculture fires consumed the most area ($5.5 \pm 6.7 \times 10^5$ ha/year), burning 50% more than agricultural fires.”

More broadly across the southeast, the 2018 National Prescribed Fire Use Survey Report conducted by the National Association of State Foresters and the Coalition of Prescribed Fire Councils reported that 77% of prescribed fire acres burned in 2017 were forestry related compared to 23% agricultural. This breakdown is comparable to previous surveys released by the organizations in 2012 and 2015.

[Printer-friendly version](#)[Discussion paper](#)

<https://www.stateforesters.org/newsroom/nasf-coalition-of-prescribed-fire-councils-release-national-survey-on-prescribed-fire-use/>

<https://www.stateforesters.org/wp-content/uploads/2018/12/2018-Prescribed-Fire-Use-Survey-Report-1.pdf>

L454-461: It would be helpful compare FiNN burned area versus GFED over CONUS. Maybe just add a sentence comparing average annual burned area. I suspect this would indicate a large difference in burned area, especially in the west, as the authors suggest. GFAS and QFED estimate/interpolate FRP between MODIS observations, essentially gap-filling for the large time periods without observations.

Section 6: While I find this analysis and interpretation valid, it leaves the impression BB smoke is not relevant wrt population exposure. I believe the 24 hour average PM_{2.5} is the metric that should be used for BB health impacts (35 ug/m³ per NAAQS). Day to week length exposures to wildfire smoke are associated with negative health impacts, see e.g. Liu et al. (2015), Fisk and Chan (2017), Moeltner et al (2013), Williamson et al. (2016). In the western US, days with high PM_{2.5} or days where PM_{2.5} exceeds the NAAQS standard, tend to be associated with BB smoke (Liu et al., 2016; Brey et al., 2018; McClure and Jaffe, 2018). Section 6 should have focused on 24 hour average PM. I do not think it is necessary for the authors to do so; however, I think it would improve the study and perhaps without significant extra effort. If the authors decide to not include an analysis based on 24 hour average PM, then they need to discuss smoke – health impact linkages associated with day – week(s) exposure.

Fisk and Chan (2017) *Indoor Air*, 27, 191–204

Liu et al. (2015) *Environmental Research*, 136, 120-132

Liu et al. (2016) *Climatic Change*, 138, 655-666

Moeltner et al. (2013) *Journal of Environmental Economics and Management*, 66, 476-496

McClure and Jaffe (2018), PNAS, 115, 7901-7906

Williamson et al. (2016) Environmental Research Letters, 11, 125009

Technical Comments

L29-32: Jumbled / missing text sentence L164: “OA” or “POA”?

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2019-589>, 2019.

ACPD

Interactive
comment

Printer-friendly version

Discussion paper

