

Supplement to "Using ship-borne observations of methane isotopic ratio in the Arctic ocean to understand methane sources in the Arctic"

Berchet Antoine^{1,*}, Isabelle Pison¹, Patrick M. Crill², Brett Thornton², Philippe Bousquet¹, Thibaud Thonat¹, Thomas Hocking¹, Joël Thanwerdas¹, Jean-Daniel Paris¹ and Marielle Saunois¹

¹Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, IPSL, Gif-sur-Yvette, France.

²Department of Geological Sciences, Stockholm University, SE-10691 Stockholm, Sweden.

Correspondence: A. Berchet (antoine.berchet@lsce.ipsl.fr)




Figure 1. Average Sea Ice cover during summer 2014 (July–August). Data from NOAA/NSIDC passive microwave sea ice concentration climate data record (Comiso and Nishio, 2008).




Figure 2. Curtains plot of vertical wind speed (top panel) and simulated concentration from the top of the domain (bottom panel)




Figure 3. Vertical distribution of isotopic ratio as simulated by the model LMDZ averaged over the Arctic ocean for the year 2014.




Figure 4. Keeling plot for the observed (black) and simulated (blue/red before/after inversion) $\delta^{13}\text{C-CH}_4$ during the campaign.

References

Comiso, J. C. and Nishio, F.: Trends in the sea ice cover using enhanced and compatible AMSR-E, SSM/I, and SMMR data, *Journal of Geophysical Research: Oceans*, 113, <https://doi.org/10.1029/2007JC004257>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JC004257>, 2008.