1 Supplement of

2 21st Century Asian air pollution impacts glacier in northwestern 3 Tibet

4 M. Roxana Sierra-Hernández¹, Emilie Beaudon¹, Paolo Gabrielli^{1,2}, and Lonnie G. Thompson^{1,2}

¹Byrd Polar and Climate Research Center, The Ohio State University, Columbus, OH, 43210, USA

7 ²School of Earth Sciences, The Ohio State University, Columbus, OH, 43210, USA

Correspondence to: M. Roxana Sierra-Hernández (sierra-hernandez.1@osu.edu)

Table S1. Limit of detection (LOD), procedural blank (TE concentrations of the water used to make the artificial ice

45 core and of the ice from the artificial ice core), accuracy, and blanks average. LOD corresponds to three times the

46 standard deviation of the concentration of 10 measurements of ultrapure water (18.3 M Ω). The concentrations of the

47 Reference Material (TMRain-95) are reported as total concentrations accounting for the dilution factor of ~20.

		Procedu	Procedural Blank		Accuracy	
Trace	IOD	Ultrapure	Artificial	TMRain-95	TMRain-95	
Element	LOD	water	ice core	Found	Certified	
Ag (pg g^{-1})	0.1	0.5 ± 0.1	1 ± 0.01			
Al (ng g^{-1})	0.03	0.6 ± 0.6	0.9 ± 1.2	2 ± 0.9	2 ± 0.9	
As (pg g ⁻¹)	0.8	3 ± 0.7	4 ± 1	1126 ± 153	1070 ± 250	
Ba (pg g^{-1})	2	27 ± 11	32 ± 19	762 ± 59	730 ± 150	
Bi (pg g ⁻¹)	0.01	0.03 ± 0	0.04 ± 0.03	802 ± 13	630 ± 260	
$Cd (pg g^{-1})$	0.1	0.9 ± 0.5	1 ± 0.2	468 ± 14	480 ± 120	
$Co (pg g^{-1})$	0.2	0.3 ± 0.1	1 ± 0.5	227 ± 10	220 ± 37	
$\operatorname{Cr}(\operatorname{pg}\operatorname{g}^{-1})$	1	3 ± 3	7 ± 5	770 ± 37	790 ± 170	
Cs (pg g ⁻¹)	0.1	1 ± 0.3	2 ± 0.5			
Cu (pg g ⁻¹)	1	21 ± 4	27 ± 5	6305 ± 101	6200 ± 930	
Fe (ng g^{-1})	0.2	0.1 ± 0.2	0.4 ± 0.3	24 ± 10	24 ± 4	
Ga (pg g ⁻¹)	0.5	2 ± 2	3 ± 3			
$\text{Li}(\text{ng g}^{-1})$	0.04	0.8 ± 0.3	0.7 ± 0.1	0.3 ± 0.2	0.4 ± 0.08	
Mg (ng g ⁻¹)	0.02	0.2 ± 0.1	0.2 ± 0.1			
Mn (pg g ⁻¹)	1	4 ± 5	5 ± 2	6013 ± 77	6100 ± 780	
Mo (pg g ⁻¹)	0.2	0.5 ± 0.2	1 ± 0.1	174 ± 7	170 ± 100	
Na (ng g ⁻¹)	0.4	1 ± 0.7	2 ± 0.7			
Nb (pg g ⁻¹)	0.2	5 ± 2	10 ± 6			
Ni (pg g^{-1})	0.8	3 ± 0.6	3 ± 0.8	845 ± 35	800 ± 170	
Pb (pg g^{-1})	0.3	0.43 ± 0.2	0.8 ± 0.5	281 ± 5	290 ± 93	
$Rb (pg g^{-1})$	1	18 ± 18	25 ± 16			
Sb (pg g^{-1})	0.1	0.1 ± 0.03	0.1 ± 0.01	322 ± 7	350 ± 100	
Sn (pg g^{-1})	4	2 ± 0.5	2 ± 0.7			
$Sr(pg g^{-1})$	5	285 ± 133	296 ± 132	1729 ± 58	1700 ± 260	
Ti (pg g ⁻¹)	10	21 ± 26	31 ± 24			
$Tl (pg g^{-1})$	0.02	0.03 ± 0.01	0.05 ± 0.02	330 ± 6	330 ± 72	
U (pg g ⁻¹)	0.03	0.07 ± 0.01	0.09 ± 0.02	262 ± 5	250 ± 60	
$V(pg g^{-1})$	1	4 ± 4	7 ± 5	678 ± 39	640 ± 120	
Zn (pg g ⁻¹)	3	8 ± 4	5 ± 1			

54 concentration data set (1971–2015).

ТЕ	Factor 1	Factor 2	Factor 3	Communality
Ag	0.92	-0.29	0.16	0.95
Al	0.96	-0.24	0.07	0.99
As	0.83	-0.50	0.02	0.94
Ba	0.85	-0.47	0.07	0.96
Bi	0.94	-0.25	0.15	0.97
Cd	0.79	-0.30	0.50	0.97
Co	0.96	-0.26	0.08	0.99
Cr	0.96	-0.26	0.08	0.99
Cs	0.95	-0.25	-0.02	0.97
Cu	0.93	-0.32	0.10	0.98
Fe	0.96	-0.25	0.06	0.99
Ga	0.96	-0.26	0.06	0.99
Li	0.79	-0.56	0.01	0.95
Mg	0.80	-0.57	0.05	0.96
Mn	0.84	-0.43	0.18	0.92
Mo	0.41	-0.85	0.08	0.90
Na	0.09	-0.96	0.03	0.93
Nb	0.94	-0.28	-0.02	0.96
Ni	0.95	-0.28	0.10	0.99
Pb	0.93	-0.24	0.24	0.97
Rb	0.94	-0.32	-0.01	0.99
Sb	0.78	-0.57	0.04	0.93
Sn	0.82	-0.36	0.30	0.89
Sr	0.20	-0.95	0.07	0.94
Ti	0.79	-0.58	0.04	0.96
Tl	0.92	-0.35	0.11	0.98
U	0.82	-0.50	0.08	0.93
V	0.95	-0.29	0.07	0.99
Zn	0.94	-0.24	0.21	0.99
Variance (%)	72.6	21.4	2.1	
Cum. Variance (%)	72.6	94.0	96.1	

59 Guliya's 1650–2015 trace element records

60 In Thompson et al. (2018), we showed the high reproducibility between the 1992 and 2015 Guliya δ^{18} O profiles. This 61 reproducibility is also observed in the TE records (Figure S1). Likewise, the Al and Fe (not shown) median 62 concentrations are 0.3 µg g⁻¹ in both records during the 1971–1991 period in which both TE records overlap.

63 Figure S1 displays 5-year median concentrations and EFs of the 1992 (1650–1991) and 2015 (1971–2015) cores for 64 TEs that showed post-1850s enrichments (Pb, Cd, Zn, and Al for comparison). The 5-year median concentrations of 65 Cd, Pb, and Zn are slightly higher in the 2015 record than the 1992 time series for the 1971–1991 period. For example, the Cd median concentrations are 5 and 6 pg g-1 in the 1992 and 2015 records, respectively. The difference in 66 67 concentrations between the 1992 and the 2015 records is not significant (Mann–Whitney test: p < 0.0005 for medians) 68 and may be due to spatial variability of ice layers between the two boreholes. Similarly, during the 1971–1991 period, 69 the EFs in the 2015 record are slightly higher than in the 1992 record. This might result from the natural signal to 70 noise ratio differences between the two records. Despite the slight EF differences between the two records during the 71 1971–1991 period, the reproducibility of TEs allows determination of temporal trends from pre-industrial times

72 (~1650) into the 21^{st} century (2015) using the 1992 and the 2015 TE records.



Figure S1. Pb, Cd, Zn, and Al shown as 5-year median concentrations, excess concentrations and enrichment factors

- 75 (EF) from the 1992 Guliya ice core (1650–1991) and the 2015 Guliya ice core (1971–2015; thick line). The
- horizontal dotted lines show the 1650–1991 concentration and EF medians for the 1992 core.



Figure S2. Comparison of Factor 1 scores (crustal contribution), dust concentrations (particles ml⁻¹) (Thompson et al., 2018), and concentrations of the typical crustal TEs Fe and Al. All data are presented as five-year running means.

81



Figure S3. Comparison of Factor 2 scores (evaporitic contribution), NO₃⁻ ion concentrations (Thompson

et al., 2018), and TEs concentrations of sodium (Na) and strontium (Sr). All data are presented as five-

85 year running means.

86



Figure S4. Cluster analysis extracted from the first three factors during the 1971–2015 period.



94

95 Figure S5. (a) Metal production in China (Zn, Pb, Cu, and Ni), Pakistan (Pb, Cu), India (Zn, Pb, Cu) and

96 Kazakhstan (Zn, Pb, Cu) (BGS, 2015) and (b) PM_{2.5} from industrial processes (including the production of cement,

97 lime, chemicals, and metal production). The Guliya EF composite (average of Cd, Pb, Zn, and Ni EF z-scores) is

shown at the bottom of each panel for comparison. The two Guliya maxima at 2000 and 2008 are shown as shaded

99 bars.



100

101 Figure S6. (a) Phosphate fertilizer consumption (FAO, 2019) and (b) PM_{2.5} from agricultural activities (including

direct soil emission, rice cultivation, and manure management) (EDGARv4.3.2, 2017; Crippa et al., 2018). The

103 Guliya EF composite (average of Cd, Pb, Zn, and Ni EF z-scores) is shown at the bottom of each panel for

104 comparison. The two Guliya maxima at 2000 and 2008 are shown as shaded bars.

105 References

106 BGS, British Geological Survey. World Mineral Statistics and World Mineral Production, 2/1/2018, 2015.

107

108 Crippa, M., D. Guizzardi, M. Muntean, E. Schaaf, F. Dentener, J. A. van Aardenne, S. Monni, U. Doering, J. G. J.

Olivier, V. Pagliari and G. Janssens-Maenhout, Gridded emissions of air pollutants for the period 1970–2012 within
EDGAR v4.3.2. Earth Syst. Sci. Data, 10, 1987–2013, 10.5194/essd-10-1987-2018, 2018.

111

EDGARv4.3.2, European Commission, Joint Research Centre (EC-JRC)/Netherlands Environmental Assessment
Agency (PBL). Emissions Database for Global Atmospheric Research (EDGAR), release EDGAR v4.3.2 (1970 –
2012), 2017.

115

FAO, Food and Agricultural Organization of the United Nations . Phosphate Fertilizer Consumption, 4/22/2019, 2019.

- 118 Thompson, L. G., T. Yao, M. E. Davis, E. Mosley-Thompson, G. Wu, S. E. Porter, B. Xu, P.-N. Lin, N. Wang, E.
- 119 Beaudon, K. Duan, M. R. Sierra-Hernández and D. V. Kenny, Ice core records of climate variability on the Third Pole
- 120 with emphasis on the Guliya ice cap, western Kunlun Mountains. Quat. Sci. Rev., 188, 1-14, 2018.
- 121