

The global impact of bacterial processes on carbon mass

Barbara Ervens and Pierre Amato

5 Université Clermont Auvergne, CNRS, Sigma-Clermont, Institut de Chimie de Clermont-Ferrand, 63000 Clermont-Ferrand, France

Correspondence to: barbara.ervens@uca.fr and pierre.amato@uca.fr

Abstract. Many recent studies have identified biological material as a major fraction of ambient aerosol loading. A
10 small fraction of these bioaerosols consist of bacteria that have attracted a lot of attention due to their role in cloud
formation and adverse health effects. Current atmospheric models consider bacteria as inert quantities and neglect cell
growth and multiplication. We provide here a framework to estimate the production of secondary biological aerosol
(SBA) mass in clouds by microbial cell growth and multiplication. The best estimate of SBA formation rates of 3.7 Tg
yr⁻¹ is comparable to previous model estimates of the primary emission of bacteria into the atmosphere, and thus might
15 represent a previously unrecognized source of biological aerosol material. We discuss in detail the large uncertainties
associated with our estimates based on the rather sparse available data on bacteria abundance, growth conditions and
properties. Additionally, the loss of water-soluble organic carbon (WSOC) due to microbial processes in cloud droplets
has been suggested to compete under some conditions with WSOC loss by chemical (OH) reactions. Our estimates
suggest that microbial and chemical processes might lead to a global loss of WSOC of 8 - 11 Tg yr⁻¹ and 8 - 20 Tg yr⁻¹,
20 respectively. While also this estimate is very approximate, the analysis of the uncertainties and ranges of all
parameters suggests that high concentrations of metabolically active bacteria in clouds might represent an efficient
sink for organics. Our estimates also highlight the urgent needs for more data concerning microbial concentrations,
fluxes and activity in the atmosphere to evaluate the role of bacterial processes as net aerosol sink or source on various
spatial and temporal scales.

25 1. Introduction

The characterization and quantification of outdoor bioaerosols is an active field of current atmospheric research since
bioaerosols have been suggested to contribute to adverse health effects and cloud formation as ice-nucleating particles
(Després et al., 2012). Biological material includes debris, pollen, bacteria, fungal spores, and viruses and is usually
considered as being directly emitted to the atmosphere (primary biological aerosol, PBA (Jaenicke, 2005)). The total
30 number and mass concentrations of PBA particles vary widely in space and time: Posfai et al. (1998) found 1% of
particles with biological material above the Southern Ocean whereas Artaxo et al. (1990) identified more than 90% of
all particles to contain biological material during the wet season in the Amazon. In an urban/remote region in Germany,
24% of all particles were found to include a biological fraction (Matthias-Maser and Jaenicke, 2000). Similar
concentrations were observed at a remote high-altitude site with 16 - 64% of the mass of particles with diameters of
35 less than 10 µm being composed of biological mass (Wiedinmyer et al., 2009) whereas the PBA number fraction was

much smaller (0.3 - 18%) in Rome, Italy (Perrino and Marcovecchio, 2016). Bacteria only comprise a small fraction of the total biological aerosol mass but they alone can contribute up to about ~20% of the total number of particles with diameters greater than 0.5 μm (Bowers et al., 2012).

40 Near the ground, typical concentrations of total airborne bacteria range from ~100 to 10^6 cells m^{-3} , depending on the emission source (Burrows et al., 2009b), and on temporal, meteorological, and other environmental conditions influencing its propensity to emit particles to the air (Carotenuto et al., 2017; Huffman et al., 2013; Lighthart, 1997; Lighthart and Shaffer, 1995). Atmospheric mixing aloft tends to homogenize the number and diversity of the various bacteria types as the distance from sources increases. In the free troposphere, concentrations of ~10,000 cells m^{-3} are reported, including in clouds (DeLeon-Rodriguez et al., 2013; Väitilingom et al., 2013). Some extent of selection
45 toward certain species of bacteria probably occurs during aerosolization and atmospheric transport (Joly et al., 2015; Michaud et al., 2018). However, such selection has not been clearly proven yet as the bacterial assemblages found at high altitude often resemble those observed near the ground (Amato et al., 2017; DeLeon-Rodriguez et al., 2013; Smith et al., 2018).

The atmosphere is a harsh environment for living microorganisms: low temperatures at high altitude, UV radiation
50 (Madronich et al., 2018) and high free radical concentrations (Haddrell and Thomas, 2017; Marinoni et al., 2011) are thought to greatly challenge living organisms (Amato et al., 2019; Joly et al., 2015; Smith et al., 2011). Additionally, the rapidly changing conditions in clouds, like condensation/evaporation and freeze/thaw cycles, can cause strong physiological shocks and physical damages to cells, which can eventually be lethal. The viability of airborne microorganisms is thus very variable in space and time depending on environmental conditions (Fahlgren et al., 2010; Hu et al., 2017; Lighthart and Shaffer, 1995; Monteil et al., 2014), but yet the fact that a fraction of bacteria cells are
55 viable was shown in many experiments of microbiological cultures from ambient aerosol samples (Amato et al., 2007b; Bovallius et al., 1978; Lighthart, 1997; Newman, 1948). This was specified and quantified more recently by direct observations and measurements of biological activity imprints (Amato et al., 2007a, 2017; Sattler et al., 2001; Wirgot et al., 2017). The multiplication of airborne bacteria was observed from aerosols generated from bacteria cultures
60 (Dimmick et al., 1979), as well as in natural polluted fog (Fuzzi et al., 1997). Thus, the estimated PBA emissions might be biased high as SBA formation provides an additional source of bacteria mass and, thus, observed bacteria concentrations represent the sum of emission fluxes that are smaller than assumed and the secondary production in the atmosphere.

Efficient bacteria cell growth and multiplication are largely constrained by the presence of liquid water (Davey, 1989; Haddrell and Thomas, 2017). One can thus assume that microbial processes in the atmosphere are limited to the time
65 microorganisms spend in clouds (*Figure 1*). Cell growth and multiplication lead to an increase of the initial cell mass and, thus, to more biological material (Kaprelyants and Kell, 1993; Norris, 2015; Si et al., 2017) whereas bacteria dormancy and death do not lead to any change in cell mass (Engelberg-Kulka et al., 2006; Kaprelyants and Kell, 1993; Price and Sowers, 2004). We introduce the term ‘secondary biological aerosol’ (SBA) mass here in order to distinguish
70 this aerosol source from directly emitted PBA. Heterotrophic bacterial processes require the uptake of organic substrates by the cells, which are subsequently converted by metabolic processes into new organic products, biochemical energy and CO_2 (‘respiration’, *Figure 1*). These substrates include organics (e.g., carboxylic acids, sugars); other elements (e.g., nitrogen, phosphorous, potassium, metals) are also needed and exist in bioavailable forms

in cloud water. The biotransformation of formate, acetate, succinate, lactate, oxalate, and formaldehyde (Ariya et al., 2002; Vaitilingom et al., 2010), phenol (Lallement et al., 2018) and methane (Šantl-Temkiv et al., 2013) by bacteria and fungi was studied in aqueous solution mimicking the typical chemical composition of cloud water, and it was suggested that under specific conditions, microbial processes might be competitive to chemical radical processes as sinks for these compounds (Delort et al., 2010; Vaitilingom et al., 2011, 2013). The efficiency of such metabolic processes strongly depends on the bacteria types, substrates and their availability within the cloud droplets. In the present study, we perform an estimate of the global importance of SBA formation and microbial WSOC loss. All parameters and their uncertainties are discussed based on the sparse data sets currently available.




Figure 1: Bacterial processes in the atmosphere leading to SBA formation and loss of water-soluble organic carbon (WSOC) in clouds.

2. Data and assumptions on bacterial processes in clouds

2.1 Atmospheric concentrations of bacteria cells

Burrows et al. (2009a, 2009b) have summarized data on number concentrations and emission fluxes of bacteria above various ecosystems on the Earth surface. Their compilation also includes the estimates of cell concentrations near the surface in a range of $10,000 \text{ m}^{-3}$ (seas) up to $650,000 \text{ m}^{-3}$ (urban). Bacteria populations aloft represent a mixture of bacteria that were emitted from different ecosystems and subsequently mixed (Burrows et al., 2009b). Despite these mixing processes, there are bacteria types that can be considered characteristic for each ecosystem (Wéry et al., 2017). **Table 1** lists cell concentrations as published by Burrows et al. (2009b) complemented by some more recent measurements. We extend this overview by data on bacteria types, suggested as predominant or characteristic for each ecosystem. In several cases, more than one predominant bacteria type is listed as specific geographical, meteorological and other environmental conditions might lead to differences in the diversity of bacteria populations for the same category of ecosystem. We also provide global average data (Category ‘All’) and define one of the most abundant

95 bacteria type (Alpha-Proteobacteria) alive in the atmosphere (Amato et al., 2019; Klein et al., 2016) as a representative type. Several studies report total concentrations of bacteria cells in the atmosphere whereas others present only the concentration of viable cells. The complexity of distinguishing viable, cultivable and dead bacteria cells in the atmosphere has been discussed in several studies (Burrows et al., 2009a; Otero Fernandez et al., 2019).

100 **Table 1:** Summary of ambient cell concentrations C_{cell} and generation rates R_{cell} for predominant bacteria types in all ecosystems

	$C_{Cell} [m^{-3}]^a$	Representative strain affiliation	Generation rate $R_{Cell} [h^{-1}]$
All	10,000	Alpha-Proteobacterium <i>Sphingomonas</i> sp. (average of 32b-11, 32b-49, 32b-57, 35b-32, 35b-38) ^e	0.06 (5°C) ^e 0.2 (17°C) ^e 0.35 (27°C) ^e 0.45 (37°C) ^e
Tundra	12,000	<i>Pseudomonas</i> spp. (<i>P. graminis</i>) ^{e, f, g}	0.12 (5°C) ^e 0.21 (17°C) ^e 0.82 (27°C) ^e 0.27 (37°C) ^e
Grassland	110,000	<i>Psychrobacter</i> sp. ^g	0.0007 (-10°C) ^g
		<i>Rhodococcus</i> sp. ^g	0.0001 (-10°C) ^g
		<i>Pseudomonas syringae</i> ^{e, h}	0.1 (5°C) ^e 0.25 (15°C) ^e 0.9 (27°C) ^e
Coastal	76,000	<i>Frigoribacterium</i> sp. ^{e, i}	1.7·10 ⁻⁴ -2.91·10 ⁻⁴ (12-18°C) ^j
Wetlands	90,000		
Crops	110,000		
Land ice	(5,000)		
Deserts	(10,000) 612 ^c		
Forests	56,000 6,323-12,748 ^d		
Shrubs	350,000		
Seas	10,000	<i>Pseudoalteromonas</i> ^k	0.25 (T unknown) ^k
Seas (estuary) ^b		Gamma-Proteobacterium (the fastest) ^l	0.17 (14°C) ^l 0.19 (24°C) ^l

^a All cell concentrations are taken from Burrows et al. (2009b) unless otherwise noted. ^b 'seas estuary' was not included as a separate ecosystem by Burrows et al. (2009b); ^c (Lighthart and Shaffer, 1994); ^d (Helin et al., 2017); ^e Amato et al. (unpublished data; strains originally reported in Amato et al., 2007c and Väitilingom et al., 2012); ^f (Männistö and Häggblom, 2006); ^g (Bakermans et al., 2003); ^h; (Morris et al., 2000) ⁱ (Copeland et al., 2015); ^j (Stibal and Elster, 2005); ^k (Middelboe, 2000) ; ^l (Fuchs et al., 2000)

We assume in **Section 3** that all bacteria cells as listed in **Table 1** are metabolically active. The atmospheric lifetime of

105

bacteria cells is limited to several minutes (Otero Fernandez et al., 2019) to hours (Amato et al., 2015). In our estimates, we neither include assumptions on the limited lifetime of bacteria cells nor on their residence time in the atmosphere as it is assumed that PBA emissions lead to a continuous replenishment of bacteria in the atmosphere resulting in a steady-state concentration of living cells. The consequences of limited cell life- and residence time on SBA formation warrant further studies in more sophisticated model approaches.

2.2 Cell generation rates R_{Cell}

Different levels of metabolic activity can be distinguished, from survival, where cells only repair molecular damages, to maintenance (dormancy), where cells do not divide but maintain biological functions, to growth, allowing the net production of biological mass (Price and Sowers, 2004) (**Figure 1**). The generation rate of a microorganism during growth is probably the most common microbiological criterion used for characterizing microbial multiplication in the laboratory; it corresponds to the time that is needed for doubling the cell number, i.e., for producing two “children” cells from one individual. This requires mass production from nutrients that provide the necessary molecular bricks and biochemical energy. The activity depends on physiological traits of the microorganism, with optima at a given temperature, pH, salinity, and other conditions that define its fitness for its habitat. The generation time in bacteria at their optimum growth conditions usually ranges from ~20 minutes (Marr, 1991) to several days or weeks; as conditions deviate from the optima, this lengthen to virtually infinite time in non-dividing cells.

The *cellular* growth rate itself, i.e. the increase of individual cell size and mass, is intimately linked with generation time: cell size increases in a predictable way as generation time decreases (Si et al., 2017), and it can vary by a factor of up to eight within a single bacteria species. Compared to generation rates, cellular growth rates are usually small and, thus, in the following only data for generation rates are used to estimate SBA mass formation rates.

Metabolic activity, in terms of carbon uptake per units of biomass and time, can range over more than ten orders of magnitude, depending on many factors of which temperature is a major one (Price and Sowers, 2004). Therefore, if available, temperature-dependent generation rates R_{cell} are listed in **Table 1** and shown in **Figure 2**. In addition, the highest expectable growth rate for bacteria as measured under laboratory conditions in culture medium is also shown in the figure for constraining an upper theoretical limit. This corresponds to the generation rate of the laboratory model *Escherichia coli* under optimal conditions (37°C). However, it can be expected that this is not representative of situations encountered in clouds.

Generally, the temperature dependence of cell generation rates can be scaled by the empirical relationship in Eq-1:

$$R_{cell}(T_2) = R_{cell}(T_1) \cdot Q_{10}^{(T_2-T_1)/10} \quad (1)$$

whereas $R(T_1)$ and $R(T_2)$ are the growth rates (h^{-1}) at two temperatures T_1 and T_2 . Q_{10} is a dimensionless scaling factor that expresses the change of these rates over an interval of 10°C and typically has values between two and three within relatively small temperature intervals (Lipson et al., 2002; Sand-Jensen et al., 2007). In general, Equation (1) can be applied for all bacteria types and is usually valid for liquid water over a temperature range up to ~ 25°C; however, the slope (Q_{10} factor) and the maximum temperature depends on the bacteria type. In **Figure 2**, the dashed lines towards

lower temperatures represent extrapolations of the generation rates at ~20°C reflecting the general agreement between measured and calculated temperature dependencies using $Q_{10} = 2$ or 3, respectively. Using generation rates measured at ~20°C might lead to an overestimate for SBA mass formation rates in colder clouds. However, we chose these values for the calculations in Section 3 as most experimentally-derived growth rates are available for temperatures of ~ 20 – 30°C. Generally, at temperatures below 0°C, cell metabolic activity is negligible in terms of carbon flux even though cells can maintain and survive under such conditions (Amato et al., 2009, 2010; Price and Sowers, 2004).




Figure 2: Temperature dependence of generation rates R_{cell} for bacteria types representative for the ecosystems in **Table 1** and *E. coli* as a likely upper limit for cell generation. Dashed lines are extrapolations of the rates at $T \sim 20^\circ\text{C}$ using $Q_{10} = 2$ or $Q_{10} = 3$, respectively.

2.2 Bacteria growth efficiency (BGE)

Chemoheterotrophs - representatives of which were shown to maintain metabolic activity in clouds (Amato et al., 2017) - take up carbon from dissolved organic material for both recovering biochemical energy and converting the substrates into CO_2 and other products. Bacteria growth efficiency (BGE) is defined as the biological mass that is produced relatively to the amount of carbon taken up from the environment, the rest being converted into CO_2 (Eiler et al., 2003):

$$\text{BGE} = \frac{d[\text{organic products}]}{d[\text{organic products}] + d[\text{CO}_2]} \quad (2)$$

Note that in the original literature BGE is defined as a measure of ‘biomass production’ instead of organic products’ in Eq (2). Given the large body of atmospheric literature on aerosol processes that discusses ‘biomass’ as material from any living matter (e.g., aerosol from forest fires), we avoid using ‘biomass’ in the current context of microbial processes. BGE for planktonic bacteria range from < 0.4% to 80% with the highest values for eutrophic conditions (Eiler et al., 2003). In turn, in the same study, it was shown that, when substrate availability is limited, values from ~7% to ~14% are generally observed. As the conditions in cloud water can be considered oligotrophic with typical concentrations of dissolved organic carbon (DOC) of less than 0.1 mM (Herckes et al., 2013), low BGEs in the range of 0.1 – 10% can be expected, i.e. DOC is efficiently converted into CO_2 .

2.4 Cloud properties relevant for microbial activity

2.4.1 Cell concentrations in cloud water

165 Bacteria cells have sizes of up to several micrometers which explains their high efficiency to act as cloud condensation
nuclei (CCN) (Bauer et al., 2003; Després et al., 2012). Assessing the hygroscopicity of biological particles is complex
since it cannot be calculated in a similar way as for chemical compounds where total hygroscopicity represents the
sum of the contributions of all components (Ariya et al., 2009). Once particles form cloud droplets, chemical
compounds dissolved in cloud water, will trigger the growth of the processed CCN and, thus, enhancing hygroscopicity
and CCN activity of aged particles in subsequent cloud cycles. The dissolution of ambient cell populations of 100 to
170 50,000 m⁻³ (**Table 1**) results in 200 to 500,000 cells mL⁻¹ for clouds with liquid water contents (LWC) of 0.5 g m⁻³ and
0.1 g m⁻³. The reasonable agreement of cell concentrations outside of clouds and those in cloud water suggests that a
large fraction of bacteria cells are scavenged and, thus, act as CCN.

Some bacteria are well-known to efficiently act as ice nuclei (Amato et al., 2015; Möhler et al., 2008; Morris et al.,
2004). In the current study, we neglect the potential role of ice clouds as media of microbial metabolic activity. In
175 addition to low temperatures resulting in very low generation rates (**Figure 2**), the substrate diffusion to the bacteria
will be limited resulting in negligible consumption of dissolved carbon.

2.4.2 Time fractions of microbial processes in clouds (F_{cloud})

As we assume that both SBA mass production and WSOC loss only occur when bacteria are suspended in cloud
droplets, we need to estimate the time bacteria spend in liquid clouds. In general, cloud contact times, i.e., the time air
180 spends in a cloud, are dependent on cloud depth and vertical velocity (Feingold et al., 2013). This small-scale
information is not consistently available for the large regions as covered by the ecosystems listed in **Table 1**. In order
to give an estimate of the cloud processing time over the various large ecosystems as identified by Olson et al. (1992),
we use the approximate cloud fractions during spring averaged for 2000 – 2011 from MODIS Terra (e.g., Figure 2b
by King et al. (2013)). This representation gives only a general view of cloudiness that varies over smaller spatial and
185 temporal scales. However, given the conceptual nature of our study that builds upon the categories as used in the
previous study by Burrows et al. (2009) for primary bacteria emissions, our approach seems sufficient to give (i) a
reasonable estimate of cloudiness above the various ecosystems and (ii) enough detail of its concept to be refined in
future studies.

Globally, a range of cloud thicknesses of 1.4 – 1.9 km has been derived (Table 1 in (Wang et al., 2000)) from which
190 we use $h = 1.5$ km as a single value for the average cloud thickness. Assuming further that globally > 90% of all liquid
clouds reside in the lowest 6 km of the atmosphere ($\Delta z = 6$ km) (Pruppacher and Jaenicke, 1995), we can convert the
cloud coverage as obtained from satellite data (F_{clc} in Table 2) into cloud volume fractions using Eq-3:

$$F_{cloud} = F_{clc} \cdot \frac{\text{cloud thickness [km]}}{\Delta z [\text{km}]} = F_{clc} \cdot \frac{1.5 \text{ km}}{6 \text{ km}} \quad (3)$$

195 Comparison of previous estimates of global cloud coverage of 60% (Pruppacher and Jaenicke, 1995) and the volume

fraction of liquid clouds within the atmosphere of 15% (Lelieveld and Crutzen, 1990) generally supports this relationship. The resulting F_{cloud} values are summarized in **Table 2** together with the percentage area fraction of each ecosystem of the Earth surface, taken from Burrows et al. (2009b) and originally obtained from Olson (1992) and the cloud coverage data (F_{clc}).

200

Table 2: Surface coverage of ecosystems on Earth surface (Burrows et al., 2009b), approximate cloud coverage F_{clc} above the ecosystems, estimated based on maps of annual cloud cover data obtained by MODIS, and estimated time fraction bacteria spend in clouds (F_{cloud})

	% of Earth surface ^{a)}	F_{clc}	F_{cloud}
All	100	0.6	0.15
Tundra	3.3	0.4	0.1
Grassland	2.2	0.7	0.2
Coastal	0.2	0.4	0.1
Wetlands	0.6	0.5	0.15
Crops	3.0	0.7	0.3
Land ice	3.1	0.4	0.1
Deserts	3.7	0.2	0.05
Forests	7.0	0.9	0.25
Shrubs	5.8	0.3	0.1
Seas	71.0	0.7	0.2

205

^{a)} Data from Burrows et al. (2009b) and Olson (1992)

3. Results and Discussion

3.1 SBA mass production

3.1.1 Calculation of SBA formation rates

We calculate the SBA mass formation rate [$\text{ng m}^{-3} \text{day}^{-1}$] above each ecosystem i , using Eq (4):

$$210 \quad \left(\frac{dm}{dt}\right)_{SBA,i,day} = R_{cell,i} \cdot F_{live} \cdot C_{cell,i} \cdot F_{cloud,i} \cdot m_{cell} \quad (4)$$

where R_{cell} is the cell generation rate [h^{-1}] (**Table 1**). For ecosystems, for which R_{cell} of the representative bacteria types is not available, we assume the average formation rate of $R_{cell} = 0.3 \text{ h}^{-1}$ as an upper limit for atmospherically-relevant conditions, corresponding to a generation time of approximately three hours. C_{cell} denotes the ambient cell concentration [cell m^{-3}] (**Table 1**), F_{live} is the fraction of living cells in total cell concentration and assumed to be unity here, F_{cloud} is the fraction of total time when bacteria are active in clouds (**Table 2**), and m_{cell} is the average mass of a single cell, independent of the bacteria type. The cell mass m_{cell} is assumed to be $52 \cdot 10^{-15} \text{ g cell}^{-1}$ (Sattler et al., 2001), equivalent of a spherical particle with diameter of 500 nm and a density of 1 g cm^{-3} .

For nearly all ecosystems, predicted SBA formation rates are in the range of ~ 0.1 to $\sim 1 \text{ ng m}^{-3} \text{ s}^{-1}$ (**Figure 3a**), with higher values for crops and shrubs where C_{cell} were found to be highest (**Table 1**). The average value ($0.6 \text{ ng m}^{-3} \text{ day}^{-1}$), calculated using the average values representative for all ecosystems (Category ‘All’ in Table 1), is similar to most of the formation rates in the individual ecosystems, suggesting that using these average data for a global estimate gives

220

a reasonable order of magnitude of SBA formation. Only above land ice where C_{cell} is small, the rate is significantly smaller. Given that the temperatures above land ice might be on average lower than above other regions, the relative importance of SBA formation there might be even smaller. According to the definition of the categories as suggested by Olson et al. (1992), the category ‘land ice’ does not include sea ice. It can be expected that above sea ice the sources and metabolic activity of bacteria are also very low (Martin et al., 2009) and thus can be likely neglected on a global scale.

To compare the mass production to other global aerosol sources, $m_{SBA,day}$ is converted into a production flux [Tg yr⁻¹] for each ecosystem i and scaled by the surface fraction of each ecosystem A_i :

$$P_{SBA,i} = \left(\frac{dm}{dt} \right)_{SBA,i,day} \cdot 365 \text{ days} \cdot A_i \cdot V_{atmos} \quad (5)$$

where V_{atmos} is the volume of atmosphere ($3 \cdot 10^{18} \text{ m}^3$) and A_i is the surface fraction of each ecosystem (**Table 2**).




Figure 3: Predicted production of secondary biological aerosol mass above the various ecosystems. The blue bar indicates the predicted production using the average values for all ecosystems (‘all’ in Table 1); a) SBA formation rates [$\text{ng m}^{-3} \text{ day}^{-1}$]; b) SBA production rates [Tg year^{-1}]. The red shaded bar represents the sum of the contributions from all ecosystems, bacteria emissions are shown for comparison, taken from Burrows et al. (2009b)

The production fluxes for each ecosystem are shown in **Figure 3b**, together with their sum for all ecosystems. The total predicted amount of SBA production is 3.7 Tg yr⁻¹ with highest contributions from bacterial activities above seas (0.5 Tg yr⁻¹) as they cover most of the globe (71%) and above shrubs (0.5 Tg yr⁻¹) since the highest bacteria concentrations have been identified there (**Table 1**). Forests with a much smaller surface area (7%) but higher cell concentration contribute the same amount (0.5 Tg yr⁻¹). Using average data instead of those for individual ecosystems results in 0.7 Tg yr⁻¹. Given the large uncertainties in all factors of Eq (4) and Eq (5), we suggest that and the value based on the weighted sum of all ecosystems (3.7 Tg yr⁻¹) might be a reasonable 'first best estimate' of total SBA contribution by bacteria on a global scale. These values are similar to the range of 1 - 10 Tg yr⁻¹ that was extrapolated by Sattler et al. (2001) based on carbon production rates of bacteria in supercooled clouds at mount Sonnblick observatory in the Austrian Alps. Our estimated SBA mass production represents the production of total bacteria mass. The carbon content of bacteria cells is roughly 50% of their dry mass, with the remainder composed of nitrogen, oxygen, phosphorous, hydrogen and other elements (Whitman et al., 1998). Thus, we suggest that SBA formation may lead to ~1.9 Tg carbon yr⁻¹ based on our best estimate that is bound in biological mass.

245 **3.1.2 Discussion of uncertainties in SBA formation**

The formation rates in **Section 3.1.1** represent an estimate of a previously unrecognized source of biological aerosol mass in the atmosphere. All parameters are associated with large uncertainties that need to be constrained in the future as they might vary depending on temporal, meteorological, spatial and geographical conditions. Ranges of observed values for all parameters in Eq (4) and Eq (5) are summarized in **Table 3** and discussed in the following:

250 (i) The cell concentrations C_{cell} in the atmosphere homogenize aloft due to mixing processes and average to concentrations of $\sim 10^4$ m⁻³ at most locations. However, spatial deviations might be present in particular locations, such as cell concentrations of $\sim 7 \cdot 10^5$ - $4 \cdot 10^6$ cells m⁻³ that were found during haze periods in China (Li et al., 2018; Xie et al., 2018), and even 10^9 m⁻³ above a wastewater storage lagoon which can be considered the highest expectable value of ambient bacteria (Paez-Rubio et al., 2005). Using the framework presented in the present study, SBA formation in such rather spatially limited areas can be estimated if growth rates of the individual bacteria types were available.

(ii) The growth rates R_{cell} assumed here likely represent an overestimate as cloud temperatures are often lower than ~ 15 - 20°C . At temperatures $> \sim 0^\circ\text{C}$, this overestimate is likely less than an order of magnitude (**Figure 2**); in supercooled cloud droplets ($< 0^\circ\text{C}$), metabolic activity R_{cell} might be some orders of magnitude lower and cell multiplication can be considered negligible.

260 (iii) While some of the studies listed by Burrows et al. (2009b) and in **Table 1** report the concentrations of viable cells, others give the total cell concentrations. In addition, the large discrepancy in reported F_{live} between $< 0.1\%$ up to nearly 100% as discussed e.g., by Lindeman et al. (1982), Lighthart and Shaffer (1994) and Gandolfi et al. (2013), might be also due to differences in the measurement techniques. Consistent experimental methodologies are needed to give comprehensive data on F_{live} and the survival rates of bacteria in aerosol particles (Otero Fernandez et al., 2019).

265 (iv) The average cell mass depends on bacteria type and their growth stage. Sattler (2001) estimated carbon mass of bacteria cells in cloud water as 17 fg carbon cell⁻¹ in agreement with values for marine and freshwater ecosystems. Approximating the total mass by doubling the carbon mass, results in 34 fg cell⁻¹, i.e. equivalent to spherical particles

with diameters of $\sim 0.4 \mu\text{m}$. Carbon mass and the carbon-to-total mass ratio can greatly differ from these values; for example, total masses of prokaryotic cell of 200 fg cell^{-1} in soil have been reported (Whitman et al., 1998). In their global study, Burrows et al. (2009b) assumed a mass of 520 fg cell^{-1} ($1 \mu\text{m}$ particle).

Table 3: Parameters used in the estimate of SBA mass formation and their possible minimum and maximum values based on literature data

Parameter	Value in Eq (4) and Eq (5)	Range		Comment
		Minimum	Maximum	
$C_{\text{cell}} [\text{m}^{-3}]$	10^4	100	10^9	$C_{\text{cell,min}}$: above desert during low RH and high radiation (Lighthart and Shaffer, 1994); $\sim 10^6$: in a highly polluted area (Xi'an, China) (Xie et al., 2018); $C_{\text{cell,max}}$ was measured above a wastewater storage lagoon (Paez-Rubio et al., 2005)
$R_{\text{cell}} [\text{h}^{-1}]$	0.3	0	3	$R_{\text{cell}} \sim 0$ might occur under stressful conditions when cells become dormant (e.g. low temperature, little water...); $R_{\text{cell}} \sim 3$ for <i>E. coli</i> under optimal conditions (37°C , appropriate culture medium).
F_{live}	1	0.0001	1	0.0001 – 0.2 based on global microbial diversity (Gandolfi et al., 2013); 0.22 above crop fields (Lindemann et al., 1982), 0.81 above desert (Lighthart and Shaffer, 1994). Some studies report concentrations of viable cells; in this case reported concentrations imply $F_{\text{live}} = 1$
$m_{\text{cell}} [\text{fg cell}^{-1}]$	52	34	520	$m_{\text{cell,min}}$ corresponds to cells in clouds assuming that cells are composed of 50% carbon (Sattler et al., 2001); masses of other prokaryotic cells might be $\leq 200 \text{ fg cell}^{-1}$ (Whitman et al., 1998); $m_{\text{cell,max}}$ corresponding to a spherical cell of a diameter $1 \mu\text{m}$ and density of 1 g cm^{-3} (Burrows et al., 2009b).
F_{cloud}	0.15	> 0	1	The average global value might be higher than 0.15 if bacterial processes also occur outside of clouds. On small scales or above individual ecosystems, the value can be smaller than the average value, depending on cloud variability.

(v) While several studies have shown that liquid water is necessary for efficient microbial activity, it is not clear yet whether bacteria maintain activity in wet aerosols. Klein et al. (2016) found indications that bacteria metabolic activity exists in aerosols but no quantitative data is reported yet. Bacteria become dormant (Kaprelyants and Kell, 1993) or have reduced viability at relative humidities of 86 - 97% (Haddrell and Thomas, 2017). In soil samples, it has been shown that cycles of drying and rewetting might enhance microbial activity compared to constantly moist samples (Meisner et al., 2017; Xiang et al., 2008); thus, it may be speculated that such effects also occur in rapidly changing

humidity conditions in atmospheric deliquesced aerosols. Under those conditions, the time fraction of microbial activity would exceed F_{cloud} . On locally smaller scales, cloud processing time might be shorter than the average F_{cloud} (Table 2) depending on cloud variability.

285 SBA mass formation calculated by Eq (4) and Eq (5) depends linearly on all parameters discussed in (i) to (v). Thus, the uncertainty of the predicted formation rates can be simply estimated by the ranges given in **Table 3**. However, in the atmosphere, all parameters might continuously change over time and thus might affect SBA mass to different extents.

3.1.3 Comparison of SBA formation to other aerosol sources

290 An estimate of aerosol emissions from the biosphere suggested a source strength of primary biological particles of 1000 Tg yr⁻¹ (Jaenicke, 2005). However, in this study, PBA was defined to include all cellular material, proteins, and their fragments. A global model study predicted total PBA emissions (bacteria, fungal spores and pollen) of 123 Tg yr⁻¹ of which bacteria comprised 0.79 Tg yr⁻¹ (Myriokefalitakis et al., 2017). This number is similar to the range of 0.4 – 1.8 Tg bacteria yr⁻¹ as given by Burrows et al (2009b). Bacteria usually comprise only a small mass fraction of total
295 PBA; a major fraction is composed of fungal spores. Thus, their emissions are generally estimated to be larger in mass than those of bacteria. For example, their global emissions (25 Tg yr⁻¹) was suggested to contribute 23% to total primary organic aerosol (Heald and Spracklen, 2009). An estimate of fungal spore emissions based on tracer compounds resulted in predicted 50 Tg yr⁻¹ (Elbert et al., 2007)). None of these estimates include microbial activity as a source of biological mass. Our predicted SBA source of 3.7 Tg year⁻¹ is similar to the model predictions for primary bacteria
300 emissions. The estimates of primary bacteria emissions are performed such that observed cell concentrations are matched by the models without considering another source of cells in the atmosphere. Our SBA estimates might be equally biased as they are based on the same ambient cell concentrations as reported by Burrows et al. (2009a, b) and Myriokefalitakis et al. (2017). The absolute values and the ratio of primary to secondary bacteria mass need to be evaluated by more complex model studies as our simple framework can provide.

305 Total organic aerosol is composed of mostly secondary mass. Best estimates based on observational and model studies of the net production rate of secondary organic aerosol (SOA) mass are on the order of 136 – 280 Tg year⁻¹ (Hodzic et al., 2016). These amounts are similar to the predicted global sulfate production of 117 Tg year⁻¹ (39 Tg S year⁻¹) (Chin et al., 2000). Thus, SBA production can be estimated to be on the order of ~1% of the secondary aerosol sources. The net aerosol mass formation due to SBA production might be even smaller if bacteria metabolize substrates that are
310 already in the particle phase. In this case, biotransformation processes lead to the conversion of non-biological into biological aerosol mass. The unique properties of biological aerosol material have been extensively discussed in the context of heterogeneous ice nucleation where it has been shown that even small amounts of biological material could have significant effects on clouds and precipitation (Möhler et al., 2008; Morris et al., 2004; Santl-Temkiv et al., 2015). Given the low ambient concentrations of ice nucleating particles and their high sensitivity to the ice/liquid partitioning
315 in mixed-phase clouds (e.g., Ervens et al., 2011), a small change in biological mass possibly translates into significant changes in the evolution of cold clouds.

3.2 Consumption of organic carbon in clouds

3.2.1 Calculation of microbial and chemical WSOC loss rates

Bacteria can be metabolically active in the aqueous phase of clouds (Delort et al., 2010) and on the surface or bulk phase of particles (Klein et al., 2016; Estillore et al., 2016). Such processes lead to a decrease of water-soluble organic carbon (WSOC) mass within cloud droplets as bacteria convert substrates into CO₂ (**Figure 1**). Also processes of biological mass production from CO₂ exist (autotrophy) and include photosynthesis (photoautotrophs). However, despite the fact that photosynthetic microorganisms were reported in the atmosphere (Tesson and Santl-Temkiv, 2018) there is neither a clear indication yet of photosynthetic activity in clouds, nor of other modes of autotrophy.

The split between carbon uptake for biological mass production and mineralization is quantified by the bacterial growth efficiency BGE (Eq (2)). Studies have shown that generally metabolic processes produce mostly CO₂ under atmospheric conditions and only a small fraction of the carbon is mineralized into organic products (< 1 – 10% of the total C utilized). We introduce here the factor F_{CO_2} as a measure of the loss of organics due to bacterial processes:

$$F_{CO_2} = 1 - BGE \quad (6)$$

Using F_{CO_2} , we can write the carbon balance as



The loss rate of carbon can be, thus, calculated as

$$R_{WSOC,Bact} \left[\frac{g}{L(aq) s} \right] = - \frac{d(WSOC)}{dt} = \frac{dCO_2}{dt} = \quad (7)$$

$$-F_{CO_2} k_{Bact} \left[\frac{L(aq)}{cell \cdot s} \right] \cdot C_{Cell,aq} \left[\frac{cell}{L(aq)} \right] \cdot F_C \cdot C_{WSOC} \left[\frac{gC}{L(aq)} \right] \cdot F_{cloud}$$

whereas the cell concentration in cloud water can be replaced by


$$C_{Cell,aq} \left[\frac{cell}{L(aq)} \right] = C_{cell,g} \left[\frac{cell}{m^3(g)} \right] / LWC \left[\frac{m^3(g)}{L(aq)} \right] \quad (8)$$

with $C_{Cell,aq}$ and C_{WSOC} being the concentrations of bacteria and water-soluble organic carbon in cloud water, respectively, and $C_{cell,g}$ ambient cell concentrations in the gas phase (e.g. **Table 1**). These concentrations are on average $\sim 10^7$ cells L⁻¹ (Vaithilingom et al., 2013) and 0.1 mM (Herckes et al., 2013) in cloud water whereas both values might differ over a few orders of magnitude locally and temporally (**Section 3.2.2**). Usually experimental loss rates of organics by bacteria in real and artificial cloud water are reported to be on the order of $\leq 10^{-17}$ mol cell⁻¹ s⁻¹ for organic substrates (e.g., formic, acetic, and succinic acids) (Vaithilingom et al., 2010, 2011). The cell activity is dependent on the bacteria type and the availability of the organic substrate. Thus, strictly, such rates [mol cell⁻¹ s⁻¹] are only valid for

345 the substrate-to-cell ratio as applied in the experiments. In order to account for the ratio as encountered in cloud water, we use here k_{Bact} [$L \text{ cell}^{-1} \text{ s}^{-1}$] (i.e. measured rate divided by the concentration of organic substrate in the experiments) that is applicable to the full range of conditions where the cells exhibit a similar microbial activity. Resulting rate constants for formic, acetic and succinate acids are on the order of $k_{Bact} \sim 10^{-13} L \text{ cell}^{-1} \text{ s}^{-1}$. Equation 7 includes an additional factor F_C that accounts for the microbial selectivity towards only some organics by each bacteria type (e.g. 350 Šantl-Temkiv et al., 2013; Bianco et al., 2019). For example, it has been shown in a single study that, upon laboratory incubation of cloud water, oxalic acid is not affected by cloud borne microorganisms, formate is only consumed after a lag time of several hours, which is much longer than the lifetime of a cloud droplet, and, on the other hand, compounds such as acetate or succinate are readily biodegraded (Vařtilingom et al., 2011). Since these compounds comprise major constituents of WSOC in cloud water, it seems reasonable to introduce a factor $F_C < 1$. Given the complexity of the organic matter in the atmosphere, the numerous organic molecules existing in cloud water and their variable susceptibility to biodegradation, F_C is hard to specify with precision. Bianco et al. (2019a) observed experimentally by FT-ICR-MS during laboratory incubation of cloud water that out of the 2178 compounds detected, 1094 were actually utilized by bacteria (~50%). Assuming that all these compounds were equally abundant, one could conclude that 50% of all cloud water organics were prone to be microbiologically consumed (i.e. $F_C = 0.5$). More quantitative support of 360 this assumption could be given based on the fact that preferably small oxygenated organics are taken up by bacteria. Compilations of speciated cloud water organics have shown that small acids, such as formic and acetic acid, comprise a large fraction (at least ~30%) of the characterized fraction of cloud water organics (e.g. Figure 6 in Herckes et al., (2013). Lab experiments have shown that these acids are degraded by bacteria (Amato et al., 2007a; Vařtilingom et al., 2013). While $F_C = 0.5$ seems a reasonable compromise, this factor is highly uncertain and strongly depends on the microbial and chemical composition of cloud water. The calculated rates $R_{Bact,WSOC}$ are summarized in **Figure 4** for $0.8 \leq F_{CO_2} \leq 0.99$ and $F_C = 0.5$.

Several studies have discussed the competition of microbial and chemical processes in cloud water as a sink of specific organic compounds (Ariya et al., 2002; Husárová et al., 2011; Vařtilingom et al., 2010, 2013). The most efficient loss reactions for organics in cloud water are initiated by OH radicals. The general rate constant of the OH radical with water-soluble organic carbon is $k_{OH} = 3.8 \cdot 10^8 M^{-1} s^{-1}$ (Arakaki et al., 2013). The reactions of WSOC with OH lead to volatile and non-volatile oxidation products. Radicals are much less selective towards organics than bacteria are; thus, the assumption of a factor equivalent to F_C in Eq (7) is not necessary as all water-soluble organics react with OH whereas the chemical reactivity mostly depends on the structure of the organic compound. The yield of volatile products (Y_{volC}) includes CO_2 but also formaldehyde and other volatile compounds that do not remain in the particle phase after cloud evaporation and thus do not contribute to the aerosol loading. We assume $0.2 \leq Y_{volC} \leq 0.5$, but in 375 general, Y_{volC} depends on the WSOC composition, with higher values for more aged organics that are more readily oxidized to volatile products.

Equivalent to reaction (R 1), we express the carbon loss by the OH radical in clouds as



380 With $WSOC_{aer}$ the WSOC fraction that remains in the aerosol phase after drop evaporation. We calculate the loss rate

accordingly:

$$R_{OH,WSOC} = -\frac{d(WSOC)_{OH}}{dt} \left[\frac{g}{L(aq)s} \right]$$

$$= k_{OH} [L mol^{-1} s^{-1}] [OH]_{aq} [mol L^{-1}] \cdot Y_{volC} C_{WSOC} \left[\frac{gC}{L} \right] \cdot LWC \left[\frac{g}{m^3} \right] \cdot F_{cloud} \quad (9)$$

OH concentrations in cloud water are in the range of $10^{-16} M < [OH]_{aq} < 10^{-14} M$ (Arakaki et al., 2013; Bianco et al., 2015; Ervens et al., 2014) and an average cloud liquid water content (LWC) of $0.15 g m^{-3}$ is assumed. The results in **Figure 4** suggest that the microbial rates may be smaller than the chemical ones under most conditions. Overall, the values shown in Figure 4 only differ by a factor of ~ 2.5 which might imply that there are conditions under which chemical and biological processes in the aqueous phase compete. This trend is in agreement with several previous studies that focused on the comparison of microbial versus chemical processes as sinks for specific organic substrates (Amato et al., 2007a; Vařtilingom et al., 2010). These loss fluxes are relatively large as compared to the predicted SBA formation (**Figure 3**).




Figure 4: Predicted loss of WSOC by bacterial utilization and by chemical (OH) processing in cloud water for different assumption on F_{CO_2} and Y_{volC} .

In a previous study, it was estimated that microbial processes in clouds lead to a total carbon loss of $\sim 10 - 50 Tg yr^{-1}$ and to a production of $\sim 100 Tg year^{-1} CO_2$ with the assumptions of complete respiration ($F_{CO_2} = 1$), microbial non-selectivity towards WSOC ($F_C = 1$) and applying the same loss rates as observed in lab experiments without correcting for differences in the ratio of bacteria cell to WSOC concentrations (Vařtilingom et al., 2013). Thus, this former estimate can be considered an upper limit whereas the one in the current study ($\sim 30 Tg yr^{-1}$) is more conservative, suggesting that the respiration of bacteria is a negligible CO_2 source as compared to the sum of anthropogenic sources ($\sim 50,000 Tg CO_2 year^{-1}$ (IPCC, 2014)).

The WSOC loss rates calculated here are also much smaller than those estimated for the loss of total SOA due to heterogeneous reactions and photolysis on particle surfaces ($\leq 50.3 Tg year^{-1}$) and by wet deposition ($\leq \sim 50 Tg year^{-1}$, (Hodzic et al., 2016)). For the parameters chosen in our estimate, they are also smaller than the predicted production

rates of aqSOA in clouds of 13.1 - 46.8 Tg year⁻¹ (Lin et al., 2014) or 20 – 30 Tg yr⁻¹ (Liu et al., 2012). This estimate represents organic carbon sources and sinks in general. It should be noted that the organics comprising WSOC and aqSOA, respectively, might not be identical. These predictions strongly depend on the representation of clouds, and particularly on the liquid water content (He et al., 2013).

3.2.2 Discussion of uncertainties of microbial and chemical WSOC loss

The calculation of microbial and chemical WSOC loss should be regarded an approximation using a set of parameters that are all associated with considerable uncertainties. Similar to the discussion in *Section 3.1.2*, we compile all parameters and minimum and maximum values based on literature data in *Table 4*:

(i) There are not as many measurements of $C_{Bact,aq}$ as for cell concentrations C_{cell} in cloud-free regions. The assumption that all particles that contain bacteria cells are activated into cloud droplets does likely not lead to a large overestimate. The sizes of bacteria-containing particles usually exceed several hundred nanometers and thus can all be considered CCN. Differences in LWC - as the conversion factor from gas to aqueous phase concentrations - are relatively small, within a factor of 2 – 3, within the categories of common cloud types (Pruppacher and Klett, 2003).

(ii) The activity of microorganisms towards organic substrates is often reported in units of ‘mol(substrate) cell⁻¹ s⁻¹’ which expresses the amount of substrate that is consumed per cell and time. For several compounds (e.g., formate, acetate, succinate) these rates differ by approximately one order of magnitude (Vaitilingom et al., 2011). However, the resulting k_{Bact} values (i.e. rate divided by substrate concentration) are all on the order of $k_{Bact} \sim 10^{-13}$ L cell⁻¹ s⁻¹ which appears to represent an upper limit for the organics that have been investigated for metabolic activity in clouds. A much lower constant was derived from experiments with less oxygenated compounds such as phenol (Lallement, 2017).

While we only consider the direct interaction of bacteria and organics, additional processes might lead to more complex chemical and microbial interactions. For example, siderophores form iron complexes (Passananti et al., 2016) and, thus, suppress Fenton reactions that affect oxidant levels in cloud droplets (e.g., Deguillaume et al., 2004). Such indirect feedbacks of microbial processes on chemical budgets require more comprehensive data sets that are currently not available for models.

(iii) The respiration of bacteria depends on many different factors such as stress due harsh conditions. It can be expected that at higher stress levels (nutritional or thermal), F_{CO_2} increases to supply elevated energy needs (Amato and Christner, 2009; Eiler et al., 2003). Values of BGE as low as < 0.4% ($F_{CO_2} = 0.996$) were observed (Eiler et al, 2003 and references therein), indicating that nearly all the carbon used was mineralized into CO₂.

(iv) The fraction of organic material metabolized by bacteria in clouds (F_C) is likely not unity for a single bacteria type (Bianco et al., 2019; Vaitilingom et al., 2011). Carboxylic acids that are preferentially metabolized by several common bacteria types often comprise a major fraction (~ 20%) of the cloud-water organics that can be speciated on a molecular level (Herckes et al., 2013). This fraction might be regarded a lower limit of F_C since the reactivity of the large fraction of unspciated organics (often ~70%) towards bacteria is not known. However, a recent qualitative study suggested that ~50% of all organics in cloud water are microbially consumed by bacteria (Bianco et al., 2019). Our comparison implies the same spatial accessibility of bacteria and OH, respectively, to WSOC. This might be an oversimplification as bacteria are unevenly distributed among cloud drop populations as statistically only one in ~10,000 droplets may

contain a single bacteria cell. OH can be expected to be present in all cloud droplets as the direct phase transfer from the gas phase represents one of the major OH sources in cloud water.

440

Table 4: Values for parameters in Eq (7) and Eq (9) used in the estimate of WSOC loss by microbial and chemical processes and their minimum and maximum values

Parameter	Value in Eq (7) and Eq (9)	Range		Comment
		Minimum	Maximum	
$C_{\text{Bact, aq}}$ [cells L ⁻¹]	10 ⁷	10 ⁶	10 ⁸	Range of total bacteria concentration observed in cloud water samples collected from a mid-altitude mountain site over several years (31 samples) (Vařtilingom et al., 2012)
LWC [g m ⁻³]	0.15	0.1	1	The minimum and maximum value describe a range for a wide variety of cloud types. The assumption of LWC is not needed if it is assumed that all bacteria-containing particles act as CCN.
k_{Bact} [L cell ⁻¹ s ⁻¹]	10 ⁻¹³	10 ⁻¹⁵	10 ⁻¹³	$k_{\text{Bact, min}}$ was derived for microbial activity towards phenol (Lallement, 2017). $k_{\text{Bact, max}}$ was derived from experiments using cloud water and is valid for the microbial activity of various highly oxygenated compounds.
F_{CO_2}	0.8 – 0.99	0.2	1	Even though BGE ranging from <0.4% to 80% ($0.996 > F_{\text{CO}_2} > 0.2$) were estimated in natural environments (Eiler et al. 2003 and references therein), at low nutrient concentrations, as encountered in clouds, high F_{CO_2} can be expected.
F_C	0.5	0.2	< 1	Herckes et al. (2013) report that ~20% of total organic carbon in clouds is composed of speciated carboxylic acids; Bianco et al. (2019) demonstrate that ~50% of all organics in cloud water are affected by bacteria. $F_C = 1$ seems unlikely due to variation in microbial and chemical cloud water composition.
k_{OH} [L mol ⁻¹ s ⁻¹]	3.8·10 ⁸	10 ⁶	10 ¹⁰	Typically, undissociated acids (low pH) and polyfunctional compounds have k_{OH} at the lower end of this range whereas the upper limit is constrained by diffusion limitation (Herrmann, 2003; Monod and Doussin, 2008)
$[\text{OH}]_{\text{aq}}$ [M]	10 ⁻¹⁵	10 ⁻¹⁷	10 ⁻¹⁴	The suggested range includes concentrations that were inferred for night-time conditions (minimum) to day-time conditions in clean air masses (low OH sinks).
Y_{volC}	0.3 – 0.5	0.2	0.8	This value has not been comprehensively quantified yet; largest values can be likely expected in aged WSOC with high O/C ratios

(v) While the absolute importance of microbial loss depends on the parameters discussed in (i) to (iv), the relative importance compared to chemical processes might be of interest in studies where the fate of individual organics in the

445

cloud droplets or in the atmospheric multiphase system is explored. $[\text{OH}]_{\text{aq}}$ depends mostly on photochemical processes as source processes and on the concentrations of WSOC as the main sinks; it ranges from 10^{-17} M (night-time) to 10^{-14} M (day time, clean air masses) (Arakaki et al., 2013).

450 (vi) Given that formate and acetate comprise major contributors to cloud water organics (Herckes et al., 2013), some fraction of WSOC will be converted into highly volatile products, such as CO_2 and CH_3CHO that will not remain in the particle phase after cloud evaporation. However, Y_{volC} likely does not approach unity since several studies have suggested that radical reactions in cloud water lead to the successive decay of dicarboxylic acids into their next smaller homologue which will remain in the aqueous phase. Within these limits, we conservatively suggest a range of 0.2 to 0.8 for Y_{volC} but point out the need for studies to refine this parameter

455 4. Summary and conclusions

We have estimated the amount of biological mass that is formed in the atmosphere by growth and multiplication of bacteria cells ('secondary biological aerosol', SBA). Data for representative bacteria strains and their generation rates have been compiled for major ecosystems. Using average values for cloudiness above the various ecosystems, we estimate that 3.7 Tg year^{-1} SBA mass is formed globally which is comparable to current estimates of direct bacteria emissions ($0.4 - 0.7 \text{ Tg yr}^{-1}$ (Burrows et al., 2009b; Myriokefalitakis et al., 2017)) which comprise a small fraction of total biological aerosol mass. While these production rates make up $\sim 1\%$ of other major secondary aerosol formation rates (secondary organics or sulfate), their importance might differ on spatial or temporal scales. In addition, SBA production leads to an increase in biological aerosol mass which might sensitively affect physicochemical particle properties (e.g. ice nucleation ability). SBA formation linearly depends on several parameters, such as the number concentration of metabolically active bacteria cells, their generation rates and the time scales during which they are assumed to grow or multiply – all of which are associated with considerable uncertainties. The ecosystem categories in Table 2 represent fairly large regions. It might be expected that SBA formation rates are different on smaller spatial and/or temporal scales. For example, it has been shown that human activities in cities lead to high bacteria concentrations; also forests have been identified as significant sources of biogenic aerosol. However, detailed data on bacteria are sparse in such regions. While several recent studies have characterized the diversity of microorganisms in forested regions (rainforest, tropics) (Gusareva et al., 2019; Souza et al., 2019), these studies did not report cell concentrations which highlights the urgent need of additional measurements.

The detailed discussion of the parameters and their uncertainties in our simplified approach highlights the likely variability of SBA formation on smaller scales and the need of future studies to refine these parameters. Similar approaches as ours may be applied to yeast growth. Yeast cells are generally larger ($\sim 2 - 10 \mu\text{m}$) than bacteria cells (Fröhlich-Nowoisky et al., 2009) and, thus, their residence time in the atmosphere is likely shorter. Detailed data on their activity in clouds are not available which currently prevents the assessment of their potential contribution to SBA. We also quantify the role of clouds as sinks of total WSOC by microbial and chemical processes, unlike previous studies that focused on microbial activity towards individual organic compounds. It is estimated that microbial processes lead to an organic mass loss of $8 - 11 \text{ Tg yr}^{-1}$ whereas chemical processes by the OH radical in clouds lead to a loss of $8 - 20 \text{ Tg yr}^{-1}$. These numbers are small compared to other sinks such as aerosol removal by deposition.

Not all of the WSOC mass even contributes to organic aerosol loading as water-soluble, volatile organics are dissolved in cloud water but evaporate during drop evaporation. Thus, the loss of organic aerosol mass due to direct microbial activity in clouds might be smaller than the predicted loss of WSOC. Large uncertainties in these estimates represent the assumptions on the fraction of carbon that is converted into volatile products. For bacteria, this fraction is quantified by the bacteria growth efficiency that depends on numerous factors, such as bacteria type, substrate availability and physical conditions in the condensed phase.

In current atmospheric models, when considered, bacteria cells are inert, i.e. they neither change their mass or number concentrations during their residence time in the atmosphere nor do they interact with other aerosol constituents. The approach presented in our study provides a first simplified estimate of SBA formation and WSOC loss due to bacteria that could be easily adapted in models. Given the current great activities in the field of atmospheric bioaerosols, it can be expected that the discussed parameters in the estimates can and should be refined in the future in order to quantify the role of bacterial processes as source of biological mass and net source or sink of organic aerosol in the atmosphere.

Data availability. All data are available upon request from the authors.

Author contributions. BE and PA planned and carried out the study and wrote the manuscript together.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgement. This work has been funded by the French National Research Agency (ANR) in the framework of the ‘Investment for the Future’ program, ANR-17-MPGA-0013.

References

- Amato, P. and Christner, B. C.: Energy metabolism response to low-temperature and frozen conditions in *Psychrobacter cryohalolentis*, *Appl. Environ. Microbiol.*, 75(3), 711–718, doi:10.1128/AEM.02193-08, 2009.
- Amato, P., Demeer, F., Melaouhi, A., Fontanella, S., Martin-Biesse, A. S., Sancelme, M., Laj, P. and Delort, A. M.: A fate for organic acids, formaldehyde and methanol in cloud water: their biotransformation by micro-organisms, *Atmos Chem Phys*, 7(15), 4159–4169, doi:10.5194/acp-7-4159-2007, 2007a.
- Amato, P., Parazols, M., Sancelme, M., Laj, P., Mailhot, G. and Delort, A.-M.: Microorganisms isolated from the water phase of tropospheric clouds at the Puy de Dôme: major groups and growth abilities at low temperatures, *FEMS Microbiol. Ecol.*, 59(2), 242–254, doi:10.1111/j.1574-6941.2006.00199.x, 2007b.
- Amato, P., Doyle, S. and Christner, B. C.: Macromolecular synthesis by yeasts under frozen conditions, *Environ. Microbiol.*, 11(3), 589–596, doi:10.1111/j.1462-2920.2008.01829.x, 2009.
- Amato, P., Doyle, S. M., Battista, J. R. and Christner, B. C.: Implications of Subzero Metabolic Activity on Long-Term Microbial Survival in Terrestrial and Extraterrestrial Permafrost,

- Astrobiology, 10(8), 789–798, doi:10.1089/ast.2010.0477, 2010.
- Amato, P., Joly, M., Schaupp, C., Attard, E., Möhler, O., Morris, C. E., Brunet, Y. and Delort, A.-M.: Survival and ice nucleation activity of bacteria as aerosols in a cloud simulation chamber, *Atmos. Chem. Phys.*, 15(11), 6455–6465, doi:10.5194/acp-15-6455-2015, 2015.
- 520 Amato, P., Joly, M., Besaury, L., Oudart, A., Taib, N., Moné, A. I., Deguillaume, L., Delort, A.-M. and Debroyas, D.: Active microorganisms thrive among extremely diverse communities in cloud water, *PLOS ONE*, 12(8), e0182869, doi:10.1371/journal.pone.0182869, 2017.
- Amato, P., Besaury, L., Joly, M., Penaud, B., Deguillaume, L. and Delort, A.-M.: Metatranscriptomic exploration of microbial functioning in clouds, *Sci. Rep.*, 9(1), 4383, 525 doi:10.1038/s41598-019-41032-4, 2019.
- Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S., Kimura, T., Tshako, A. and Miyagi, Y.: A general scavenging rate constant for reaction of hydroxyl radical with organic carbon in atmospheric waters, *Env. Sci Technol*, 47(15), 8196–8203, doi:10.1021/es401927b, 2013.
- 530 Ariya, P. A., Nepotchatykh, O., Ignatova, O. and Amyot, M.: Microbiological degradation of atmospheric organic compounds, *Geophys. Res. Lett.*, 29(22), doi: 10.1029/2002GL015637, 2002.
- Ariya, P. A., Sun, J., Eltouny, N. A., Hudson, E. D., Hayes, C. T. and Kos, G.: Physical and chemical characterization of bioaerosols - Implications for nucleation processes, *Int. Rev. Phys. Chem.*, 28(1), 1–32, doi:10.1080/01442350802597438, 2009.
- 535 Artaxo, P., Maenhaut, W., Storms, H. and Van Grieken, R.: Aerosol characteristics and sources for the Amazon Basin during the wet season, *J. Geophys. Res. Atmospheres*, 95(D10), 16971–16985, doi:10.1029/JD095iD10p16971, 1990.
- Bakermans, C., Tsapin, A. I., Souza-Egipsy, V., Gilichinsky, D. A. and Neilson, K. H.: Reproduction and metabolism at -10°C of bacteria isolated from Siberian permafrost, *Environ. Microbiol.*, 5(4), 321–326, doi:10.1046/j.1462-2920.2003.00419.x, 2003.
- 540 Bauer, H., Giebl, H., Hitzemberger, R., Kasper-Giebl, A., Reischl, G., Zibuschka, F. and Puxbaum, H.: Airborne bacteria as cloud condensation nuclei, *J Geophys Res*, 108(D21), 4658, doi:10.1029/2003jd003545, 2003.
- Bianco, A., Passananti, M., Perroux, H., Vuyard, G., Mouchel-Vallon, C., Chaumerliac, N., 545 Mailhot, G., Deguillaume, L. and Brigante, M.: A better understanding of hydroxyl radical photochemical sources in cloud waters collected at the puy de Dôme station – experimental versus modelled formation rates, *Atmos Chem Phys*, 15(16), 9191–9202, doi:10.5194/acp-15-9191-2015, 2015.
- 550 Bianco, A., Deguillaume, L., Chaumerliac, N., Vaïtilingom, M., Wang, M., Delort, A.-M. and Bridoux, M. C.: Effect of endogenous microbiota on the molecular composition of cloud water: a study by Fourier-transform ion cyclotron resonance mass spectrometry (FT-ICR MS), *Sci. Rep.*, 9(1), 7663, doi:10.1038/s41598-019-44149-8, 2019.

- 555 Bovallius, A., Bucht, B., Roffey, R. and Anäs, P.: Three-year investigation of the natural airborne bacterial flora at four localities in Sweden., *Appl. Environ. Microbiol.*, 35(5), 847, doi:0099-2240/78/0035-0847, 1978.
- Bowers, R. M., McCubbin, I. B., Hallar, A. G. and Fierer, N.: Seasonal variability in airborne bacterial communities at a high-elevation site, *Atmos. Environ.*, 50, 41–49, doi:10.1016/j.atmosenv.2012.01.005, 2012.
- 560 Burrows, S. M., Elbert, W., Lawrence, M. G. and Pöschl, U.: Bacteria in the global atmosphere, Part 1: Review and synthesis of literature data for different ecosystems, *Atmos Chem Phys*, 9(23), 9263–9280, doi:10.5194/acp-9-9263-2009, 2009a.
- Burrows, S. M., Butler, T., Jöckel, P., Tost, H., Kerkweg, A., Pöschl, U. and Lawrence, M. G.: Bacteria in the global atmosphere, Part 2: Modeling of emissions and transport between different ecosystems, *Atmos Chem Phys*, 9(23), 9281–9297, doi:10.5194/acp-9-9281-2009, 2009b.
- 565 Carotenuto, F., Georgiadis, T., Gioli, B., Leyronas, C., Morris, C. E., Nardino, M., Wohlfahrt, G. and Miglietta, F.: Measurements and modeling of surface–atmosphere exchange of microorganisms in Mediterranean grassland, *Atmos Chem Phys*, 17(24), 14919–14936, doi:10.5194/acp-17-14919-2017, 2017.
- 570 Chin, M., Rood, R. B., Lin, S.-J., Müller, J.-F. and Thompson, A. M.: Atmospheric sulfur cycle simulated in the global model GOCART: Model description and global properties, *J Geophys Res - Atmos*, 105(D20), 24671–24687, doi:10.1029/2000jd900384, 2000.
- Copeland, J. K., Yuan, L., Layeghifard, M., Wang, P. W. and Guttman, D. S.: Seasonal Community Succession of the Phyllosphere Microbiome, *Mol. Plant. Microbe Interact.*, 28(3), 274–285, doi:10.1094/MPMI-10-14-0331-FI, 2015.
- 575 Davey, K. R.: A predictive model for combined temperature and water activity on microbial growth during the growth phase, *J. Appl. Bacteriol.*, 67(5), 483–488, doi:10.1111/j.1365-2672.1989.tb02519.x, 1989.
- 580 Deguillaume, L., Leriche, M., Monod, A. and Chaumerliac, N.: The role of transition metal ions on HO_x radicals in clouds: a numerical evaluation of its impact on multiphase chemistry, *Atmos Chem Phys*, 4, 95–110, 2004.
- 585 DeLeon-Rodriguez, N., Latham, T. L., Rodriguez-R, L. M., Barazesh, J. M., Anderson, B. E., Beyersdorf, A. J., Ziemba, L. D., Bergin, M., Nenes, A. and Konstantinidis, K. T.: Microbiome of the upper troposphere: Species composition and prevalence, effects of tropical storms, and atmospheric implications, *Proc. Natl. Acad. Sci.*, 110(7), 2575, doi:10.1073/pnas.1212089110, 2013.
- Delort, A.-M., Vaïtilingom, M., Amato, P., Sancelme, M., Parazols, M., Mailhot, G., Laj, P. and Deguillaume, L.: A short overview of the microbial population in clouds: Potential roles in atmospheric chemistry and nucleation processes, *Atmos Res*, 98(2–4), 249–260, 2010.
- Després, V. R., Huffman, J. A., Burrows, S. M., Hoose, C., Safatov, A. S., Buryak, G., Fröhlich-

- 590 Nowoisky, J., Elbert, W., Andreae, M. O., Pöschl, U. and Jaenicke, R.: Primary biological aerosol particles in the atmosphere: a review, *Tellus B*, 64, 15598, doi:10.3402/tellusb.v64i0.15598, 2012.
- Dimmick, R. L., Wolochow, H. and Chatigny, M. A.: Evidence that bacteria can form new cells in airborne particles, *Appl. Environ. Microbiol.*, 37(5), 924–927, 1979.
- 595 Eiler, A., Langenheder, S., Bertilsson, S. and Tranvik, L. J.: Heterotrophic bacterial growth efficiency and community structure at different natural organic carbon concentrations, *Appl. Environ. Microbiol.*, 69(7), 3701–3709, doi:10.1128/AEM.69.7.3701-3709.2003, 2003.
- Elbert, W., Taylor, P. E., Andreae, M. O. and Pöschl, U.: Contribution of fungi to primary biogenic aerosols in the atmosphere: wet and dry discharged spores, carbohydrates, and inorganic ions, *Atmos Chem Phys*, 7(17), 4569–4588, doi:10.5194/acp-7-4569-2007, 2007.
- 600 Engelberg-Kulka, H., Amitai, S., Kolodkin-Gal, I. and Hazan, R.: Bacterial Programmed Cell Death and Multicellular Behavior in Bacteria, *PLOS Genet.*, 2(10), e135, doi:10.1371/journal.pgen.0020135, 2006.
- 605 Ervens, B., Feingold, G., Sulia, K. and Harrington, J.: The impact of microphysical parameters, ice nucleation mode, and habit growth on the ice/liquid partitioning in mixed-phase Arctic clouds, *J Geophys Res*, 116(D17), D17205, doi:10.1029/2011jd015729, 2011.
- Ervens, B., Sorooshian, A., Lim, Y. B. and Turpin, B. J.: Key parameters controlling OH-initiated formation of secondary organic aerosol in the aqueous phase (aqSOA), *J Geophys Res - Atmos*, 119(7), 3997–4016, doi:10.1002/2013JD021021, 2014.
- 610 Estillore, A. D., Trueblood, J. V. and Grassian, V. H.: Atmospheric chemistry of bioaerosols: heterogeneous and multiphase reactions with atmospheric oxidants and other trace gases, *Chem. Sci.*, 7(11), 6604–6616, doi:10.1039/c6sc02353c, 2016.
- Fahlgren, C., Hagström, Å., Nilsson, D. and Zweifel, U. L.: Annual Variations in the Diversity, Viability, and Origin of Airborne Bacteria, *Appl. Environ. Microbiol.*, 76(9), 3015–3025, doi:10.1128/AEM.02092-09, 2010.
- 615 Feingold, G., McComiskey, A., Rosenfeld, D. and Sorooshian, A.: On the relationship between cloud contact time and precipitation susceptibility to aerosol, *J. Geophys. Res. Atmospheres*, 118(18), 10,544–10,554, doi:10.1002/jgrd.50819, 2013.
- 620 Fröhlich-Nowoisky, J., Pickersgill, D. A., Després, V. R. and Pöschl, U.: High diversity of fungi in air particulate matter, *Proc. Natl. Acad. Sci.*, 106(31), 12814, doi:10.1073/pnas.0811003106, 2009.
- Fuchs, B. M., Zubkov, M. V., Sahn, K., Burkill, P. H. and Amann, R.: Changes in community composition during dilution cultures of marine bacterioplankton as assessed by flow cytometric and molecular biological techniques, *Environ. Microbiol.*, 2(2), 191–201, doi:10.1046/j.1462-2920.2000.00092.x, 2000.
- 625 Fuzzi, S., Mandrioli, P. and Perfetto, A.: Fog droplets—an atmospheric source of secondary

- biological aerosol particles, *Atmos. Environ.*, 31(2), 287–290, doi:10.1016/1352-2310(96)00160-4, 1997.
- Gandolfi, I., Bertolini, V., Ambrosini, R., Bestetti, G. and Franzetti, A.: Unravelling the bacterial diversity in the atmosphere, *Appl. Microbiol. Biotechnol.*, 97(11), 4727–4736, doi:10.1007/s00253-013-4901-2, 2013.
- 630 Gusareva, E. S., Acerbi, E., Lau, K. J. X., Luhung, I., Premkrishnan, B. N. V., Kolundžija, S., Purbojati, R. W., Wong, A., Houghton, J. N. I., Miller, D., Gaultier, N. E., Heinle, C. E., Clare, M. E., Vettath, V. K., Kee, C., Lim, S. B. Y., Chénard, C., Phung, W. J., Kushwaha, K. K., Nee, A. P., Putra, A., Panicker, D., Yanqing, K., Hwee, Y. Z., Lohar, S. R., Kuwata, M., Kim, H. L., Yang, L., Uchida, A., Drautz-Moses, D. I., Junqueira, A. C. M. and Schuster, S. C.: Microbial communities in the tropical air ecosystem follow a precise diel cycle, *Proc. Natl. Acad. Sci.*, 201908493, doi:10.1073/pnas.1908493116, 2019.
- Haddrell, A. E. and Thomas, R. J.: *Aerobiology: Experimental Considerations, Observations, and Future Tools*, *Appl. Environ. Microbiol.*, 83(17), e00809-17, doi:10.1128/AEM.00809-17, 2017.
- 640 He, C., Liu, J., Carlton, A. G., Fan, S., Horowitz, L. W., Levy II, H. and Tao, S.: Evaluation of factors controlling global secondary organic aerosol production from cloud processes, *Atmos Chem Phys*, 13(4), 1913–1926, doi:10.5194/acp-13-1913-2013, 2013.
- Heald, C. L. and Spracklen, D. V.: Atmospheric budget of primary biological aerosol particles from fungal spores, *Geophys. Res. Lett.*, 36(9), doi:10.1029/2009GL037493, 2009.
- 645 Helin, A., Sietiö, O.-M., Heinonsalo, J., Bäck, J., Riekkola, M.-L. and Parshintsev, J.: Characterization of free amino acids, bacteria and fungi in size-segregated atmospheric aerosols in boreal forest: seasonal patterns, abundances and size distributions, *Atmos Chem Phys*, 17(21), 13089–13101, doi:10.5194/acp-17-13089-2017, 2017.
- 650 Herckes, P., Valsaraj, K. T. and Collett Jr, J. L.: A review of observations of organic matter in fogs and clouds: Origin, processing and fate, *Atmos Res*, 132–133, 434–449, doi:org/10.1016/j.atmosres.2013.06.005, 2013.
- Herrmann, H.: Kinetics of aqueous phase reactions relevant for atmospheric chemistry, *Chem. Rev.*, 103(12), 4691–4716, 2003.
- 655 Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C. D., Jimenez, J. L., Madronich, S. and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos Chem Phys*, 16(12), 7917–7941, doi:10.5194/acp-16-7917-2016, 2016.
- 660 Hu, W., Murata, K., Toyonaga, S. and Zhang, D.: Bacterial abundance and viability in rainwater associated with cyclones, stationary fronts and typhoons in southwestern Japan, *Atmos. Environ.*, 167, 104–115, doi:10.1016/j.atmosenv.2017.08.013, 2017.
- Huffman, J. A., Prenni, A. J., DeMott, P. J., Pöhlker, C., Mason, R. H., Robinson, N. H., Fröhlich-Nowoisky, J., Tobo, Y., Després, V. R., Garcia, E., Gochis, D. J., Harris, E., Müller-Germann, I.,

665 Ruzene, C., Schmer, B., Sinha, B., Day, D. A., Andreae, M. O., Jimenez, J. L., Gallagher, M., Kreidenweis, S. M., Bertram, A. K. and Pöschl, U.: High concentrations of biological aerosol particles and ice nuclei during and after rain, *Atmos Chem Phys*, 13(13), 6151–6164, doi:10.5194/acp-13-6151-2013, 2013.

670 Husárová, S., Vařtilingom, M., Deguillaume, L., Traikia, M., Vinatier, V., Sancelme, M., Amato, P., Matulová, M. and Delort, A.-M.: Biotransformation of methanol and formaldehyde by bacteria isolated from clouds. Comparison with radical chemistry, *Atmos. Environ.*, 45(33), 6093–6102, doi:org/10.1016/j.atmosenv.2011.06.035, 2011.

675 IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2014.

Jaenicke, R.: Abundance of Cellular Material and Proteins in the Atmosphere, *Science*, 308(5718), 73, doi:10.1126/science.1106335, 2005.

680 Joly, M., Amato, P., Sancelme, M., Vinatier, V., Abrantes, M., Deguillaume, L. and Delort, A.-M.: Survival of microbial isolates from clouds toward simulated atmospheric stress factors, *Atmos. Environ.*, 117, 92–98, doi:10.1016/j.atmosenv.2015.07.009, 2015.

Kaprelyants, A. S. and Kell, D. B.: Dormancy in Stationary-Phase Cultures of *Micrococcus luteus*: Flow Cytometric Analysis of Starvation and Resuscitation, *Appl. Environ. Microbiol.*, 59(10), 3187, doi:0099-2240/93/103187-10\$02.00/0, 1993.

685 King, S. Platnick, W. P. Menzel, S. A. Ackerman and P. A. Hubanks: Spatial and Temporal Distribution of Clouds Observed by MODIS Onboard the Terra and Aqua Satellites, *IEEE Trans. Geosci. Remote Sens.*, 51(7), 3826–3852, doi:10.1109/TGRS.2012.2227333, 2013.

690 Klein, A. M., Bohannan, B. J. M., Jaffe, D. A., Levin, D. A. and Green, J. L.: Molecular Evidence for Metabolically Active Bacteria in the Atmosphere, *Front. Microbiol.*, 7, 772, doi:10.3389/fmicb.2016.00772, 2016.

Lallement, A.: Impact des processus photochimiques et biologiques sur la composition chimique du nuage, PhD Thesis, Université Clermont Auvergne, Clermont-Ferrand (France)., 2017.

695 Lallement, A., Besaury, L., Tixier, E., Sancelme, M., Amato, P., Vinatier, V., Canet, I., Polyakova, O. V., Artaev, V. B., Lebedev, A. T., Deguillaume, L., Mailhot, G. and Delort, A.-M.: Potential for phenol biodegradation in cloud waters, *Biogeosciences*, 15(18), 5733–5744, doi:10.5194/bg-15-5733-2018, 2018.

Lelieveld, J. and Crutzen, P. J.: Influences of cloud photochemical processes on tropospheric ozone, *Nature*, 343, 227–233, 1990.

Li, W., Yang, J., Zhang, D., Li, B., Wang, E. and Yuan, H.: Concentration and Community of

- 700 Airborne Bacteria in Response to Cyclical Haze Events During the Fall and Midwinter in Beijing, China, *Front. Microbiol.*, 9, 1741, doi:10.3389/fmicb.2018.01741, 2018.
- Lighthart, B.: The ecology of bacteria in the al fresco atmosphere, *FEMS Microbiol. Ecol.*, 23(4), 263–274, 1997.
- Lighthart, B. and Shaffer, B. T.: Bacterial flux from chaparral into the atmosphere in mid-summer at a high desert location, *Atmos. Environ.*, 28(7), 1267–1274, doi:10.1016/1352-2310(94)90273-9, 1994.
- Lighthart, B. and Shaffer, B. T.: Viable bacterial aerosol particle size distributions in the midsummer atmosphere at an isolated location in the high desert chaparral, *Aerobiologia*, 11(1), 19–25, 1995.
- 710 Lin, G., Sillman, S., Penner, J. E. and Ito, A.: Global modeling of SOA: the use of different mechanisms for aqueous phase formation, *Atmos Chem Phys*, 14, 5451–5475, doi:10.5194/acp-14-5451-2014, 2014.
- Lindemann, J., Constantinidou, H. A., Barchet, W. R. and Upper, C. D.: Plants as sources of airborne bacteria, including ice nucleation-active bacteria, *Appl. Environ. Microbiol.*, 44(5), 1059–1063, 1982.
- 715 Lipson, D. A., Schadt, C. W. and Schmidt, S. K.: Changes in Soil Microbial Community Structure and Function in an Alpine Dry Meadow Following Spring Snow Melt, *Microb. Ecol.*, 43(3), 307–314, doi:10.1007/s00248-001-1057-x, 2002.
- Liu, J., Horowitz, L. W., Fan, S., Carlton, A. G. and Levy, H.: Global in-cloud production of secondary organic aerosols: Implementation of a detailed chemical mechanism in the GFDL atmospheric model AM3, *J Geophys Res - Atmos*, 117(D15), D15303, doi:10.1029/2012jd017838, 2012.
- 720 Madronich, S., Bjorn, L. O. and McKenzie, R. L.: Solar UV radiation and microbial life in the atmosphere, *Photochem. Photobiol. Sci.*, 17(12), 1918–1931, doi:10.1039/C7PP00407A, 2018.
- 725 Marinoni, A., Parazols, M., Brigante, M., Deguillaume, L., Amato, P., Delort, A.-M., Laj, P. and Mailhot, G.: Hydrogen peroxide in natural cloud water: Sources and photoreactivity, *Atmos Res*, 101(1–2), 256–263, doi:10.1016/j.atmosres.2011.02.013, 2011.
- Marr, A. G.: Growth rate of *Escherichia coli*, *Microbiol. Rev.*, 55(2), 316–333, 1991.
- Martin, A., Hall, J. and Ryan, K.: Low Salinity and High-Level UV-B Radiation Reduce Single-Cell Activity in Antarctic Sea Ice Bacteria, *Appl. Environ. Microbiol.*, 75(23), 7570, doi:10.1128/AEM.00829-09, 2009.
- 730 Matthias-Maser, S. and Jaenicke, R.: The size distribution of primary biological aerosol particles in the multiphase atmosphere, *Aerobiologia*, 16(2), 207–210, doi:10.1023/A:1007607614544, 2000.

- 735 Meisner, A., Leizeaga, A., Rousk, J. and Bååth, E.: Partial drying accelerates bacterial growth recovery to rewetting, *Soil Biol. Biochem.*, 112, 269–276, doi:10.1016/j.soilbio.2017.05.016, 2017.
- Michaud, J. M., Thompson, L. R., Kaul, D., Espinoza, J. L., Richter, R. A., Xu, Z. Z., Lee, C., Pham, K. M., Beall, C. M., Malfatti, F., Azam, F., Knight, R., Burkart, M. D., Dupont, C. L. and
740 Prather, K. A.: Taxon-specific aerosolization of bacteria and viruses in an experimental ocean-atmosphere mesocosm, *Nat. Commun.*, 9(1), 2017, doi:10.1038/s41467-018-04409-z, 2018.
- Middelboe, M.: Bacterial Growth Rate and Marine Virus–Host Dynamics, *Microb Ecol*, 114–124, 2000.
- Möhler, O., Georgakopoulos, D. G., Morris, C. E., Benz, S., Ebert, V., Hunsmann, S., Saathoff, H.,
745 Schnaiter, M. and Wagner, R.: Heterogeneous ice nucleation activity of bacteria: new laboratory experiments at simulated cloud conditions, *Biogeosciences*, 5(5), 1425–1435, doi:10.5194/bg-5-1425-2008, 2008.
- Monod, A. and Doussin, J. F.: Structure-activity relationship for the estimation of OH-oxidation rate constants of aliphatic organic compounds in the aqueous phase: alkanes, alcohols, organic
750 acids and bases, *Atmos. Environ.*, 42(33), 7611–7622, doi:DOI: 10.1016/j.atmosenv.2008.06.005, 2008.
- Monteil, C. L., Bardin, M. and Morris, C. E.: Features of air masses associated with the deposition of *Pseudomonas syringae* and *Botrytis cinerea* by rain and snowfall, *ISME J.*, 8(11), 2290–2304, doi:10.1038/ismej.2014.55, 2014.
- 755 Morris, C. E., Glaux, C., Latour, X., Gardan, L., Samson, R. and Pitrat, M.: The Relationship of Host Range, Physiology, and Genotype to Virulence on Cantaloupe in *Pseudomonas syringae* from Cantaloupe Blight Epidemics in France, *Phytopathology*, 90(6), 636–646, doi:10.1094/PHYTO.2000.90.6.636, 2000.
- Morris, C. E., D. G. Georgakopoulos and D. C. Sands: Ice nucleation active bacteria and their
760 potential role in precipitation, *J Phys IV Fr.*, 121, 87–103, 2004.
- Myriokefalitakis, S., Fanourgakis, G. and Kanakidou, M.: The Contribution of Bioaerosols to the Organic Carbon Budget of the Atmosphere, in *Perspectives on Atmospheric Sciences*, edited by T. Karacostas, A. Bais, and P. T. Nastos, pp. 845–851, Springer International Publishing., 2017.
- Newman, I. V.: Aerobiology on commercial air routes, *Nature*, 161(4086), 275, 1948.
- 765 Norris, V.: Why do bacteria divide?, *Front. Microbiol.*, 6, 322, doi:10.3389/fmicb.2015.00322, 2015.
- Olson, J.: World ecosystems (WE1.4): Digital raster data on a 10 minute geographic 1080 (2160 grid square), *Global Ecosystem Database*, Version 1, 1992.
- 770 Otero Fernandez, M., Thomas, R., Garton, N., Hudson, A., Haddrell, A. and Reid, J.: Assessing the Airborne Survival of Bacteria in Populations of Aerosol Droplets with a Novel Technology, *J. R.*

- Soc. Interface, 16(150), doi:10.1098/rsif.2018.0779, 2019.
- 775 Paez-Rubio, T., Viau, E., Romero-Hernandez, S. and Peccia, J.: Source Bioaerosol Concentration and rRNA Gene-Based Identification of Microorganisms Aerosolized at a Flood Irrigation Wastewater Reuse Site, *Appl. Environ. Microbiol.*, 71(2), 804, doi:10.1128/AEM.71.2.804-810.2005, 2005.
- Passananti, M., Vinatier, V., Delort, A.-M., Mailhot, G. and Brigante, M.: Siderophores in Cloud Waters and Potential Impact on Atmospheric Chemistry: Photoreactivity of Iron Complexes under Sun-Simulated Conditions, *Environ. Sci. Technol.*, 50(17), 9324–9332, doi:10.1021/acs.est.6b02338, 2016.
- 780 Perrino, C. and Marcovecchio, F.: A new method for assessing the contribution of Primary Biological Atmospheric Particles to the mass concentration of the atmospheric aerosol, *Environ. Int.*, 87, 108–115, doi:10.1016/j.envint.2015.11.015, 2016.
- 785 Posfai, M., Xu, H., Anderson, J. R. and Buseck, P. R.: Wet and dry sizes of atmospheric aerosol particles: An AFM-TEM study, *Geophys. Res. Lett.*, 25(11), 1907–1910, doi:10.1029/98GL01416, 1998.
- Price, P. B. and Sowers, T.: Temperature dependence of metabolic rates for microbial growth, maintenance, and survival, *Proc. Natl. Acad. Sci. U. S. A.*, 101(13), 4631–4636, doi:10.1073/pnas.0400522101, 2004.
- 790 Pruppacher, H. R. and Jaenicke, R.: The processing of water vapor and aerosols by atmospheric clouds, a global estimate, *Atmos Res*, 38(1–4), 283–295, doi:http://dx.doi.org/10.1016/0169-8095(94)00098-X, 1995.
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of clouds and precipitation*, 2nd revised and enlarged., Kluwer Academic Publishers, Dordrecht (NL), 2003.
- 795 Sand-Jensen, K., Pedersen, N. L. and Sandergaard, M.: Bacterial metabolism in small temperate streams under contemporary and future climates, *Freshw. Biol.*, 52(12), 2340–2353, doi:10.1111/j.1365-2427.2007.01852.x, 2007.
- Šantl-Temkiv, T., Finster, K., Hansen, B. M., Pašić, L. and Karlson, U. G.: Viable methanotrophic bacteria enriched from air and rain can oxidize methane at cloud-like conditions, *Aerobiologia*, 29(3), 373–384, doi:10.1007/s10453-013-9287-1, 2013.
- 800 Santl-Temkiv, T., Sahyoun, M., Finster, K., Hartmann, S., Augustin-Bauditz, S., Stratmann, F., Wex, H., Clauss, T., Nielsen, N. W., Sørensen, J. H., Korsholm, U. S., Wick, L. Y. and Karlson, U. G.: Characterization of airborne ice-nucleation-active bacteria and bacterial fragments, *Atmos. Environ.*, 109, 105–117, doi:10.1016/j.atmosenv.2015.02.060, 2015.
- 805 Sattler, B., Puxbaum, H. and Psenner, B.: Bacterial growth in supercooled cloud droplets, *Geophys. Res. Lett.*, 28(2), 239–242, doi:10.1029/2000GL011684, 2001.
- Si, F., Li, D., Cox, S. E., Sauls, J. T., Azizi, O., Sou, C., Schwartz, A. B., Erickstad, M. J., Jun, Y.,

- Li, X. and Jun, S.: Invariance of Initiation Mass and Predictability of Cell Size in *Escherichia coli*, *Curr. Biol.*, 27(9), 1278–1287, doi:10.1016/j.cub.2017.03.022, 2017.
- 810 Smith, D. J., Griffin, D. W., McPeters, R. D., Ward, P. D. and Schuerger, A. C.: Microbial survival in the stratosphere and implications for global dispersal, *Aerobiologia*, 27(4), 319–332, doi:10.1007/s10453-011-9203-5, 2011.
- 815 Smith, D. J., Ravichandar, J. D., Jain, S., Griffin, D. W., Yu, H., Tan, Q., Thissen, J., Lusby, T., Nicoll, P., Shedler, S., Martinez, P., Osorio, A., Lechniak, J., Choi, S., Sabino, K., Iverson, K., Chan, L., Jaing, C. and McGrath, J.: Airborne Bacteria in Earth's Lower Stratosphere Resemble Taxa Detected in the Troposphere: Results From a New NASA Aircraft Bioaerosol Collector (ABC), *Front. Microbiol.*, 9, 1752, doi:10.3389/fmicb.2018.01752, 2018.
- 820 Souza, F. F. C., Rissi, D. V., Pedrosa, F. O., Souza, E. M., Baura, V. A., Monteiro, R. A., Balsanelli, E., Cruz, L. M., Souza, R. A. F., Andreae, M. O., Reis, R. A., Godoi, R. H. M. and Huergo, L. F.: Uncovering prokaryotic biodiversity within aerosols of the pristine Amazon forest, *Sci. Total Environ.*, 688, 83–86, doi:10.1016/j.scitotenv.2019.06.218, 2019.
- Stibal, M. and Elster, J.: Growth and morphology variation as a response to changing environmental factors in two Arctic species of *Raphidonema* (Trebouxiophyceae) from snow and soil, *Polar Biol.*, 28(7), 558–567, doi:10.1007/s00300-004-0709-y, 2005.
- 825 Tesson, S. V. M. and Santl-Temkiv, T.: Ice Nucleation Activity and Aeolian Dispersal Success in Airborne and Aquatic Microalgae, *Front. Microbiol.*, 9, 2681, doi:10.3389/fmicb.2018.02681, 2018.
- Vaïtilingom, M., Amato, P., Sancelme, M., Laj, P., Leriche, M. and Delort, A.-M.: Contribution of Microbial Activity to Carbon Chemistry in Clouds, *Appl. Environ. Microbiol.*, 76(1), 23–29, doi:10.1128/AEM.01127-09, 2010.
- 830 Vaïtilingom, M., Charbouillot, T., Deguillaume, L., Maisonobe, R., Parazols, M., Amato, P., Sancelme, M. and Delort, A. M.: Atmospheric chemistry of carboxylic acids: microbial implication versus photochemistry, *Atmos Chem Phys*, 11(16), 8721–8733, doi:10.5194/acp-11-8721-2011, 2011.
- 835 Vaïtilingom, M., Attard, E., Gaiani, N., Sancelme, M., Deguillaume, L., Flossmann, A. I., Amato, P. and Delort, A.-M.: Long-term features of cloud microbiology at the puy de Dôme (France), *Atmos. Environ.*, 56(0), 88–100, doi:http://dx.doi.org/10.1016/j.atmosenv.2012.03.072, 2012.
- Vaïtilingom, M., Deguillaume, L., Vinatier, V., Sancelme, M., Amato, P., Chaumerliac, N. and Delort, A.-M.: Potential impact of microbial activity on the oxidant capacity and organic carbon budget in clouds, *Proc. Natl. Acad. Sci.*, 110(2), 559–564, doi:10.1073/pnas.1205743110, 2013.
- 840 Wang, J., Rossow, W. B. and Zhang, Y.: Cloud Vertical Structure and Its Variations from a 20-Yr Global Rawinsonde Dataset, *J. Clim.*, 13(17), 3041–3056, doi:10.1175/1520-0442(2000)013<3041:CVSAIV>2.0.CO;2, 2000.
- Wéry, N., Gales, A. and Brunet, Y.: Bioaerosol sources, in *Microbiology of Aerosols*, pp. 115–

135, A.M. Delort and P. Amato, Hoboken, NJ: John Wiley & Sons, Inc., 2017.

845 Whitman, W. B., Coleman, D. C. and Wiebe, W. J.: Prokaryotes: the unseen majority, *Proc. Natl. Acad. Sci. U. S. A.*, 95(12), 6578–6583, doi:10.1073/pnas.95.12.6578, 1998.

Wiedinmyer, C., Bowers, R. M., Fierer, N., Horanyi, E., Hannigan, M., Hallar, A. G., McCubbin, I. and Baustian, K.: The contribution of biological particles to observed particulate organic carbon at a remote high altitude site, *Atmos. Environ.*, 43(28), 4278–4282, doi:10.1016/j.atmosenv.2009.06.012, 2009.

Wirgot, N., Vinatier, V., Deguillaume, L., Sancelme, M. and Delort, A.-. M.: H₂O₂ modulates the energetic metabolism of the cloud microbiome, *Atmos Chem Phys*, 17, doi:10.5194/acp-17-14841-2017, 2017.

855 Xiang, S.-R., Doyle, A., Holden, P. A. and Schimel, J. P.: Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils, *Soil Biol. Biochem.*, 40(9), 2281–2289, doi:10.1016/j.soilbio.2008.05.004, 2008.

Xie, Z., Li, Y., Lu, R., Li, W., Fan, C., Liu, P., Wang, J. and Wang, W.: Characteristics of total airborne microbes at various air quality levels, *J. Aerosol Sci.*, 116, 57–65, doi:10.1016/j.jaerosci.2017.11.001, 2018.

860