



# The global impact of bacterial processes on carbon mass

**Barbara Ervens and Pierre Amato**

5 Université Clermont Auvergne, CNRS, Sigma-Clermont, Institut de Chimie de Clermont-Ferrand, 63000 Clermont-Ferrand, France

*Correspondence to:* barbara.ervens@uca.fr and pierre.amato@uca.fr

**Abstract.** Many recent studies have identified biological material as a major fraction of ambient aerosol loading. A  
10 small fraction of these bioaerosols consist of bacteria that have attracted a lot of attention due to their role in cloud  
formation and adverse health effects. Current atmospheric models consider bacteria as inert quantities and neglect cell  
growth and multiplication. We provide here a framework to estimate the production of secondary biological aerosol  
(SBA) mass in clouds by microbial cell growth and multiplication. The best estimate of SBA formation rates of 3.7 Tg  
15 yr<sup>-1</sup> are comparable to previous model estimates of the primary emission of bacteria into the atmosphere, and thus  
might represent a previously unrecognized source of biological aerosol material. We discuss in detail the large  
uncertainties associated with our estimates based on the rather sparse available data on bacteria abundance, growth  
conditions and properties. Additionally, the loss of water-soluble organic carbon (WSOC) due to microbial processes  
in cloud droplets has been suggested to compete under some conditions with WSOC loss by chemical (OH) reactions.  
Our estimates suggest that microbial and chemical processes might lead to a global loss of WSOC of 8 - 11 Tg yr<sup>-1</sup> and  
20 8 - 20 Tg yr<sup>-1</sup>, respectively. While also this estimate is very approximate, the analysis of the uncertainties and ranges  
of all parameters gives hints about the conditions under which microbial processes cannot be neglected as organic  
carbon sinks in clouds. Our estimates also highlight the urgent needs for more data concerning microbial  
concentrations, fluxes and activity in the atmosphere to evaluate the role of bacterial processes as net aerosol sink or  
source on various spatial and temporal scales.

## 25 1. Introduction

The characterization and quantification of outdoor bioaerosols is an active field of current atmospheric research since  
bioaerosols have been suggested to contribute to adverse health effects and cloud formation as ice-nucleating particles  
(Després et al., 2012). Biological material includes debris, pollen, bacteria, fungal spores, and viruses and is usually  
considered as being directly emitted to the atmosphere (primary biological aerosol, PBA (Jaenicke, 2005)). The total  
30 number and mass concentrations of PBA particles vary widely in space and time: Posfai et al. (1998) found 1% of  
particles with biological material above the Southern Ocean whereas Artaxo et al. (1990) identified more than 90% of  
all particles to contain biological material during the wet season in the Amazon. In an urban/remote region in Germany,  
24% of all particles were found to include a biological fraction (Matthias-Maser and Jaenicke, 2000). Similar  
concentrations were observed at a remote high-altitude site with 16 - 64% of the mass of particles with diameters of  
35 less than 10 µm being composed of biological mass (Wiedinmyer et al., 2009) whereas the PBA number fraction was




much smaller (0.3 - 18%) in Rome, Italy (Perrino and Marcovecchio, 2016). Bacteria only comprise a small fraction of the total biological aerosol mass but they alone can contribute up to about ~20% of the total number of particles with diameters greater than 0.5  $\mu\text{m}$  (Bowers et al., 2012).

Near the ground, typical concentrations of total airborne bacteria range from ~100 to  $10^6$  cells  $\text{m}^{-3}$ , depending on the emission source (Burrows et al., 2009b), and on temporal, meteorological, and other environmental conditions influencing its propensity to emit particles to the air (Carotenuto et al., 2017; Huffman et al., 2013; Lighthart, 1997; Lighthart and Shaffer, 1995). Atmospheric mixing aloft tends to homogenize the number and diversity of the various bacteria types as the distance from sources increases. In the free troposphere, concentrations of ~10,000 cells  $\text{m}^{-3}$  are reported, including in clouds (DeLeon-Rodriguez et al., 2013; Vařtilingom et al., 2013). Some extent of selection toward certain species of bacteria probably occurs during aerosolization and atmospheric transport (Joly et al., 2015; Michaud et al., 2018). However, such selection has not been clearly proven yet as the bacterial assemblages found at high altitude often resemble those observed near the ground (Amato et al., 2017; DeLeon-Rodriguez et al., 2013; Smith et al., 2018).

The atmosphere is a harsh environment for living microorganisms: low temperatures at high altitude, UV radiation (Madronich et al., 2018) and high free radical concentrations (Haddrell and Thomas, 2017; Marinoni et al., 2011) are thought to greatly challenge living organisms (Amato et al., 2019; Joly et al., 2015; Smith et al., 2011). Additionally, the rapidly changing conditions in clouds, like condensation/evaporation and freeze/thaw cycles, can cause strong physiological shocks and physical damages to cells, which can eventually be lethal. The viability of airborne microorganisms is thus very variable in space and time depending on environmental conditions (Fahlgren et al., 2010; Hu et al., 2017; Lighthart and Shaffer, 1995; Monteil et al., 2014), but yet the fact that a fraction of bacteria cells are viable was shown in many experiments of microbiological cultures from ambient aerosol samples (Amato et al., 2007c; Bovallius et al., 1978; Lighthart, 1997; Newman, 1948). This was specified and quantified more recently by direct observations and measurements of biological activity imprints (Amato et al., 2007b, 2017; Sattler et al., 2001; Wirgot et al., 2017). The multiplication of airborne bacteria was observed from aerosols generated from bacteria cultures (Dimmick et al., 1979), as well as in natural polluted fog (Fuzzi et al., 1997). Thus, the estimated emissions might be biased high as SBA formation provides an additional source of bacteria mass and, thus, observed bacteria concentrations represent the sum of emission fluxes that are smaller than assumed and the secondary production in the atmosphere. Efficient bacteria cell growth and multiplication are largely constrained by the presence of liquid water (Davey, 1989; Haddrell and Thomas, 2017). One can thus assume that microbial processes in the atmosphere are limited to the time microorganisms spend in clouds (*Figure 1*). Cell growth and multiplication lead to an increase of the initial cell mass and, thus, to more biological material. We introduce the term ‘secondary biological aerosol’ (SBA) mass here in order to distinguish this aerosol source from directly emitted PBA. Heterotrophic bacterial processes require the uptake of organic substrates by the cells, which are subsequently converted by metabolic processes into new organic products, biochemical energy and  $\text{CO}_2$  (‘respiration’, *Figure 1*). These substrates include organics (e.g., carboxylic acids, sugars); other elements (e.g., nitrogen, phosphorous, potassium, metals) are also needed and exist in bioavailable forms in cloud water. The biotransformation of formate, acetate, succinate, lactate, oxalate, and formaldehyde (Ariya et al., 2002; Vařtilingom et al., 2010), phenol (Lallement et al., 2018) and methane (Šantl-Temkiv et al., 2013) by bacteria and fungi was studied in aqueous solution mimicking the typical chemical composition of



75 cloud water, and it was suggested that under specific conditions, microbial processes might be competitive to chemical radical processes as sinks for these compounds (Delort et al., 2010; Vaitilingom et al., 2011, 2013). The efficiency of such metabolic processes strongly depends on the bacteria types, substrates and their availability within the cloud droplets. In the present study, we perform an estimate of the global importance of SBA formation and microbial WSOC loss. All parameters and their uncertainties are discussed based on the sparse data sets currently available.



**Figure 1:** Bacterial processes in the atmosphere leading to SBA formation and loss of water-soluble organic carbon (WSOC) in clouds

## 80 2. Data and assumptions on bacterial processes in clouds

### 2.1 Atmospheric concentrations of bacteria cells

Burrows et al. (2009a, 2009b) have summarized data on number concentrations and emission fluxes of bacteria above various ecosystems on the Earth surface. Their compilation also included the estimates of cell concentrations near the surface in a range of  $10,000 \text{ m}^{-3}$  (seas) up to  $650,000 \text{ m}^{-3}$  (urban). Bacteria populations aloft represent a mixture of  
85 bacteria that were emitted from different ecosystems and subsequently mixed (Burrows et al., 2009b). Despite these mixing processes, there are bacteria types that can be considered characteristic for each ecosystem (Wéry et al., 2017). **Table 1** lists cell concentrations as published by Burrows et al. (2009b) complemented by some more recent measurements. We extend this overview by data on bacteria types, suggested as predominant or characteristic for each ecosystem. In several cases, more than one predominant bacteria type is listed as specific geographical, meteorological  
90 and other environmental conditions might lead to differences in the diversity of bacteria populations for the same category of ecosystem. We also provide global average data (Category ‘All’) and define one of the most abundant bacteria type (Alpha-Proteobacteria) alive in the atmosphere (Amato et al., 2019; Klein et al., 2016) as a representative type.

The atmospheric lifetime of bacteria cells is limited to several minutes (Otero Fernandez et al., 2019) to hours (Amato



95 et al., 2015). Several studies report total concentrations of bacteria cells in the atmosphere whereas others present only  
 the concentration of viable cells. The complexity of distinguishing viable, cultivable and dead bacteria cells in the  
 atmosphere has been discussed in several studies (Burrows et al., 2009a; Otero Fernandez et al., 2019).

100 **Table 1:** Summary of ambient cell concentrations  $C_{cell}$  and generation rates  $R_{cell}$  for predominant bacteria types in all  
 ecosystems

	$C_{cell}$ [ $m^{-3}$ ] <sup>a</sup>	Representative strain affiliation	Generation rate $R_{cell}$ [ $h^{-1}$ ]
All	10,000	Alpha-Proteobacterium <i>Sphingomonas</i> sp. (average of 32b-11, 32b-49, 32b-57, 35b-32, 35b-38) <sup>e</sup>	0.06 (5°C) <sup>e</sup> 0.2 (17°C) <sup>e</sup> 0.35 (27°C) <sup>e</sup> 0.45 (37°C) <sup>e</sup>
Tundra	12,000	<i>Pseudomonas</i> spp. ( <i>P. graminis</i> ) <sup>e, f, g</sup>	0.12 (5°C) <sup>e</sup> 0.21 (17°C) <sup>e</sup> 0.82 (27°C) <sup>e</sup> 0.27 (37°C) <sup>e</sup>
		<i>Psychrobacter</i> sp. <sup>g</sup>	0.0007 (-10°C) <sup>g</sup>
		<i>Rhodococcus</i> sp. <sup>g</sup>	0.0001 (-10°C) <sup>g</sup>
Grassland	110,000	<i>Pseudomonas syringae</i> <sup>e, h</sup>	0.1 (5°C) <sup>e</sup> 0.25 (15°C) <sup>e</sup> 0.9 (27°C) <sup>e</sup>
Coastal	76,000		
Wetlands	90,000		
Crops	110,000	<i>Frigoribacterium</i> sp. <sup>e, i</sup>	
Land ice	(5,000)	<i>Raphidonema</i> spp. <sup>j</sup>	$1.7 \cdot 10^{-4}$ - $2.91 \cdot 10^{-4}$ (12-18°C) <sup>j</sup>
Deserts	(10,000) 612 <sup>c</sup>		
Forests	56,000 6,323-12,748 <sup>d</sup>		
Shrubs	350,000		
Seas	10,000	<i>Pseudoalteromonas</i> <sup>k</sup>	0.25 (T unknown) <sup>k</sup>
Seas (estuary) <sup>b</sup>		Gamma-Proteobacterium (the fastest) <sup>l</sup>	0.17 (14°C) <sup>l</sup> 0.19 (24°C) <sup>l</sup>

<sup>a</sup> All cell concentrations are taken from Burrows et al. (2009b) unless otherwise noted. <sup>b</sup> 'seas estuary' was not included as a separate ecosystem by Burrows et al. (2009b); <sup>c</sup> (Lighthart and Shaffer, 1994); <sup>d</sup> (Helin et al., 2017); <sup>e</sup> Amato et al. (unpublished data; strains originally reported in Amato et al., 2007c or Väitilingom et al., 2012); <sup>f</sup> (Männistö and Häggblom, 2006); <sup>g</sup> (Bakermans et al., 2003); <sup>h</sup>; (Morris et al., 2000) <sup>i</sup> (Copeland et al., 2015); <sup>j</sup> (Stibal and Elster, 2005); <sup>k</sup> (Middelboe, 2000); <sup>l</sup> (Fuchs et al., 2000)

105



We assume in **Section 3** that all bacteria cells as listed in **Table 1** are metabolically active. In our estimates, we neither include assumptions on the limited lifetime of bacteria cells nor on their residence time in the atmosphere as it is assumed that PBA emissions lead to a continuous replenishment of bacteria in the atmosphere resulting in a steady-state concentration of living cells. The consequences of limited cell life- and residence time on SBA formation warrant further studies in more sophisticated model approaches.

## 2.2 Cell generation rates $R_{cell}$

Different levels of metabolic activity can be distinguished, from survival, where cells only repair molecular damages, to maintenance (dormancy), where cells do not divide but maintain biological functions, to growth, allowing the net production of biological mass (Price and Sowers, 2004) (**Figure 1**). The generation rate of a microorganism during growth is probably the most common microbiological criterion used for characterizing microbial multiplication in the laboratory; it corresponds to the time that is needed for doubling the cell number, i.e., for producing two “children” cells from one individual. This requires mass production from nutrients that provide the necessary molecular bricks and biochemical energy. The activity depends on physiological traits of the microorganism, with optima at a given temperature, pH, salinity, and other conditions that define its fitness for its habitat. The generation time in bacteria at their optimum growth conditions usually ranges from ~20 minutes (Marr, 1991) to several days or weeks; as conditions deviate from the optima, this lengthen to virtually infinite time in non-dividing cells.

The *cellular* growth rate itself, i.e. the increase of individual cell size and mass, is intimately linked with generation time: cell size increases in a predictable way as generation time decreases (Si et al., 2017), and it can vary of a factor of up to eight within a single bacteria species. Compared to generation rates, cellular growth rates are usually small and, thus, in the following only data for generation rates are used to estimate SBA mass formation rates.

Metabolic activity, in terms of carbon uptake per units of biomass and time, can range over more than ten orders of magnitude, depending on many factors of which temperature is a major one (Price and Sowers, 2004). Therefore, if available, temperature-dependent generation rates  $R_{cell}$  are listed in **Table 1** and shown in **Figure 2**. In addition, the highest expectable growth rate for bacteria as measured under laboratory conditions in culture medium is also shown in the figure for constraining an upper theoretical limit. This corresponds to the generation rate of the laboratory model *Escherichia coli* under optimal conditions (37°C). However, it can be expected that this is not representative of situations encountered in clouds.


Generally, the temperature dependence of cell generation rates can be scaled by the empirical relationship

$$R_{cell}(T_2) = R_{cell}(T_1) \cdot Q_{10}^{(T_2-T_1)/10} \quad (1)$$

whereas  $R(T_1)$  and  $R(T_2)$  are the growth rates ( $\text{h}^{-1}$ ) at two temperatures  $T_1$  and  $T_2$ .  $Q_{10}$  is a dimensionless scaling factor that expresses the change of these rates over an interval of 10°C and typically has values between two and three within relatively small temperature intervals (Lipson et al., 2002; Sand-Jensen et al., 2007). In general, Equation (1) can be applied for all bacteria types and is usually valid for liquid water over a temperature range up to ~ 25°C; however, the slope ( $Q_{10}$  factor) and the maximum temperature depends on the bacteria type.



In **Figure 2**, the dashed lines towards lower temperatures represent extrapolations of the generation rates at  $\sim 20^{\circ}\text{C}$  reflecting the general agreement between measured and calculated temperature dependencies using  $Q_{10} = 2$  or 3, respectively. Using generation rates measured at  $\sim 20^{\circ}\text{C}$  might lead to an overestimate for SBA mass formation rates in colder clouds. However, we chose these values for the calculations in **Section 3** as most experimentally-derived growth rates are available for temperatures of 20 – 30°C. Generally, at temperatures below 0°C, cell metabolic activity is negligible in terms of carbon flux even though cells can maintain and survive under such conditions (Amato et al., 2009, 2010; Price and Sowers, 2004).



**Figure 2:** Temperature dependence of generation rates  $R_{cell}$  for bacteria types representative for the ecosystems in **Table 1** and *E. coli* as a likely upper limit for cell generation. Dashed lines are extrapolations of the rates at  $T \sim 20^{\circ}\text{C}$  using  $Q_{10} = 2$  or  $Q_{10} = 3$ , respectively.

## 2.2 Bacteria growth efficiency (BGE)

150 Chemoheterotrophs - representatives of which were shown to maintain metabolic activity in clouds (Amato et al., 2017) - take up carbon from dissolved organic material for both recovering biochemical energy and converting the substrates into  $\text{CO}_2$  and other products. Bacteria growth efficiency (BGE) is defined as the biological mass that is produced relatively to the amount of carbon taken up from the environment, the rest being converted into  $\text{CO}_2$  (Eiler et al., 2003):

$$\text{BGE} = \frac{d[\text{organic products}]}{d[\text{organic products}] + d[\text{CO}_2]} \quad (2)$$

155 Note that in the original literature BGE is defined as a measure of ‘biomass production’ instead of organic products’ in *Eq (2)*. Given the large body of atmospheric literature on aerosol processes that discusses ‘biomass’ as material from any living matter (e.g., aerosol from forest fires), we avoid using ‘biomass’ in the current context of microbial processes. BGE for planktonic bacteria range from  $< 0.4\%$  to 80% with the highest values for eutrophic conditions (Eiler et al., 2003). In turn, when substrate availability is limited, values from  $\sim 7\%$  to  $\sim 14\%$  are generally observed.

160 As the conditions in cloud water can be considered oligotrophic with typical concentrations of dissolved organic carbon (DOC) of less than 0.1 mM (Herckes et al., 2013), low BGEs in the range of 0.1 – 10% can be expected, i.e. DOC is efficiently converted into  $\text{CO}_2$ .



## 2.4 Cloud properties relevant for microbial activity

### 2.4.1 Cell concentrations in cloud water

165 Bacteria cells have sizes of up to several micrometers which explains their high efficiency to act as cloud condensation  
nuclei (CCN) (Bauer et al., 2003; Després et al., 2012). Assessing the hygroscopicity of biological particles is complex  
since it cannot be calculated in a similar way as for chemical compounds where total hygroscopicity represents the  
sum of the contributions of all components (Ariya et al., 2009). Once particles form cloud droplets, chemical  
compounds dissolved in cloud water, will trigger the growth of the processed CCN and, thus, enhancing hygroscopicity  
170 and CCN activity of aged particles in subsequent cloud cycles. The dissolution of ambient cell populations of 100 to  
50,000  $\text{m}^{-3}$  (**Table 1**) results in 200 to 500,000 cells  $\text{mL}^{-1}$  for clouds with liquid water contents (LWC) of 0.5  $\text{g m}^{-3}$  and  
0.1  $\text{g m}^{-3}$  (Vaithilingom et al., 2010). The reasonable agreement of cell concentrations outside of clouds and those in  
cloud water suggests that a large fraction of bacteria cells are scavenged and, thus, act as CCN. Some bacteria are well-  
known to efficiently act as ice nuclei (Amato et al., 2015; Möhler et al., 2008; Morris et al., 2004). In the current study,  
175 we neglect the potential role of ice clouds as media of microbial metabolic activity. In addition to low temperatures  
resulting in very low generation rates (**Figure 2**), the substrate diffusion to the bacteria will be limited resulting in  
negligible consumption of dissolved carbon.

### 2.4.2 Time fractions of microbial processes in clouds ( $F_{cloud}$ )

As we assume that both SBA mass production and WSOC loss only occur when bacteria are suspended in cloud  
180 droplets, we need to estimate the time bacteria spend in liquid clouds. In general, cloud contact times, i.e., the time air  
spends in a cloud, are dependent on cloud depth and vertical velocity (Feingold et al., 2013). This small-scale  
information is not consistently available for the large regions as covered by the ecosystems listed in **Table 1**. Instead,  
the fraction of cloud contact time  $F_{cloud}$  is here equaled to the atmospheric volume fraction that is filled with clouds  
(Lelieveld and Crutzen, 1990).

185 Global cloud coverage data based on satellite images derived from MODIS data (King et al., 2013) show cloud  
coverage of  $\sim 0.2 - 0.9$  above various ecosystems. In order to convert these two-dimensional values of cloud coverage  
into three-dimensional values that describe the volume fraction of the atmosphere that is filled with clouds, we assume  
an average cloud thickness of 1.5 km within the lowest 6 km of the atmosphere where  $> 90\%$  of all liquid clouds reside  
(Pruppacher and Jaenicke, 1995; Wang et al., 2000). This results in a conversion factor of 0.25 (1.5 km / 6 km) to  
190 obtain an average cloud volume, and thus  $F_{cloud}$ , above each ecosystem.

Comparison of previous estimates of global cloud coverage of 60% (Pruppacher and Jaenicke, 1995) and the volume  
fraction of liquid clouds within the atmosphere of 15% (Lelieveld and Crutzen, 1990) generally supports this  
relationship. The resulting  $F_{cloud}$  values are summarized in **Table 2** together with the percentage area fraction of each  
ecosystem of the Earth surface, taken from Burrows et al. (2009b) and originally obtained from Olson (1992).



195 **Table 1:** Surface coverage of ecosystems on Earth surface (Burrows et al., 2009b) and estimated time fraction bacteria spend in clouds ( $F_{cloud}$ )

	% of Earth surface <sup>a)</sup>	$F_{cloud}$
All	100	0.15
Tundra	3.3	0.1
Grassland	2.2	0.15
Coastal	0.2	0.1
Wetlands	0.6	0.15
Crops	3.0	0.25
Land ice	3.1	0.1
Deserts	3.7	0.05
Forests	7.0	0.2
Shrubs	5.8	0.2
Seas	71.0	0.17

<sup>a)</sup> Data from Burrows et al. (2009b) and Olson (1992)

## 200 3. Results and Discussion

### 3.1 SBA mass production

#### 3.1.1 Calculation of SBA formation rates

We calculate the SBA mass formation rate [ $\text{ng m}^{-3} \text{day}^{-1}$ ] above each ecosystem  $i$ , using Eq (3):

$$\left(\frac{dm}{dt}\right)_{SBA,i,day} = R_{cell,i} \cdot F_{live} \cdot C_{cell,i} \cdot F_{cloud,i} \cdot m_{cell} \quad (3)$$

205 where  $R_{cell}$  is the cell generation rate [ $\text{h}^{-1}$ ] (**Table 1**). For ecosystems, for which  $R_{cell}$  of the representative bacteria types is not available, we assume the average value of  $R_{cell} = 0.3 \text{ h}^{-1}$  as an upper limit for atmospherically-relevant conditions, corresponding to a generation time of approximately three hours.  $C_{cell}$  denotes the ambient cell concentration [ $\text{cell m}^{-3}$ ] (**Table 1**),  $F_{live}$  is the fraction of living cells in total cell concentration and assumed to be unity here,  $F_{cloud}$  is the fraction of total time bacteria when are active in clouds (**Table 2**), and  $m_{cell}$  is the average mass of a single cell, independent of the bacteria type. The cell mass  $m_{cell}$  is assumed to be  $52 \cdot 10^{-15} \text{ g cell}^{-1}$  (Sattler et al., 2001), equivalent of a spherical particle with diameter of 500 nm and a density of  $1 \text{ g cm}^{-3}$ .

For nearly all ecosystems, predicted SBA formation rates are in the range of  $\sim 0.1$  to  $\sim 1 \text{ ng m}^{-3} \text{ s}^{-1}$  (**Figure 3a**), with higher values for crops and shrubs where  $C_{cell}$  were found to be highest (**Table 1**). The average formation rate ( $0.6 \text{ ng m}^{-3} \text{ day}^{-1}$ ), calculated using the average values representative for all ecosystems (Category ‘All’ in **Table 1**), is similar to most of the other formation rates, suggesting that using these average data for a global estimate gives a reasonable order of magnitude of SBA formation. Only above land ice where  $C_{cell}$  is small, the rate is significantly smaller. Given that the temperatures above land ice might be on average lower than above other regions, the relative importance of SBA formation there might be even smaller.


215 To compare the mass production to other global aerosol sources,  $m_{SBA,day}$  is converted into a production flux [ $\text{Tg yr}^{-1}$ ]  
 220 for each ecosystem  $i$  and scaled by the surface fraction of each ecosystem  $A_i$ :





$$P_{SBA,i} = \left( \frac{dm}{dt} \right)_{SBA,i,day} \cdot 365 \text{ days} \cdot A_i \cdot V_{atmos} \quad (4)$$

where  $V_{atmos}$  is the volume of atmosphere ( $3 \cdot 10^{18} \text{ m}^3$ , 0 – 6 km) and  $A_i$  is the surface fraction of each ecosystem (*Table 2*).



**Figure 3:** Predicted production of secondary biological aerosol mass above the various ecosystems. The blue bar indicates the predicted production using the average values for all ecosystems ('all' in Table 1); a) SBA formation rates [ $\text{ng m}^{-3} \text{ day}^{-1}$ ]; b) SBA production rates [ $\text{Tg year}^{-1}$ ]. The red shaded bar represents the sum of the contributions from all ecosystems, bacteria emissions are shown for comparison, taken from Burrows

225

The production fluxes for each ecosystem are shown in *Figure 3b*, together with their sum for all ecosystems. The total predicted amount of SBA production is  $3.7 \text{ Tg yr}^{-1}$  with highest contributions from bacterial activities above seas ( $0.5 \text{ Tg yr}^{-1}$ ) as they cover most of the globe (71%) and above shrubs ( $0.5 \text{ Tg yr}^{-1}$ ) since the highest bacteria concentrations have been identified there (*Table 1*). Forests with a much smaller surface area (7%) but higher cell concentration contribute the same amount ( $0.5 \text{ Tg yr}^{-1}$ ). We suggest that this formation rate of  $3.7 \text{ Tg yr}^{-1}$  can be

230



considered the best estimate of SBA formation based on the available sparse data set presented in **Tables 1 and 2**. Using average data (Category ‘All’ in **Tables 1 and 2**) instead of those for individual ecosystems results in  $0.7 \text{ Tg yr}^{-1}$ , i.e. on a similar order of magnitude as the more sophisticated best estimate. These values are similar to the range of  $1 - 10 \text{ Tg yr}^{-1}$  that was extrapolated by Sattler et al. (2001) based on carbon production rates of bacteria in supercooled clouds at mount Sonnblick observatory in the Austrian Alps. Our estimated SBA mass production represents the production of total bacteria mass. The carbon content of bacteria cells is roughly 50% of their dry mass, with the remainder composed of nitrogen, oxygen, phosphorous, hydrogen and other elements (Whitman et al., 1998). Thus, we suggest that SBA formation may lead to  $1.9 \text{ Tg carbon yr}^{-1}$  ( $0.4 \text{ Tg carbon yr}^{-1}$  if the average value is applied, respectively) that is bound in biological mass.

### 240 3.1.2 Discussion of uncertainties in SBA formation

The formation rates in **Section 3.1.1** represent an estimate of a previously unrecognized source of biological aerosol mass in the atmosphere. All parameters are associated with large uncertainties that need to be constrained in the future as they might vary depending on temporal, meteorological, spatial and geographical conditions. Ranges of observed values for all parameters in Eq (3) and Eq (4) are summarized in **Table 3** and discussed in the following:

245 (i) The cell concentrations  $C_{cell}$  in the atmosphere homogenize aloft due to mixing processes and average to concentrations of  $\sim 10,000 \text{ m}^{-3}$  at most locations. However, spatial deviations might be present in particular locations, such as cell concentrations of  $> 10^6 \text{ m}^{-3}$  that were found in highly polluted areas (Xie et al., 2018), and even  $10^9 \text{ m}^{-3}$  near the ground (Paez-Rubio et al., 2005).

250 (ii) The growth rates  $R_{cell}$  assumed here likely represent an overestimate as cloud temperatures are likely lower than  $\sim 15\text{-}20^\circ\text{C}$ . At temperatures  $> \sim 0^\circ\text{C}$ , this overestimate is likely less than an order of magnitude (**Figure 2**); in supercooled clouds ( $< 0^\circ\text{C}$ ), metabolic activity  $R_{cell}$  might be restricted to occur in liquid droplets whereas in ice particles, it is some orders of magnitude lower and cell multiplication can be considered negligible.

255 (iii) While some of the studies listed by Burrows et al. (2009b) and in **Table 1** report the concentrations of viable cells, others give the total cell concentrations. In addition, the large discrepancy in reported  $F_{live}$  between  $< 0.1\%$  up to nearly 100% as discussed e.g., by Lindeman et al. (1982), Lighthart and Shaffer (1994) and Gandolfi et al. (2013), might be also due to differences in the measurement techniques. Consistent experimental methodologies are needed to give comprehensive data on  $F_{live}$  and the survival rates of bacteria in aerosol particles (Otero Fernandez et al., 2019).

260 (iv) The average cell mass depends on bacteria type and their growth stage. Sattler et al. (2001) estimated carbon mass of bacteria cells in cloud water as  $17 \text{ fg carbon cell}^{-1}$  in agreement with values for marine and freshwater ecosystems. Approximating the total mass by doubling the carbon mass, results in  $34 \text{ fg cell}^{-1}$ , i.e. equivalent to spherical particles with diameters of  $\sim 0.4 \mu\text{m}$ . Carbon mass and the carbon-to-total mass ratio can greatly differ from these values; for example, total masses of prokaryotic cell of  $200 \text{ fg cell}^{-1}$  in soil have been reported (Whitman et al., 1998). In their global study, Burrows et al. (2009b) assumed a mass of  $520 \text{ fg cell}^{-1}$  ( $1 \mu\text{m}$  particle).

265 (v) While several studies have shown that liquid water is necessary for efficient microbial activity, it is not clear yet whether bacteria maintain activity in wet aerosols. Klein et al. (2016) found indications that bacteria metabolic activity exists in aerosols but no quantitative data is reported yet. Bacteria become dormant or have reduced viability at relative



humidities of 86 - 97% (Haddrell and Thomas, 2017). In soil samples, it has been shown that cycles of drying and rewetting might enhance microbial activity compared to constantly moist samples (Meisner et al., 2017; Xiang et al., 2008); thus, it may be speculated that such effects also occur in rapidly changing humidity conditions in atmospheric deliquesced aerosols. Under those conditions, the time fraction of microbial activity would exceed  $F_{cloud}$ . On locally smaller scales, cloud processing time might be shorter than the average  $F_{cloud}$  (Table 2). SBA mass formation calculated by Eq (3) and Eq (4) depends linearly on all parameters discussed in (i) to (v). Thus, the sensitivity of the predicted formation rates can be simply estimated by ranges given in **Table 3**. However, in the atmosphere, all parameters might continuously change over time and thus might affect SBA mass to different extents.

275

**Table 3:** Parameters used in the estimate of SBA mass formation and their possible minimum and maximum values based on literature data

Parameter	Value in Eq (3) and Eq (4)	Range		Comment
		Minimum	Maximum	
$C_{cell}$ [ $m^{-3}$ ]	$10^4$	100	$10^9$	$C_{cell,min}$ : above desert during low RH and high radiation (Lighthart and Shaffer, 1994); $\sim 10^6$ : in a highly polluted area (Xi'an, China) (Xie et al., 2018); $C_{cell,max}$ was measured near the ground (Paez-Rubio et al., 2005)
$R_{cell}$ [ $h^{-1}$ ]	0.3	0	3	$R_{cell} \sim 0$ might occur under stressful conditions when cells become dormant (e.g. low temperature, little water...); $R_{cell} \sim 3$ for <i>E. coli</i> under optimal conditions (37°C, appropriate culture medium).
$F_{live}$	1	0.0001	1	0.0001 – 0.2 based on global microbial diversity (Gandolfi et al., 2013); 0.22 above crop fields (Lindemann et al., 1982), 0.81 above desert (Lighthart and Shaffer, 1994). Some studies report concentrations of viable cells; in this case reported concentrations imply $F_{live} = 1$
$m_{cell}$ [ $fg\ cell^{-1}$ ]	52	34	520	$m_{cell,min}$ corresponds to cells in clouds assuming that they are composed of 50% carbon (Sattler et al., 2001); masses of other prokaryotic cells might be up to 200 $fg\ cell^{-1}$ (Whitman et al., 1998); $m_{cell,max}$ is for a spherical cell of a diameter $1\mu m$ and density of $1\ g\ cm^{-3}$ (Burrows et al., 2009b).
$F_{cloud}$	0.15	> 0	1	The average global value might be higher than 0.15 if bacterial processes also occur outside of clouds. On small scales or above individual ecosystems, the value can be smaller than the average value.

### 3.1.3 Comparison of SBA formation to other aerosol sources

An estimate of aerosol emissions from the biosphere suggested a source strength of primary biological particles of 1000 Tg  $yr^{-1}$  (Jaenicke, 2005). However, in this study, PBA was defined to include all cellular material, proteins, and

280



their fragments. A global model study predicted total PBA emissions (bacteria, fungal spores and pollen) of  $123 \text{ Tg yr}^{-1}$  of which bacteria comprised  $0.79 \text{ Tg yr}^{-1}$  (Myriokefalitakis et al., 2017). This number is similar to the range of  $0.4 - 1.8 \text{ Tg bacteria yr}^{-1}$  as given by Burrows et al (2009b). Fungal spores are considered a major fraction of PBA. Their global emissions ( $25 \text{ Tg yr}^{-1}$ ) was suggested to contribute 23% to total primary organic aerosol (Heald and Spracklen, 2009). An estimate of fungal spore emissions based on tracer compounds resulted in predicted  $50 \text{ Tg yr}^{-1}$  (Elbert et al., 2007)). None of these estimates include microbial activity as a source of biological mass. Our predicted SBA source of  $3.7 \text{ Tg year}^{-1}$  is on the same order of magnitude as the model predictions for primary bacteria emissions. The estimates of bacteria emissions are performed such that observed cell concentrations are matched by the models without considering another source of cells in the atmosphere. Our SBA estimates might be equally biased as they are based on the same ambient cell concentrations as used by Burrows et al. (2009a, b) and Myriokefalitakis et al. (2017). The absolute values and the ratio of primary to secondary bacteria mass need to be evaluated by more complex model studies as our simple framework can provide.

Total organic aerosol is composed of mostly secondary mass. Best estimates based on observational and model studies of the net production rate of secondary organic aerosol (SOA) mass are on the order of  $136 - 280 \text{ Tg year}^{-1}$  (Hodzic et al., 2016). These amounts are similar to the predicted global sulfate production of  $117 \text{ Tg year}^{-1}$  ( $39 \text{ Tg S year}^{-1}$ ) (Chin et al., 2000). Thus, SBA production can be estimated to be on the order of  $\sim 1\%$  of the total aerosol sources. The net aerosol mass formation due to SBA production might be even smaller if bacteria metabolize substrates that are already in the particle phase. In this case, biotransformation processes lead to the conversion of non-biological into biological aerosol mass. The unique properties of biological aerosol material have been extensively discussed in the context of heterogeneous ice nucleation where it has been shown that even small amounts of biological material could have significant effects on clouds and precipitation (Möhler et al., 2008; Morris et al., 2004; Santl-Temkiv et al., 2015). Given the low ambient concentrations of ice nucleating particles and their high sensitivity to the ice/liquid partitioning in mixed-phase clouds (e.g., Ervens et al., 2011), a small change in biological mass possibly translates into significant changes in the evolution of cold clouds.

## 3.2 Consumption of organic carbon in clouds

### 3.2.1 Calculation of microbial and chemical WSOC loss rates

Bacteria can be metabolically active in the aqueous phase of clouds (Delort et al., 2010) and on the surface or bulk phase of particles (Klein et al., 2016; Estillore et al., 2016). Such processes lead to a decrease of water-soluble organic carbon (WSOC) mass within cloud droplets as bacteria convert substrates into  $\text{CO}_2$  (Figure 1). Also processes of biological mass production from  $\text{CO}_2$  exist (autotrophy) and include photosynthesis (photoautotrophs). However, despite the fact that photosynthetic microorganisms were reported in the atmosphere (Tesson and Santl-Temkiv, 2018) there is neither a clear indication yet of photosynthetic activity in clouds, nor of other modes of autotrophy.

The split between carbon uptake for biological mass production and mineralization is quantified by the bacterial growth efficiency BGE (Eq (2)). Studies have shown that generally metabolic processes produce mostly  $\text{CO}_2$  under atmospheric conditions and only a small fraction of the carbon is mineralized into organic products ( $< 1 - 10\%$  of the



total C utilized). We introduce here the factor  $F_{CO_2}$  as a measure of the loss of organics due to bacterial processes:

$$F_{CO_2} = 1 - BGE \quad (5)$$

Using  $F_{CO_2}$ , we can write the carbon balance as



320 The loss rate of carbon can be, thus, calculated as

$$R_{WSOC,Bact} = -\frac{d(WSOC)}{dt} = \frac{dCO_2}{dt} = \quad (6)$$

$$-F_{CO_2} k_{Bact} \left[ \frac{L}{cell \cdot s} \right] \cdot C_{Cell, aq} \left[ \frac{cell}{L} \right] \cdot F_C \cdot C_{WSOC} \left[ \frac{gC}{L} \right] \cdot LWC \left[ \frac{g}{m^3} \right] \cdot F_{cloud}$$


where  $C_{Cell, aq}$  and  $C_{WSOC}$  are the concentrations of bacteria and water-soluble organic carbon in cloud water, respectively. These concentrations are on average  $\sim 10^7$  cells  $L^{-1}$  (Vařtilingom et al., 2013) and 0.1 mM (Herckes et al.,  
325 2013) in cloud water whereas both values might differ over a few orders of magnitude locally and temporally (**Section 3.2.2**). Usually experimental loss rates of organics by bacteria in real and artificial cloud water are reported to be on the order of  $\leq 10^{-17}$  mol  $cell^{-1} s^{-1}$  for organic substrates (e.g. formic, acetic, and succinic acids) (Vařtilingom et al., 2010, 2011). The cell activity is dependent on the bacteria type and the availability of the organic substrate. Thus, strictly, such rates [ $mol\ cell^{-1} s^{-1}$ ] are only valid for the substrate-to-cell ratio as applied in the experiments. In order to account  
330 for the ratio as encountered in cloud water, we use here  $k_{Bact}$  [ $L\ cell^{-1} s^{-1}$ ] (i.e. measured rate divided by the concentration of organic substrate in the experiments) that is applicable to the full range of conditions where the cells exhibit a similar microbial activity. Resulting rate constants for formic, acetic and succinate acids are on the order of  $k_{Bact} \sim 10^{-13}$  L  $cell^{-1} s^{-1}$ .

Bacteria are selective towards organic compounds and not all organics are metabolized by every bacteria type. For  
335 example, it has been shown in a single study that, upon laboratory incubation of cloud water oxalic acid is not affected by cloud borne microorganisms, formate is only consumed after a lag time of several hours, which is much longer than the lifetime of a cloud droplet, and, on the other hand, compounds such as acetate or succinate are readily degraded (Vařtilingom et al., 2011). Since these compounds comprise major constituents of WSOC in cloud water, it seems reasonable to introduce an additional factor  $F_C$  in Eq (5) that accounts for the microbial selectivity towards only some  
340 organics by each bacteria type (e.g. Šantl-Temkiv et al., 2013; Bianco et al., 2019). Obviously, this factor is highly uncertain and strongly depends on the microbial and chemical composition of cloud water. The calculated rates  $R_{Bact, WSOC}$  are summarized in **Figure 4** for  $0.8 \leq F_{CO_2} \leq 0.99$  (Eiler et al., 2003) and  $F_C = 0.5$  (based on the characterization of metabolic activity in cloud water towards predominant cloud-water organics).

Several studies have discussed the competition of microbial and chemical processes in cloud water as a sink of specific  
345 organic compounds (Ariya et al., 2002; Husárová et al., 2011; Vařtilingom et al., 2010, 2013). The most efficient loss reactions for organics in cloud water are initiated by OH radicals. The general rate constant of the OH radical with water-soluble organic carbon is  $k_{OH} = 3.8 \cdot 10^8$   $M^{-1} s^{-1}$  (Arakaki et al., 2013). The reactions of WSOC with OH lead to




volatile and non-volatile oxidation products. Radicals are much less selective towards organics than bacteria are; thus, the assumption of a factor equivalent to  $F_C$  in Eq (6) is not necessary as all water-soluble organics react with OH  
 350 whereas the chemical reactivity mostly depends on the structure of the organic compound. The yield of volatile products ( $Y_{volC}$ ) includes  $CO_2$  but also formaldehyde and other volatile compounds that do not remain in the particle phase after cloud evaporation and thus do not contribute to the aerosol loading. We assume  $0.2 \leq Y_{volC} \leq 0.5$  (cf. **Section 3.2** and **Table 4** for the discussion of the assumptions and uncertainties of this parameter). In general,  $Y_{volC}$  depends on the WSOC composition, with higher values for more aged organics that are more readily oxidized to volatile products.  
 355 Equivalent to reaction (R 1), we express the carbon loss by the OH radical in clouds as



With  $WSOC_{aer}$  being the WSOC fraction that remains in the aerosol phase after drop evaporation. We calculate the loss rate accordingly:

$$R_{OH,WSOC} = -\frac{d(WSOC)_{OH}}{dt} = k_{OH} [L mol^{-1} s^{-1}] [OH]_{aq} [mol L^{-1}] \cdot Y_{volC} C_{WSOC} \left[\frac{gC}{L}\right] \cdot LWC \left[\frac{g}{m^3}\right] \cdot F_{cloud} \quad (7)$$

OH concentrations in cloud water are in the range of  $10^{-16} M < [OH]_{aq} < 10^{-14} M$  (Arakaki et al., 2013; Bianco et al., 2015; Ervens et al., 2014) and an average cloud liquid water content (LWC) of  $0.15 g m^{-3}$  is assumed. The results in **Figure 4** show that the predicted rates  $R_{OH,WSOC}$  ( $8 - 20 Tg yr^{-1}$ ) and  $R_{Bact,WSOC}$  ( $8 - 11 Tg yr^{-1}$ ) are for most parameter sets on a similar order of magnitude with possibly a slightly higher contribution of chemical reactions to WSOC loss.  
 365 This trend is in agreement with several previous studies that focused on the comparison of microbial versus chemical processes as sinks for specific organic substrates (Amato et al., 2007a; Vätilingom et al., 2010). These loss fluxes are



**Figure 4:** Predicted loss of WSOC by bacterial utilization and by chemical (OH) processing in cloud water for different assumption on  $F_{CO_2}$  and  $Y_{volC}$ ,



relatively large as compared to the predicted SBA formation (**Figure 3**).

In a previous study, it was estimated that microbial processes in clouds lead to a total carbon loss of  $\sim 10 - 50 \text{ Tg yr}^{-1}$  and to a production of  $\sim 100 \text{ Tg year}^{-1} \text{ CO}_2$  with the assumptions of complete respiration ( $F_{\text{CO}_2} = 1$ ), microbial non-selectivity towards WSOC ( $F_C = 1$ ) and applying the same loss rates as observed in lab experiments without correcting for differences in the ratio of bacteria cell to WSOC concentrations (Vaithilingom et al., 2013). Thus, this former estimate can be considered an upper limit whereas the one in the current study ( $\sim 50 \text{ Tg yr}^{-1}$ ) is more conservative, both suggesting that the respiration of bacteria is a negligible  $\text{CO}_2$  source as compared to the sum of anthropogenic sources ( $\sim 50,000 \text{ Tg CO}_2 \text{ year}^{-1}$  (IPCC, 2014)).

The WSOC loss rates calculated here are also much smaller than those estimated for the loss of total SOA due to heterogeneous reactions and photolysis on particle surfaces ( $\leq 50.3 \text{ Tg year}^{-1}$ ) and by wet deposition ( $\leq 50 \text{ Tg year}^{-1}$ , (Hodzic et al., 2016)). They are also smaller than the predicted production rates of aqSOA in clouds of  $13.1 - 46.8 \text{ Tg year}^{-1}$  (Lin et al., 2014) or  $20 - 30 \text{ Tg yr}^{-1}$  (Liu et al., 2012). These predictions strongly depend on the representation of clouds, and particularly on the liquid water content (He et al., 2013) but overall emphasize that clouds can be considered a net source rather than a sink for organic aerosol.

### 3.2.2 Discussion of uncertainties of microbial and chemical WSOC loss

The calculation of microbial and chemical WSOC loss should be regarded an approximation using a set of parameters that are all associated with considerable uncertainties. Similar to the discussion in **Section 3.1.2**, we compile all parameters and minimum and maximum values based on literature data in **Table 4**:

(i) There are not as many measurements of  $C_{\text{Bact,aq}}$  as for cell concentrations  $C_{\text{cell}}$  in cloud-free regions. The assumption that all particles that contain bacteria cells are activated into cloud droplets does likely not lead to a large overestimate. The sizes of bacteria-containing particles usually exceed several hundred nanometers and thus can all be considered CCN. Differences in LWC as the conversion factor from gas to aqueous phase concentrations are relatively small, within a factor of  $2 - 3$ , within the categories of common cloud types (Pruppacher and Klett, 2003).

(ii) The activity of microorganisms towards organic substrates is often reported in units of 'mol(substrate) cell $^{-1}$  s $^{-1}$ ', which expresses the amount of substrate that is consumed per cell and time. For several compounds (e.g., formate, acetate, succinate) these rates differ by approximately one order of magnitude (Vaithilingom et al., 2011). However, the resulting  $k_{\text{Bact}}$  (i.e. rate divided by substrate concentration) are all on the order of  $k_{\text{Bact}} \sim 10^{-13} \text{ L cell}^{-1} \text{ s}^{-1}$  which appears to represent an upper limit for the organics that have been investigated for metabolic activity in clouds. A much lower constant was derived from experiments with less oxygenated compounds such as phenol (Lallement, 2017).

While we only consider the direct interaction of bacteria and organics, additional processes might lead to more complex chemical and microbial interactions. For example, siderophores form iron complexes (Passananti et al., 2016) and, thus, suppress Fenton reactions that affect oxidant levels in cloud droplets (e.g., Deguillaume et al., 2004). Such indirect feedbacks of microbial processes on chemical budgets require more comprehensive data sets that are currently not available for models.

(iii) The respiration of bacteria depends on many different factors such as stress due harsh conditions. It can be expected that at higher stress levels (nutritional or thermal),  $F_{\text{CO}_2}$  increases to supply elevated energy needs (Amato and



Christner, 2009; Eiler et al., 2003). Values of BGE as low as  $< 0.4\%$  ( $F_{CO_2} = 0.996$ ) were observed (Eiler et al, 2003 and references therein), indicating that nearly all the carbon used was mineralized into  $CO_2$ .

405

**Table 4:** Values for parameters in Eq (6) and Eq (7) used in the estimate of WSOC loss by microbial and chemical processes and their likely minimum and maximum values

Parameter	Value in Eq (6) and Eq (7)	Range		Comment
		Minimum	Maximum	
$C_{Bact, aq}$ [cells $L^{-1}$ ]	$10^7$	$10^6$	$10^8$	Range of total bacteria concentration observed in cloud water samples collected from a mid-altitude mountain site over several years (31 samples) (Vaitilingom et al., 2012)
LWC [g $m^{-3}$ ]	0.15	0.1	1	The minimum and maximum value describe a range for a wide variety of cloud types. The assumption of LWC is not needed if it is assumed that all bacteria-containing particles act as CCN.
$k_{Bact}$ [ $L \text{ cell}^{-1} \text{ s}^{-1}$ ]	$10^{-13}$	$10^{-15}$	$10^{-13}$	$k_{Bact, min}$ was derived for microbial activity towards phenol (Lallement, 2017). $k_{Bact, max}$ was derived from experiments using cloud water and is valid for the microbial activity various highly oxygenated compounds.
$F_{CO_2}$	0.8 – 0.99	0.2	1	Even though BGE ranging from $< 0.4\%$ to 80% ( $0.996 > F_{CO_2} > 0.2$ ) were estimated in natural environments (Eiler et al. 2003 and references therein), at low nutrient concentrations, as encountered in clouds, high $F_{CO_2}$ can be expected.
$F_C$	0.5	0.2	$< 1$	Herckes et al. (2013) report that $\sim 20\%$ of total organic carbon in clouds is composed of speciated carboxylic acids. $F_C = 1$ seems unlikely due to variation in microbial and chemical cloud water composition.
$k_{OH}$ [ $L \text{ mol}^{-1} \text{ s}^{-1}$ ]	$3.8 \cdot 10^8$	$10^6$	$10^{10}$	Typically, undissociated acids (low pH) and polyfunctional compounds have $k_{OH}$ at the lower end of this range whereas the upper limit is constrained by diffusion limitation (Herrmann, 2003; Monod and Doussin, 2008)
$[OH]_{aq}$ [M]	$10^{-15}$	$10^{-17}$	$10^{-14}$	The suggested range includes concentrations that were inferred for night-time conditions (minimum) to day-time conditions in clean air masses (low OH sinks).
$Y_{volC}$	0.3 – 0.5	0.2	0.8	This value has not been comprehensively quantified yet; largest values can be likely expected in aged WSOC with high O/C ratios

(iv) The fraction of organic material metabolized by bacteria in clouds ( $F_C$ ) is likely not unity for a single bacteria type (Bianco et al., 2019; Vaitilingom et al., 2011). Carboxylic acids that are preferentially metabolized by several common bacteria types often comprise a major fraction ( $\sim 20\%$ ) of the cloud-water organics that can be speciated on a molecular





level (Herckes et al., 2013). This fraction might be regarded a lower limit of  $F_C$  since the reactivity of the large fraction of unspeciated organics (often ~70%) towards bacteria is not known. Our comparison implies the same spatial accessibility of bacteria and OH, respectively, to WSOC. This might be an oversimplification as bacteria are unevenly distributed among cloud drop populations as statistically only one in ~10,000 droplets may contain a single bacteria cell. OH can be expected to be present in all cloud droplets as the direct phase transfer from the gas phase represents one of the major OH sources in cloud water.

(v) While the absolute importance of microbial loss depends on the parameters discussed in (i) to (iv), the relative importance compared to chemical processes might be of interest in studies where the fate of individual organics in the cloud droplets or in the atmospheric multiphase system is explored.  $[\text{OH}]_{\text{aq}}$  depends mostly on photochemical processes as source processes and on the concentrations of WSOC as the main sinks; it ranges from  $10^{-17}$  M (night-time) to  $10^{-14}$  M (day time, clean air masses) (Arakaki et al., 2013).

(vi) Given that formate and acetate comprise major contributors to cloud water organics (Herckes et al., 2013), some fraction of WSOC will be converted into highly volatile products, such as  $\text{CO}_2$  and  $\text{CH}_3\text{CHO}$  that will not remain in the particle phase after cloud evaporation. However,  $Y_{\text{vol}C}$  likely does not approach unity since several studies have suggested that radical reactions in cloud water lead to the successive decay of dicarboxylic acids into their next smaller homologue which will remain in the aqueous phase. Within these limits, we conservatively suggest a range of 0.2 to 0.8 for  $Y_{\text{vol}C}$  but point out the need for studies to refine this parameter

#### 4. Summary and conclusions

We have estimated the amount of biological mass that is formed in the atmosphere by growth and multiplication of bacteria cells ('secondary biological aerosol', SBA). Data for representative bacteria strains and their generation rates have been compiled for major ecosystems. Using average values for cloudiness above the various ecosystems, we estimate that  $0.7 \text{ Tg yr}^{-1}$  SBA mass is formed globally which is comparable to current estimates of direct bacteria emissions ( $0.4 - 0.7 \text{ Tg yr}^{-1}$  (Burrows et al., 2009b; Myriokefalitakis et al., 2017)) which comprise a small fraction of total biological aerosol mass. While these production rates make up ~1% of formation rates of other major secondary aerosol components (secondary organics or sulfate), their importance might differ on spatial or temporal scales. In addition, SBA production leads to an increase in biological aerosol mass which might sensitively affect physicochemical particle properties (e.g. ice nucleation ability). SBA formation linearly depends on several parameters, such as the number concentration of metabolically active bacteria cells, their generation rates and the time scales during which they are assumed to grow or multiply – all of which are associated with considerable uncertainties. Our detailed discussion of the parameters and their uncertainties in our simplified approach highlight the likely variability of SBA formation on smaller scales and the need of future studies to refine these parameters. Similar approaches as ours may be applied to yeast growth. Yeast cells are generally larger (~2 – 10  $\mu\text{m}$ ) than bacteria cells (Fröhlich-Nowoisky et al., 2009) and, thus, their residence time in the atmosphere is likely shorter. Detailed data on their activity in clouds are not available which currently prevents the assessment of their potential contribution to SBA.

We also quantify the role of clouds as sinks of total WSOC by microbial and chemical processes, unlike previous studies that focused on microbial activity towards individual organic compounds. It is estimated that microbial



processes lead to an organic mass loss of 8 – 11 Tg yr<sup>-1</sup> whereas chemical processes by the OH radical in clouds lead to a loss of 8 – 20 Tg yr<sup>-1</sup>. These numbers are small compared to other sinks such as aerosol removal by deposition.

450 Not all of the organic mass even contributes to organic aerosol loading as water-soluble, volatile organics are dissolved in cloud water but evaporate during drop evaporation. Thus, the loss of organic aerosol mass due to direct microbial activity in clouds is smaller than the predicted loss of organic carbon. Large uncertainties in these estimates represent the assumptions on the fraction of carbon that is converted into volatile products. For bacteria, this fraction is quantified by the bacteria growth efficiency that depends on numerous factors, such as bacteria type, substrate availability and

455 physical conditions in the condensed phase.

In current atmospheric models, when considered, bacteria cells are inert, i.e. they neither change their mass or number concentrations during their residence time in the atmosphere nor do they interact with other aerosol constituents. The approach presented in our study is provides a first simplified estimate of SBA formation and WSOC loss due to bacteria that could be easily adapted in models. Given the current great activities in the field of atmospheric bioaerosols, it can

460 be expected that the discussed parameters in the estimates can and should be refined in the future in order to quantify the role of bacterial processes as source of biological mass and net source or sink of organic aerosol in the atmosphere.

**Data availability.** All data are available upon request from the authors.

**Author contributions.** BE and PA planned and carried out the study and wrote the manuscript together.

465 **Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgement.** This work has been funded by the French National Research Agency (ANR) in the framework of the ‘Investment for the Future’ program, ANR-17-MPGA-0013.

## References

- 470 Amato, P. and Christner, B. C.: Energy metabolism response to low-temperature and frozen conditions in *Psychrobacter cryohalolentis*, *Appl. Environ. Microbiol.*, 75(3), 711–718, doi:10.1128/AEM.02193-08, 2009.
- Amato, P., Demeer, F., Melahoui, A., Fontanella, S., Martin-Biesse, A.-S., Sancelme, M., Laj, P. and Delort, A.-M.: A fate for organic acids, formaldehyde and methanol in cloud water: their biotransformation by micro-organisms, *Atmos Chem Phys*, 7, 4159–4169, 2007a.
- 475 Amato, P., Parazols, M., Sancelme, M., Mailhot, G., Laj, P. and Delort, A.-M.: An important oceanic source of micro-organisms for cloud water at the Puy de Dôme (France), *Atmos. Environ.*, 41(37), 8253–8263, doi:10.1016/j.atmosenv.2007.06.022, 2007b.
- Amato, P., Parazols, M., Sancelme, M., Laj, P., Mailhot, G. and Delort, A.-M.: Microorganisms isolated from the water phase of tropospheric clouds at the Puy de Dôme: major groups and growth abilities at low temperatures, *FEMS Microbiol. Ecol.*, 59(2), 242–254, doi:10.1111/j.1574-6941.2006.00199.x, 2007c.
- 480 Amato, P., Doyle, S. and Christner, B. C.: Macromolecular synthesis by yeasts under frozen conditions, *Environ. Microbiol.*, 11(3), 589–596, doi:10.1111/j.1462-2920.2008.01829.x, 2009.
- Amato, P., Doyle, S. M., Battista, J. R. and Christner, B. C.: Implications of Subzero Metabolic Activity on Long-



- Term Microbial Survival in Terrestrial and Extraterrestrial Permafrost, *Astrobiology*, 10(8), 789–798, doi:10.1089/ast.2010.0477, 2010.
- 485 Amato, P., Joly, M., Schaupp, C., Attard, E., Möhler, O., Morris, C. E., Brunet, Y. and Delort, A.-M.: Survival and ice nucleation activity of bacteria as aerosols in a cloud simulation chamber, *Atmospheric Chem. Phys.*, 15(11), 6455–6465, doi:10.5194/acp-15-6455-2015, 2015.
- Amato, P., Joly, M., Besaury, L., Oudart, A., Taib, N., Moné, A. I., Deguillaume, L., Delort, A.-M. and Debroas, D.: Active microorganisms thrive among extremely diverse communities in cloud water, *PLOS ONE*, 12(8), e0182869, doi:10.1371/journal.pone.0182869, 2017.
- 490 Amato, P., Besaury, L., Joly, M., Penaud, B., Deguillaume, L. and Delort, A.-M.: Metatranscriptomic exploration of microbial functioning in clouds, *Sci. Rep.*, 9(1), 4383, doi:10.1038/s41598-019-41032-4, 2019.
- Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S., Kimura, T., Tshuhako, A. and Miyagi, Y.: A general scavenging rate constant for reaction of hydroxyl radical with organic carbon in atmospheric waters, *Env. Sci Technol*, 47(15), 8196–8203, doi:10.1021/es401927b, 2013.
- 495 Ariya, P. A., Nepotchaykh, O., Ignatova, O. and Amyot, M.: Microbiological degradation of atmospheric organic compounds, *Geophys. Res. Lett.*, 29(22), doi: 10.1029/2002GL015637, 2002.
- Ariya, P. A., Sun, J., Eltouny, N. A., Hudson, E. D., Hayes, C. T. and Kos, G.: Physical and chemical characterization of bioaerosols - Implications for nucleation processes, *Int. Rev. Phys. Chem.*, 28(1), 1–32, doi:10.1080/01442350802597438, 2009.
- 500 Artaxo, P., Maenhaut, W., Storms, H. and Van Grieken, R.: Aerosol characteristics and sources for the Amazon Basin during the wet season, *J. Geophys. Res. Atmospheres*, 95(D10), 16971–16985, doi:10.1029/JD095iD10p16971, 1990.
- Bakermans, C., Tsapin, A. I., Souza-Egipsy, V., Gilichinsky, D. A. and Neelson, K. H.: Reproduction and metabolism at -10°C of bacteria isolated from Siberian permafrost, *Environ. Microbiol.*, 5(4), 321–326, doi:10.1046/j.1462-2920.2003.00419.x, 2003.
- 505 Bauer, H., Giebl, H., Hitznerberger, R., Kasper-Giebl, A., Reischl, G., Zibuschka, F. and Puxbaum, H.: Airborne bacteria as cloud condensation nuclei, *J Geophys Res*, 108(D21), 4658, doi:10.1029/2003jd003545, 2003.
- Bianco, A., Passananti, M., Perroux, H., Vuyard, G., Mouchel-Vallon, C., Chaumerliac, N., Mailhot, G., Deguillaume, L. and Brigante, M.: A better understanding of hydroxyl radical photochemical sources in cloud waters collected at the puy de Dôme station – experimental versus modelled formation rates, *Atmos Chem Phys*, 15(16), 9191–9202, doi:10.5194/acp-15-9191-2015, 2015.
- 510 Bianco, A., Deguillaume, L., Chaumerliac, N., Vařtilingom, M., Wang, M., Delort, A.-M. and Bridoux, M. C.: Effect of endogenous microbiota on the molecular composition of cloud water: a study by Fourier-transform ion cyclotron resonance mass spectrometry (FT-ICR MS), *Sci. Rep.*, 9(1), 7663, doi:10.1038/s41598-019-44149-8, 2019.
- 515 Bovallius, A., Bucht, B., Roffey, R. and Anäs, P.: Three-year investigation of the natural airborne bacterial flora at four localities in Sweden., *Appl. Environ. Microbiol.*, 35(5), 847, 1978.
- Bowers, R. M., McCubbin, I. B., Hallar, A. G. and Fierer, N.: Seasonal variability in airborne bacterial communities at a high-elevation site, *Atmos. Environ.*, 50, 41–49, doi:10.1016/j.atmosenv.2012.01.005, 2012.
- 520 Burrows, S. M., Elbert, W., Lawrence, M. G. and Pöschl, U.: Bacteria in the global atmosphere, Part 1: Review and synthesis of literature data for different ecosystems, *Atmos Chem Phys*, 9(23), 9263–9280, doi:10.5194/acp-9-9263-2009, 2009a.
- Burrows, S. M., Butler, T., Jöckel, P., Tost, H., Kerkweg, A., Pöschl, U. and Lawrence, M. G.: Bacteria in the global



- atmosphere, Part 2: Modeling of emissions and transport between different ecosystems, *Atmos Chem Phys*, 9(23), 9281–9297, doi:10.5194/acp-9-9281-2009, 2009b.
- 525 Carotenuto, F., Georgiadis, T., Gioli, B., Leyronas, C., Morris, C. E., Nardino, M., Wohlfahrt, G. and Miglietta, F.: Measurements and modeling of surface–atmosphere exchange of microorganisms in Mediterranean grassland, *Atmos Chem Phys*, 17(24), 14919–14936, doi:10.5194/acp-17-14919-2017, 2017.
- Chin, M., Rood, R. B., Lin, S.-J., Müller, J.-F. and Thompson, A. M.: Atmospheric sulfur cycle simulated in the global model GOCART: Model description and global properties, *J Geophys Res - Atmos*, 105(D20), 24671–24687, doi:10.1029/2000jd900384, 2000.
- 530 Copeland, J. K., Yuan, L., Layeghifard, M., Wang, P. W. and Guttman, D. S.: Seasonal Community Succession of the Phyllosphere Microbiome, *Mol. Plant. Microbe Interact.*, 28(3), 274–285, doi:10.1094/MPMI-10-14-0331-FI, 2015.
- Davey, K. R.: A predictive model for combined temperature and water activity on microbial growth during the growth phase, *J. Appl. Bacteriol.*, 67(5), 483–488, doi:10.1111/j.1365-2672.1989.tb02519.x, 1989.
- 535 Deguillaume, L., Leriche, M., Monod, A. and Chaumerliac, N.: The role of transition metal ions on HO<sub>x</sub> radicals in clouds: a numerical evaluation of its impact on multiphase chemistry, *Atmos Chem Phys*, 4, 95–110, 2004.
- DeLeon-Rodriguez, N., Latham, T. L., Rodriguez-R, L. M., Barazesh, J. M., Anderson, B. E., Beyersdorf, A. J., Ziemba, L. D., Bergin, M., Nenes, A. and Konstantinidis, K. T.: Microbiome of the upper troposphere: Species composition and prevalence, effects of tropical storms, and atmospheric implications, *Proc. Natl. Acad. Sci.*, 110(7), 2575, doi:10.1073/pnas.1212089110, 2013.
- 540 Delort, A.-M., Vaïtilingom, M., Amato, P., Sancelme, M., Parazols, M., Mailhot, G., Laj, P. and Deguillaume, L.: A short overview of the microbial population in clouds: Potential roles in atmospheric chemistry and nucleation processes, *Atmos Res*, 98(2–4), 249–260, 2010.
- Després, V. R., Huffman, J. A., Burrows, S. M., Hoose, C., Safatov, A. S., Buryak, G., Fröhlich-Nowoisky, J., Elbert, W., Andreae, M. O., Pöschl, U. and Jaenicke, R.: Primary biological aerosol particles in the atmosphere: a review, *Tellus B*, 64, 15598, doi:10.3402/tellusb.v64i0.15598, 2012.
- 545 Dimmick, R. L., Wolochow, H. and Chatigny, M. A.: Evidence that bacteria can form new cells in airborne particles, *Appl. Environ. Microbiol.*, 37(5), 924–927, 1979.
- Eiler, A., Langenheder, S., Bertilsson, S. and Tranvik, L. J.: Heterotrophic bacterial growth efficiency and community structure at different natural organic carbon concentrations, *Appl. Environ. Microbiol.*, 69(7), 3701–3709, doi:10.1128/AEM.69.7.3701-3709.2003, 2003.
- 550 Elbert, W., Taylor, P. E., Andreae, M. O. and Pöschl, U.: Contribution of fungi to primary biogenic aerosols in the atmosphere: wet and dry discharged spores, carbohydrates, and inorganic ions, *Atmos Chem Phys*, 7(17), 4569–4588, doi:10.5194/acp-7-4569-2007, 2007.
- 555 Ervens, B., Feingold, G., Sulia, K. and Harrington, J.: The impact of microphysical parameters, ice nucleation mode, and habit growth on the ice/liquid partitioning in mixed-phase Arctic clouds, *J Geophys Res*, 116(D17), D17205, doi:10.1029/2011jd015729, 2011.
- 560 Ervens, B., Sorooshian, A., Lim, Y. B. and Turpin, B. J.: Key parameters controlling OH-initiated formation of secondary organic aerosol in the aqueous phase (aqSOA), *J Geophys Res - Atmos*, 119(7), 3997–4016, doi:10.1002/2013JD021021, 2014.
- Estillore, A. D., Trueblood, J. V. and Grassian, V. H.: Atmospheric chemistry of bioaerosols: heterogeneous and multiphase reactions with atmospheric oxidants and other trace gases, *Chem. Sci.*, 7(11), 6604–6616, doi:10.1039/c6sc02353c, 2016.



- 565 Fahlgren, C., Hagström, Å., Nilsson, D. and Zweifel, U. L.: Annual Variations in the Diversity, Viability, and Origin of Airborne Bacteria, *Appl. Environ. Microbiol.*, 76(9), 3015–3025, doi:10.1128/AEM.02092-09, 2010.
- Feingold, G., McComiskey, A., Rosenfeld, D. and Sorooshian, A.: On the relationship between cloud contact time and precipitation susceptibility to aerosol, *J. Geophys. Res. Atmospheres*, 118(18), 10,544–10,554, doi:10.1002/jgrd.50819, 2013.
- 570 Fröhlich-Nowoisky, J., Pickersgill, D. A., Després, V. R. and Pöschl, U.: High diversity of fungi in air particulate matter, *Proc. Natl. Acad. Sci.*, 106(31), 12814, doi:10.1073/pnas.0811003106, 2009.
- Fuchs, B. M., Zubkov, M. V., Sahm, K., Burkill, P. H. and Amann, R.: Changes in community composition during dilution cultures of marine bacterioplankton as assessed by flow cytometric and molecular biological techniques, *Environ. Microbiol.*, 2(2), 191–201, doi:10.1046/j.1462-2920.2000.00092.x, 2000.
- 575 Fuzzi, S., Mandrioli, P. and Perfetto, A.: Fog droplets—an atmospheric source of secondary biological aerosol particles, *Atmos. Environ.*, 31(2), 287–290, doi:10.1016/1352-2310(96)00160-4, 1997.
- Gandolfi, I., Bertolini, V., Ambrosini, R., Bestetti, G. and Franzetti, A.: Unravelling the bacterial diversity in the atmosphere, *Appl. Microbiol. Biotechnol.*, 97(11), 4727–4736, doi:10.1007/s00253-013-4901-2, 2013.
- Haddrell, A. E. and Thomas, R. J.: *Aerobiology: Experimental Considerations, Observations, and Future Tools*, *Appl. Environ. Microbiol.*, 83(17), e00809-17, doi:10.1128/AEM.00809-17, 2017.
- 580 He, C., Liu, J., Carlton, A. G., Fan, S., Horowitz, L. W., Levy II, H. and Tao, S.: Evaluation of factors controlling global secondary organic aerosol production from cloud processes, *Atmos Chem Phys*, 13(4), 1913–1926, doi:10.5194/acp-13-1913-2013, 2013.
- Heald, C. L. and Spracklen, D. V.: Atmospheric budget of primary biological aerosol particles from fungal spores, *Geophys. Res. Lett.*, 36(9), doi:10.1029/2009GL037493, 2009.
- 585 Helin, A., Sietiö, O.-M., Heinonsalo, J., Bäck, J., Riekkola, M.-L. and Parshintsev, J.: Characterization of free amino acids, bacteria and fungi in size-segregated atmospheric aerosols in boreal forest: seasonal patterns, abundances and size distributions, *Atmos Chem Phys*, 17(21), 13089–13101, doi:10.5194/acp-17-13089-2017, 2017.
- Herckes, P., Valsaraj, K. T. and Collett Jr, J. L.: A review of observations of organic matter in fogs and clouds: Origin, processing and fate, *Atmos Res*, 132–133, 434–449, doi:10.1016/j.atmosres.2013.06.005, 2013.
- 590 Herrmann, H.: Kinetics of aqueous phase reactions relevant for atmospheric chemistry, *Chem. Rev.*, 103(12), 4691–4716, 2003.
- Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C. D., Jimenez, J. L., Madronich, S. and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos Chem Phys*, 16(12), 7917–7941, doi:10.5194/acp-16-7917-2016, 2016.
- 595 Hu, W., Murata, K., Toyonaga, S. and Zhang, D.: Bacterial abundance and viability in rainwater associated with cyclones, stationary fronts and typhoons in southwestern Japan, *Atmos. Environ.*, 167, 104–115, doi:10.1016/j.atmosenv.2017.08.013, 2017.
- 600 Huffman, J. A., Prenni, A. J., DeMott, P. J., Pöhlker, C., Mason, R. H., Robinson, N. H., Fröhlich-Nowoisky, J., Tobo, Y., Després, V. R., Garcia, E., Gochis, D. J., Harris, E., Müller-Germann, I., Ruzene, C., Schmer, B., Sinha, B., Day, D. A., Andreae, M. O., Jimenez, J. L., Gallagher, M., Kreidenweis, S. M., Bertram, A. K. and Pöschl, U.: High concentrations of biological aerosol particles and ice nuclei during and after rain, *Atmos Chem Phys*, 13(13), 6151–6164, doi:10.5194/acp-13-6151-2013, 2013.
- Husárová, S., Vaitilingom, M., Deguillaume, L., Traikia, M., Vinatier, V., Sancelme, M., Amato, P., Matulová, M. and



- 605 Delort, A.-M.: Biotransformation of methanol and formaldehyde by bacteria isolated from clouds. Comparison with radical chemistry, *Atmos. Environ.*, 45(33), 6093–6102, doi:org/10.1016/j.atmosenv.2011.06.035, 2011.
- 610 IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2014.
- Jaenicke, R.: Abundance of Cellular Material and Proteins in the Atmosphere, *Science*, 308(5718), 73, doi:10.1126/science.1106335, 2005.
- 615 Joly, M., Amato, P., Sancelme, M., Vinatier, V., Abrantes, M., Deguillaume, L. and Delort, A.-M.: Survival of microbial isolates from clouds toward simulated atmospheric stress factors, *Atmos. Environ.*, 117, 92–98, doi:10.1016/j.atmosenv.2015.07.009, 2015.
- King, S. Platnick, W. P. Menzel, S. A. Ackerman and P. A. Hubanks: Spatial and Temporal Distribution of Clouds Observed by MODIS Onboard the Terra and Aqua Satellites, *IEEE Trans. Geosci. Remote Sens.*, 51(7), 3826–3852, doi:10.1109/TGRS.2012.2227333, 2013.
- 620 Klein, A. M., Bohannon, B. J. M., Jaffe, D. A., Levin, D. A. and Green, J. L.: Molecular Evidence for Metabolically Active Bacteria in the Atmosphere, *Front. Microbiol.*, 7, 772, doi:10.3389/fmicb.2016.00772, 2016.
- Lallement, A.: Impact des processus photochimiques et biologiques sur la composition chimique du nuage, PhD Thesis, Université Clermont Auvergne, Clermont-Ferrand (France), 2017.
- 625 Lallement, A., Besaury, L., Tixier, E., Sancelme, M., Amato, P., Vinatier, V., Canet, I., Polyakova, O. V., Artaev, V. B., Lebedev, A. T., Deguillaume, L., Mailhot, G. and Delort, A.-M.: Potential for phenol biodegradation in cloud waters, *Biogeosciences*, 15(18), 5733–5744, doi:10.5194/bg-15-5733-2018, 2018.
- Lelieveld, J. and Crutzen, P. J.: Influences of cloud photochemical processes on tropospheric ozone, *Nature*, 343, 227–233, 1990.
- Lighthart, B.: The ecology of bacteria in the alfresco atmosphere, *FEMS Microbiol. Ecol.*, 23(4), 263–274, 1997.
- 630 Lighthart, B. and Shaffer, B. T.: Bacterial flux from chaparral into the atmosphere in mid-summer at a high desert location, *Atmos. Environ.*, 28(7), 1267–1274, doi:10.1016/1352-2310(94)90273-9, 1994.
- Lighthart, B. and Shaffer, B. T.: Viable bacterial aerosol particle size distributions in the midsummer atmosphere at an isolated location in the high desert chaparral, *Aerobiologia*, 11(1), 19–25, 1995.
- Lin, G., Sillman, S., Penner, J. E. and Ito, A.: Global modeling of SOA: the use of different mechanisms for aqueous phase formation, *Atmos Chem Phys*, 14, 5451–5475, doi:10.5194/acp-14-5451-2014, 2014.
- 635 Lindemann, J., Constantinidou, H. A., Barchet, W. R. and Upper, C. D.: Plants as sources of airborne bacteria, including ice nucleation-active bacteria, *Appl. Environ. Microbiol.*, 44(5), 1059–1063, 1982.
- Lipson, D. A., Schadt, C. W. and Schmidt, S. K.: Changes in Soil Microbial Community Structure and Function in an Alpine Dry Meadow Following Spring Snow Melt, *Microb. Ecol.*, 43(3), 307–314, doi:10.1007/s00248-001-1057-x, 2002.
- 640 Liu, J., Horowitz, L. W., Fan, S., Carlton, A. G. and Levy, H.: Global in-cloud production of secondary organic aerosols: Implementation of a detailed chemical mechanism in the GFDL atmospheric model AM3, *J Geophys Res - Atmos*, 117(D15), D15303, doi:10.1029/2012jd017838, 2012.



- Madronich, S., Bjorn, L. O. and McKenzie, R. L.: Solar UV radiation and microbial life in the atmosphere, *Photochem. Photobiol. Sci.*, 17(12), 1918–1931, doi:10.1039/C7PP00407A, 2018.
- 645 Marinoni, A., Parazols, M., Brigante, M., Deguillaume, L., Amato, P., Delort, A.-M., Laj, P. and Mailhot, G.: Hydrogen peroxide in natural cloud water: Sources and photoreactivity, *Atmos Res*, 101(1–2), 256–263, doi:10.1016/j.atmosres.2011.02.013, 2011.
- Marr, A. G.: Growth rate of *Escherichia coli*, *Microbiol. Rev.*, 55(2), 316–333, 1991.
- 650 Matthias-Maser, S. and Jaenicke, R.: The size distribution of primary biological aerosol particles in the multiphase atmosphere, *Aerobiologia*, 16(2), 207–210, doi:10.1023/A:1007607614544, 2000.
- Meisner, A., Leizeaga, A., Rousk, J. and Bååth, E.: Partial drying accelerates bacterial growth recovery to rewetting, *Soil Biol. Biochem.*, 112, 269–276, doi:10.1016/j.soilbio.2017.05.016, 2017.
- 655 Michaud, J. M., Thompson, L. R., Kaul, D., Espinoza, J. L., Richter, R. A., Xu, Z. Z., Lee, C., Pham, K. M., Beall, C. M., Malfatti, F., Azam, F., Knight, R., Burkart, M. D., Dupont, C. L. and Prather, K. A.: Taxon-specific aerosolization of bacteria and viruses in an experimental ocean-atmosphere mesocosm, *Nat. Commun.*, 9(1), 2017, doi:10.1038/s41467-018-04409-z, 2018.
- Middelboe, M.: Bacterial Growth Rate and Marine Virus–Host Dynamics, *Microb Ecol*, 114–124, 2000.
- 660 Möhler, O., Georgakopoulos, D. G., Morris, C. E., Benz, S., Ebert, V., Hunsmann, S., Saathoff, H., Schnaiter, M. and Wagner, R.: Heterogeneous ice nucleation activity of bacteria: new laboratory experiments at simulated cloud conditions, *Biogeosciences*, 5(5), 1425–1435, doi:10.5194/bg-5-1425-2008, 2008.
- Monod, A. and Doussin, J. F.: Structure-activity relationship for the estimation of OH-oxidation rate constants of aliphatic organic compounds in the aqueous phase: alkanes, alcohols, organic acids and bases, *Atmos. Environ.*, 42(33), 7611–7622, doi:DOI: 10.1016/j.atmosenv.2008.06.005, 2008.
- 665 Monteil, C. L., Bardin, M. and Morris, C. E.: Features of air masses associated with the deposition of *Pseudomonas syringae* and *Botrytis cinerea* by rain and snowfall, *ISME J.*, 8(11), 2290–2304, doi:10.1038/ismej.2014.55, 2014.
- Morris, C. E., Glaux, C., Latour, X., Gardan, L., Samson, R. and Pitrat, M.: The Relationship of Host Range, Physiology, and Genotype to Virulence on Cantaloupe in *Pseudomonas syringae* from Cantaloupe Blight Epidemics in France, *Phytopathology*, 90(6), 636–646, doi:10.1094/PHYTO.2000.90.6.636, 2000.
- 670 Morris, C. E., D. G. Georgakopoulos and D. C. Sands: Ice nucleation active bacteria and their potential role in precipitation, *J Phys IV Fr.*, 121, 87–103, 2004.
- Myriokefalitakis, S., Fanourgakis, G. and Kanakidou, M.: The Contribution of Bioaerosols to the Organic Carbon Budget of the Atmosphere, in *Perspectives on Atmospheric Sciences*, edited by T. Karacostas, A. Bais, and P. T. Nastos, pp. 845–851, Springer International Publishing., 2017.
- Newman, I. V.: Aerobiology on commercial air routes, *Nature*, 161(4086), 275, 1948.
- 675 Olson, J.: World ecosystems (WE1.4): Digital raster data on a 10 minute geographic 1080 (2160 grid square), *Global Ecosystem Database*, Version 1, 1992.
- Otero Fernandez, M., Thomas, R., Garton, N., Hudson, A., Haddrell, A. and Reid, J.: Assessing the Airborne Survival of Bacteria in Populations of Aerosol Droplets with a Novel Technology, *J. R. Soc. Interface*, 16(150), doi:10.1098/rsif.2018.0779, 2019.
- 680 Paez-Rubio, T., Viau, E., Romero-Hernandez, S. and Peccia, J.: Source Bioaerosol Concentration and rRNA Gene-Based Identification of Microorganisms Aerosolized at a Flood Irrigation Wastewater Reuse Site, *Appl. Environ.*



- Microbiol., 71(2), 804, doi:10.1128/AEM.71.2.804-810.2005, 2005.
- 685 Passananti, M., Vinatier, V., Delort, A.-M., Mailhot, G. and Brigante, M.: Siderophores in Cloud Waters and Potential Impact on Atmospheric Chemistry: Photoreactivity of Iron Complexes under Sun-Simulated Conditions, *Environ. Sci. Technol.*, 50(17), 9324–9332, doi:10.1021/acs.est.6b02338, 2016.
- Perrino, C. and Marcovecchio, F.: A new method for assessing the contribution of Primary Biological Atmospheric Particles to the mass concentration of the atmospheric aerosol, *Environ. Int.*, 87, 108–115, doi:10.1016/j.envint.2015.11.015, 2016.
- 690 Posfai, M., Xu, H., Anderson, J. R. and Buseck, P. R.: Wet and dry sizes of atmospheric particles: An AFM-TEM study, *Geophys. Res. Lett.*, 25(11), 1907–1910, 1998.
- Price, P. B. and Sowers, T.: Temperature dependence of metabolic rates for microbial growth, maintenance, and survival, *Proc. Natl. Acad. Sci. U. S. A.*, 101(13), 4631–4636, doi:10.1073/pnas.0400522101, 2004.
- Pruppacher, H. R. and Jaenicke, R.: The processing of water vapor and aerosols by atmospheric clouds, a global estimate, *Atmos Res.*, 38(1–4), 283–295, doi:http://dx.doi.org/10.1016/0169-8095(94)00098-X, 1995.
- 695 Pruppacher, H. R. and Klett, J. D.: *Microphysics of clouds and precipitation*, 2nd revised and enlarged., Kluwer Academics Publisher, Dordrecht (NL), 2003.
- Sand-Jensen, K., Pedersen, N. L. and Sandergaard, M.: Bacterial metabolism in small temperate streams under contemporary and future climates, *Freshw. Biol.*, 52(12), 2340–2353, doi:10.1111/j.1365-2427.2007.01852.x, 2007.
- 700 Šantl-Temkiv, T., Finster, K., Hansen, B. M., Pašić, L. and Karlson, U. G.: Viable methanotrophic bacteria enriched from air and rain can oxidize methane at cloud-like conditions, *Aerobiologia*, 1–12, 2013.
- Santl-Temkiv, T., Sahyoun, M., Finster, K., Hartmann, S., Augustin-Bauditz, S., Stratmann, F., Wex, H., Clauss, T., Nielsen, N. W., Sørensen, J. H., Korsholm, U. S., Wick, L. Y. and Karlson, U. G.: Characterization of airborne ice-nucleation-active bacteria and bacterial fragments, *Atmos. Environ.*, 109, 105–117, doi:10.1016/j.atmosenv.2015.02.060, 2015.
- 705 Sattler, B., Puxbaum, H. and Psenner, B.: Bacterial growth in supercooled cloud droplets, *Geophys. Res. Lett.*, 28(2), 239–242, 2001.
- Si, F., Li, D., Cox, S. E., Sauls, J. T., Azizi, O., Sou, C., Schwartz, A. B., Erickstad, M. J., Jun, Y., Li, X. and Jun, S.: Invariance of Initiation Mass and Predictability of Cell Size in *Escherichia coli*, *Curr. Biol.*, 27(9), 1278–1287, doi:10.1016/j.cub.2017.03.022, 2017.
- 710 Smith, D. J., Griffin, D. W., McPeters, R. D., Ward, P. D. and Schuerger, A. C.: Microbial survival in the stratosphere and implications for global dispersal, *Aerobiologia*, 27(4), 319–332, doi:10.1007/s10453-011-9203-5, 2011.
- 715 Smith, D. J., Ravichandar, J. D., Jain, S., Griffin, D. W., Yu, H., Tan, Q., Thissen, J., Lusby, T., Nicoll, P., Shedler, S., Martinez, P., Osorio, A., Lechniak, J., Choi, S., Sabino, K., Iverson, K., Chan, L., Jaing, C. and McGrath, J.: Airborne Bacteria in Earth's Lower Stratosphere Resemble Taxa Detected in the Troposphere: Results From a New NASA Aircraft Bioaerosol Collector (ABC), *Front. Microbiol.*, 9, 1752, doi:10.3389/fmicb.2018.01752, 2018.
- Stibal, M. and Elster, J.: Growth and morphology variation as a response to changing environmental factors in two Arctic species of *Raphidonema* (Trebouxiophyceae) from snow and soil, *Polar Biol.*, 28(7), 558–567, doi:10.1007/s00300-004-0709-y, 2005.
- 720 Tesson, S. V. M. and Santl-Temkiv, T.: Ice Nucleation Activity and Aeolian Dispersal Success in Airborne and Aquatic Microalgae, *Front. Microbiol.*, 9, 2681, doi:10.3389/fmicb.2018.02681, 2018.





- Vaïtilingom, M., Amato, P., Sancelme, M., Laj, P., Leriche, M. and Delort, A.-M.: Contribution of Microbial Activity to Carbon Chemistry in Clouds, *Appl. Environ. Microbiol.*, 76(1), 23–29, doi:10.1128/AEM.01127-09, 2010.
- Vaïtilingom, M., Charbouillot, T., Deguillaume, L., Maisonobe, R., Parazols, M., Amato, P., Sancelme, M. and Delort, A. M.: Atmospheric chemistry of carboxylic acids: microbial implication versus photochemistry, *Atmos Chem Phys*, 11(16), 8721–8733, doi:10.5194/acp-11-8721-2011, 2011.
- 725
- Vaïtilingom, M., Attard, E., Gaiani, N., Sancelme, M., Deguillaume, L., Flossmann, A. I., Amato, P. and Delort, A.-M.: Long-term features of cloud microbiology at the puy de Dôme (France), *Atmos. Environ.*, 56(0), 88–100, doi:http://dx.doi.org/10.1016/j.atmosenv.2012.03.072, 2012.
- Vaïtilingom, M., Deguillaume, L., Vinatier, V., Sancelme, M., Amato, P., Chaumerliac, N. and Delort, A.-M.: Potential impact of microbial activity on the oxidant capacity and organic carbon budget in clouds, *Proc. Natl. Acad. Sci.*, 110(2), 559–564, doi:10.1073/pnas.1205743110, 2013.
- 730
- Wang, J., Rossow, W. B. and Zhang, Y.: Cloud Vertical Structure and Its Variations from a 20-Yr Global Rawinsonde Dataset, *J. Clim.*, 13(17), 3041–3056, doi:10.1175/1520-0442(2000)013<3041:CVSAIV>2.0.CO;2, 2000.
- Wéry, N., Gales, A. and Brunet, Y.: Bioaerosol sources, in *Microbiology of Aerosols*, pp. 115–135, A.M. Delort and P. Amato, Hoboken, NJ: John Wiley & Sons, Inc., 2017.
- 735
- Whitman, W. B., Coleman, D. C. and Wiebe, W. J.: Prokaryotes: the unseen majority, *Proc. Natl. Acad. Sci. U. S. A.*, 95(12), 6578–6583, doi:10.1073/pnas.95.12.6578, 1998.
- Wiedinmyer, C., Bowers, R. M., Fierer, N., Horanyi, E., Hannigan, M., Hallar, A. G., McCubbin, I. and Baustian, K.: The contribution of biological particles to observed particulate organic carbon at a remote high altitude site, *Atmos. Environ.*, 43(28), 4278–4282, doi:10.1016/j.atmosenv.2009.06.012, 2009.
- 740
- Wirgot, N., Vinatier, V., Deguillaume, L., Sancelme, M. and Delort, A.-M.: H<sub>2</sub>O<sub>2</sub> modulates the energetic metabolism of the cloud microbiome, *Atmos Chem Phys*, 17(24), 14841–14851, doi:10.5194/acp-17-14841-2017, 2017.
- Xiang, S.-R., Doyle, A., Holden, P. A. and Schimel, J. P.: Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils, *Soil Biol. Biochem.*, 40(9), 2281–2289, doi:10.1016/j.soilbio.2008.05.004, 2008.
- 745
- Xie, Z., Li, Y., Lu, R., Li, W., Fan, C., Liu, P., Wang, J. and Wang, W.: Characteristics of total airborne microbes at various air quality levels, *J. Aerosol Sci.*, 116, 57–65, doi:10.1016/j.jaerosci.2017.11.001, 2018.