

1 Occurrence and source apportionment of perfluoroalkyl acids 2 (PFAAs) in the atmosphere in China

3 Deming Han¹, Yingge Ma², Cheng Huang², Xufeng Zhang¹, Hao Xu¹, Yong Zhou¹, Shan Liang¹,
4 Xiaojia Chen¹, Xiqian Huang¹, Haoxiang Liao¹, Shuang Fu¹, Xue Hu¹, Jinping Cheng¹

5 ¹ School of Environmental Science and Engineering, Shanghai Jiao Tong University, Shanghai 200240, China

6 ² State Environmental Protection Key Laboratory of the Formation and Prevention of Urban Air Pollution Complex,
7 Shanghai Academy of Environmental Sciences, Shanghai 200233, China

8 *Correspondence to:* Jinping Cheng (jpcheng@sjtu.edu.cn)

9 **Abstract:**

10 Perfluoroalkyl acids (PFAAs) are a form of toxic pollutant that can be transported across the globe and accumulated in
11 the bodies of wildlife and humans. A nationwide geographical investigation considering atmospheric PFAAs [via](#)
12 [XAD–Passive Air Sampler](#) was conducted [in 23 different provinces/municipalities/autonomous regions](#) in China, which
13 provides an excellent chance to investigate [their occurrences](#), spatial trends, and potential sources. The total [atmospheric](#)
14 concentrations of thirteen PFAAs ([n=268](#)) were 6.19–292.57 pg/m³, with an average value of 39.84±28.08 pg/m³, which
15 were higher than other urban levels but lower than point source measurements. Perfluorooctanoic acid (PFOA) was the
16 dominant PFAAs (20.6%), followed by [perfluorohexanoic acid](#) (PFHxA), perfluorooctane sulfonate (PFOS), and
17 [perfluoroheptanoic acid](#) (PFPeA). An increasing seasonal trend of PFAAs concentrations was shown as summer <
18 autumn < spring < winter, which may be initiated by stagnant meteorological conditions. Spatially, the content of PFAAs
19 displayed a declining gradient trend of [central China> northern China> eastern China> northeast of China> southwest of](#)
20 [China> northwest of China> southern China areas](#), and Henan contributed as the largest proportion of PFAAs. Four
21 sources of PFAAs were identified using a positive matrix factorization (PMF) model, including PFOS–based products
22 (26.1%), PFOA–based, and PFNA–based products (36.6%), degradation products of [fluorotelomere–based](#) products
23 (15.5%), and an unknown source (21.8%).

25 **1.Introduction**

26 Perfluoroalkyl acids (PFAAs) are [one](#) class of ionic polyfluoroalkyl substances (PFASs), which have excellent
27 characteristics in terms of chemical and thermal stability, high surface activity, and water and oil repulsion (Lindstrom et
28 al., 2011; Wang et al., 2014). They are applied to a wide variety of domestic and industrial products such as textiles, oil

29 and liquid repellents, firefighting foam, pesticides, and food packaging materials (Xie et al., 2013;Wang et al., 2014).
30 PFAAs can be released to the surrounding environment during manufacturing and use of PFAAs containing products,
31 which are ubiquitous in the environment (e.g., in the atmosphere, water, or snow) [\(Dreyer et al., 2009;Wang et al.,](#)
32 [2017;Hu et al., 2016\)](#), in wildlife [\(Sedlak et al., 2017\)](#), and even in the human body [\(Cardenas et al., 2017;Tian et al.,](#)
33 [2018\)](#). PFAAs can change adult thyroid hormone levels, reduce newborn birth weight, and [biomagnify up](#) the food chain,
34 which can be extremely toxic to animals and humans (Hu et al., 2016;Jian et al., 2017;Baard Ingegerdsson et al., 2010).
35 Of the PFAAs, the long-chain [\(C >8\)](#) perfluoroalkyl carboxylic acids (PFCAs) and [\(C >7\)](#) perfluoroalkyl sulfonic acids
36 (PFSAAs) are more toxic and bio-accumulative than their short-chain analogues [\(Buck et al., 2011\)](#). This especially
37 applies to perfluorooctanoic acid (PFOA), [perfluorooctane sulfonate \(PFOS\) and perfluorohexane sulfonate \(PFHxS\), in](#)
38 [which PFOS and PFOA have been added to Annex B and Annex A of the Stockholm Convention in 2009 and 2019,](#)
39 [respectively, while PFHxS was under review by the Persistent Organic Pollutants Review Committee \(Johansson et al.,](#)
40 [2008; UNEP Stockholm Convention, 2019\)](#).

41 PFAAs can originate from direct sources of products' emissions as well as indirect sources of incomplete degradation of
42 their precursors. It is estimated that the global historical emission quantities of C4–C14 PFCAs were 2610–21400 t in the
43 period of 1951–2015, of which PFOA-based and perfluorononanoic-acid (PFNA)-based products contributed the most
44 (Wang et al., 2014). A trend of geographical distribution of major fluorochemical manufacturing sites has shifted from
45 Western Europe, US, and Japan to the emerging economies in the Asia Pacific area over the past decades. This is
46 especially true for China, which was the world's largest industrial contributor of PFOAs (50–80 t) and PFOS-related
47 compounds (~1800 t) in 2009 (Xie et al., 2013). PFOA- and PFOS- based products were added to the Catalogue for the
48 Guidance of Industrial Structure Adjustment in China in 2011, and restricted elimination of PFOA/PFOS substances
49 production were conducted. With a large quantity of PFAAs and their products manufacturing and consumption, China
50 has become the emerging contamination hotspots in the world. In spite of several studies on atmospheric PFAAs levels
51 having been conducted in a few cities (Liu et al., 2015) and point sources (Yao et al., 2016a;Tian et al., 2018) in China,
52 due to the imbalanced urbanization and industrialization levels, there is still a lack of systemic research on atmospheric
53 PFAAs quantification and trends in China.

54 Additionally, the long range or mesoscale transport was also suggested to have a contribution to PFAAs in the air (Dreyer
55 et al., 2009;Cai et al., 2012a). In general, three pathways/hypotheses for the transportation of PFAAs were suggested:
56 transport associated with particles, degradation from precursor, and sea salts from current bursting in coastal areas. The
57 PFAAs precursors such as fluorotelomere alcohols (FTOHs), which can form the corresponding PFAAs through
58 oxidation reactions initiated by hydroxyl radicals (OH·) in the atmosphere (Thackray and Selin, 2017), are more volatile

59 than PFAAs and can reach remote areas via long-range transportation (Martin et al., 2006; Wang et al., 2018). Due to the
60 lower acid dissociation coefficient (pK_A), 0–3.8 for PFCAs and –3.3 for PFSA, PFAAs are expected to be mainly
61 associated with aerosols in the non-volatile anionic form (Lai et al., 2018; [Karásková et al., 2018](#)). However, recent field
62 studies have confirmed their occurrence in gaseous phase (Cassandra et al., 2018; Ahrens et al., 2013), e.g. [Fang et al.,
63 \(2018\) found the total concentrations of C2, C4–C10 PFCAs and C6 and C8 PFSA in the gas phase were 0.076–4.0
64 \$\text{pg}/\text{m}^3\$ in the air above the Bohai and Yellow Seas, China](#). Investigating the transport pathways of PFAAs in nationwide
65 region via active air sampler (AAS) is challenging, due to their electronic power supply and high cost. Fortunately, a
66 number of reports showed that the XAD (a styrene–divinylbenzene copolymer) impregnated sorbent based passive air
67 sampler (SIP–PAS) and XAD based PAS (XAD–PAS), were proven to be an ideal alternative sampling tool for
68 monitoring PFAAs in a wide region. [Despite several publications suggested XAD-PAS collects primarily gaseous PFAAs
69 \(Melymuk et al., 2014; Lai et al., 2018\) in the ambient, current findings were not consistent. Due to the unimpeded
70 movements of particles into the sampler, XAD–PAS was indicated to collect a representative sample of both gas and
71 particle phases \(Ahrens et al., 2013; Okeme et al., 2016; Karásková et al., 2018\). Moreover, the dominant sorbent for
72 fluorinated compounds was reported as XAD resin in the XAD impregnated SIP–PAS, instead of PUF themselves
73 \(Krogseth et al., 2013\)](#). XAD–PAS give PFASs profiles that were more closely resembled to those from AAS in
74 comparing with PUF–PAS, have sufficient uptake rates for the PFCAs and PFSA to be [deployed](#) for short time duration
75 (Lai et al., 2018).

76 Given the factors mentioned above, we conducted a nationwide survey of PFAAs in China at a provincial level using a
77 XAD–PAS from January to December in 2017. The objective of this research was: (1) to examine the tempo–spatial
78 variations of PFAAs, and (2) to identify their potential affecting factors and evaluate the affecting pathways. To the best
79 of our knowledge, this is the first research paper analyzing both a long-term and nationwide atmospheric PFAAs data set
80 complemented by a comprehensive investigation in China.

81 **2. Material and methods**

82 **2.1 Chemicals and reagents**

83 The PFAAs standards used were Wellington Laboratories (Guelph, ON, Canada) PFAC–MXB standard materials,
84 including C5–C14 PFCAs analogues ([Perfluoropentanoic acid \(PFPeA\), Perfluorohexanoic acid \(PFHxA\),
85 Perfluoroheptanoic acid \(PFHpA\), PFOA, Perfluorononanoic acid \(PFNA\), Perfluorodecanoic acid \(PFDA\),
86 Perfluoroundecanoic acid \(PFUdA\), Perfluorododecanoic acid \(PFDoA\), Perfluorotridecanoic acid \(PFTrDA\), and](#)

87 [Perfluorotetradecanoic acid \(PFTeDA\)](#)), as well as C4, C6, and C8 PFSA analogues ([Perfluorobutane sulfonic acid](#)
88 [\(PFBS\)](#), PFHxS, and PFOS). The mass-labeled 1,2-¹³C₂-PFHxA, 1,2,3,4-¹³C₄-PFOA, 1,2,3,4,5-¹³C₅-PFNA,
89 1,2-¹³C₂-PFDA, 1,2-¹³C₂-PFUDA, 1,2-¹³C₂-PFDaA, ¹⁸O₂-PFHxS, and 1,2,3,4-¹³C₄-PFOS were used as internal
90 standards (ISs, MPFAC-MXA, Wellington Laboratories Inc.) in high-performance liquid chromatography (HPLC)
91 coupled with a tandem mass spectrometer (MS/MS). HPLC-grade reagents that were used include methanol, ethyl
92 acetate, ammonia acetate, acetone, methylene dichloride, n-hexane, and Milli-Q water. Detailed sources of the target
93 PFAAs and their ISs are listed in Table S1 in the Supplementary Materials.

94 **2.2 Sample collection**

95 Sampling campaigns were carried out at 23 different provinces/municipalities/autonomous regions in China
96 simultaneously from January to December 2017, of which 20 were urban sites and three were rural sites (Zhejiang,
97 Shanxi, and Liaoning). Urban samples typically came from urban residential areas, and the rural samples were obtained
98 from villages. These sampling sites were divided into seven administrative divisions: [norther China \(NC, n=3 sites\)](#),
99 [southern China \(SC, n=2\)](#), [central China \(CC, n=3\)](#), [eastern China \(EC, n=7\)](#), [northwest of China \(NW, n=3\)](#), [northeast of](#)
100 [China \(NE, n=2\)](#), and [southwest of China \(SW, n=3\)](#). A geographical map of the sampling sites is displayed in Figure S1,
101 and the detailed information on sampling sites such as elevation, meteorological parameters, local resident population
102 and gross domestic product were listed in Table S2 [and Figure S1](#).

103 Samples were collected with Amberlite XAD-2 resin using XAD-PAS, which have been successfully monitored PFCAs
104 (C4-C16) and PFSAs (C4-C10) in the atmosphere (Krogseth et al., 2013; Armitage et al., 2013). Briefly, the mesh
105 cylinder (L× I.D.: 10 cm × 2 cm) was prebaked at 450°C for 3 h, filled with ~10 g XAD-2 resin, and capped with an
106 aluminum cap. [The particle size of XAD-2 is ~20-60 mesh, with water content of 20%-45%, its specific surface area](#)
107 [≥430 m²/g, and the reference adsorption capacity ≥35 mg/g. We should keep in mind that the unimpeded movement of](#)
108 [particle bound PFAAs would be captured during sampling using XAD-PAS, which cannot differentiate PFAAs between](#)
109 [gas and particle phases. Despite some research suggest the sampling efficiency of gas and particle phase PFAAs were](#)
110 [similar \(Karásková et al., 2018\). In the present study, the two phases PFAAs sampled by XAD-PAS were treated as the](#)
111 [whole atmosphere PFAAs concentration](#). The sampling program for each sample lasted approximately a month (30 days),
112 and the error of the sampling time was controlled within 3 d. At the end of each deployment period, the atmosphere
113 samples were retrieved, resealed in their original solvent-cleaned aluminum tins at the sampling location, and transported
114 by express post to Shanghai Jiao Tong University. On receipt, they were stored and frozen ([-20 °C](#)) until extraction.
115 The sampling rate of XAD-PAS is a crucial factor to derive the chemical concentrations accumulated in the XAD resin.

116 Ahrens et al. (2013) found that sampling rate of PFCAs and PFASs ranged 1.80–5.50 m³/d with XAD impregnated
117 sorbent, and the sampling rate increased as the carbon chain adding, while Karásková et al. (2018) suggested that the
118 sampling rate of XAD–PAS of 0.21–15.00 m³/d for PFAAs. The loss of depuration compounds could be used to calculate
119 the sampling rate, assessing the impacts from meteorological factors like temperature and wind speed. According to
120 Ahrens et al. (2013) the 1,2,3,4-¹³C₄–PFOA was used to calculate the sampling rates of PFAAs at Shanghai sampling site
121 (Shanghai Jiao Tong University) in the present study, by assessing 1,2,3,4-¹³C₄–PFOA abundance loss. The specific
122 description of the sampling rate calculation in this study is shown in Section S1 in the Supplementary Materials.

123 **2.3 Sample preparation and instrument analysis**

124 The sample preparation and analysis were according to the method described by previous researches (Liu et al.,
125 2015; Tian et al., 2018). The MPFAC–MXA ISs mixture surrogates (10 ng) were added to each spiked sample prior to
126 extraction. This was done to account for the loss of substances from the samples associated with instrument instability
127 caused by the changes in laboratory environmental conditions. The XAD resin samples were Soxhlet–extracted for 24 h
128 using a Soxhlet extraction system, with n–hexane: acetone (1:1, V:V) as a solvent in a 300 mL polypropylene (PP) bottle,
129 following extracted with methanol for 4 h. These two extracts were combined and reduced to ~5 mL via a rotary
130 evaporator (RE–52AA, Yarong Biochemical Instrument Inc., Shanghai, China) at a temperature below 35 °C, and then
131 transferred to a 10 mL PP tube for centrifugation (10 min, 8,000 rpm). The supernatant was transferred to another PP tube,
132 filtered three times through a 0.22 µm nylon filter, with an addition of 1 mL methanol each time. The extracts were
133 further condensed under a gentle stream of nitrogen (99.999%, Shanghai Liquid Gas Cor.) at [35 °C](#) to a final 200 µL for
134 instrument analysis.

135 The separation and detection of PFAAs were performed using a HPLC system (Thermo Ultra 3000⁺, Thermo Scientific,
136 USA) coupled with a triple quadrupole negative electrospray ionization MS/MS (Thermo API 3000, Thermo Scientific,
137 USA). An Agilent Eclipse XDB C18 (3.5 µm, 2.1 mm, 150 mm) was used to separate the desorbed substances. The
138 column temperature was set to [40 °C](#), and the flow rate was 0.30 mL/min. The injection volume was 20 µL. The gradient
139 elution program of the mobile phase [A \(5 mmol/L aqueous ammonium acetate\)](#) and [B \(methanol\)](#) was [80% A + 20% B](#) at
140 the start, [5% A + 95% B](#) at 8 min, 100% a at 13 min, [80% A + 20% B](#) at 14 min, and was maintained for 6 min. The
141 MS/MS was operated in a negative ion scan and multiple reaction monitoring (MRM) mode, and the electrospray voltage
142 was set to 4500 V. The ion source temperature was [450 °C](#). The flow rates of the atomization gas and air curtain gas was
143 10 and 9 L/min, respectively. Species identification was achieved by comparing the mass spectra and retention time of
144 the chromatographic peaks with the corresponding authentic standards.

145 2.4 Quality assurance and quality control

146 To avoid exogenous contamination, the XAD-2 resin was precleaned using a Soxhlet extraction system with acetone and
147 petroleum ether at extraction times of 24 h and 4 h, respectively. The extracted XAD resin was dried under a vacuum
148 desiccator, wrapped in an aluminum foil and zip-lock bags, and stored at -20 °C to avoid contamination. All laboratory
149 vessels were PP, and these vessels were washed with ultrapure water and methanol three times, respectively.

150 For quantification, six-point calibration curves of PFAAs were constructed by adopting different calibration solutions
151 with values of 1, 3, 6, 15, 30, and 60 ng/mL. The same concentration for the internal calibration (10 ng/mL) was used for
152 each level of the calibration solution. Recovery standards were added to each of the samples to monitor procedural
153 performance, and the mean spiked PFAAs recoveries ranged from 81%±25% to 108%±22%. All the analyzed PFAAs
154 were normalized against the recovery of the corresponding mass-labeled ISs. Field blanks were prepared at all sampling
155 sites, transported, and analyzed in the same way as the samples. Laboratory blanks were obtained by taking amounts of
156 solvent via extraction, cleanup, and analysis. A total of 8 field blanks and 26 laboratory blanks were analyzed, with
157 individual blank values of BDL (below detection limit)-1.12 pg/m³ and BDL-1.29 pg/m³, respectively. All the results
158 were corrected according to the blank and recovery results. The method detection limit (MDL) was derived from three
159 times standard deviation of the field blank values. The limit of detection (LOD) and the limit of quantification (LOQ)
160 were determined as a signal-to-noise ratio of 3:1 and 10:1, respectively (Rauert et al., 2018;Liu et al., 2015). To convert
161 MDLs, LODs and LOQs values to pg/m³, the mean volume of sampling air (m³) was applied. For the analytes that were
162 not detected or were below the LOQs in field blanks, MDLs were derived directly from three times the corresponding
163 LODs. More detailed information on the individual compounds of PFASs on MDL, LOD, LOQ, the recovery values, and
164 blank values are listed in Table S3.

165 2.5 Statistical and geostatistical analysis

166 Statistical analyses were carried out by SPSS Statistics 22 (IBM Inc. US), and the values of 1/2 MDL were used to
167 replace these measured results of BDL. The statistics figures were depicted using technical software of SigmaPlot 14.0
168 (Systat Software, US). And the geographical variations of atmospheric PFAAs were analyzed with ArcGIS 10.4 (ESRI,
169 US). The Hybrid Single-Particle Lagrangian Integrated Trajectory (Hysplit) back trajectory model (NOAA, US) was used
170 to study the long range transport of air masses in the sampling locations (Zhen et al., 2014). Positive matrix factorization
171 (PMF) is considered an advanced algorithm among various receptor models, which has been successfully applied for
172 source identification of environmental pollutants (Han et al., 2018;Han et al., 2019). PMF (5.0, US EPA) was adopted to
173 cluster the PFAAs with similar behaviors to identify potential sources, and a more detailed description of PMF can be

174 seen in Section S2.

175 3.Results and discussion

176 3.1 Abundances and compositions

177 The descriptive statistics of all targeted atmosphere PFAAs (n=268) are presented in Table 1 and Table S4. The total
178 concentrations of Σ_{13} PFAAs analogues varied between 6.19 and 292.57 pg/m^3 , with an average value of 39.84 ± 28.08
179 pg/m^3 . The commonly concerned PFCAs analogues (C5–C14) occupied 79.6% of the total PFAAs, at a level of
180 4.50–247.23 pg/m^3 , whereas the PFSAAs concentrations were 1.04–42.61 pg/m^3 . [The long-chain PFCAs concentrations](#)
181 [were 17.96 \$\pm\$ 13.71 \$\text{pg}/\text{m}^3\$, which were significantly higher than the short-chain concentrations \(13.74 \$\pm\$ 12.19 \$\text{pg}/\text{m}^3\$ \)](#)
182 [\(p<0.05\). Similarly, a recent PFAAs measurement conducted in the landfill atmosphere in Tianjin, China \(Tian et al.,](#)
183 [2018\), found the long chain PFCAs were much higher than the short species.](#) Specifically, PFOA was the dominant
184 PFAAs (accounting 20.6%), and was detected in all atmospheric samples with an average value of 8.19 ± 8.03 pg/m^3 . This
185 phenomenon could occur since PFOA is widely used in the manufacturing of polytetrafluoroethylene (PTFE),
186 perfluorinated ethylene propolymer (FEP), and perfluoroalkoxy polymers (PFA) (Wang et al., 2014). The domestic
187 demand for and the industrial production of PFOA-based products have been increasing in China since the late 1990s
188 (Wang et al., 2014), and direct emissions of FOSA-based products may contribute to the relative high level of PFOA.
189 Meanwhile, one major variation of PFOA precursor, 8:2 FTOH, [was reported to rank](#) as the highest concentration among
190 [neural PFASs in air of China](#) (De Silva, 2004; Martin et al., 2006). Among PFAAs' composition profile, it was followed
191 by PFHxA, PFOS, and PFPeA, with mean concentrations of 5.36, 5.20, and 4.95 pg/m^3 , respectively. The detection
192 frequencies of PFCAs decreased gradually as the carbon chain length increased – for instance, the PFPeA and PFTrDA
193 were detected in 84.8% and 37.3%, respectively.

194 Compared with other gaseous PFAAs measurements, Liu et al. (2015) reported that PFAAs in the urban atmosphere
195 sampled with XAD-containing sorbent in Shenzhen city in China was 15 ± 8.8 pg/m^3 , which contributed to nearly half of
196 this study. Wong et al. (2018) found that a much lower PFAAs levels in the remote Arctic area than this study, with mean
197 value of 1.95 pg/m^3 . This study found generally higher PFAAs abundances compared to measurement in Canada
198 (Gewurtz et al., 2013), which may be attributed to the relative high abundance of industrial and domestic emissions in
199 China. However, the PFAAs concentrations in urban/rural areas in this study were far lower than the measurements at
200 point sources, for example, landfill atmosphere (Tian et al., 2018) ($360\text{--}820$ pg/m^3) and fluorochemical manufacturing
201 facility (Chen et al., 2018) (4900 ± 4200 pg/m^3), suggesting that PFAAs were susceptible to being affected by local source

202 emissions. [Although there existed inherent differences of PFAAs levels between regions, the impacts from differences in](#)
 203 [sampling techniques and sorbents between XAD-PAS and SIP-PAS could not be neglected. As indicated by previous](#)
 204 [researches, XAD has much higher sorptive capacity of PFASs than PUF, wind speed and temperature displayed different](#)
 205 [degrees of impact on their sampling capacity among different regions. Additionally, UV radiation has the potential to](#)
 206 [degrade PFAAs due to O₃, OH·, and other atmospheric oxidants during sampling.](#)

207
 208 **Table 1.** Comparison of PFAAs levels in the present research with measurements in other areas (pg/m³)

Sampling sites	Duration	Sampling location	Sampler type ^a	PFAAs ^b	PFCAs ^c	Reference
23 provinces in China	2017.1–12	Urban and rural areas	XAD–PAS	6.19–292.57; 39.84±28.08	4.50–247.23; 31.69±23.88; C5–C14	This study
Shenzhen, China	2011.9–11	Urban area	SIP–PAS	3.4–34; 15±8.8	11.59±8.74; C4–C12	(Liu et al., 2015)
Fuxin, China	2016.9–10	Fluorochemical manufacturing facilities	SIP–PAS	4900±4200	4900±4200; C4–C12	(Chen et al., 2018)
Tianjin, China	2013	Waste water treatment plant	SIP–PAS	87.9–227; 123	87.9–227; 123; C6–C12	(Yao et al., 2016a)
Tianjin, China	2016.5–6	Landfill	SIP–PAS	280–820	280–820; C4–C12	(Tian et al., 2018)
Canada	2006–2011	Remote and urban areas	SIP–PAS	0.014–0.44	0.014–0.44; C8–C12	(Gewurtz et al., 2013)
Alert, Arctic	2006.8–2015	Remote area	SIP–PAS	1.95	1.95; C4–C8	(Wong et al., 2018)
Toronto, Canada	2010.3–10	Semi–urban site	SIP–PAS	11.24±7.95	11.24±7.95 ; C4–C18	(Ahrens et al., 2013)
Brno, Czech Republic	2013.4–9	Suburban background site	XAD–PAS	30–153	26–147.6; C4–C14	(Karásková et al., 2018)

209 ^a: SIP-PAS represent XAD impregnated PUF sorbent based PAS, which is composed of PUF, mashed XAD-4, and PUF;

210 ^b: [represent the total concentration ranges of PFCAs and PFASs; mean concentrations of the total PFCAs and PFASs;](#)


211 ^c: represent concentration range; mean value; carbon length of PFCAs.

212 3.2 Temporal variations

213 Monthly and seasonal variations of the mean PFAAs concentrations are depicted in Figure 1. In general, an increasing
 214 seasonal mean of PFAAs concentrations [from 23 sampling sites](#) existed for summer (31.35 pg/m³) < autumn (35.63

215 pg/m^3) < spring (42.40 pg/m^3) < winter (52.83 pg/m^3). The winter maxima abundance of PFAAs could be attribute to the
216 stagnant atmospheric conditions, in which atmospheric contaminants were trapped in the air with a weak diluting effect.
217 XAD–PAS showed similar efficiency of capturing gas and particle phases PFASs, while the unimpeded particle gathering
218 efficiency is challenging to quantify. In addition, despite the increase in atmospheric oxidation of precursors in summer
219 may lead to PFCAs rise (Li et al., 2011; Yao et al., 2016a), the abundant rainfall would enhance their scavenging activities
220 (Table S5), ultimately leading to the relatively low concentrations of PFAAs in the summer. Specifically, the PFAAs
221 showed much higher concentrations in spring than other seasons in Shanghai, which was different from Tianjin and
222 Xinjiang (Figure S2). An extreme high level of PFAAs of 135.51 pg/m^3 was occurred in November in Beijing, which was
223 2–4.5 times higher than in other month, indicating the potential point source of PFAAs contamination in this site. In fact,
224 numerous fluoride related products manufacturers were distributed in EC, NC (including Beijing) and CC areas, see
225 detail in Figure S3. As gaseous PFAAs measurements were majorly reported at a relative short time (several weeks to
226 several months), it is somewhat difficult to compare their temporal trends.

227 Interestingly, the evolution of PFAAs showed a dramatic monthly variation, and the monthly mean levels varied from
228 25.92 to 60.57 pg/m^3 , with the lowest and the highest abundances being present in September and December, respectively.
229 For the specific composition profile of PFAAs, the average concentrations of PFOA, PFHxA, PFPeA, and PFOS were
230 10.36 , 8.42 , 6.55 , and 6.44 pg/m^3 in winter, respectively, which were nearly two times higher than in the summer. The
231 seasonal variation trend of PFOS was summer < spring \approx autumn < winter, while PFNA appeared to show winter maxima
232 with concentrations 4 and 3 times higher than in the summer and spring, respectively. However, Wong et al. (2018)
233 reported that PFBS showed the maximal value in winter but found no consistent seasonality for PFOS in the Arctic area.
234 The differences may be explained as the PFAAs in air in the remote Arctic area were originated from long–range
235 transport and volatilization from snow or sea, but not affected by local direct anthropogenic emission.



236
237 **Fig. 1.** Monthly mean concentrations of PFAAs in China from January to December 2017
238

239 **3.3 Geographical distributions**

240 Due to the stark differences in topography and socioeconomic development of Chinese provinces, municipalities, or
 241 autonomous regions, as well as the enormous differences in industrialization and emissions, PFAAs showed significantly
 242 different distribution patterns in China (Figure 2). Overall, the predominant declining gradient of PFAAs' contents was
 243 CC> NC> EC> NE> SW> NW> SC areas in China, which was similar to previous research that the outdoor
 244 dust-bound PFAAs were relatively enriched in the eastern part of mainland China (Yao et al., 2016b). This trend was
 245 not surprising since numerous PFAAs related photoelectric industries, chemical industries, and mechanical industries are
 246 dispersed across CC, EC and NC areas, e.g., Shanghai, Zhejiang, Fujian, Henan, and Jiangsu. As expected, the western
 247 mountain and highland areas, e.g., Xinjiang and Yunnan (20.88 pg/m³), with relatively low population densities and high
 248 latitudes, displayed significantly lower PFAAs concentrations. It was reported that high orographic conditions have a
 249 cold trapping effect on atmospheric PFASs, the transportation of PFAAs involving particles or not should be dramatically
 250 reduced (Konstantinos et al., 2010; Yao et al., 2016a). Given that altitudes increase gradually from several meters in EC,
 251 NC and SC coastal areas to nearly 2,000 meters in SW and NW highland regions in China, the high altitude blocking
 252 effect for atmospheric PFAAs transportation should not be neglected.

253 The annual average concentrations of PFAAs at the provincial level ranged from 12.38 pg/m³ in Xinjiang to 90.88 pg/m³
 254 in Henan, and the composition patterns varied widely. Henan contributed the largest proportion of PFAAs in China, and
 255 showed the highest PFOA level (19.07 pg/m³), which is a typical, heavily-industrialized province characterized by textile

256 treatments, metal plating, and firefighting foam manufacturing, and a large amount of PFAAs emulsifier fluoropolymers
257 were used in industrial production. Special attention should be paid to Zhejiang, the level of which (61.68 pg/m^3) ranked
258 second in PFAAs abundances in spite of its sampling site being located in a village. As well as this, several
259 painting–packaging plants, mechanical plants, and electrical equipment manufacturers were dispersed around this
260 sampling site (see Figure S4), which would contribute to the PFAAs variations in this site. In fact, the GDP of Zhejiang
261 ranked fourth in China, specializing in mechanical manufacture, textiles, and chemical industry. Moreover, the top six
262 sites with abundant of PFAAs were located in the most economically–developed and populated areas (the Yangtze River
263 Delta area, the Circum–Bohai Sea Region), and in the rapidly–developing regions (Henan, Sichuan) in China. In line
264 with this result, a sampling campaign conducted in Asia, including 18 sites in China, found very high levels of PFAAs
265 precursors (8:2 FTOH, 10:2 FTOH) existed in Beijing, Tianjin, and Zhejiang (Li et al., 2011). But meanwhile we should
266 keep in mind that the production of PFCA in the atmosphere from gaseous precursors degradation may be impaired in
267 urban areas, due to the high abundance of NO_x compete for $\text{OH}\cdot$ radicals.

268 Furthermore, PFOA concentrations were apparently high in Henan, Zhejiang, Beijing, Tianjin, and Hubei, where mean
269 values ranged of $11.65\text{--}19.14 \text{ pg/m}^3$ compared with in other provinces ($2.93\text{--}8.54 \text{ pg/m}^3$). PFOA and PFOA–related
270 products have not been banned for use in various industrial and domestic applications (Konstantinos et al., 2010; Wang et
271 al., 2014), which were manufactured extensively in EC and NC areas and were used widely. However, the highest
272 concentration of PFOS was found in Zhejiang (14.13 pg/m^3), which may be affected by local manufacturing of PFOS
273 based products, e.g. leather, paper and metal plating. It was followed by Beijing (8.98 pg/m^3) and Fujian (9.09 pg/m^3),
274 while Xinjiang and Yunnan shared the lowest levels ($1.20\text{--}3.57 \text{ pg/m}^3$). This spatial variation patterns of PFOS in the
275 present study, matched well with a previous national survey that found most PFOS and its derivative facilities in China
276 are suited in EC, CC and NC areas, with emission density ranged from $1\text{--}500 \text{ g}/(\text{km}^2\cdot\text{a})$ (Konstantinos et al., 2010; Wang
277 et al., 2014).




Fig. 2. The spatial distributions of PFAAs in China (annual average of PFAAs, created by ArcGIS 10.4).


3.4 Geographical distributions transport pathway

The PFAAs variations in the atmosphere depended on their local source emissions as well as regional atmosphere transportation. In order to give readers a direct impression of factors affecting the geographical variations of PFAAs in China, here we analyzed PFAAs variations along three pathway transects and one coastal line to determine how PFAAs distribute spatially.

As shown in Figure 3a, PFAAs concentrations were enriched in southeastern areas ($40.58\text{--}47.17\text{ pg/m}^3$) at low altitudes (2–30 m), but relatively low abundances ($12.31\text{--}29.44\text{ pg/m}^3$) existed in the northwestern part of China (397–1,517 m in altitude). As discussed above, the EC areas (e.g. Fujian) were the most intensively industrialized regions, direct emissions from PFAAs manufacturing processes would enhance their atmospheric abundances. However, high altitudes existed in NW areas would have a blocking effect to the transportation of PFAAs from eastern polluted areas.

In terms of the SW–NE transect (Figure 3b), Yunnan and Liaoning showed much lower PFAAs concentrations (20.88 and

292 24.99 pg/m^3) than other areas (44.76–52.58 pg/m^3). Notably, a steady increasing trend of PFAAs concentrations existed
 293 across the W–E transect (Figure 3c), which escalated from 20.88 pg/m^3 in Yunnan to 61.68 pg/m^3 in Zhejiang. The
 294 composition profiles of PFAAs along this transect differed from each other; for instance, PFOA occupied 28.5% of the
 295 total PFAAs in Zhejiang, while it only accounted for 15.6%–21.8% in other areas. Note that PFAAs released from point
 296 sources would be eliminated by deposition, degradation, or dilution during transportation in the atmosphere, e.g., PFOA
 297 could decrease by ~90% within 5 km of its point source (Chen et al., 2018). However, the long range transport of PFAAs
 298 bounded with particles also have been explored in previous research (Pickard et al., 2018). As illustrated in Figure S5,
 299 the 48 hours back trajectories were generally associated with air masses originating from the surrounding areas of the
 300 sampling locations, the trajectories which overlapped with urban areas in Zhejiang, Jiangxi and Shanghai, which
 301 confirmed that the air mass origins was a driving factor for PFAAs variation.
 302 Interestingly, with the exclusion of the site directly affected by surrounding sources in Zhejiang, PFAAs were rather
 303 uniformly distributed among the coastal areas, with concentrations ranging from 24.92–45.76 pg/m^3 (Figure 3d).
 304 Excluded industrial and domestic emissions as well as secondary formation, the PFAAs containing sea spray aerosols
 305 could contribute the variations of PFAAs in coastal atmosphere (Cai et al., 2012b; Pickard et al., 2018).



306
307 **Fig. 3.** Transects of PFAAs concentrations across three different directions and one coastal line

308

309 3.5 Source identification

310 Understanding the sources of PFAAs and their corresponding importance would enable elucidation of the levels of

311 PFAAs in the environment. As discussed above, the observations from tempo-spatial variations of PFAAs suggest that
312 several factors may have a combined effect on the variations of PFAAs. Hence, a PMF model was adopted to extract the
313 potential factors affecting PFAAs variations, and four sources were extracted in this study (see Figure 4).

314 High percentages (~90.0%) of PFPeA and PFBS were found in factor 1, and were moderately loaded with PFOS (62.6%).
315 Three major types of PFOS-related chemicals; namely PFOS salts, PFOS substances and PFOS polymers, are used in
316 industrial products in China (Xie et al., 2013). PFOS salts are usually used in metal plating, firefighting foams, and
317 pesticides, while PFOS substances are adopted in paper treatment and the semiconductor industry. PFOS polymers are
318 employed for textile and leather treatment. These PFOS-related products would lead to direct emissions of PFOS during
319 their industrial and domestic activities. PFPeA and PFBS are the main substitutes for long-chain PFAAs in China, which
320 would release as impurities or by-products when manufacturing PFOS-based products (Liu et al., 2017). Hence, this
321 factor was regarded as the direct source of PFOS-based products. This was consistent with the spatial observations that
322 high PFOS concentrations were shown in Zhejiang, Fujian, Guangdong, and Shanghai, where manufacturing facilities are
323 distributed.


324 Factor 2 was characterized by PFHxA, PFOA, PFNA, and PFDA, each representing over 60.0% of their explained
325 variations. Their rather strong positive correlations ($r=0.54-0.84$, $p<0.01$) suggested that they may have originated from a
326 similar source (Table S6). PFOA was considered as the marker for the emulsification of plastics, rubber products, flame
327 retardants for textiles, paper surface treatments, fire foams and [PTFE emulsifiers](#) (Liu et al., 2015;Konstantinos et al.,
328 2010). It has been reported that there was an increase in PFCAs emissions at the manufacturing sites of PFOA-based
329 products in China between 2002 and 2012 due to a rapid increase in domestic demand and production of PFOA-related
330 products (Wang et al., 2014). PFNA and its derivatives have similar physicochemical properties to PFOA and its
331 derivatives, and both can be emitted through exhaust gases. The PFNA-based production was found to be related to
332 polyvinylidene fluoride (PVDF) production, and it has been suggested that PVDF production increased in China after
333 2008 (Wang et al., 2014). Therefore, factor 2 represents direct sources of PFOA-based and PFNA-based products.

334 The compositions of factor 3 were characterized by a high loading of PFHpA and PFHxS, with loading factor values of
335 84.9% and 81.7%, respectively. The historical production and uses of PFHpA and its derivatives remain unidentified.
336 Factor with PFHxS alone did not indicate a specific source, so this factor may be classified as an unknown source, which
337 may be affected by atmosphere air mass transport, sea aerosol bursting and/or other origins.

338 The final factor was dominated by PFUdA, PFDaA, PFTrDA, and PFTeDA, with loading factor values larger than 80%.
339 These long-chain PFAAs (C11-C14) analogues have been interpreted as degradation products of fluorotelomer-based
340 products in previous research (Liu et al., 2017;Wang et al., 2014; Thackray and Selin, 2017). Based on the life-cycle

341 usage and release from fluorotelomer and other fluorinated products, the global cumulative estimation of PFUdA,
 342 PFDoA, PFTrDA, and PFTeDA from quantified sources was estimated to be 9–230 tons in the period of 2003–2015, and
 343 projected to be between 0–84 tons between 2016–2030 (Wang et al., 2014). It was reported that the manufacturing of
 344 fluorotelomer-based substances would increase in China. In addition, these four analogues showed apparent positive
 345 correlations to each other ($r = 0.59\text{--}0.79$, $p < 0.01$). Thus, this factor was explained as the degradation products of
 346 fluorotelomer-based products, which could be proven by their higher abundances caused by an enhanced atmospheric
 347 oxidation ability in the summer than other seasons.

348 Direct emission sources, including PFOS-based products, PFOA-based products, and PFNA-based products were
 349 estimated to represent 62.7% of the total PFAAs sources. Indirect sources of degradation products of
 350 fluorotelomer-based products played a minor role, contributing 15.5%, and there are 21.8% of variances that could still
 351 not be explained and need further detailed investigation. This source apportionment result was similar to one recent piece
 352 of research that found that industrial PFOA emissions were the major sources of atmospheric PFAAs in Shenzhen, China
 353 (Liu et al., 2015), and the long-distance transportation of pollutants also made a contribution.



354

355

Fig. 4. Factor profiles of PFAAs extracted by the PMF model

356

4. Conclusion

357

In the present study, PFAAs were ubiquitously detected in the atmosphere across China over the length of a year. Results

358 indicated that the measured PFAAs [in the present study](#) were several times to several magnitudes higher than [the levels](#)
359 [conducted in most other urban locations, while far lower than the measurements implemented at point sources. In which,](#)
360 [the C5–C14 PFCA analogues occupied 79.6% of the total PFAAs variations, PFOA, PFHxA and PFOS ranked the top](#)
361 [three species. Additionally,](#) much higher abundances of PFAAs existed in winter compared with in summer. In terms of
362 spatial distribution, the PFAAs concentrations were higher in central and eastern China, where dense residential and
363 industrial manufacturing facilities were distributed. Correlation analysis, Hysplit backward trajectories, and PMF
364 receptor model, [have combined to suggest](#) that the direct sources of PFOS-based, PFOA-based, and PFNA-based
365 products made a predominant contribution to variations in PFAAs, while indirect degradation played a minor role.

366 **Acknowledgements**

367 This study was financially supported by National Key Research & Development Plan (2016YFC0200104), National
368 Natural Science Foundation of China (No. 21577090 and No. 21777094), and China Postdoctoral Innovative Talent
369 Support Project (BX20190169). We thank Lei Ye (Xi'an University of Architecture and Technology), Fengxia Wang
370 (Hainan University), Linrui Jia (Beijing Normal University), Songfeng Chu (Tongji University), and other 18 volunteers,
371 for coordinating the sampling process and for their valuable contribution to field measurement. We appreciate senior
372 engineer Xiaofang Hu (Instrumental Analysis Center, SESE, Shanghai Jiao Tong University) for her assistance in
373 experiment analysis.

374 **Appendix A: Supplementary material**

375 **References**

376 Ahrens, L., Harner, T., Shoeib, M., Koblizkova, M., and Reiner, E. J.: Characterization of Two Passive Air Samplers for
377 Per- and Polyfluoroalkyl Substances, *Environ. Sci. Technol.*, 47, 14024-14033, <https://doi.org/10.1021/es4048945>, 2013.
378 Armitage, J. M., Hayward, S. J., and Frank, W.: Modeling the Uptake of Neutral Organic Chemicals on XAD Passive Air
379 Samplers under Variable Temperatures, External Wind Speeds and Ambient Air Concentrations (PAS-SIM), *Environ. Sci.*
380 *Technol.*, 47, 13546-13554, <https://doi.org/10.1021/es402978a>, 2013.
381 Baard Ingegerdsson, F., Line Sm. Stuen, H., Raymond, O., Hanne Line, D., Merete, H., Cathrine, T., Syvert, T., Georg, B.,
382 Paal, M., and Ellingsen, D. G.: Occupational exposure to airborne perfluorinated compounds during professional ski
383 waxing, *Environ. Sci. Technol.*, 44, 7723-7728, <https://doi.org/10.1021/es102033k>, 2010.

384 [Buck, R. C., Franklin, J., Berger, U., Conder, J. M., Cousins, I. T., Voogt, P. D.: Perfluoroalkyl and polyfluoroalkyl](#)
385 [substances in the environment: terminology, classification, and origins. *Int. Environ. Assess. Manag.*, 7\(4\), 513-541,](#)
386 <https://doi.org/10.1002/ieam.258>, 2011.

387 Cai, M., Xie, Z., Moeller, A., Yin, Z., Huang, P., Cai, M., Cai M. Yang, H., Sturm, R., and He, J.: Polyfluorinated
388 compounds in the atmosphere along a cruise pathway from the Japan Sea to the Arctic Ocean, *Chemosphere*, 87(9),
389 989-997 <https://doi.org/10.1016/j.chemosphere.2011.11.010>, 2012a.

390 Cai M., Zheo, Z., Yin Z., Ahrens L., Huang P., Cai M., Yang H., He J., Sturm R., Ebinghaus R., Xie Z.: Occurrence of
391 perfluoroalkyl compounds in surface waters from the North Pacific to the Arctic Ocean, *Environ. Sci. Technol.*, 46,
392 661-668, <https://doi.org/10.1021/es2026278>, 2012b.

393 Cardenas, A., Gold, D. R., Hauser, R., Kleinman, K. P., Hivert, M. F., Calafat, A. M., Ye, X., Webster, T. F., Horton, E. S.,
394 and Oken, E.: Plasma Concentrations of Per- and Polyfluoroalkyl Substances at Baseline and Associations with Glycemic
395 Indicators and Diabetes Incidence among High-Risk Adults in the Diabetes Prevention Program Trial, *Environ Health*
396 *Perspect*, 125, 107001, <https://doi.org/10.1289/EHP1612>, 2017.

397 Cassandra, R., Tom, H., K, S. J., Anita, E., Gilberto, F., Eugenia, C. L., Oscar, F., Martin, V. I., S.B., M. K., and Isabel, M.
398 R.: Atmospheric concentrations of new POPs and emerging chemicals of concern in the Group of Latin America and
399 Caribbean (GRULAC) region, *Environ. Sci. Technol.*, 52(13), 7240-7249, <https://doi.org/10.1021/acs.est.8b00995>, 2018.

400 Chen, H., Yao, Y., Zhao, Z., Wang, Y., Wang, Q., Ren, C., Wang, B., Sun, H., Alder, A. C., and Kannan, K.: Multimedia
401 Distribution and Transfer of Per- and Polyfluoroalkyl Substances (PFASs) Surrounding Two Fluorochemical
402 Manufacturing Facilities in Fuxin, China, *Environ. Sci. Technol.*, 52, 8263-8271, <https://doi.org/10.1021/acs.est.8b00544>,
403 2018.

404 De Silva, A. O.: Degradation of fluorotelomer alcohols: a likely atmospheric source of perfluorinated carboxylic acids,
405 *Environ. Sci. Technol.*, 38, 3316-33121, <https://doi.org/10.1021/es049860w>, 2004.

406 Dreyer, A., Weinberg, I., Temme, C., and Ebinghaus, R.: Polyfluorinated Compounds in the Atmosphere of the Atlantic
407 and Southern Oceans: Evidence for a Global Distribution, *Environ. Sci. Technol.*, 43, 6507-6514, <https://doi.org/10.1021/es9010465>, 2009.

409 [Fang, X., Wang, Q., Zhao, Z., Tang, J., Tian, C., Yao, Y., Yu, J., and Sun, H.: Distribution and dry deposition of](#)
410 [alternative and legacy perfluoroalkyl and polyfluoroalkyl substances in the air above the Bohai and Yellow Seas, China,](#)
411 [Atmos. Environ.](#), 192, 128-135, <https://doi.org/10.1016/j.atmosenv.2018.08.052>, 2018.

412 Gewurtz, S. B., Backus, S. M., Silva, A. O., De, Lutz, A., Alain, A., Marlene, E., Susan, F., Melissa, G., Paula, G., and
413 Tom, H.: Perfluoroalkyl acids in the Canadian environment: multi-media assessment of current status and trends, *Environ.*

414 Int., 59, 183-200, <https://doi.org/10.1016/j.envint.2013.05.008>, 2013.

415 Han, D., Fu, Q., Gao, S., Li, L., Ma, Y., Qiao, L., Xu, H., Liang, S., Cheng, P., Chen, X., Zhou, Y., Yu, J. Z., and Cheng,
416 J.: Non-polar organic compounds in autumn and winter aerosols in a typical city of eastern China: size distribution and
417 impact of gas–particle partitioning on PM_{2.5} source apportionment, *Atmos. Chem. Phys.*, 18, 9375-9391, [https://doi.org/](https://doi.org/10.5194/acp-18-9375-2018)
418 10.5194/acp-18-9375-2018, 2018.

419 Han, D., Fu, Q., Gao, S., Zhang, X., Feng, J., Chen, X., Huang, X., Liao, H., Cheng, J., and Wang, W.: Investigate the
420 impact of local iron–steel industrial emission on atmospheric mercury concentration in Yangtze River Delta, China,
421 *Environ. Sci. Pollut. Res.*, 26(6), 5862–5872, <https://doi.org/10.1007/s11356-018-3978-7>, 2019.

422 Hu, X. C., Andrews, D. Q., and Lindstrom, A. B.: Detection of Poly- and Perfluoroalkyl Substances (PFASs) in U.S.
423 Drinking Water Linked to Industrial Sites, Military Fire Training Areas, and Wastewater Treatment Plants, *Environ. Sci.*
424 *Technol. Lett.*, 3, 344-350, <https://doi.org/10.1021/acs.estlett.6b00260>, 2016.

425 Jian, J. M., Guo, Y., Zeng, L., Liu, L. Y., Lu, X., Wang, F., and Zeng, E. Y.: Global distribution of perfluorochemicals
426 (PFCs) in potential human exposure source—A review, *Environ. Int.*, 108, 51-62, <https://doi.org/>, 2017.

427 Johansson, N., Fredriksson, A., and Eriksson, P.: Neonatal exposure to perfluorooctane sulfonate (PFOS) and
428 perfluorooctanoic acid (PFOA) causes neurobehavioural defects in adult mice, *Neurotoxicology*, 29, 160-169,
429 <https://doi.org/10.1016/j.neuro.2007.10.008>, 2008.

430 Karásková, P., Codling, G., Melymuk, L., and Klánová, J.: A critical assessment of passive air samplers for per- and
431 polyfluoroalkyl substances, *Atmos. Environ.*, 185, 186-195, <https://doi.org/10.1016/j.atmosenv.2018.05.030>, 2018.

432 Konstantinos, P., Cousins, I. T., Buck, R. C., and Korzeniowski, S. H.: Sources, fate and transport of
433 perfluorocarboxylates, *Environ. Sci. Technol.*, 40(1), 32-44, <https://doi.org/10.1002/chin.200611255>, 2010.

434 Krogseth, I. S., Xianming, Z., Ying, D., Lei, Frank, W., and Knut, B.: Calibration and application of a passive air sampler
435 (XAD-PAS) for volatile methyl siloxanes, *Environ. Sci. Technol.*, 47, 4463-4470, <https://doi.org/10.1021/es400427h>,
436 2013.

437 Lai, F. Y., Rauert, C., Gobelius, L., and Ahrens, L.: A critical review on passive sampling in air and water for per- and
438 polyfluoroalkyl substances (PFASs), *TrAC Trends Anal. Chem.*, Available online 23 Nov. 2018, [https://doi.org/](https://doi.org/10.1016/j.trac.2018.11.009)
439 10.1016/j.trac.2018.11.009, 2018.

440 Li, J., Vento, S. D., Schuster, J., Gan, Z., Chakraborty, P., Kobara, Y., and Jones, K. C.: Perfluorinated Compounds in the
441 Asian Atmosphere, *Environ. Sci. Technol.*, 45, 7241-7428, <https://doi.org/10.1021/es201739t>, 2011.

442 Lindstrom, A. B., Strynar, M. J., and Libelo, E. L.: Polyfluorinated Compounds: Past, Present, and Future, *Environ. Sci.*
443 *Technol.*, 45, 7954-7961, <https://doi.org/10.1021/es2011622>, 2011.

444 Liu, B., Zhang, H., Yao, D., Li, J., Xie, L., Wang, X., Wang, Y., Liu, G., and Yang, B.: Perfluorinated compounds (PFCs)
445 in the atmosphere of Shenzhen, China: Spatial distribution, sources and health risk assessment, *Chemosphere*, 138,
446 511-518, <https://doi.org/10.1016/j.chemosphere.2015.07.012>, 2015.

447 Liu, Z., Lu, Y., Wang, P., Wang, T., Liu, S., Johnson, A. C., Sweetman, A. J., and Baninla, Y.: Pollution pathways and
448 release estimation of perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) in central and eastern China,
449 *Sci. Total Environ.*, 580, 1247-1256, <https://doi.org/10.1016/j.scitotenv.2016.12.085>, 2017.

450 Martin, J. W., Ellis, D. A., Mabury, S. A., Hurley, M. D., and Wallington, T. J.: Atmospheric chemistry of
451 perfluoroalkanesulfonamides: kinetic and product studies of the OH radical and Cl atom initiated oxidation of N-ethyl
452 perfluorobutanesulfonamide, *Environ. Sci. Technol.*, 40, 864-872, <https://doi.org/10.1021/es051362f>, 2006.

453 [Melymuk, L., Bohlin, P., Sáňka, O., Pozo, K., and Klánová, J.: Current challenges in air sampling of semivolatile organic
454 contaminants: sampling artifacts and their influence on data comparability. *Environ. Sci. Technol.*, 48\(24\), 14077-91,
455 <https://doi.org/10.1021/es502164r>, 2014.](https://doi.org/10.1021/es502164r)

456 [Okeme, J. O., Saini, A., Yang, C., Zhu, J., Smedes, F. and Klánová, J.: Calibration of polydimethylsiloxane and
457 XAD-pocket passive air samplers \(PAS\) for measuring gas- and particle-phase SVOCs. *Atmos. Environ.*, 143, 202-208,
458 <https://doi.org/10.1016/j.atmosenv.2016.08.023>, 2016.](https://doi.org/10.1016/j.atmosenv.2016.08.023)

459 Pickard H. M., Criscitiello A. S., Spencer C., Sharp M. J., Muir D.C. G., De Silva A. O., Young C. J.: Continuous
460 non-marine inputs of per- and polyfluoroalkyl substances to the High Arctic: a multi-decadal temporal record, *Atmos.*
461 *Chem. Phys.*, 18, 5045–5058, <https://doi.org/10.5194/acp-18-5045-2018>, 2018.

462 Rauert, C., Harner, T., Schuster, J. K., Eng, A., Fillmann, G., Castillo, L. E., Fentanes, O., Villa, M. I., Miglioranza, K.,
463 and Moreno, I. R.: Atmospheric Concentrations of New Persistent Organic Pollutants and Emerging Chemicals of
464 Concern in the Group of Latin America and Caribbean (GRULAC) Region, *Environ. Sci. Technol.*, 52(13), 7240-7249,
465 <https://doi.org/10.1021/acs.est.8b00995>, 2018.

466 Sedlak, M. D., Benskin, J. P., Wong, A., Grace, R., and Greig, D. J.: Per- and polyfluoroalkyl substances (PFASs) in San
467 Francisco Bay wildlife: Temporal trends, exposure pathways, and notable presence of precursor compounds,
468 *Chemosphere*, 185, 1217-1226, <https://doi.org/10.1016/j.chemosphere.2017.04.096>, 2017.

469 Thackray, C. P., Selin N. E.: Uncertainty and variability in atmospheric formation of PFCA from fluorotelomer
470 precursors, *Atmos. Chem. Phys.*, 17, 4585–4597, <https://doi.org/10.5194/acp-17-4585-2017>, 2017.

471 Tian, Y., Yao, Y., Chang, S., Zhao, Z., Zhao, Y., Yuan, X., Wu, F., and Sun, H.: Occurrence and Phase Distribution of
472 Neutral and Ionizable Per- and Polyfluoroalkyl Substances (PFASs) in the Atmosphere and Plant Leaves around Landfills:
473 A Case Study in Tianjin, China, *Environ. Sci. Technol.*, 52, 1301-1310, <https://doi.org/10.1021/acs.est.7b05385>, 2018.

474 Tian, Y., Zhou, Y., Miao, M., Wang, Z., Yuan, W., Liu, X., Wang, X., Wang, Z., Wen, S., and Liang, H.: Determinants of
475 plasma concentrations of perfluoroalkyl and polyfluoroalkyl substances in pregnant women from a birth cohort in
476 Shanghai, China, *Environment International*, 119, 165-173, <https://doi.org/10.1016/j.envint.2018.06.015>, 2018.

477 Wang, Q. W., Yang, G. P., Zhang, Z. M., and Jian, S.: Perfluoroalkyl acids in surface sediments of the East China Sea,
478 *Environ. Pollut.*, 231, 59-67, <https://doi.org/10.1016/j.envpol.2017.07.078>, 2017.

479 Wang, Z., Cousins, I. T., Scheringer, M., Buck, R. C., and Hungerbühler, K.: Global emission inventories for C4-C14
480 perfluoroalkyl carboxylic acid (PFCA) homologues from 1951 to 2030, Part I: production and emissions from
481 quantifiable sources, *Environ. Int.*, 70, 62–75, <https://doi.org/10.1016/j.envint.2014.04.013>, 2014.

482 Wang X., Schuster J., Jones K. C., Gong P.: Occurrence and spatial distribution of neutral perfluoroalkyl substances and
483 cyclic volatile methylsiloxanes in the atmosphere of the Tibetan Plateau, *Atmos. Chem. Phys.*, 18, 8745–8755,
484 <https://doi.org/10.5194/acp-18-8745-2018>, 2018.

485 Wong, F., Shoeib, M., Katsoyiannis, A., Eckhardt, S., Stohl, A., Bohlinizzetto, P., Li, H., Fellin, P., Su, Y., and Hung, H.:
486 Assessing temporal trends and source regions of per- and polyfluoroalkyl substances (PFASs) in air under the Arctic
487 Monitoring and Assessment Programme (AMAP), *Atmos. Environ.*, 172, 65-73, <https://doi.org/10.1016/j.atmosenv.2017.10.028>, 2018.

488

489 Xie, S., Wang, T., Liu, S., Jones, K. C., Sweetman, A. J., and Lu, Y.: Industrial source identification and emission
490 estimation of perfluorooctane sulfonate in China, *Environ. Int.*, 52, 1-8, <https://doi.org/10.1016/j.envint.2012.11.004>,
491 2013.

492 Yao, Y., Chang, S., Sun, H., Gan, Z., Hu, H., Zhao, Y., and Zhang, Y.: Neutral and ionic per- and polyfluoroalkyl
493 substances (PFASs) in atmospheric and dry deposition samples over a source region (Tianjin, China), *Environ Pollut*, 212,
494 449-456, <https://doi.org/10.1016/j.envpol.2016.02.023>, 2016a.

495 Yao, Y., Sun, H., Gan, Z., Hu, H., Zhao, Y., Chang, S., and Zhou, Q.X.: A Nationwide Distribution of Per- and
496 Polyfluoroalkyl Substances (PFASs) in Outdoor Dust in Mainland China From Eastern to Western Areas, *Environ. Sci.*
497 *Technol.*, 50(7), 3676-3685, <https://doi.org/10.1021/acs.est.6b00649>, 2016b.

498 [Zhen, W., Xie, Z., Möller, A., Mi, W., Wolschke, H., and Ebinghaus, R.: Atmospheric concentrations and gas/particle](https://doi.org/10.1016/j.atmosenv.2014.06.036)
499 [partitioning of neutral poly- and perfluoroalkyl substances in northern german coast. *Atmos. Environ.*, 95\(1\), 207-213,](https://doi.org/10.1016/j.atmosenv.2014.06.036)
500 <https://doi.org/10.1016/j.atmosenv.2014.06.036>, 2014.