

Dear editor:

We would like to thank again the referees for their helpful comments and suggestions, which have been fully taken into account upon manuscript revision. A point-by-point response to the comments and a revised manuscript were uploaded.

Sincerely yours,

Nan Ma and co-authors

Response to comments of referee #2

The manuscript has been extensively edited with several new sections, resulting in a clear and comprehensive work with only some minor issues to be solved.

Response:

Thanks for the comment.

(1) The section comparing emissions with detected decrease would profit from a description of the emission data. Where are the emission data from, what are they based on etc.?

Response:

Thanks for the suggestion. The emission data used in the comparison Sect. 3.2 is from German Informative Inventory Report (UBA, 2020). This report was produced by the national co-ordination agency for the National System of Emissions Inventories (Nationales Systems Emissionsinventare; NaSE), sited within the German Federal Environmental Agency (UBA). The dataset is based on a large number of sources and publications and the detailed information is available at <http://iir-de.wikidot.com/>. In Sect. 3.2, we have added the following description:

“In this section, the long-term variation of total emission of major pollutants in Germany are compared with the trends of aerosol concentrations measured at the six regional background and mountain sites. The emission data is from German Informative Inventory Report (UBA, 2020) produced by the national co-ordination agency for the National System of Emissions Inventories (Nationales Systems Emissionsinventare; NaSE) sited within the German Federal Environmental Agency (UBA), and is based on a large number of sources and publications. Detailed information is available in UBA (2020).”

Reference:

German Federal Environmental Agency, UBA: German Informative Inventory Report 2020 (IIR 2020). available at <http://iir-de.wikidot.com/>; access: 1 March 2020.

(2) Have the differences before and after LEZ / between individual LEZ levels been tested for statistical significance? (L378+L389)

Response:

Thanks for the comment. Two-sample t-test was used to test the differences between the concentration levels before and after LEZ. And we find that all differences are statistically significant at 95 % significance level. To make it clearer, the sentences have been revised as:

“Figure 6a and b show that the eBC mass concentration and $N_{[30-200]}$ have gradually decreased after the implementation of each of new stages of LEZ. And the differences between the concentration levels at different stages are statistically significant at 95 % significance level. However, the difference between stage 2 and 3 is relatively negligible.”

“Statistically significant differences are found between the concentration levels before and after the implementation of LEZ for both $N_{[30-200]}$ and eBC mass concentration. But the decrease in $N_{[30-200]}$ after 2010 (Fig. 6c) is much larger than that in eBC mass concentration (Fig. 6d).”

(3) L273 The large contribution of SOA on PM would be expected from a close relationship between decrease in precursors and decrease in PM? Since this is not the case, the sentence is confusing.

Response:

Thanks for the comment. The sentences have been revised as:

“Secondary aerosol formation processes are highly complex and nonlinear, determined not only by the concentrations of precursors but also many other factors such as solar radiation, temperature, humidity, and diffusion condition etc. Thus, the concentration of secondary aerosol might not follow the variation of precursor concentrations. It was found that secondary aerosol contributes a large fraction in particulate matters in the regional background settings (Castro et al., 1999). Therefore, discrepancies are observed between the emissions and particle concentrations although decrease trends are found in both of them.”

(4) L325 It is not clear what other parameters show similar diurnal pattern as N_{10-30} at UB?

Response:

Thanks for the comment. The sentence has been revised as “It is interesting that at the roadside and urban background sites the trend of N_{10-30} has similar diurnal pattern as those for N_{30-200} and eBC mass concentration, but at regional background and low mountain range sites they look quite different.”

(5) Table 5 Why there is no percentage change in temperature data if it is done for precipitation and wind speed?

Response:

Since the average ambient temperature (Celsius) may be close to 0° or below 0° in cold seasons, the calculated percentage change may be a very large number or opposite to the absolute change of temperature, which does not indicate the real trend. And absolute temperature (K) is not usually used for ambient temperature. Therefore, we decided to give only the absolute temperature change in Table 5.

(6) In figure captions, it is sometimes not clear what data are plotted; for example, in Fig. 3 the text says it is only from regional background and mountain stations but the figure caption states just Germany... The same issue is also with Fig. 10.

Response:

Thanks for the comment. The captions of Fig. 3 and Fig. 10 have been revised as:

Figure 3: Long-term change index (% of average in 2009) of measured parameters at the regional background and mountain stations sites and total emissions in Germany: (a) measured eBC mass concentration and BC emission; (b) measured N_{10-30} , N_{30-200} , $N_{200-800}$, and V_{20-800} , and emission of $PM_{2.5}$, SO_2 , NMVOC, NH_3 and NO_x .

Figure 10: Annual average eBC mass concentration and $N_{200-800}$ measured at the regional background and low mountain range sites for the polluted and clean air mass categories. Green bars show the frequency of polluted air masses in each year.

(7) In Fig. 7, maybe ΔeBC etc. would be a better y-axis label instead of only eBC? Similarly, the same applies on b and c parts as well.

Response:

Thanks for the suggestion. Figure 7 has been modified, as shown below. Accordingly, the description of Fig. 7 in the text has been revised as: “Figure 7 illustrates the annually averaged diurnal cycles of the increments of eBC mass concentration (ΔeBC), $N_{[30-200]}$ ($\Delta N_{[30-200]}$) and $N_{[200-800]}$ ($\Delta N_{[200-800]}$). Before and after the implementation of LEZ (2010 and 2011), the increments show a sudden decrease of up to 40 % during daytime. The average ΔeBC , $\Delta N_{[30-200]}$ and $\Delta N_{[200-800]}$ during working hours (06:00 to 18:00 local time) in 2010 are respectively about 1.63, 1.33, and 1.58 times higher than those in 2011.”




Figure 7: Average diurnal cycles of the increment of eBC mass concentration (ΔeBC , panel a), $N_{[30-200]}$ ($\Delta N_{[30-200]}$, panel b) and $N_{[200-800]}$ ($\Delta N_{[200-800]}$, panel c).

(8) In Fig. 4, the RS, UB etc. abbreviations could be explained in the caption?

Response:

Thanks for the suggestion. The caption of Fig. 4 has been revised as:

Figure 4: Diurnal variations of the trends of PNCs and eBC mass concentration for the four site categories (RS: roadside; UB: urban background; RB: regional background; LMT&HA: low mountain range and high Alpine).

Decreasing Trends of Particle Number and Black Carbon Mass Concentrations at 16 Observational Sites in Germany from 2009 to 2018

Jia Sun¹, Wolfram Birmili^{2,1}, Markus Hermann¹, Thomas Tuch¹, Kay Weinhold¹, Maik Merkel¹, Fabian Rasch¹, Thomas Müller¹, Alexander Schladitz^{3,a}, Susanne Bastian³, Gunter Löschau³, Josef Cyrus^{4,5}, Jianwei Gu^{4,5,b}, Harald Flentje⁶, Björn Briel⁶, Christoph Asbach⁷, Heinz Kaminski⁷, Ludwig Ries², Ralf Sohmer², Holger Gerwig², Klaus Wirtz², Frank Meinhardt², Andreas Schwerin², Olaf Bath², Nan Ma^{8,1}, Alfred Wiedensohler¹

¹Leibniz Institute for Tropospheric Research (TROPOS), Leipzig, Germany

²German Environment Agency (UBA), Dessau-Roßlau, Germany

³Saxon State Office for Environment, Agriculture and Geology (LfULG), Dresden, Germany

⁴Helmholtz Zentrum München (HMGU), Institute of Epidemiology II, Neuherberg, Germany

⁵University of Augsburg (UA), Wissenschaftszentrum Umwelt, Augsburg, Germany

⁶Deutscher Wetterdienst (DWD), Meteorologisches Observatorium Hohenpeißenberg, Germany

⁷Institute of Energy and Environmental Technology (IUTA), Duisburg, Germany

⁸Institute for Environmental and Climate Research, Jinan University, Guangzhou, Guangdong 511443, China

^anow at: SICK Engineering GmbH, Ottendorf-Okrilla, Germany

^bnow at: Fraunhofer Wilhelm-Klauditz-Institut (WKI), Braunschweig, Germany

Correspondence to: Nan Ma (nan.ma@jnu.edu.cn) and Alfred Wiedensohler (alfred.wiedensohler@tropos.de)

Abstract

Anthropogenic emissions are dominant contributors to air pollution. Consequently, mitigation policies have been attempted since the 1990s in Europe to reduce pollution by anthropogenic emissions. To evaluate the effectiveness of these mitigation policies, the German Ultrafine Aerosol Network (GUAN) was established in 2008, focusing on black carbon (BC) and sub-micrometre aerosol particles. In this study, long-term trends of atmospheric particle number concentrations (PNCs) and equivalent BC (eBC) mass concentration over a 10-year period (2009–2018) were determined for 16 GUAN sites ranging from roadside to high Alpine environment. Overall, statistically significant decreasing trends are found for most of these parameters and environments in Germany. The annual relative slope of eBC mass concentration varies between –13.1 % and –1.7 % per year. The slopes of the PNCs vary from –17.2 % to –1.7 %, –7.8 % to –1.1 %, and –11.1 % to –1.2 % per year for 10–30 nm, 30–200 nm, and 200–800 nm size ranges, respectively. The reductions in various anthropogenic emissions are found to be the dominant factors responsible for the decreasing trends of eBC mass concentration and PNCs. The diurnal and seasonal variations in the trends clearly show the effects of the mitigation policies for road transport and residential emissions. The influences of other factors such as air masses, precipitation and temperature were also examined and found to be less important or negligible. This study proves that a combination of emission mitigation policies can effectively improve the air quality on large spatial scales. It also suggests that a long-term aerosol measurement network in multi-type sites is an efficient and necessary tool for evaluating emission mitigation policies.

1 Introduction

Epidemiological studies have shown that increased particulate air pollution due to anthropogenic emissions leads to adverse health effects (Seaton et al., 1995) and increases global disease burden (Cohen et al., 2017). Among the ambient sub-micrometre aerosol particles (diameter $< 1\ \mu\text{m}$), ultrafine particles (UFP, diameter $< 100\ \text{nm}$) share the largest number fraction. The UFP can penetrate deep into lungs and translocate to other organs such as brain, causing health problems such as respiratory and cardiovascular diseases (Kreyling et al., 2006; Schmid and Stoeger, 2016). In urban areas, black carbon (BC) produced by incomplete combustion of fossil fuel and biomass accounts for a significant fraction of UFP mass (Chen et al., 2014; Cheng et al., 2013; Pérez et al., 2010). As BC could operate as a universal carrier of a wide variety of toxins such as polycyclic aromatic hydrocarbons (PAH) and heavy metals into human body, exposure to BC could cause acute health effects such as cardiopulmonary morbidity and mortality (Janssen et al., 2012).

The European Union (EU) was one of the earliest regions around the world to implement emission reduction policies to reduce the harmful effects of air pollution. EU emission mitigation legislations are directly formulated based on emission sources. In Europe, the main anthropogenic sources of primary aerosol particles are fuel combustions from industrial installations (power generation, industry, etc.), non-road and road transport, and domestic heating (European Environment Agency, 2017). The Member States of EU were required to draft National Programmes to the Commission (<http://ec.europa.eu/environment/air/reduction/implementation.htm>). In Germany, the Federal Environment Ministry issued the Federal Emission Control Regulations (Bundes-Immissionsschutzverordnung, BImSchV). To reduce the emission from industrial installations, the BImSchV regulates permits for construction and operation of industrial installations. The emission limits for large combustion, such as for that from power plants, are defined as well. For domestic heating emission, the unsuitable fuels are listed and their emission values are defined. For traffic emission, low emission zones (LEZs) were set up to limit the emission of nitrogen oxide and aerosol particles from traffic exhaust.

It is important to evaluate the effectiveness of the implemented emission mitigation policies through long-term observations of pollutants such as sub-micrometre particles and BC. There have been many studies on the long-term trends of particle number concentration (PNC) and/or BC mass concentration since the 1990s. These studies have concluded that emission mitigation policies may reduce human exposure to the pollutants and the results were important for the policy makers (Barnpadimos et al., 2011; Masiol et al., 2018; Kutzner et al., 2018; Putaud et al., 2014). Most of these studies were conducted at roadside or urban background, which are largely dominated by traffic emissions. Few of them have focused on long-term trends at the regional background setting (Asmi et al., 2013; Barnpadimos et al., 2011; Murphy et al., 2011). Murphy et al. (2011) found that between 1990 and 2004, the elemental carbon (EC) mass concentration decreased in several of the national parks and other remote sites in the US. This result was an indication that the emission control policies were effective in reducing the EC mass concentration in the background air across the US. Asmi et al. (2013) analysed the long-term change of PNC at the regional background and remote sites in Europe, North America, Antarctica, and Pacific Ocean islands during 2001–2010. The results showed that decreased PNCs could likely be explained by the reduction of anthropogenic emissions. Kutzner et al. (2018) evaluated the trend of BC over Germany based on measurements at traffic, urban background, and rural sites for the period of 2005–2014, and concluded that the observed decreasing trends in BC are likely owing largely to mitigation measures in the traffic sector. However, there is still a lack of a thorough investigation of the connections between the long-term trends of PNCs/BC and the change of different

anthropogenic emissions. A better understanding of the influence of the inter-annual variation of meteorological conditions on the observed trends is also needed.

Based on a unique dataset from the German Ultrafine Aerosol Network (GUAN), this study investigates the long-term variation in the regional PNC and BC mass concentration, to understand the effectiveness of the emission mitigation policies in reducing the PNC and BC in Germany. The study was conducted for the period of 2009–2018 with data from 16 observational sites representing different types of environment (roadside, urban background, regional background, low mountain, and high Alpine). The overall, diurnal, and seasonal trends of PNCs and BC are evaluated and the role of potential decisive factors, including not only emission mitigation policies, but also other potential drivers (i.e. inter-annual change in meteorological conditions and long-range transport patterns) are discussed.

2 Data and methods

2.1 The German Ultrafine Aerosol Network (GUAN)

The GUAN is a specialized network in Germany that provides continuous measurements including sub-micrometre particle number size distribution (PNSD) and the equivalent BC (eBC) mass concentration in diverse environments including roadside, urban background, regional background, low mountain range and high Alpine. The GUAN combines federal and state air quality monitoring stations as well as atmospheric observatories from research institutes, aiming at a better understanding of sub-micrometre PNC and BC with respect to human health and climate impact (Birmili et al., 2016).

Table 1 lists the basic information of the GUAN sites evaluated in this study. The locations of all 16 sites are illustrated in Fig. 1. A summarised description of the sites is given here and more details are available in Birmili et al. (2016). Among the 16 sites, seven are located in the state of Saxony: Leipzig-Mitte (LMI), Leipzig-Eisenbahnstraße (LEI), Leipzig-Leibniz Institute for Tropospheric Research (TROPOS) (LTR), Melpitz (MEL), Dresden-Nord (DDN), Dresden-Winkelmann-straße (DDW), and Annaberg-Buchholz (ANA). LMI and LEI are two roadside stations in Leipzig. LMI is located on roadside in an open area in the city centre, while LEI is a street canyon station. The traffic volumes at these two sites are 44000 and 12000 vehicles per day, respectively. LTR is an atmospheric research station situated on the roof of the main building of TROPOS. Station MEL is located in a farmland about 50 km from Leipzig. Previous studies have showed that MEL can well represent the regional background atmosphere of Central Europe (Spindler et al., 2013). Two stations are located in the city of Dresden: a roadside site DDN with a traffic volume of about 36000 vehicles per day and an urban background site DDW, 1.7 km away from the city centre. ANA is an urban background station located in the city of Annaberg-Buchholz in the Ore mountain area about 10 km away from the German-Czech border (Schladitz et al., 2015).

Three stations are located in the lowlands of Northern Germany: Bösel (BOS), Neuglobsow (NEU) and Waldhof (WAL). The urban background site BOS is located in the village of Bösel, about 100 km from the North Sea. It is, therefore, partly influenced by maritime air masses. NEU and WAL are located in forests, representing regional background conditions of the Northern Germany lowlands.

Two stations, Langen (LAN) and Mülheim-Styrum (MST), are located in the western part of Germany. LAN is an urban background site located in the city of Langen at the edge of a residential area and a forest. Emissions from the Frankfurt's Rhein-Main airport (about 5 km to the southeast) might sometimes influence the observations at LAN. MST is situated in the western end of the Ruhr area, the largest urban cluster area in Germany.

Four stations are located in the southern part of Germany, including one urban background site Augsburg (AUG), two low mountain range sites Schauinsland (SCH) and Hohenpeißenberg (HPB), and one high Alpine site Zugspitze (ZSF). AUG is located on the premises of Augsburg's University of Applied Sciences, about 1 km southeast of Augsburg city centre. The two low mountain range sites SCH and HPB are surrounded mainly by forests and agricultural pastures, and are located at the elevations of 1205 and 980 m above sea level (a.s.l.), respectively. The high Alpine site ZSF is located at 2670 m a.s.l. and 300 m below the summit of the Zugspitze, ~~at the south side of~~ the highest mountain in Germany.

2.2 Instrumentation

The technical details of the PNSD and the eBC mass concentration measurements at each GUAN site are summarised in Table 2. More detailed information on the instrumentation and data processing are provided in Birmili et al. (2016). Depending on individual set-up, PNSD are measured either by the Mobility Particle Size Spectrometers (MPSS, Wiedensohler et al., 2012) or by the Dual Mobility Particle Size Spectrometers (D-MPSS). Aerosol dryers are used to dry the aerosol sample to a relative humidity below 40 % (Swietlicki et al., 2008). The PNSD is retrieved from the raw mobility distribution using an inversion algorithm (Pfeifer et al., 2014), including the commonly used bipolar charge distribution (Wiedensohler, 1988). The corrections for particle losses in instruments and inlets are made based on Wiedensohler et al. (2012).

Mass concentrations of eBC are measured using the Multi-Angle Absorption Photometers (MAAP, Thermo Scientific, model 5012), except for AUG, where an Aethalometer (Type 8100, Thermo Fisher Scientific Inc.) is used. For the MAAP measurement, eBC mass concentration is obtained using a mass absorption cross section of $6.6 \text{ m}^2 \text{ g}^{-1}$ at the wavelength of 637 nm (Petzold and Schönlinner, 2004; Müller et al., 2011). eBC data is not available for LAN and MST.

Quality assurance of PNSD measurements in GUAN are periodically done to ensure that measurements remain stable both instrument to instrument (or site to site) and instrument to standard. Monthly maintenance and onsite/laboratory inter-comparisons with a reference MPSS with a frequency between one to four times per year as recommended by Wiedensohler et al. (2018) are done by the World Calibration Centre for Aerosol Physics (WCCAP, <http://www.wmo-gaw-wcc-aerosol-physics.org/>). These procedures can ensure an accuracy of $\pm 10 \%$ for PNCs over the entire measurement period (Birmili et al., 2016). Although the uncertainty of PNCs is comparable or higher than their annual trends (Sect. 3.1), with the application of periodical quality assurance procedures, there should be no monotonicity change or systematic bias in the measurement uncertainties. Therefore, the influence of the measurement uncertainty on the detection of long-term trends of PNCs is assumed to be negligible.

To condense the information provided by PNSD, three particle size ranges, 10–30 nm, 30–200 nm, and 200–800 nm, are chosen to obtain integrated PNCs. $N_{[10-30]}$ represents the nucleation mode in which particles are freshly formed by homogeneous nucleation from either photochemical processes or downstream of traffic exhausts. $N_{[30-200]}$ represents the Aitken mode in which particles are either directly emitted from incomplete combustion or grown by condensational growth. $N_{[200-800]}$ represents the accumulation mode in which particles have undergone condensational growth or cloud processing during transport. As the particles below 20 nm are not measured from 2009 to 2018 at ZSF and MST, we use $N_{[20-800]}$ to represent total PNC in this study instead of $N_{[10-800]}$. Figure 2 illustrates the time span for which data are available at the 16 stations in GUAN during the study period of 2009

to 2018. As a sum-up, Table S1 in the Supplemental Material (SM) gives the number of stations used in trend analysis.

2.3 Trend analysis methods

Environmental data are usually not normally distributed. Therefore, non-parametric methods are often used to detect the long-term trends (e.g., Asmi et al., 2013; Bigi and Ghermandi, 2014; Collaud Coen et al., 2013). Detection of long-term linear trends can be affected by several factors such as time span and time resolution, magnitude of variability, autocorrelation and periodicity of the time series (Weatherhead et al., 1998). To analyse the long-term trends of the PNCs and the eBC mass concentrations, two trend evaluation methods, a customised Sen-Theil trend estimator and the generalized least-square-regression (GLS), were used in this study. Moreover, the regional Mann-Kendall test was used to detect the overall trends over the whole study region. To ensure the comparability of trend slopes among the different sites, relative slopes (absolute slope divided by the median value of the whole time series) in % per year were used.

2.3.1 Customised Sen-Theil trend estimator

The customised Sen-Theil trend estimator (customised Sen's estimator, hereafter) is a modified non-parametric procedure based on the ordinary Sen's slope estimator (Sen, 1968; Theil, 1992; Birmili et al., 2015). This approach can give the true slope of atmospheric parameters by fully considering the effect of their periodic variations, such as, seasonal, weekly, and diurnal cycles, and avoid the influence of outliers, missing values, and statistical distribution of the data. Based on the hourly or daily time series $x(i)$, firstly, the rates of change $m_{i,k}$ of each data pair $[x(i), x(i + k \times 364 \text{ days})]$ is calculated as

$$m_{i,k} = \frac{(x(i+\Delta t) - x(i))}{\Delta t} \quad (1)$$

$\Delta t = k \times 364 \text{ days}$

where k is the integer. Δt ensures that each data point is compared only with data points that are separated by a multiple of 52 weeks (364 days), that is, data points from two different years are compared only if they both were measured on the same hour of the day, day of the week, and season of the year. For our dataset, more than 10000 slopes $m_{i,k}$ are calculated for each time series. The median of those slopes is taken as the true slope for the whole period. Significance and confidence interval (CI) of the trends are determined at 95 % significance level from the distribution of $m_{i,k}$.

2.3.2 Generalized least-square-regression and auto-regressive bootstrap confidence intervals

The second method used to detect the trends is the generalized least-square-regression (GLS) (Mudelsee, 2010; Asmi et al., 2013). A brief introduction of the method is given here, for details refer to Mudelsee (2010) and Asmi et al. (2013). GLS is an approach for estimating the linear parameters in a linear regression model. For a time series of an observed parameter $x(i)$, compactly written as $\{t(i), x(i)\}_{i=1}^n$, the linear regression model can be defined as:

$$x(i) = \beta_1 + \beta_2 t(i) + \Omega(t(i)) + S(i)e(i) \quad (2)$$

where β_1 and β_2 are the two trend parameters (intercept and slope). $\Omega(t(i))$ is a seasonal component. $S(i)$ is a variability function scaling the random noise term $e(i)$. The GLS regresses the trend and seasonal parameters (denoted as β , hereinafter) by minimizing the sum of squares of the residuals ($SSQG$).

$$SSQG(\beta) = (x - T\beta)'V^{-1}(x - T\beta) \quad (3)$$

where T is time, x is observation data, and V is the covariance matrix that can be estimated by Eq. (6) in Asmi et al. (2013).

To evaluate the robustness of the derived slopes, the auto-regressive bootstrap (ARB) method is used to construct the CIs (Mudelsee, 2010, algorithm 3.5). The ARB resamples the white-noise residuals of data using the auto-regressive persistence model, adds the resampled residuals to the fitted data, and re-calculates the slopes. The resampling is repeated 1000 times and the CIs are estimated from these 1000 resampled slopes.

2.3.3 Regional Mann-Kendall test

To detect if an overall increase or decrease exists in the whole region, the regional Mann-Kendall test was also applied in this study. Mann-Kendall test is a commonly used method for detection of long-term trends (Mann 1945; Kendall 1938). It detects the trend using the Kendall's tau test which is known as a rank correlation test and evaluates if a monotonic increasing or decreasing trend exists. If a significant monotonic increase or decrease is detected, the Sen's slope estimator is further used to determine the slope and CI of the time series (Gilbert, 1987). The regional Mann-Kendall test (Helsel and Frans, 2006) is a method adapted from the seasonal Kendall test to determine whether a monotonic increase or decrease can be observed across a large area including multiple locations. For details about the regional Mann-Kendall test refer to Helsel and Frans (2006). Considering the dataset size and calculation efficiency, the monthly median time series was used for the regional Mann-Kendall test in this study.

3 Results and discussion

3.1 Overall trends over the time period 2009–2018

The temporal trends of the observed PNCs and eBC mass concentrations were evaluated using the customised Sen's estimator and GLS/ARB. The daily median time series was used for the customized Sen's estimator, and the monthly median time series was used for GLS/ARB. The relative annual slopes are listed in Table 3. The trend is marked as "statistically significant" (s.s.) in the table if it is statistically significant at 95 % significance level. For the five parameters at the 16 sites (76 trends in total, see Table S1 in the SM), the trends detected by the two methods agree well with each other with seven exceptions that we conclude as no s.s. trends ($N_{[20-800]}$ at MEL, $N_{[10-30]}$ at LMI, LAN, HPB and SCH, and $N_{[200-800]}$ at LAN and BOS). The s.s. negative slopes were found in 14 out of 16 sites for $N_{[30-200]}$ and $N_{[200-800]}$, and $N_{[20-800]}$, in 8 out of 14 sites for $N_{[10-30]}$, and in all sites for eBC mass concentration. The annual slope of the eBC mass concentration varies between -13.1 % and -1.7 % per year. The slopes of the PNCs vary from -17.2 % to -1.7 %, -7.8 % to -1.1 %, and -11.1 % to -1.2 % per year (only the s.s. trends) for 10–30 nm, 30–200 nm, and 200–800 nm particle diameter, respectively. Robustness analysis (see Sect. 1 in SM) suggests that the time span of our dataset is long enough for slope detection.

To evaluate the overall trends of PNCs and eBC mass concentration over Germany, the regional Mann-Kendall test was applied to our dataset and the results are also listed in Table 3. It should be noted that roadside sites might bias the regional Mann-Kendall trends because of their prominent local influence. Moreover, the 13 non-roadside sites in GUAN are not evenly distributed in spatial scale as there are five sites located in the state of Saxony and HPB is only 42 km away from ZSF. To ensure the representativeness of the regional trends, three roadside sites (DDN, LMI and LEI) as well as LTR, ANA, and ZSF were excluded in the regional Mann-Kendall test. The

highest regional decrease of 5 % per year appears in the eBC mass concentration of which anthropogenic emissions are the major source. The regional trends of $N_{[30-200]}$ and $N_{[200-800]}$ are both s.s. negative. In the urban area, $N_{[30-200]}$ and eBC mass concentration are found to be closely related to the emissions from incomplete diesel combustion (Cheng et al., 2013; Krecl et al., 2015). Therefore, the s.s. regional decreases in $N_{[30-200]}$ and eBC mass concentration are very likely to stem from the decrease of anthropogenic emissions in Germany. However, insignificant regional trend was detected for $N_{[10-30]}$. A plausible explanation is that anthropogenic emissions have only minor or negligible influence on $N_{[10-30]}$ in the regional background area due to the short lifetime and high spatial variability of nucleation mode particles (Sun et al., 2019).

The trends of the PNCs and eBC mass concentration in this study are consistent with results from other studies conducted in Europe. Table 4 compares the long-term trends of aerosol concentrations between the present and other studies. Since 2001, the s.s. decrease in BC, PNCs, and $PM_{2.5}$ have been detected for most of the evaluated sites in Table 4. The implementation of emission mitigation policies have been thought to be the dominant impact factors in these studies. Especially, there is one similar study that evaluated the trend of BC mass concentration in Germany for the time period 2005–2014 (Kutzner et al., 2018), in which decreased BC mass concentration was detected in 12 sites. Comparing the two studies, the absolute decreasing trend of BC mass for 2005–2014 is stronger than our results for 2009–2018, which might stem from the difference between the effects of emission mitigation policies in the two study periods.

3.2 Emission change in Germany

Long-term trends of aerosol concentrations on regional scales could occur due to several factors such as emission changes and changes in long-range transport and vertical diffusion due to inter-annual variations of weather and climate. In areas strongly affected by human activities (e.g. traffic, domestic heating, industry etc.), changes in emissions are usually the main cause of the trend of aerosol concentrations. In this section, the long-term variation of total emission of major pollutants in Germany are compared with the trends of aerosol concentrations measured at the six regional background and mountain sites. The emission data is from German Informative Inventory Report (UBA, 2020) produced by the national co-ordination agency for the National System of Emissions Inventories (Nationales Systems Emissionsinventare; NaSE) sited within the German Federal Environmental Agency (UBA), and is based on a large number of sources and publications. Detailed information is available in UBA (2020).

From 2005, the German Federal Government has established German Sustainability Strategy with the goal of reducing the emissions of SO_2 , NO_x , NH_3 , non-methane volatile organic compounds (NMVOC), and $PM_{2.5}$. Figure 3a illustrates the variation of BC total emission in Germany from 2009 to 2017 (black line) and the annual mean eBC mass concentration index (defined as the percentage of the concentration for the year of 2009) for the six regional background and mountain sites (magenta line). From 2009 to 2017, the total emission of BC in Germany decreases about -3.4 % per year and highly agrees with the trend of mean eBC mass concentration. eBC mass concentration is influenced by emission, transport, and scavenging simultaneously. The agreement between the trend of BC total emission and eBC mass concentration suggests that emission reduction is very likely to be the dominant factor for the decreasing of eBC mass concentration over Germany, while other factors do not show a clear contribution.

Other than primary emission, another important process controlling the PNC is the formation of secondary particulate matters. The decreased anthropogenic emissions might reduce the concentration of precursor gases and thus inhibit secondary aerosol formation. Figure 3b illustrates the variation of total emission of $PM_{2.5}$ and some

带格式的: 缩进: 首行缩进: 0.48 厘米

selected precursor gasses, as well as the annual mean PNCs index for the six regional background and mountain sites. It should be noted that PNC is not a conserved parameter and might change rapidly by particle coagulation. Thus, another parameter, particle volume concentration (PVC) in the size range of 20–800 nm ($V_{[200-800]}$), is also shown in Fig. 3b for a better comparability. Total emissions of all precursors and $PM_{2.5}$ except NH_3 decreased around $-2.2 \sim -0.9$ % per year during 2009–2018. However, the measured $N_{[30-200]}$, $N_{[200-800]}$ and $V_{[200-800]}$ decreases about -2.5 % per year, which is stronger than the decreases in the emissions. ~~This might be because the secondary aerosol contributes a large fraction in particulate matters in the regional background settings (Castro et al., 1999).~~ Secondary aerosol formation processes are highly complex and nonlinear, determined not only by the concentrations of precursors but also many other factors such as solar radiation, temperature, humidity, and diffusion conditions etc. ~~Thus, the concentration of secondary aerosol might not follow the variation of precursor concentrations. It was found that secondary aerosol contributes a large fraction in particulate matters in the regional background settings (Castro et al., 1999). The change in secondary aerosol particle concentration might not follow the change of precursor emission.~~ Therefore, larger discrepancies are observed between the emissions and particle concentrations although decrease trends are found in both of them. ~~This might be because the secondary aerosol contributes a large fraction in particulate matters in the regional background settings (Castro et al., 1999).~~

Based on the above results, we believe that the observed trends of PNCs and eBC mass concentration are mainly due to the reduction in emissions. The annual changes of meteorological conditions might have an impact on PNCs, but are not likely to be the decisive impact factor. Detailed discussion on the possible influence of meteorological conditions will be discussed in Sect. 4. The decreased pollutant concentrations are highly associated with the reduced risk of human health. Pope et al. (2009) demonstrated that a decrease of $10 \mu g m^{-3}$ in the $PM_{2.5}$ mass concentration is related with an increase of life expectancy of 0.61 ± 0.20 year in 211 countries. The improved health effects because of decreased UFP and BC would be even greater compared with that of $PM_{2.5}$ mass concentration. As of 2018, 97 % of cities in low- and mid-income countries do not meet the World Health Organization (WHO) air quality guidelines (WHO, 2018). Our result demonstrates that the implementation of proper emission mitigation policies can largely reduce the BC mass concentration and PNC, thus may effectively reduce the health risk in polluted regions.

3.3 Diurnal variation of trends

The emission intensities of some emission sources have distinct diurnal variations, such as that of traffic and residential activities (e.g. domestic heating). The trends based on the subsets of the time series might reflect the impact of these changes. In this section we will analyse the diurnal variations of trends and investigate their connection to the sources.

Figure 4 shows the Sen's slopes of the measured PNCs and eBC mass concentration at each hour of day for each site category. To evaluate the diurnal trends, data pairs belonging to the same hour of day were selected for the calculation of Sen's slope. Figure 4 shows that all the four parameters show distinct diurnal variations at the roadside and urban background sites. At the roadside sites, the decrease in eBC mass concentration, $N_{[10-30]}$ and $N_{[30-200]}$ are much stronger during daytime than during night-time. At the urban background sites, the diurnal trends also show stronger decrease in the morning and evening. Such diurnal patterns are consistent with the diurnal variation of vehicle volume in the urban area where traffic emission is the dominant source of BC and UFP (Ma and Birmili, 2015). As shown in Fig. S2 in the SM, road transport contributes to the highest reduction in total emission of BC, $PM_{2.5}$, and precursors NO_x and NMVOC, from 2009 to 2017. Therefore, it can be concluded that

the daytime strong reduction in PNCs and eBC mass concentration in the urban area is a result of the decrease in road transport emissions. In Germany, the government has made great effort to reduce the emissions due to road transport. The 10th BImSchV (first issued in 1994 and entered into force of recast on 14 December, 2010) regulates the emission requirement for petrol, diesel, and bio-diesel. And the 28th BImSchV (issued in 2004 and amended every year) regulates the type of engines for mobile machinery that can be marketed commercially, ensuring low emissions from new commercial vehicles. Meanwhile, the implementation of the European Emission Standard (EURO standards, https://en.wikipedia.org/wiki/European_emission_standards, last access: 18 September, 2019) starting from 1990s has significantly reduced the emissions from gasoline and diesel engines. Moreover, another regulation LEZ (35th BImSchV) has effectively reduced the traffic emissions by restricting highly polluting vehicles in defined area in the city. Resulting from the above policies, the road transport emissions of BC, PM_{2.5}, NO_x, and NMVOC have significantly decreased during 2009–2017 as shown in Fig. S2. Our results confirm that the reduction in traffic emissions plays a main role in the decreasing trends of eBC mass concentration in Germany, especially in the urban area.

For regional background and low mountain range sites that are far from the road traffic and not directly affected by traffic emissions, trends of eBC mass concentration and PNCs do not show distinct diurnal patterns. However, downward trends are still visible, which stems mainly from the decrease in background concentration in the whole region caused by the reduction in emissions as shown in Fig. S2. For the low mountain range and high Alpine sites, trends of eBC mass concentration, $N_{[30-200]}$ and $N_{[200-800]}$ show weakly diurnal patterns with slightly more reduction in the afternoon. This is mainly because the high-altitude sites might have more chance to merge into the planetary boundary layer (PBL) in the afternoon, resulting in a much stronger influence of anthropogenic emissions.

It is interesting that at the roadside and urban background sites the trend of $N_{[10-30]}$ has similar diurnal pattern as those for $N_{[30-200]}$ and eBC mass concentration, but at regional background and low mountain range sites they look quite different. It is interesting that at the urban sites $N_{[10-30]}$ has similar diurnal pattern as other parameters, but at regional background and low mountain range sites they look quite different. At the regional background sites, $N_{[10-30]}$ shows a maximum average reduction rate of around -1.5 % per year in the afternoon but basically zero trend during the night. New particle formation (NPF) is a dominant source of $N_{[10-30]}$ in the non-urban areas. Ma and Birmili (2015) reported that the annual average contribution of NPF on $N_{[5-20]}$ is 54 % at regional background sites. Also, the contribution of NPF has a diurnal pattern with higher levels in the afternoon and no influence during night. Thus, the diurnal variations in $N_{[10-30]}$ trend in the regional background sites is likely to have resulted from the inter-annual changes in the regional NPF events. At the low mountain range sites, statistically insignificant positive trends are observed. One possible reason is that the low mountain range sites are far from emission sources of nucleation mode particles (e.g. traffic) and NPF is also rare in the areas. Thus the trend of $N_{[10-30]}$ might be more influenced by metrological conditions.

3.4 Seasonal variation of trends

Figure 5 shows the trends of the PNCs and eBC mass concentration for each season. In general, negative trends are found for the five parameters in all seasons. Similar seasonal trend patterns with stronger decreasing rate in winter are detected for all PNCs, which is likely to have been caused by factors that have strong seasonal variation such as domestic heating and/or meteorological conditions. The emission of domestic heating is much stronger during cold season. The 1st BImSchV limited the emission for medium and small combustions (e.g. domestic

heating). Although domestic heating (residential sector) contributes only a minor fraction of the total emission of BC, PM_{2.5}, SO₂ and NMVOC, its absolute decrease from 2009 to 2017 is large and comparable with other sectors (Fig. S2). The least decreasing rates in the PNCs were found in summer. Other than the low residential emission in warm seasons, another reason might be the strong seasonal variations in biogenic emissions (Asmi et al., 2013). Biogenic emissions contribute considerable secondary organic aerosol (SOA) precursors in summer and thus a higher contribution on PNCs. Without any strong long-term variation, the stable contribution of biogenic emissions on PNCs might lower the relative decreasing rates in PNCs in summer. No clear seasonal pattern could be observed for eBC mass concentration because its emission decrease is mostly contributed by the road transport that has no obvious seasonal variation. Long-term change of meteorological parameters might also affect the seasonal trends as well and will be discussed in Sect. 4.

3.5 Evaluation of low emission zones

As discussed in the previous sections, the reduction in total emissions could be reflected directly in the long-term trends of the aerosol observation in Germany, suggesting that long-term observation network with different types of site is an effective tool to verify the effectiveness of emission control policies. However, the observed decrease in trends is a combined result of the various emission mitigation policies. A question raised is that can such long-term observation network reflect the effect of a specific emission mitigation policy. LEZ is believed to be a good candidate for such an evaluation due to several reasons. Firstly, LEZ has usually a clear introduced date in each city. Secondly, traffic emissions are the major source of aerosol particles in the urban area. Thirdly, traffic emission sources are basically evenly distributed in the urban area and therefore, its contribution of particulate matter in the urban area will not be strongly influenced by wind direction. In this section, we will analyse the effects of two LEZs in the urban area based on our dataset.

LEZ is an urban access regulation in Europe and is one of the key ways to reduce traffic emissions in urban areas. In Germany, high-, medium-, and low-emitting vehicles are required to be marked with red, yellow or green colour stickers on the front window shield. The green sticker denotes the diesel vehicles with at least Euro 4 or Euro 3 engines with a particular filter and petrol vehicles meeting at least Euro 1 standard. Vehicles with green stickers have lowest emissions and can enter all LEZs. Vehicles with other stickers, meaning higher emissions, are restricted. In Germany, the first LEZ was launched in 2008 in Berlin. As of November 2019, LEZs are implemented in over 60 cities. Short-term studies have shown that LEZ can immediately reduce the pollutant levels after the implementation (Rasch et al., 2013; Qadir et al., 2013; Jones et al., 2012).

We select two cities that have implemented LEZ during 2009–2018 and have measurements of both PNCs and eBC mass concentration: Leipzig and Augsburg. Figure 6 illustrates the deseasonalised time series of monthly averaged parameters measured at the two urban background sites AUG and LTR, by subtracting the seasonal cycle from the mean monthly time series. The horizontal dashed lines in Fig. 6 denote the mean values with respect to different LEZ stages. In Augsburg, the first, second, and third stages of the LEZ were implemented respectively on 1 July, 2009 (red dashed line), 1 January, 2011 (yellow dashed line), and 1 June, 2016 (green dashed line). Figure 6a and b show that the eBC mass concentration and $N_{[30-200]}$ have gradually decreased after the implementation of each of new stages of LEZ. And the differences between the concentration levels at different stages are statistically significant at 95% significance level. However, the difference between stage 2 and 3 is relatively negligible. A possible reason is that the third stage of LEZ came into force in June 2016 in Augsburg. By June 2016, around 52 cities in Germany had already implemented the third stage of LEZ, which accelerates

the fleet update in the whole country (also in Augsburg). Thus, no large difference could be seen between the second and third stage of LEZ in the city of Augsburg. It is worth noticing that even with the seasonal variation subtracted from the time series, the amplitude of short-term variations of eBC mass concentration and $N_{[30-200]}$ are still very large mostly due to variations in meteorological conditions. Sometimes it is even larger than the concentration decrease caused by the implementation of LEZ. This indicates that short-term measurement might be influenced largely by the variations in meteorological conditions and long-term measurements are necessary for a trustworthy verification of LEZs.

The LEZ was entered into force in the city of Leipzig directly on third level on 1 March, 2011. Statistically significant differences are found between the concentration levels before and after the implementation of LEZ for both $N_{[30-200]}$ and eBC mass concentration. But the decrease in $N_{[30-200]}$ after 2010 (Fig. 6c) is much larger than that in eBC mass concentration (Fig. 6d). Clear decrease is observed in $N_{[30-200]}$ after 2010 (Fig. 6c) but nearly invisible in eBC mass concentration (Fig. 6d). In Leipzig, the PNCs and eBC mass concentrations were measured at both roadside and urban background sites simultaneously, which provide us the possibility to directly detect the traffic contribution by evaluating the increment of the aerosol concentration (the difference of the concentrations between the traffic and the background sites). The effects of background variation, other sources and meteorological factors can be ignored. Figure 7 illustrates the annually averaged diurnal cycles of the increments of eBC mass concentration (ΔeBC), $N_{[30-200]}$ ($\Delta N_{[30-200]}$) and $N_{[200-800]}$ ($\Delta N_{[200-800]}$). Before and after the implementation of LEZ (2010 and 2011), the ~~increments PNCs and eBC mass concentration~~ show a sudden decrease of up to 40 % during daytime. ~~The average ΔeBC , $\Delta N_{[30-200]}$ and $\Delta N_{[200-800]}$. The mean concentration of eBC mass, $N_{[30-200]}$ and $N_{[200-800]}$~~ during working hours (06:00 to 18:00 local time) in 2010 are respectively about 1.63, 1.33, and 1.58 times higher than those in 2011. After 2011, these aerosol variables continued to decrease, mainly due to the continuous update of vehicle fleet. This result suggests that with a multiple-site network, the effect of emission control policy could be better detected from the increments between near-source and background sites.

4 Meteorological influences on the trends of particle number and eBC mass concentration

Meteorological conditions also influence the concentration of aerosol particles (Birmili et al., 2001; Spindler et al., 2013; von Bismarck-Osten et al., 2013; Hussein et al., 2006) and their inter-annual changes might modify the trends of the parameters studied. In this section, the potential influence of meteorological conditions, including precipitation, temperature, wind speed, and air mass types are discussed.

4.1 Influence of precipitation, temperature, wind speed on the detected trends

Table 5 provides the long-term trends of precipitation, ambient temperature, and wind speed during 2009–2018 based on the 76 measuring sites distributed all over Germany. The trends of the parameters were evaluated by the customised Sen's estimator. The data was obtained from Germany's National Meteorological Service (Deutscher Wetterdienst, DWD). The 76 DWD sites are grouped into three categories: urban background, regional background, and mountain area. The trends in all three meteorological parameters agree well among the three categories. Temperature shows a negligible change in spring, a slight increase in summer, and a larger increase up to 0.43 °C per year in autumn and winter. Increased temperature during cold seasons might have led to lower anthropogenic emissions from domestic heating and power generation, and further led to a decrease in PNCs and eBC mass

设置了格式: 字体: 倾斜

设置了格式: 下标

设置了格式: 字体: 倾斜

设置了格式: 下标

concentrations. Precipitation presents a s.s. decreasing trend up to about -6% per year in summer, which might inhibit the wet deposition of aerosol particles and diminish the reduction of eBC mass concentrations and $N_{[200-800]}$ to a certain extent. No obvious trend is observed in wind speed. In summary, increased ambient temperature might contribute to the decrease of the PNCs and eBC mass concentrations shown in Sect. 3.1 and 3.4 by indirectly influencing anthropogenic emissions. While decreased precipitation in summer might diminish the decrease of the PNCs and eBC mass concentration by inhibiting aerosol wet deposition.

4.2. Influence of air mass condition on the detected trends

Synoptic-scale air mass condition, including origin region and pathways is an important factor driving regional pollutant concentration (Ma et al., 2014; Hussein et al., 2006). Atmospheric stability is also important as it dominates the vertical dilution of pollutants. Based on a self-developed back-trajectory cluster method (BCLM), the influences of the inter-annual changes in air mass conditions and atmospheric stability on the detected trends are investigated.

BCLM is based on a joint cluster analysis considering air mass backward trajectories, profiles of pseudo-potential temperature, and PM_{10} mass concentration over Germany (Birmili et al., 2010; Engler et al., 2007; Ma et al., 2014). In BCLM, 15 air mass types are defined to represent different meteorological conditions on a scale of Germany. Detailed information about data preparation, cluster processing, and the data procedures and data products is described in Sect. 3 of SM and in Ma et al. (2014). Figure 8 shows the average trajectories and the normalized profiles of pseudo potential temperature (θ_p) for the 15 air mass types. The 15 air mass types are named by seasons (CS: cold season; TS: transition season; and WS: warm season) and synoptic patterns (ST: Stagnant; A1: Anti-cyclonic with air mass originating from Eastern Europe; A2: Anti-cyclonic with air mass originating from west; C1: cyclonic with air mass originating from relatively south; C2: cyclonic with air mass originating from the north). Table 6 lists the basic statistical information of the 15 air mass types.

Figure 9 illustrates the statistics of PNCs and eBC mass concentrations for each air mass type at the regional background site category (MEL, WAL and NEU). Large differences in the mean values of $N_{[200-800]}$ and eBC mass concentrations are observed among different air mass types; whereas $N_{[10-30]}$ and $N_{[30-200]}$ show less significant difference as $N_{[10-30]}$ and $N_{[30-200]}$ represent more local information and are not as sensitive as $N_{[200-800]}$ and eBC mass concentration. In the following discussion, only $N_{[200-800]}$ and eBC mass concentration are used.

Due to the high sensitivity of $N_{[200-800]}$ and eBC mass concentration on the air mass types, frequency changes of air mass types might lead to changes in their long-term trends. However, it is difficult to investigate the influence for each air mass type since the frequencies of the air mass types are quite low ($3.0\% \sim 12.0\%$). Therefore, the 15 air mass types are grouped into two categories according to pollution level. If both the median eBC mass concentration and $N_{[200-800]}$ are higher than their overall median concentration, the air mass is grouped into polluted air mass category, and vice versa.

(1): Polluted air mass category that includes CS-ST, CS-A1, CS-A2, CS-C1, TS-A1, WS-ST, WS-A1, and WS-C1;

(2): Cleaner air mass category that includes CS-C2a, CS-C2b, TS-A2, TS-C1, TS-C2, WS-A2, and WS-C2.

The annual occurrence of polluted air mass category together with the annual mean values of $N_{[200-800]}$ and eBC mass concentration at the regional background and low mountain range sites are shown in Fig. 10. No clear trend of the occurrence of polluted air mass category could be found. However, large difference is visible in the occurrences between different years. If the change in air mass frequency plays an important role in the variations

in $N_{[30-200]}$ and eBC mass concentration, low polluted air mass occurrences should be associated with relatively low $N_{[200-800]}$ and eBC mass concentration. However, such a relationship is not visible in Fig. 10. The annual mean values of $N_{[200-800]}$ and eBC mass concentration (black lines in Fig. 10) decrease consistently. Therefore, it could be concluded that the inter-annual changes in synoptic-scale air mass conditions are not the reason for the decrease of pollutant concentrations shown in Sect. 3.

5 Conclusions

In this study, long-term trends of atmospheric PNCs and eBC mass concentration over a 10 years period (2009–2018) are determined for 16 sites in the GUAN, ranging from roadside to high Alpine environments. Overall, statistically significant decreasing trends are found for 85 % of the parameters and observation sites. Concretely, the annual slope of the eBC mass concentration varies between -13.1 % and -1.7 % per year. The slopes of the PNCs vary from -17.2 % to -1.7 %, -7.8 % to -1.1 %, and -11.1 % to -1.2 % per year for 10–30 nm, 30–200 nm, and 200–800 nm size ranges, respectively. The regional Mann-Kendall test yields regional-scale trends of eBC mass concentration, $N_{[30-200]}$, and $N_{[200-800]}$ of -5 %, -2.5 % and -2.9 % per year, respectively, indicating an overall decreasing trend in sub-micrometre PNC (except $N_{[10-30]}$) and eBC mass concentration all over Germany.

Comparing the trends of measured parameters with the long-term change in total emission, we believe that the observed trends of PNCs and eBC mass concentrations are mainly due to the emission reduction. The decreasing trend of eBC mass concentration agrees well with the variation of BC total emission in Germany, suggesting reduction in emissions is the dominant factor for the reduction in eBC mass concentrations over the country. The decreasing rates of $N_{[30-200]}$, $N_{[200-800]}$, and $V_{[200-800]}$ are stronger than the decrease in the total emissions of all precursors and $PM_{2.5}$, which could have been caused by the highly complex and nonlinear processes of secondary aerosol formation.

The highest decrease in eBC mass concentration was observed during daytime at the roadside and urban background, implying a strong evidence of reduced traffic emissions in the urban area. When there are fewer motor vehicles at night, the PNCs and eBC mass concentration in the urban sites also show a significant decrease, which could be due to the background concentration decrease caused by the reduction in other emissions such as domestic heating, industry, etc. Stronger reductions in PNCs are found in winter, which is likely to be caused by the decreased emissions from domestic heating combustion.

Meteorological conditions are also able to influence the temporal variation of aerosol concentration. The inter-annual changes in precipitation and temperature might have some influences on the detected trends by indirectly influencing anthropogenic emissions and inhibiting aerosol wet deposition, but they seem to be not the dominant factors for the long-term decrease of the measured parameters. The influence of long-range transport pattern is also evaluated and the inter-annual changes in synoptic-scale air mass conditions are found to be not the reason for the decrease in pollutant concentrations.

This study suggests that a combination of emission mitigation policies can effectively improve the air quality on large spatial scales such as in Germany. Given the relative novelty of the long-term measurements (PNSD, BC) in a network such as GUAN, the results proved to be robust and comprehensive. Our study also shows that long-term measurements of aerosol parameters in different environments could be very instrumental in detecting and understanding the long-term effects of emission mitigation policies.

511
512 **Acknowledgement.** We acknowledge funding by the German Federal Environment Ministry (BMU) grants F&E
513 370343200 (German title: Erfassung der Zahl feiner und ultrafeiner Partikel in der Außenluft) from 2008 to 2010,
514 and F&E 371143232 (German title: Trendanalysen gesundheitsgefährdender Fein- und Ultrafeinstaubfraktionen
515 unter Nutzung der im German Ultrafine Aerosol Network (GUAN) ermittelten Immissionsdaten durch Fortführung
516 und Interpretation der Messreihen) from 2012 to 2014. For the MST (Mülheim-Styrum) measurements, we thank
517 the co-funding by the North Rhine-Westphalia Agency for Nature, Environment and Consumer Protection
518 (LANUV). Measurements at Annaberg-Buchholz were supported by the EU-Ziel3 project UltraSchwarz (German
519 title: Ultrafeinstaub und Gesundheit im Erzgebirgskreis und Region Usti), grant 100083657. Measurements at
520 DDW (Dresden-Winkelmannstraße) were co-funded by the European Regional Development Fund Financing
521 Programme Central Europe, grant No. 3CE288P (UFIREG). Measurements in AUG (Augsburg) were funded
522 partly also by UFIREG and by the Helmholtz-Zentrum.

523 The authors would like to thank the technical and scientific staff members of the stations included in this study.
524 André Sonntag and Stephan Nordmann (TROPOS/UBA) contributed to data processing. Prof. Dr. Thomas A.J.
525 Kuhlbusch and Dr. Ulrich Quass contributed the data quality assurance and data analysis at MST. Horst-Günther
526 Kath (State Dept. for Environmental and Agricultural Operations in Saxony, Betriebsgesellschaft für Umwelt und
527 Landwirtschaft – BfUL), Andreas Hainsch (Labour Inspectorate of Lower Saxony, Staatliches
528 Gewerbeaufsichtsamt Hildesheim – GAA), and Dieter Glatke (Agency for Nature Protection, the Environment,
529 and Customer Protection in North Rhine-Westphalia, Landesamt für Natur, Umwelt und Verbraucherschutz
530 Nordrhein-Westfalen – LANUV) made the GUAN measurements possible at their respective observations sites.
531 We also thank Werner Wunderlich in Hessian State Office for Nature Conservation, Environment and Geology
532 for the eBC mass concentration data at Raunheim and Karin Uhse at German Environment Agency (UBA) for the
533 PNCs data quality check at LAN. We thank Andreas Rudolph, Dustin Konzack and Andreas Fischer at TOPAS
534 GmbH, Dresden, for kindly providing the UFP-monitor TSI 3031 at LAN (data 2015 – 2018), and yearly quality
535 assurance checks.

536 This work was also accomplished in the frame of the project ACTRIS-2 (Aerosols, Clouds, and Trace gases
537 Research InfraStructure) under the European Union—Research Infrastructure Action in the frame of the H2020
538 program for “Integrating and opening existing national and regional research infrastructures of European interest”
539 under Grant Agreement N654109 (Horizon 2020). Additionally, we acknowledge the WCCAP (World Calibration
540 Centre for Aerosol Physics) as part of the WMO-GAW program base-funded by the UBA.

541

References

- Asmi, A., Collaud Coen, M., Ogren, J. A., Andrews, E., Sheridan, P., Jefferson, A., Weingartner, E., Baltensperger, U., Bukowiecki, N., Lihavainen, H., Kivekas, N., Asmi, E., Aalto, P. P., Kulmala, M., Wiedensohler, A., Birmili, W., Hamed, A., O'Dowd, C., Jennings, S. G., Weller, R., Flentje, H., Fjaeraa, A. M., Fiebig, M., Myhre, C. L., Hallar, A. G., Swietlicki, E., Kristensson, A., and Laj, P.: Aerosol decadal trends - Part 2: In-situ aerosol particle number concentrations at GAW and ACTRIS stations, *Atmos. Chem. Phys.*, 13, 895-916, 10.5194/acp-13-895-2013, 2013.
- Barnpadimos, I., Hueglin, C., Keller, J., Henne, S., and Prévôt, A. S. H.: Influence of meteorology on PM₁₀ trends and variability in Switzerland from 1991 to 2008, *Atmos. Chem. Phys.*, 11, 1813-1835, 10.5194/acp-11-1813-2011, 2011.
- Bigi, A., and Ghermandi, G.: Long-term trend and variability of atmospheric PM₁₀ concentration in the Po Valley, *Atmos. Chem. Phys.*, 14, 4895-4907, 10.5194/acp-14-4895-2014, 2014.
- Birmili W, Wiedensohler A, Heintzenberg J, Lehmann K. Atmospheric particle number size distribution in central Europe: Statistical relations to air masses and meteorology. *Journal of Geophysical Research: Atmospheres*. 106(D23), 32005-18, 2001.
- Birmili, W., Heinke, K., Pitz, M., Matschullat, J., Wiedensohler, A., Cyrys, J., Wichmann, H. E., and Peters, A.: Particle number size distributions in urban air before and after volatilisation, *Atmos. Chem. Phys.*, 10, 4643-4660, 10.5194/acp-10-4643-2010, 2010.
- Birmili, W., Sun, J., Weinhold, K., Merkel, M., Rasch, F., Wiedensohler, A., Bastian, S., Löschau, G., Schladitz, A., Quass, U., Kuhlbusch, T. A. J., Kaminski, H., Cyrys, J., Pitz, M., Gu, J., Peters, A., Flentje, H., Meinhardt, F., Schwerin, A., Bath, O., Ries, L., Gerwig, H., Wirtz, K., and Weber, S.: Atmospheric aerosol measurements in the German Ultrafine Aerosol Network (GUAN) – Part 3: Black Carbon mass and particle number concentrations 2009 to 2014, *Gefährst. Reinh. Luft*, 75, 2015.
- Birmili, W., Weinhold, K., Rasch, F., Sonntag, A., Sun, J., Merkel, M., Wiedensohler, A., Bastian, S., Schladitz, A., Löschau, G., Cyrys, J., Pitz, M., Gu, J., Kusch, T., Flentje, H., Quass, U., Kaminski, H., Kuhlbusch, T. A. J., Meinhardt, F., Schwerin, A., Bath, O., Ries, L., Gerwig, H., Wirtz, K., and Fiebig, M.: Long-term observations of tropospheric particle number size distributions and equivalent black carbon mass concentrations in the German Ultrafine Aerosol Network (GUAN), *Earth Syst. Sci. Data*, 8, 355-382, 10.5194/essd-8-355-2016, 2016.
- Castro, L.M., Pio, C.A., Harrison, R.M., Smith, D.J.T.: Carbonaceous aerosol in urban and rural European atmospheres: estimation of secondary organic carbon concentrations, *Atmospheric Environment*, 33, 2771-2781, [https://doi.org/10.1016/S1352-2310\(98\)00331-8](https://doi.org/10.1016/S1352-2310(98)00331-8), 1999.
- Chen, X., Zhang, Z., Engling, G., Zhang, R., Tao, J., Lin, M., Sang, X., Chan, C., Li, S., and Li, Y.: Characterization of fine particulate black carbon in Guangzhou, a megacity of South China, *Atmospheric Pollution Research*, 5, 361-370, <https://doi.org/10.5094/APR.2014.042>, 2014.
- Cheng, Y.H., Shiu, B.T., Lin, M.H. and Yan, J.W.: Levels of black carbon and their relationship with particle number levels—observation at an urban roadside in Taipei City. *Environmental Science and Pollution Research*, 20(3), 1537-1545, 2013.
- Cohen, A. J., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., and Dandona, R.: Estimates and 25-year trends of the global burden of disease attributable to

582 ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015, *The Lancet*, 389,
 583 1907-1918, 2017.

584 Collaud Coen, M., Andrews, E., Asmi, A., Baltensperger, U., Bukowiecki, N., Day, D., Fiebig, M., Fjaeraa, A. M.,
 585 Flentje, H., Hyvarinen, A., Jefferson, A., Jennings, S. G., Kouvarakis, G., Lihavainen, H., Myhre, C. L., Malm,
 586 W. C., Mihalopoulos, N., Molenaar, J. V., O'Dowd, C., Ogren, J. A., Schichtel, B. A., Sheridan, P., Virkkula,
 587 A., Weingartner, E., Weller, R., and Laj, P.: Aerosol decadal trends - Part 1: In-situ optical measurements at
 588 GAW and IMPROVE stations, *Atmos. Chem. Phys.*, 13, 869-894, 10.5194/acp-13-869-2013, 2013.

589 Engler, C., Rose, D., Wehner, B., Wiedensohler, A., Brüggemann, E., Gnauk, T., Spindler, G., Tuch, T., and
 590 Birmili, W.: Size distributions of non-volatile particle residuals ($D_p < 800$ nm) at a rural site in Germany and
 591 relation to air mass origin, *Atmos. Chem. Phys.*, 7, 5785-5802, 10.5194/acp-7-5785-2007, 2007.

592 European Environment Agency, EEA: Air quality in Europe – 2017 report, Luxembourg, 74 pp., available at:
 593 <https://www.eea.europa.eu/publications/air-quality-in-europe-2017>, 2017.

594 Gilbert, R. O.: Statistical methods for environmental pollution monitoring, John Wiley & Sons, 1987.

595 Helsel, D. R., and Frans, L. M.: Regional Kendall Test for Trend, *Environmental Science & Technology*, 40, 4066-
 596 4073, 10.1021/es051650b, 2006.

597 Hussein, T., Karppinen, A., Kukkonen, J., Härkönen, J., Aalto, P.P., Hämeri, K., Kerminen, V.M. and Kulmala,
 598 M.: Meteorological dependence of size-fractionated number concentrations of urban aerosol particles.
 599 *Atmospheric Environment*, 40(8), 1427-1440, 2006.

600 Janssen, N. A., Gerlofs-Nijland, M. E., Lanki, T., Salonen, R. O., Cassee, F., Hoek, G., Fischer, P., Brunekreef,
 601 B., and Krzyzanowski, M.: Health effects of black carbon, WHO, 86, 2012.

602 Jones, A. M., Harrison, R. M., Barratt, B., and Fuller, G.: A large reduction in airborne particle number
 603 concentrations at the time of the introduction of “sulphur free” diesel and the London Low Emission Zone,
 604 *Atmos. Environ.*, 50, 129-138, <http://dx.doi.org/10.1016/j.atmosenv.2011.12.050>, 2012.

605 Kendall, M. G.: A new measure of rank correlation, *Biometrika*, 30, 81-93, 1938.

606 Krecl, P., Targino, A. C., Johansson, C., and Ström, J.: Characterisation and source apportionment of submicron
 607 particle number size distributions in a busy street canyon, *Aerosol Air Qual. Res.*, 15, 220-233, 2015.

608 Kreyling, W. G., Semmler-Behnke, M., and Möller, W.: Health implications of nanoparticles, *Journal of*
 609 *Nanoparticle Research*, 8, 543-562, 10.1007/s11051-005-9068-z, 2006.

610 Kutzner, R. D., von Schneidmesser, E., Kuik, F., Quedenau, J., Weatherhead, E. C., and Schmale, J.: Long-term
 611 monitoring of black carbon across Germany, *Atmos. Environ.*, 185, 41-52,
 612 <https://doi.org/10.1016/j.atmosenv.2018.04.039>, 2018.

613 Ma, N., and Birmili, W.: Estimating the contribution of photochemical particle formation to ultrafine particle
 614 number averages in an urban atmosphere, *Science of The Total Environment*, 512-513, 154-166,
 615 <http://dx.doi.org/10.1016/j.scitotenv.2015.01.009>, 2015.

616 Ma, N., Birmili, W., Müller, T., Tuch, T., Cheng, Y. F., Xu, W. Y., Zhao, C. S., and Wiedensohler, A.:
 617 Tropospheric aerosol scattering and absorption over central Europe: a closure study for the dry particle state,
 618 *Atmos. Chem. Phys.*, 14, 6241-6259, 10.5194/acp-14-6241-2014, 2014.

619 Mann, H. B.: Nonparametric tests against trend, *Econometrica: Journal of the Econometric Society*, 245-259, 1945.

620 Masiol, M., Squizzato, S., Chalupa, D. C., Utell, M. J., Rich, D. Q., and Hopke, P. K.: Long-term trends in
 621 submicron particle concentrations in a metropolitan area of the northeastern United States, *Science of The*
 622 *Total Environment*, 633, 59-70, <https://doi.org/10.1016/j.scitotenv.2018.03.151>, 2018.

623 Mudelsee, M.: *Climate Time Series Analysis: Classical Statistical and Bootstrap Methods.*, Springer, 2010.

624 Murphy, D. M., Chow, J. C., Leibensperger, E. M., Malm, W. C., Pitchford, M., Schichtel, B. A., Watson, J. G.,
625 and White, W. H.: Decreases in elemental carbon and fine particle mass in the United States, *Atmos. Chem.*
626 *Phys.*, 11, 4679-4686, 10.5194/acp-11-4679-2011, 2011.

627 Müller, T., Henzing, J. S., de Leeuw, G., Wiedensohler, A., Alastuey, A., Angelov, H., Bizjak, M., Collaud Coen,
628 M., Engström, J. E., Gruening, C., Hillamo, R., Hoffer, A., Imre, K., Ivanow, P., Jennings, G., Sun, J. Y.,
629 Kalivitis, N., Karlsson, H., Komppula, M., Laj, P., Li, S. M., Lunder, C., Marinoni, A., Martins dos Santos, S.,
630 Moerman, M., Nowak, A., Ogren, J. A., Petzold, A., Pichon, J. M., Rodriguez, S., Sharma, S., Sheridan, P. J.,
631 Teinilä, K., Tuch, T., Viana, M., Virkkula, A., Weingartner, E., Wilhelm, R., and Wang, Y. Q.:
632 Characterization and intercomparison of aerosol absorption photometers: result of two intercomparison
633 workshops, *Atmos. Meas. Tech.*, 4, 245-268, 10.5194/amt-4-245-2011, 2011.

634 Pérez, N., Pey, J., Cusack, M., Reche, C., Querol, X., Alastuey, A., and Viana, M.: Variability of particle number,
635 black carbon, and PM₁₀, PM_{2.5}, and PM₁ levels and speciation: influence of road traffic emissions on urban air
636 quality, *Aerosol Science and Technology*, 44, 487-499, 2010.

637 Petzold, A., and Schönlinner, M.: Multi-angle absorption photometry—a new method for the measurement of
638 aerosol light absorption and atmospheric black carbon, *Journal of Aerosol Science*, 35, 421-441, 2004.

639 Pfeifer, S., Birmili, W., Schladitz, A., Müller, T., Nowak, A., and Wiedensohler, A.: A fast and easy-to-implement
640 inversion algorithm for mobility particle size spectrometers considering particle number size distribution
641 information outside of the detection range, *Atmos. Meas. Tech.*, 7, 95-105, 10.5194/amt-7-95-2014, 2014.

642 Pope, C. A., Ezzati, M., and Dockery, D. W.: Fine-Particulate Air Pollution and Life Expectancy in the United
643 States, *New England Journal of Medicine*, 360, 376-386, 10.1056/NEJMsa0805646, 2009.

644 Putaud, J., Cavalli, F., Martins dos Santos, S., and Dell'Acqua, A.: Long-term trends in aerosol optical
645 characteristics in the Po Valley, Italy, *Atmos. Chem. Phys.*, 14, 9129-9136, 2014.

646 Qadir, R. M., Abbaszade, G., Schnelle-Kreis, J., Chow, J. C., and Zimmermann, R.: Concentrations and source
647 contributions of particulate organic matter before and after implementation of a low emission zone in Munich,
648 Germany, *Environ. Pollut.*, 175, 158-167, <http://dx.doi.org/10.1016/j.envpol.2013.01.002>, 2013.

649 Rasch, F., Birmili, W., Weinhold, K., Nordmann, S., Sonntag, A., Spindler, G., Herrmann, H., Wiedensohler, A.,
650 and Löschau, G.: Significant reduction of ambient black carbon and particle number in Leipzig as a result of
651 the low emission zone, *Gefährst. Reinh. Luft*, 73, 2013.

652 Schladitz, A., Leníček, J., Beneš, I., Kováč, M., Skorkovský, J., Soukup, A., Jandlová, J., Poulain, L., Plachá, H.,
653 Löschau, G., and Wiedensohler, A.: Air quality in the German–Czech border region: A focus on harmful
654 fractions of PM and ultrafine particles, *Atmos. Environ.*, 122, 236-249,
655 <http://dx.doi.org/10.1016/j.atmosenv.2015.09.044>, 2015.

656 Schmid, O., and Stoeger, T.: Surface area is the biologically most effective dose metric for acute nanoparticle
657 toxicity in the lung, *Journal of Aerosol Science*, 99, 133-143, 2016.

658 Seaton, A., Godden, D., MacNee, W., and Donaldson, K.: Particulate air pollution and acute health effects, *The*
659 *Lancet*, 345, 176-178, [https://doi.org/10.1016/S0140-6736\(95\)90173-6](https://doi.org/10.1016/S0140-6736(95)90173-6), 1995.

660 Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall's Tau, *Journal of the American Statistical*
661 *Association*, 63, 1379-1389, 10.1080/01621459.1968.10480934, 1968.

662 Singh, A., Bloss, W. J., and Pope, F. D.: 60 years of UK visibility measurements: impact of meteorology and
663 atmospheric pollutants on visibility, *Atmos. Chem. Phys.*, 17, 2085-2101, 10.5194/acp-17-2085-2017, 2017.

664 Spindler, G., Gr ner, A., M ller, K., Schlimper, S., and Herrmann, H.: Long-term size-segregated particle (PM₁₀,
665 PM_{2.5}, PM₁) characterization study at Melpitz -- influence of air mass inflow, weather conditions and season,
666 J Atmos Chem, 70, 165-195, 10.1007/s10874-013-9263-8, 2013.

667 Sun, J., Birmili, W., Hermann, M., Tuch, T., Weinhold, K., Spindler, G., Schl ditz, A., Bastian, S., L schau, G.,
668 Cyrys, J., Gu, J., Flentje, H., Briel, B., Asbach, C., Kaminski, H., Ries, L., Sohmer, R., Gerwig, H., Wirtz, K.,
669 Meinhardt, F., Schwerin, A., Bath, O., Ma, N., and Wiedensohler, A.: Variability of black carbon mass
670 concentrations, sub-micrometer particle number concentrations and size distributions: results of the German
671 Ultrafine Aerosol Network ranging from city street to High Alpine locations, Atmos. Environ., 202, 256-268,
672 <https://doi.org/10.1016/j.atmosenv.2018.12.029>, 2019.

673 Swietlicki, E., Hansson, H.-C., H meri, K., Svenningsson, B., Massling, A., McFiggans, G., McMurry, P., Pet j ,
674 T., Tunved, P., and Gysel, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles
675 measured with H-TDMA instruments in various environments—a review, Tellus B: Chemical and Physical
676 Meteorology, 60, 432-469, 2008.

677 Theil, H.: A Rank-Invariant Method of Linear and Polynomial Regression Analysis, in: Henri Theil's
678 Contributions to Economics and Econometrics: Econometric Theory and Methodology, edited by: Raj, B., and
679 Koerts, J., Springer Netherlands, Dordrecht, 345-381, 1992.

680 [German Federal Environmental Agency, UBA: German Informative Inventory Report 2020 \(IIR 2020\). available](http://iir-de.wikidot.com/)
681 [at <http://iir-de.wikidot.com/>; access: 1 March 2020.](http://iir-de.wikidot.com/)

682 von Bismarck-Osten, C., Birmili, W., Ketzel, M., Massling, A., Pet j , T., and Weber, S.: Characterization of
683 parameters influencing the spatio-temporal variability of urban particle number size distributions in four
684 European cities, Atmos. Environ., 77, 415-429, <http://dx.doi.org/10.1016/j.atmosenv.2013.05.029>, 2013.

685 Weatherhead, E. C., Reinsel, G. C., Tiao, G. C., Meng, X. L., Choi, D., Cheang, W. K., Keller, T., DeLuisi, J.,
686 Wuebbles, D. J., and Kerr, J. B.: Factors affecting the detection of trends: Statistical considerations and
687 applications to environmental data, Journal of Geophysical Research: Atmospheres (1984–2012), 103, 17149-
688 17161, 1998.

689 World Health Organization, WHO: WHO Global Ambient Air Quality Database (update 2018). available at:
690 <https://www.who.int/airpollution/data/cities/en/>; last access: 12 December 2019.

691 Wiedensohler, A.: An approximation of the bipolar charge distribution for particles in the submicron range, J.
692 Aerosol Sci., 19, 387-389, 1988.

693 Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer,
694 S., Fiebig, M., F j raa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., L j, P., Aalto, P., Ogren,
695 J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., H glin, C., Fierz-Schmidhauser, R., Gysel, M.,
696 Weingartner, E., Riccobono, F., Santos, S., Gr ning, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C.,
697 Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H. G., Keck, L., Jiang, J., Scheckman, J., McMurry, P.
698 H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., L schau, G., and Bastian, S.: Mobility
699 particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality
700 long-term observations of atmospheric particle number size distributions, Atmos. Meas. Tech., 5, 657-685,
701 10.5194/amt-5-657-2012, 2012.

702 Wiedensohler, A., Wiesner, A., Weinhold, K., Birmili, W., Hermann, M., Merkel, M., M ller, T., Pfeifer, S.,
703 Schmidt, A., and Tuch, T.: Mobility particle size spectrometers: Calibration procedures and measurement
704 uncertainties, Aerosol Science and Technology, 52, 146-164, 2018.

设置了格式: 字体: 非加粗

705 Table 1: Basic information on the 16 sites in the German Ultrafine Aerosol Network (GUAN), in alphabetic order.

No.	Site name	Abbreviation	Status (Until 2017)	Site category	Elevation (m)	Location
1	Annaberg-Buchholz	ANA	In operation	Urban background	545	50°34'18" N, 12°59'56" E
2	Augsburg	AUG	In operation	Urban background	485	48°21'29" N, 10°54'25" E
3	Bösel	BOS	Terminated end of 2014	Urban background	17	52°59'53" N, 07°56'34" E
4	Dresden-Nord	DDN	In operation	Roadside	116	51°03'54" N, 13°44'29" E
5	Dresden- Winckelmann-straße	DDW	In operation	Urban background	120	51°02'10" N, 13°43'50" E
6	Hohenpeißenberg	HPB	In operation	Low mountain range	980	47°48'06" N, 11°00'34" E
7	Langen	LAN	In operation	Urban background	130	50°00'18" N, 08°39'05" E
8	Leipzig- Eisenbahnstraße	LEI	In operation	Roadside	120	51°20'45" N, 12°24'23" E
9	Leipzig-Mitte	LMI	In operation	Roadside	111	51°20'39" N, 12°22'38" E
10	Leipzig-TROPOS	LTR	In operation	Urban background	126	51°21'10" N, 12°26'03" E
11	Melpitz	MEL	In operation	Regional background	86	51°31'32" N, 12°55'40" E
12	Mülheim-Styrum	MST	In operation	Urban background	37	51°27'17" N, 06°51'56" E
13	Neuglobsow	NEU	In operation	Regional background	70	53°08'28" N, 13°01'52" E
14	Schauinsland	SCH	In operation	Low mountain range	1205	47°54'49" N, 07°54'29" E
15	Waldhof	WAL	In operation	Regional background	75	52°48'04" N, 10°45'23" E
16	Zugspitze (Schneefemerhaus)	ZSF	In operation	High alpine	2670	47°25'00" N, 10°58'47" E

706

707 Table 2: Technical details of the instrumentations of the 16 GUAN sites.

NO.	Name	Type	Inlet height above ground (m)	Particle mobility size spectrometer type	Size range (nm)	eBC instrument	eBC cut-off
1	ANA	portable cabin	4	MPSS	10–800	MAAP	PM ₁
2	AUG	portable cabin	4	D-MPSS	5–800	Aethalometer (Type 8100)	PM _{2.5}
3	BOS	portable cabin	4	MPSS	10–800	MAAP	PM ₁₀
4	DDN	portable cabin	4	D-MPSS	5–800	MAAP	PM ₁
5	DDW	portable cabin	4	MPSS	10–800	MAAP	PM ₁
6	HPB	building	12	MPSS	10–800	MAAP	PM ₁₀
7	LAN	portable cabin	14	MPSS (TSI 3936)	10–600	–	PM ₁
8	LEI	building	6	TDMPSS	5–800	MAAP	PM ₁
9	LMI	portable cabin	4	TDMPSS	5–800	MAAP	PM ₁₀
10	LTR	portable cabin	16	TDMPSS	5–800	MAAP	PM ₁₀
11	MEL	portable cabin	4	D-MPSS	5–800	MAAP	PM ₁₀
12	MST	portable cabin	4	MPSS (TSI 3936)	14–750	–	PM ₁₀
13	NEU	building	6	MPSS	10–800	MAAP	PM ₁₀
14	SCH	building	6	MPSS	10–800	MAAP	PM ₁₀
15	WAL	building	6	MPSS	10–800	MAAP	PM ₁₀
16	ZSF	building	6	MPSS (TSI 3936)	20–600	MAAP	PM ₁₀

708

709

Table 3: Multi-annual trends of the PNCs and eBC mass concentration in percentage per year, calculated using the customised Sen's estimator and generalized least-square-regression with autoregression bootstrap (GLS /ARB). Statistically significant slopes at 95 % significance level are marked as bold numbers. Five site categories in the left column are roadside (RS), urban background (UB), regional background (RB), low mountain range (LMT), and high Alpine (HA).

Category	Site	eBC mass concentration		$N_{[20-800]}$		$N_{[10-30]}$		$N_{[30-200]}$		$N_{[200-800]}$	
		Sen slope	GLS /ARB slope	Sen slope	GLS /ARB slope	Sen slope	GLS /ARB slope	Sen slope	GLS /ARB slope	Sen slope	GLS /ARB slope
RS	DDN	-11.3	-13.1	-7.3	-6.0	-8.0	-7.3	-6.7	-5.6	-9.7	-8.6
	LEI	-5.0	-6.3	-2.9	-4.0	-5.0	-4.8	-2.9	-3.8	-1.2	-3.3
	LMI	-5.5	-6.8	-4.8	-6.9	-0.2	-1.4	-5.5	-7.3	-4.8	-6.7
UB	MST	---	---	-2.6	-1.5	---	---	-3.2	-1.8	-6.1	-5.6
	LTR	-4.1	-6.0	-4.3	-6.7	-4.7	-17.2	-4.1	-5.4	-4.6	-5.2
	ANA	-6.9	-13.1	-5.5	-7.7	-6.5	-8.4	-5.4	-7.5	-11.1	-8.2
	AUG	-2.3	-3.9	-6.3	-7.9	-6.0	-8.8	-6.3	-7.8	-3.7	-5.1
	DDW	-8.1	-11.0	-4.8	-8.9	-3.2	-5.8	-5.0	-5.4	-8.8	-8.2
	LAN	---	---	-3.4	-2.6	-1.4	-1.1	-4.3	-3.4	-2.5	-2.0
	BOS	-4.9	-6.2	-5.5	-5.5	-1.7	-3.9	-5.9	-6.0	-6.3	-3.6
RB	MEL	-4.4	-7.6	-0.2	-3.5	1.9	0.4	-0.2	-0.3	-2.9	-3.8
	WAL	-3.2	-3.7	-4.2	-3.6	-3.3	-3.5	-4.4	-3.7	-5.2	-6.2
	NEU	-7.8	-7.8	-1.0	-0.2	-0.6	-0.3	-0.5	0.0	-3.9	-4.8
LMT	HPB	-2.8	-6.3	-1.2	-3.1	1.7	-0.6	-1.2	-1.1	-3.9	-5.6
	SCH	-1.7	-3.6	-1.5	-3.0	3.8	-1.9	-2.0	-3.0	-3.8	-3.8
HA	ZSF	-4.0	-6.6	-4.2	-4.9	---	---	-4.1	-4.3	-4.2	-9.7
Regional Mann-Kendall		-5.0		-2.1		-1.4		-2.5		-2.9	

716 Table 4. Comparison of long-term trend studies of BC, PNC, and PM in Europe.

Study	Time period	Region	Parameters	Annual slope (numbers in brackets are the absolute slope, in $\mu\text{g m}^{-3} \text{ year}^{-1}$)
This study	2009–2018	Germany	BC	Traffic (3 sites): $-11.3\% \sim -5.0\%$, $(-0.19 \sim -0.08)$; UB (5 sites): $-8.1\% \sim -2.3\%$ $(-0.08 \sim -0.03)$; RB to high Alpine (6 sites): $-7.8\% \sim -1.7\%$ $(-0.03 \sim 0.00)$
			$N_{[20-800]}$	Traffic (3 sites): $-7.3\% \sim -2.9\%$; UB (7 sites): $-6.3\% \sim -2.6\%$; RB to high Alpine (6 sites): $-4.2\% \sim -0.2\%$
Kutzner et al., 2018	2005–2014	Germany	BC	Traffic (7 sites): $(-0.31, -0.15)$; UB (4 sites): $(-0.1, -0.02)$; Rural (1 site): 0.00
Asmi et al., 2013	2001–2010	Europe	$N_{[20-800]}$	Rural to remote (4 sites): $-4.6\% \sim 1.6\%$
Collaud Coen et al., 2013	2001–2010	Europe	Absorption coef.	Rural to remote (4 sites): $-1.6\% \sim 0.0\%$
Bigi and Ghermandi, 2016	2005–2014	Italy, Po valley	$\text{PM}_{2.5}$	Traffic (2 sites): $-6.4\% \sim -4.6\%$; UB (17 sites): $-8.1\% \sim -0.4\%$; RB (4 sites): $-4.9\% \sim 0.0\%$
Singh et al., 2018	2009–2016	United Kingdom	BC	Traffic (1 site): -8.0% ; UB (2 sites): $-5.0\% \sim -4.7\%$; Rural (1 site): -7.7%

717

Table 5: Trends of meteorological parameters for the three site categories in Germany. The bold numbers are the statistically significant slopes at the 95 % significance level. The daily meteorological data are from Germany's National Meteorological Service (Deutscher Wetterdienst, DWD).


season		Urban background	Regional background	Mountain area
Spring	Precipitation mm year ⁻¹ (% year ⁻¹)	-0.02 (-1.0)	0.00 (0.0)	-0.02 (-0.5)
	Temperature °C year ⁻¹	-0.04	-0.03	-0.02
	Wind speed m s ⁻¹ year ⁻¹ (% year ⁻¹)	0.01 (0.2)	0.02 (0.3)	0.04 (0.6)
Summer	Precipitation mm year ⁻¹ (% year ⁻¹)	-0.14 (-5.5)	-0.15 (-5.8)	-0.20 (-4.7)
	Temperature °C year ⁻¹	0.15	0.13	0.16
	Wind speed m s ⁻¹ year ⁻¹ (% year ⁻¹)	0.00 (0.0)	0.02 (0.4)	-0.08 (-1.4)
Autumn	Precipitation mm year ⁻¹ (% year ⁻¹)	-0.07 (-3.9)	-0.05 (-2.5)	-0.07 (-1.9)
	Temperature °C year ⁻¹	0.37	0.36	0.29
	Wind speed m s ⁻¹ year ⁻¹ (% year ⁻¹)	-0.02 (-0.8)	-0.01 (-0.3)	-0.09 (-1.2)
Winter	Precipitation mm year ⁻¹ (% year ⁻¹)	0.02 (1.3)	0.04 (1.8)	0.14 (3.1)
	Temperature °C year ⁻¹	0.41	0.43	0.34
	Wind speed m s ⁻¹ year ⁻¹ (% year ⁻¹)	0.02 (0.5)	0.05 (0.9)	0.13 (1.5)

723 Table 6: Basic statistical information of the 15 air mass types.

Air mass type	Wind direction	Source region	Frequency 2009–2018 (%)	Mean PM ₁₀ 2009–2018(μg m ⁻³)
CS-ST	Stagnant	Central Europe	3.0	34.6
CS-A1	East	Eastern Europe	3.4	34.8
CS-A2	West	North Atlantic	5.8	23.1
CS-C1	South West	Southwest Europe	5.3	24.4
CS-C2a	South West	North Atlantic	4.0	11.5
CS-C2b	West	North Atlantic	5.8	11.3
TS-A1	North East	Subpolar	7.4	17.5
TS-A2	West	North Atlantic	6.5	16.9
TS-C1	South West	Southwest Europe	5.0	14.4
TS-C2	North West	Arctic	10.2	12.9
WS-ST	Stagnant	Central Europe	7.4	20.5
WS-A1	South East	Eastern Europe	5.9	24.8
WS-A2	North West	North Atlantic	12.0	16.7
WS-C1	West	North Atlantic	9.1	16.4
WS-C2	West	North Atlantic	9.0	12.1

724
725

726



727

728

729

Figure 1: Map of the 16 atmospheric measurement stations in the GUAN.



Figure 2: Data coverage of the PNSD and eBC mass concentrations from 2009 to 2018 at the 16 GUAN sites.





Figure 3: Long-term change index (% of average in 2009) of measured parameters at the regional background and mountain stations sites and total emissions in Germany: (a) measured eBC mass concentration and BC emission; (b) measured $N_{[10-30]}$, $N_{[30-200]}$, $N_{[200-800]}$, and $V_{[20-800]}$, and emission of PM_{2.5}, SO₂, NMVOC, NH₃ and NO_x. Figure 3: Comparison of the long-term changes in measured parameters and total emissions in Germany.

741



742

743

744


745

746

747

Figure 4: Diurnal variations of the trends of PNCs and eBC mass concentration for each of the four site categories (RS: roadside; UB: urban background; RB: regional background; LMT&HA: low mountain range and high Alpine).

748
749




750

751 Figure 5: Seasonal statistics of the trends of PNCs and eBC mass concentration. Dots denote the mean slope for
752 all sites, black line inside the box denotes the median slope, and the top and bottom of the box denotes the 75th
753 and 25th percentiles. Spring: March to May (MAM); summer: June to August (JJA); autumn: September to
754 November (SON); and winter: December to February (DJF).

755

756



757

758

759

760

761

762

Figure 6: De-seasonalised monthly time series of eBC mass concentration and $N_{[30-200]}$ at the two urban background sites AUG and LTR. The vertical dashed lines refer to the start dates of different stages of LEZ of different stages in the city of Augsburg and Leipzig. The horizontal dashed lines refer to the mean concentration levels of measured parameters during the corresponding time period.






Figure 7: Average diurnal cycles of the increment (defined as the difference between LMI and LTR) of Δ eBC mass concentration (Δ eBC, panel a), $N_{[30-200]}$ ($\Delta N_{[30-200]}$, panel b) and $N_{[200-800]}$ ($\Delta N_{[200-800]}$, panel c) eBC mass concentrations, $N_{[30-200]}$ and $N_{[200-800]}$.

769
770
771



772
773 Figure 8: (a) 15 back-trajectory clusters terminated at MEL. (b) Average normalised profiles of pseudo potential
774 temperature (θ_v) for the 15 clusters.
775

776




777

778 Figure 9: Average **concentrations of eBC mass concentration** (a) and PNCs (b to e) for the 15 air mass types at
 779 the regional background sites category (MEL, WAL and NEU). In each panel, the boxes and whiskers denote the
 780 5th, 25th, 50th, 75th and 95th percentiles, while the dots denote the mean values. The solid red line shows the
 781 overall median values.

782

783
784



785

786

787

788

789

Figure 10: Annual ~~concentrations of the~~average eBC mass concentrations and $N_{[200-800]}$ measured at the regional background and low mountain range sites for the polluted and clean air mass categories. Green bars show the frequency of polluted air masses in each year.