

## Vertical profiles of light absorption and scattering associated with black-carbon particle fractions in the springtime Arctic above 79°N

W. Richard Leaitch<sup>1</sup>, John K. Kodros<sup>2</sup>, Megan D. Willis<sup>3, a</sup>, Sarah Hanna<sup>4</sup>, Hannes Schulz<sup>5</sup>,  
5 Elisabeth Andrews<sup>6,8</sup>, Heiko Bozem<sup>7</sup>, Julia Burkart<sup>3, b</sup>, Peter Hoor<sup>7</sup>, Felicia Kolonjari<sup>1</sup>, John  
A. Ogren<sup>8</sup>, Sangeeta Sharma<sup>1</sup>, Meng Si<sup>4</sup>, Knut von Salzen<sup>9</sup>, Allan K. Bertram<sup>4</sup>, Andreas  
Herber<sup>5</sup>, Jonathan P. D. Abbatt<sup>3</sup>, Jeffrey R. Pierce<sup>2</sup>

<sup>1</sup> - Environment and Climate Change Canada, Toronto, ON, Canada

10 <sup>2</sup> – Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA

<sup>3</sup> – Department of Chemistry, University of Toronto, Toronto, ON, Canada

<sup>a</sup> – now: Lawrence Berkeley National Laboratory, Chemical Sciences Division, Berkeley, CA, USA

<sup>b</sup> - now: Aerosol Physics and Environmental Physics, University of Vienna, Austria

<sup>4</sup> – Department of Chemistry, University of British Columbia, Vancouver, BC, Canada

15 <sup>5</sup> – Alfred Wegener Institute, Helmholtz Center for POLAR and Marine Research, Bremerhaven,  
Germany

<sup>6</sup> - University of Colorado, Boulder, CO, USA

<sup>7</sup> – Institute of Atmospheric Physics, Johannes Gutenberg-University, Mainz, Germany

<sup>8</sup> - National Oceanic and Atmospheric Administration (NOAA), Boulder, CO, USA

20 <sup>9</sup> - Environment and Climate Change Canada, Victoria, BC, Canada

*Correspondence to:* W. Richard Leaitch ([leaitchs@gmail.com](mailto:leaitchs@gmail.com)); Jeffrey R. Pierce ([jeffrey.pierce@colostate.edu](mailto:jeffrey.pierce@colostate.edu))

25 **Abstract.** Despite the potential importance of black carbon (BC) to radiative forcing of the Arctic atmosphere, vertically-resolved measurements of the particle light scattering coefficient ( $\sigma_{sp}$ ) and light absorption coefficient ( $\sigma_{ap}$ ) in the springtime Arctic atmosphere are infrequent, especially measurements at latitudes at or above 80°N. Here, relationships among vertically-distributed aerosol optical properties ( $\sigma_{ap}$ ,  $\sigma_{sp}$ , and single scattering albedo or SSA), particle microphysics and particle chemistry are examined for a region of the Canadian

30 archipelago between 79.9°N and 83.4°N from near the surface to 500 hPa. Airborne data collected during April, 2015, are combined with ground-based observations from the observatory at Alert, Nunavut and simulations from the GEOS-Chem-TOMAS model (Kodros et al., 2018) to further our knowledge of the effects of BC on light absorption in the Arctic troposphere. The results are constrained for  $\sigma_{sp}$  less than 15 Mm<sup>-1</sup>, which represent 98% of the observed  $\sigma_{sp}$ , because the single scattering albedo (SSA) has a tendency to be lower at lower  $\sigma_{sp}$ ,

35 resulting in a larger relative contribution to Arctic warming. At 18.4 m<sup>2</sup> g<sup>-1</sup>, the average BC mass absorption coefficient (MAC) from the combined airborne and Alert observations is substantially higher than the two averaged modelled MAC values (9.5 m<sup>2</sup> g<sup>-1</sup> and 7.0 m<sup>2</sup> g<sup>-1</sup>) for two different internal mixing assumptions, the latter of which is based on previous observations. The higher observed MAC value may be explained by an underestimation of BC and/or possible differences in BC microphysics and morphologies between the

40 observations and model. We present  $\sigma_{ap}$  and SSA, as measured, and  $\sigma_{ap}$  and SSA adjusted to the modelled MAC values, since the latter are more consistent with accepted MAC values. Median values of the measured  $\sigma_{ap}$ , rBC and organic component of particles all increase by a factor of 1.8±0.1 going from near-surface to 750 hPa, and values higher than the surface persist to 600 hPa. Modelled BC, organics, and  $\sigma_{ap}$  agree with the near-surface measurements, but do not reproduce the higher values observed between 900 hPa and 600 hPa. The

45 differences between modelled and observed optical properties follow the same trend as the differences between the modelled and observed concentrations of the carbonaceous components (black and organic). Discrepancies between the model and observations are not due to the use of a relatively low imaginary refractive index of BC, based on simulations for two different refractive indices. Model-observation

50 discrepancies may be mostly due to the modelled ejection of biomass burning particles only into the boundary layer at sources. For the assumption of the higher observed MAC value, the SSA range between 0.88 and 0.94, which is significantly lower than other recent estimates for the Arctic, in part reflecting the constraint of  $\sigma_{sp} < 15$  Mm<sup>-1</sup>. The large uncertainties in measuring optical properties and BC as well as the large differences between measured and modelled values, here and in the literature, argue for improved measurements of BC and light

absorption by BC as well as more vertical profiles of aerosol chemistry, microphysics, and other optical  
55 properties in the Arctic.

## 1 Introduction

Aerosol particles responsible for Arctic haze (Mitchell, 1956) originate from mid-latitude pollution sources  
during winter and spring (e.g. Holmgren et al., 1974; Rahn et al., 1977; Rahn, 1981; Shaw, 1983; Barrie and Hoff,  
60 1985; Radke et al., 1984; Schnell and Raatz, 1984; Barrie, 1986). During late spring and summer, the Arctic is  
cleansed of haze particles by increased levels of precipitation (e.g. Garrett et al., 2011) resulting in a much  
cleaner, sometimes pristine, troposphere. At lower altitudes, Europe and Northern Asia are thought to be the  
dominant source regions of Arctic haze with contributions from south/central Asian sources dominating at  
higher altitudes (e.g. Stohl, 2006; Fisher et al., 2011; Sharma et al., 2013; Qi et al., 2017; Xu et al., 2017).

65 Although surface-based concentrations of Arctic haze components have declined since studies of Arctic haze  
first began (Heidam et al., 1999; Sirois and Barrie, 1999; Sharma et al., 2004; 2006; Quinn et al., 2009; Hirdman  
et al., 2010; Sinha et al., 2017), it is unclear how total atmospheric burdens have changed (e.g. Sharma et al.,  
2013) and horizontal inhomogeneities with scales of 50-100 km (Hansen and Rosen, 1985) complicate this issue.

Radiative forcing by Arctic haze particles may be an important regional driver of Arctic climate change  
70 (e.g. Law and Stohl, 2007; Quinn et al., 2008). Characterized by a unimodal number distribution centered  
between 200 and 300 nm diameter (e.g. Bigg, 1980; Heintzenberg, 1980; Radke et al., 1984; Leaitch et al., 1989;  
Staebler, 1994), the effectiveness of Arctic haze particles at scattering light (e.g. Delene and Ogren, 2002;  
Schmeisser et al., 2017) is one reason why the net effect of the Arctic aerosol has been estimated to cool the  
Arctic atmosphere (Najafi et al., 2015; Navarro et al., 2016). Black carbon (BC) is the primary anthropogenic  
75 light-absorbing component of Arctic haze, (e.g. Leighton, 1983; Valero et al., 1984; Hansen and Rosen, 1984;  
Blanchet and List, 1987; Valero et al., 1989; Pueschel and Kinne, 1995; Hansen and Nazarenko, 2004; Flanner et  
al., 2007; McConnell et al., 2007; Law and Stohl, 2007; Quinn et al., 2008; Schindell and Faluvegi, 2009; Brock et  
al., 2011). Mitigation of BC emissions has been proposed as a provisional means of slowing Arctic warming  
(Shindell and Faluvegi, 2008; Kopp and Mauzerall, 2010; Stohl et al., 2015; Sand et al., 2016), but considerable  
80 uncertainty surrounds radiative forcing by BC in the Arctic. Despite profiles of black carbon and optical  
properties in recent years (e.g. Brock et al., 2011; McNaughton et al., 2011; Schwarz et al., 2010; Spackman et  
al., 2010; Wofsy et al., 2011; Ferrero et al., 2016; Markowicz et al., 2017) there remains a shortage of such

85 observational data that limits evaluation of models of Arctic BC and light absorption (e.g. Samset et al., 2013),  
because the Arctic is subject to transport from many pollution sources at southern latitudes during winter and  
spring, and variability exists with altitude, with location and from year-to-year.

Here, we use a combination of airborne and ground-based observations plus modelling to examine  
particle light scattering coefficients ( $\sigma_{sp}$ ), light absorption coefficients ( $\sigma_{ap}$ ), and single scatter albedos (SSA) at  
the green wavelength of 550 nm from near the surface to 500 hPa over a region of the Arctic Ocean between  
79.9°N and 83.4°N. Recent Arctic studies, ground-based and airborne, indicate SSA values ranging from 0.92 to  
90 0.97 during the month of April for a wavelength of 550 nm (Delene and Ogren, 2002; Brock et al., 2011;  
McNaughton et al., 2011; Schmeisser et al., 2017), or roughly 5% of the light incident on a population of Arctic  
haze particles is absorbed. The present work, part of the Canadian Network on Climate and Aerosols:  
Addressing Key Uncertainties in Remote Canadian Environments or NETCARE, contributes knowledge  
concerning Arctic optical properties and warming of the Arctic lower troposphere by BC during April at higher  
95 latitudes.

Airborne measurements of  $\sigma_{ap}$  that are based on transmission of light through a filter, as used here, are  
constrained by instabilities during changes in pressure (i.e. altitude) and generally higher detection limits (DL)  
associated with flight conditions. To reduce the skewness of observations imposed by these constraints, we  
expand our dataset of  $\sigma_{ap}$  values by employing a linear relationship between measured  $\sigma_{ap}$  and the refractory  
100 black carbon mass concentrations (rBC) of Schulz et al. (2018). This approach enables a more comprehensive  
representation of values below DL. In the airborne component of this dataset, over 98% of one-minute  
averages of  $\sigma_{sp}$  exceeding detection limit ( $0.9 \text{ Mm}^{-1}$ ) are below  $15 \text{ Mm}^{-1}$ . By constraining the discussion to  
values of  $\sigma_{sp}$  less than  $15 \text{ Mm}^{-1}$ , we address the largest component of Arctic haze exclusive of the direct  
influence from strong plumes. Since most Arctic pollution in April is from long-range transport, the lower  $\sigma_{sp}$   
105 suggests that these particles on average spent longer times in the Arctic atmosphere and thus are more  
indicative of the “chronic” Arctic haze discussed by Brock et al. (2011). Further, the SSA for particle populations  
that fall within this constraint have been found to decrease more sharply with decreasing  $\sigma_{sp}$  (e.g. Targino et al.,  
2005; Andrews et al., 2011), making these populations more efficient at warming of the atmosphere. Our  
primary objective is to demonstrate the large uncertainty in our knowledge of the absorption by BC at lower  $\sigma_{sp}$   
110 in an area of the high Arctic with a low frequency of observations. Details of the methods employed are

described in Section 2. In Section 3, results are presented and vertical profiles of  $\sigma_{ap}$  and SSA are compared with simulated values. Section 4 is a broader discussion of the results, and conclusions are given in Section 5.

## 2 Methods

### 115 2.1 Observations overview

Ten research flights were conducted in the Arctic using the Alfred Wegener Institute (AWI) POLAR 6 aircraft beginning on April 5, 2015 and ending on April 21, 2015. The first and only flight from Longyearbyen, Svalbard, Norway is not discussed here as it dealt with some instrument and sampling issues. Subsequently, four flights were conducted from Alert, Nunavut, Canada during April 7-9 (two flights on April 8), two flights from Eureka, Nunavut, Canada (April 11 and 13) and three flights from Inuvik, Northwest Territories, Canada on April 20-21 (two flights on April 20). Tracks of the flights out of Alert and Eureka can be found in Kodros et al. (2018), Schulz et al. (2018) and Willis et al. (2018). The flights out of Inuvik were conducted within a 300 km radius of Inuvik (68.4°N; 133.7°W). Sampling of submicron particles from the POLAR 6 during NETCARE is discussed by Leaitch et al. (2016), Schulz et al. (2018) and Willis et al. (2018). All airborne and model data presented here are referenced to a temperature of 20°C and pressure of 1013.25 hPa. The POLAR 6 data are restricted to between 2 minutes after takeoff and 2 minutes prior to landing to avoid local contamination.

Ground-based observations of  $\sigma_{sp}$ ,  $\sigma_{ap}$ , rBC and particle microphysics are routinely conducted at the Dr. Neil Trivett Global Atmospheric Watch Observatory at Alert, Nunavut (82.5°N, 62.5°W). The site is approximately 7 km south-east of the main Alert station that is operated by the Canadian Department of National Defense. The impact of the camp on observations at the Alert Observatory (hereafter, referred to as Alert) has been found to be insignificant for particles larger than about 50 nm diameter (Leaitch et al., 2018), and the observations are filtered for air arriving from within a 45° arc centred on the station. The Alert optical properties data reported here are for particles less than 1  $\mu\text{m}$  diameter.

### 135 2.2 Aerosol optical measurements

Particle light scattering coefficients ( $\sigma_{sp}$ ) at wavelengths of 450 nm, 550 nm and 700 nm were measured with a TSI Model 3563 three-wavelength volume integrating nephelometer on the POLAR 6 and at Alert. To account for the absence of a forward scattering measurement where the collimated light beam is dumped (i.e., truncation of the total scattering between 0-7° and 170-180°), the correction described by Anderson and Ogren

140 (1998) was applied. A span check of the nephelometer calibration is routinely performed at Alert using CO<sub>2</sub>, and span checks of the nephelometer used on the POLAR 6 were conducted before and after the study.

Values of  $\sigma_{ap}$  at Alert are derived from a three-wavelength Particle Soot Absorption Photometer (PSAP), as discussed by Sharma et al. (2017). The  $\sigma_{ap}$  on the POLAR 6 were measured using a Continuous Light Absorption Photometer (CLAP), a three-wavelength filter-based instrument operating at 467 nm, 528 nm and  
145 652 nm. Based on the PSAP design, the CLAP, designed and built by NOAA, samples consecutively on one of eight spots on **one 47 mm glass-fibre filter (Pallflex type E70-2075W)**, which improves functionality in situations where filter changes are more difficult, such as remote sites and aircraft (Ogren et al., 2017). For both the PSAP and CLAP, the  $\sigma_{ap}$  are derived from the change in light transmission through a spot on the filter accumulating particles relative to light transmission through a reference spot. Both the PSAP and CLAP data were corrected  
150 for multiple scattering arising from the filter medium and accumulating particles using an **empirical algorithm** described by Bond et al. (1999) and Ogren (2010). To avoid artifacts associated with filter disturbances due to rapid pressure fluctuations, the POLAR 6 absorption data are limited to in-flight pressure variations, as recorded within the nephelometer, of less than 2 hPa over a two-minute period encompassing each one-minute sample. **Recently, Düsing et al. (2019) found that changes in water uptake by filter material used in particle absorption**  
155 **measurements can influence the light absorption measurement. In the present case, because the ambient temperatures were -20°C or colder and the temperatures at the intake of the nephelometer, which was situated next to the CLAP, ranged from +14°C to +30°C, we expect little influence of relative humidity on our results. As**  
**discussed in Section 3.2, in some circumstances these corrections may fail to correct overestimates of  $\sigma_{ap}$  due to multiple scattering (Cappa et al., 2008a; 2008b; Lack et al., 2008; Sinha et al., 2017).**

160 The estimated detection limits (DL) for  $\sigma_{ap}$  and  $\sigma_{sp}$  on the POLAR 6 are 0.75 Mm<sup>-1</sup> and 0.9 Mm<sup>-1</sup>, respectively, at 550 nm wavelength and for one-minute averages. The DL are calculated from the maximum of two times the standard deviation of the raw measurement during in-flight zeroes, which are also constrained to pressure variations of 2 hPa or less. The  $\sigma_{sp}$  DL was adjusted for the truncation correction, and the  $\sigma_{ap}$  DL was adjusted for multiple scattering. Of approximately 2000 one-minute averaged data points, collected over nine  
165 flights, 220 absorption data points remain after scrutiny for DL and pressure variations. Uncertainties in the POLAR 6  $\sigma_{ap}$  are considered in Section 3.3.

The DLs for  $\sigma_{ap}$  and  $\sigma_{sp}$  measured at Alert are 0.1 Mm<sup>-1</sup> for and 0.4 Mm<sup>-1</sup>, respectively, and represent DLs for the one-hour averages of Alert data used here. Based on Sherman et al. (2015) and Ogren et al. (2017),

170 uncertainties in  $\sigma_{ap}$  at  $1 \text{ Mm}^{-1}$  and  $\sigma_{sp}$  at  $10 \text{ Mm}^{-1}$  are 60% and 24% respectively. After discriminating for local influences and removal of compromised data (e.g. zeroes), the Alert data set comprises 1505 one-hour averages, or approximately 70% of the February-April, 2015 time period. The months of February and March are included to broaden the comparison with the model results.

175 The  $\sigma_{ap}$  from the POLAR 6 and from Alert were adjusted to the nephelometer wavelengths assuming a  $1/\lambda$  relationship, and the  $\sigma_{ap}$ ,  $\sigma_{sp}$  and SSA for both platforms are reported for wavelengths of 450 nm (blue), 550 nm (green) and 700 nm (red). The SSA, given by  $\sigma_{sp}/(\sigma_{sp}+\sigma_{ap})$ , were calculated for the green values of  $\sigma_{sp}$  and  $\sigma_{ap}$ .

### 2.3 rBC and Physical Measurements

180 Size distributions for particles with diameters in the range of 85 - 1000 nm were measured on the POLAR 6 using an Ultra High Sensitivity Aerosol Spectrometer (UHSAS, Droplet Measurement Technology Inc); hereafter, all particle sizes are given as diameters. The UHSAS is a laser-based aerosol spectrometer. Checks of concentration and sizing were done with nearly monodisperse particles of ammonium sulphate. Details of the measurements used here are discussed by Schulz et al. (2018). The size distributions from the UHSAS are used to estimate submicron particle volume concentrations. Submicron particle volume concentrations at Alert are estimated from size distribution measurements with a TSI 3034 SMPS, previously discussed by Leaitch et al. (2013; 2018). As discussed by Willis et al. (2018), size distributions from the UHSAS were compared with those from the TSI 3034 SMPS at the Alert from four flights when the POLAR 6 flew between 60 m and 200 m above the ocean (sea ice) surface a few kilometres north of Alert. The volume concentrations calculated from the two measurements for the 85-500 nm size range compared within 20% for all flights.

190 Refractory black carbon (rBC) was measured on the POLAR 6 using a Droplet Measurement Technologies Inc. Single Particle Soot Photometer (SP2). The SP2 detects individual particles using an intra-cavity Nd:YAG laser operating at 1064 nm. Incandescence from components of particles absorbing at 1064 nm (i.e. BC) is detected by a pair of photomultiplier tubes, and the peak amplitude of the thermal radiation is proportional to the mass of refractory material (Moteki and Kondo, 2007; Slowik et al., 2007). The detection range of the SP2 used here is 0.60 fg rBC to 330 fg rBC, or approximately 85-700 nm mass equivalent diameter for a rBC density of  $1.8 \text{ g cm}^{-3}$  (Bond and Bergstrom, 2006; Bond et al., 2013). Mass calibrations are based on Fullerene soot particles that were size selected using a differential mobility analyser. The mobility diameters

were converted to rBC mass concentrations following Gysel et al. (2011). Schulz et al. (2018) describe the dataset used here, and they estimate the uncertainty at  $\pm 15\%$ . The rBC measurements conducted at Alert, also made using a SP2, are discussed by Sharma et al. (2017).

Non-refractory aerosol mass concentrations were measured aboard POLAR 6 with an Aerodyne High Resolution Time-of-Flight Aerosol Mass Spectrometer (ToF-AMS) (DeCarlo et al., 2006). These measurements are described in detail by Willis et al. (2019). Briefly, the ToF-AMS measured non-refractory aerosol between about 70 – 700 nm vacuum aerodynamic diameter. Detection limits for sulfate and total organic aerosol, based on 10 s time resolution, were  $0.01 \mu\text{g}/\text{m}^3$  and  $0.08 \mu\text{g}/\text{m}^3$ , respectively.

Coarse particles ( $>2 \mu\text{m}$  diameter) on the POLAR 6 were measured with a GRIMM Model 1.129 Optical Particle Counter. Model 1.129 measures particles larger than  $0.25 \mu\text{m}$ , but only the coarse particle concentrations are used here. As shown by comparisons with a Particle Measuring Systems FSSP-300 probe operated under one wing of the POLAR 6, the coarse particles tend to be sampled less effectively than the submicron particles, but they are still an indicator of the presence of coarse particles, and, more importantly, the coarse particles entering the POLAR 6 sample manifold.

## 2.4 Modelling

The observations of  $\sigma_{\text{sp}}$ ,  $\sigma_{\text{ap}}$  and SSA are compared to version 10.01 of the GEOS-Chem chemical-transport model (geos-chem.com), driven by MERRA re-analysis meteorology fields. The February-April 2015 simulation was preceded by a 2-month spin-up. This version of GEOS-Chem uses a horizontal resolution of 4 degrees latitude by 5 degrees longitude (approximately  $440 \text{ km} \times 72 \text{ km}$  in the flight region near Alert and Eureka, and  $440 \text{ km} \times 200 \text{ km}$  near Inuvik) with 47 vertical layers. Relevant to the observations presented here, 24 vertical layers cover the pressure region from the surface to 500 hPa, 12 of which are from the surface to 850 hPa spaced equally by 15.5 hPa. Aerosol microphysics is simulated using Two Moment Aerosol Sectional (TOMAS) microphysics scheme (Adams and Seinfeld, 2002) coupled with GEOS-Chem (known as “GEOS-Chem-TOMAS”), including tracers for sulfate, BC, organic aerosol, sea salt, dust, and aerosol water. More details are found in Adams and Seinfeld (2002), Lee et al. (2013), Lee and Adams (2012), and Kodros et al. (2018). Here, particle water is excluded from the calculations because the observed particles were relatively dry: during the flights, the outside air temperature was  $-15^\circ\text{C}$  or less and the measured temperature of air entering the nephelometer, which was situated near the CLAP, ranged between  $+14^\circ\text{C}$  or greater, indicating very low values of relative humidity at the point of measurement; also, no significant particle water was indicated in the ToF-AMS data.



Aerosol optical properties are calculated using monthly averaged aerosol mass and number concentrations with refractive indices from the Global Aerosol Dataset (GADS; Kopke et al., 1997). To calculate aerosol optical properties, we assume two of the BC mixing states discussed by Kodros et al. (2018): 1) “Allcore”, in which BC is fully mixed with other chemical species in a core-shell morphology within each size section, where BC forms the core of the particle and hydrophilic aerosol species form a concentric shell around the BC core; 2) “Rshell”, in which BC is mixed within a particle, again as a core surrounded by hydrophilic species, but the size-dependent fraction of BC-containing particles core sizes and hydrophilic coating thicknesses are constrained by the observed sizes and the modelled BC mass concentration as described by Kodros et al. (2018). These two states are illustrated in Fig. 2 of Kodros et al. (2018), where they are referred to as “rshell-constrained” (Fig. 2b) and “allCoreShell” (Fig. 2d). Particle mass, including BC mass, is conserved. The Allcore state is less realistic because every particle contains a BC core, whereas Arctic observations identify BC in roughly 10-20% of the particles (e.g. Sharma et al., 2017). As a result, the Allcore mixing assumption overestimates absorption (e.g. Alvarado et al., 2016). Rshell, which is based on observations, has a smaller fraction of coating material participating in absorption enhancement, resulting in lower absorption compared with Allcore; Rshell absorption is higher than that for the externally mixed assumption (Kodros et al., 2018). The Mie code of Bohren and Huffman (1983) for two concentric spheres is used to calculate  $\sigma_{\text{spt}}$ ,  $\sigma_{\text{ap}}$  and SSA.

Emissions are derived from the Emissions Database for Global Atmospheric Research (EDGAR) Hemispheric Transport of Air Pollution (HTAP) version 2.2 (Janssens-Maenhout et al., 2015). Following Xu et al. (2017), BC and organic carbon emissions from gas flaring derived from the Evaluating the Climate and Air Quality Impacts of short-lived Pollutants (ECLIPSE) emission inventory (Klimont et al., 2017) are included. Biomass burning emissions are from the Fire Inventory from NCAR (FINN) for the year 2015 (Wiedinmyer et al., 2011), but the particles are only injected into the boundary layer in this case. Dust emissions follow the DEAD scheme (Zender et al., 2003), and sea-salt aerosol emissions are based on Jaeglé et al. (2011). The model assumes a refractive index for BC of  $1.75+0.45i$  in the mid visible (Hess et al., 1998). Because the imaginary part of the index for BC is lower than some estimates (e.g.  $0.65i$  from Pluchino et al., 1980; Kim et al., 2015), model simulations were also conducted with a BC refractive index of  $1.9+0.79i$ , as recommended by Bond and Bergstrom (2006). See Liu et al. (2020) for a review of the refractive index of BC. More details of the application of this version of the model to the Arctic as well as a detailed discussion of the role of mixing state on absorption by BC are given by Kodros et al (2018).

The data used here are from two vertical columns of the model: one column includes Axel Heiberg Island and the Arctic Ocean west of the island; the second is all Arctic Ocean to the west and slightly north of Alert. Hereafter, the two grids are referred to as “Axel” and “NW Alert”. They are centred on 78°N, 95°W and 86°N, 75°W, respectively, as indicated by the stars in Fig. 1. The model grid that includes Alert is not used in order to avoid problems that might be associated with modelling of the significant terrain of Ellesmere Island.

### 3 Results

In Section 3.1, we identify the strongest influences of dust on the  $\sigma_{ap}$  observed from the POLAR 6. In section 3.2, we consider the mass absorption coefficients (MAC) for BC based on the observations (Alert and POLAR 6) and the model, including MAC at low BC mass concentrations. In Section 3.3, we present the profile data of rBC, median  $\sigma_{ap}$ , median SSA,  $\sigma_{sp}$  as well as sulphate and organic mass concentrations, and compare with like quantities from the GEOS-Chem-TOMAS model. The regression between  $\sigma_{ap}$  and rBC is used to increase the number of  $\sigma_{ap}$  and SSA values, from which the medians are derived, enabling the influence of values below DL and during pressure changes to be better represented.

#### 3.1 Dust absorption

Figure 2 shows the blue (450 nm), green (550 nm) and red (700 nm)  $\sigma_{ap}$  plotted as a function of rBC for the nine POLAR 6 flights. A number of points in the rBC range of 0.15-0.3  $\mu\text{g m}^{-3}$  lie well above their respective linear regression lines. Those points represent measurements from two of the flights conducted out of Inuvik in plumes above 3 km with trajectories that trace back over Northern China and Mongolia. In Fig. 3, the scattering Ångström exponents for green-red wavelengths are plotted versus the number concentrations of particles larger than 2  $\mu\text{m}$ . The Inuvik points are identified separately from the Alert and Eureka points, and the coarse particle influence associated with the Inuvik points is evident. The pattern of the lower Ångström exponent points (<0.8) versus increasing number concentrations of coarse particles agrees with the results of Hallar et al. (2015) for Asian dust measured at a mountain top in Colorado and with our general knowledge of aerosol optical properties (e.g. Aryal et al., 2014; Clarke et al., 2007; Russell et al., 2010). We assume the Inuvik points with a scattering Ångström exponent of less than 0.8 and coarse particle number concentrations greater than 0.5  $\text{cm}^{-3}$  are strongly influenced by dust; although there is some evidence of dust associated with all the Inuvik points.

285 The SSA from both the POLAR 6 and Alert are plotted against  $\sigma_{sp}$  at 550 nm, in Fig. 4. The POLAR 6  
points are from the POLAR 6 CLAP and Nephelometer measurements, and the dust-influenced points associated  
with the Inuvik flights are highlighted. The Alert data cover the period February, 2015 to April 14, 2015, when  
the POLAR 6 left that region. Alert data from February and March of 2015 are included in part to provide a  
larger comparison base with the model and in part to demonstrate consistency of the POLAR 6 data with those  
290 from Alert, since the latter only reaches into the  $0.75 \text{ Mm}^{-1} < \sigma_{ap} < 1.6 \text{ Mm}^{-1}$  region during February. As in  
previous observations (e.g Targino et al., 2005; Andrews et al., 2011; Schmeisser et al., 2018), lower values of  
SSA are more frequent at these smaller  $\sigma_{sp}$ . Hereafter, we only consider data that fall into the region of green  
 $\sigma_{sp} < 15 \text{ Mm}^{-1}$ , which represent 98% of the above-DL POLAR 6 data collected over Alert and Eureka and 85% of  
the above-DL Alert data (February to April 14, inclusive). With these constraints, significant contributions to  $\sigma_{ap}$   
295 from dust are removed. As next discussed, this improves the linear regression of  $\sigma_{ap}$  with rBC.

### 3.2 Mass Absorption

The regression of  $\sigma_{ap}$  with rBC for the combined POLAR 6 and Alert data (April 1-14), constrained as just  
discussed, has a slope of  $18.4 \text{ m}^2 \text{ g}^{-1}$  and intercept of  $+0.15$  (Fig. 5). In Fig. 5, separate regressions through the  
300 Alert data and through the POLAR 6 data overlap at a confidence level of greater than 95%. The  $\sigma_{ap}$ -rBC slope  
represents the average MAC of rBC in the measured particles, but this value is about twice that typically found  
for Arctic measurements at the green wavelength (e.g.  $9.5 \text{ m}^2 \text{ g}^{-1}$ , McNaughton et al., 2011;  $8 \text{ m}^2 \text{ g}^{-1}$ , Sharma et  
al., 2017;  $9.8 \text{ m}^2 \text{ g}^{-1}$ , Zanatta et al., 2018). In Fig. 6a and 6b, modelled  $\sigma_{ap}$ , which include a significant dust  
presence in many cases, are plotted against modelled BC for the NW Alert and Axel grid boxes, respectively.  
305 Modelled MAC values when dust is included are  $11.7$  and  $11.1 \text{ m}^2 \text{ g}^{-1}$  for Allcore and Rshell, respectively.  
Removal of points with modelled dust concentrations greater than  $1.5 \mu\text{g m}^{-3}$  (arbitrary value) leaves modelled  
MAC values in the Allcore case of  $9.30$  and  $9.62 \text{ m}^2 \text{ g}^{-1}$  for NW Alert and Axel grids, respectively, and  $6.94$  and  
 $7.08 \text{ m}^2 \text{ g}^{-1}$  for NW Alert and Axel, respectively, in the Rshell case. The modelled Allcore results are consistent  
with the previous Arctic MAC values reported above. The lower Rshell results are more consistent with the  
310 MAC value estimated for “freshly emitted” BC (e.g.  $7.5 \text{ m}^2 \text{ g}^{-1}$ , Bond et al., 2006; Kahnert, 2010), due to a  
smaller influence of the coating material on absorption enhancement relative to the Allcore case.

The  $\sigma_{ap}$ -rBC regressions of the observations and the simulations have positive intercepts (regression-based confidence levels exceed 99.9%; the standard error of the intercepts, based on the measurement error, is 0.034). The intercepts are the result of increases in individual MAC values with decreasing concentrations of BC, as shown in Fig. 7a for the model and 7b for the observations; in these plots, MAC should be constant across all BC if absorption is solely due to BC without enhancements, such as lensing. In the case of the modelled MAC values, the persistence of dust concentrations at low BC concentrations (Fig. 7a) may have contributed to the increase in MAC at lower BC. As the BC concentrations decrease up to 100 times less than the dust concentrations, absorption by dust with its lower imaginary refractive index (0.0065 in the model) may approach absorption by BC, which would increase the apparent MAC for BC. Because the modelled organic aerosol (OA) concentrations decrease proportionately with decreasing BC (Fig. 7a) and the imaginary refractive index is lower (0.0055), absorbing OA cannot explain the increased MAC at lower BC. Also, the higher modelled MAC at lower BC may have a contribution from an increase in the coating enhancement factor as the BC core decreases in size.

In terms of the observations, absorbing dust may be present at lower BC concentrations, as shown in Fig. 7b. The mass concentrations of coarse particles, estimated from the POLAR 6 size distributions assuming a density of  $2 \text{ g cm}^{-3}$ , are present across all rBC concentrations. However, we cannot distinguish whether the composition of these coarse particles is dust or sea salt. In Fig. 7b, the 14 points with coarse particle mass concentrations of zero suggest nothing other than an increase in MAC with decreasing BC (Fig. 7b), implying that dust may not explain the higher MAC at lower BC in this case. Other potential explanations include a greater deficiency in the rBC measurement at the lower BC concentrations, the presence of smaller BC components, as shown in Fig. 8, and/or more complex morphology of BC within the particles. According to Yu et al. (2019), the latter enhances absorption by factors of 3-16 within Arctic summer particles, when BC is also very low (e.g. Schulz et al., 2018).

The MAC-vs-BC curves in Fig. 7a and 7b have asymptotes in line with the MAC values determined from Fig. 5 and 6. Reasons for the higher observation-based MAC, based on the regression in Fig. 5, are unclear. There is no indication from the blue versus green slopes (Fig. 2) to suggest brown carbon is a significant factor for the Alert and Eureka data, and dust will not explain the differences in observed and modelled MAC. We consider three possibilities for the higher MAC:

- 340 1) The higher MAC is reasonable. This is suggested by the recent observations in smoke plumes and in background particles over the continental U.S.A. (Mason et al., 2018), and by Yu et al (2019) as discussed above. Another factor suggesting the higher MAC is reasonable is the close agreement between the present  $\sigma_{ap}$  and SSA for Alert with the April summary of Schmeisser et al. (2018) for Alert: the  $\sigma_{ap}$  are  $0.50 \text{ Mm}^{-1}$  and  $0.45 \text{ Mm}^{-1}$  for the present Alert analysis and for Schmeisser et al. (2018),  
345 respectively; the SSA are 0.95 and 0.95, respectively.
- 2) The higher observed MAC results from the overestimation of absorption by the filter-based measurements, as suggested by a number of observations. Lack et al. (2008) and Cappa et al. (2008a; 2008b) found that  $\sigma_{abs}$  measured with a PSAP ranged from 1.3 to over two times higher than  $\sigma_{abs}$  measured with a photoacoustic technique for organic mass (OM) concentrations above  $2.5 \mu\text{g m}^{-3}$  and the ratio of OM to the light absorbing component of the carbonaceous components exceeded  
350 about 15. To explain the absence of a similar effect on their results by ammonium sulphate particles, they suggested that the liquid nature of non-absorbing OM enhanced multiple scattering across the filter, further increasing absorption by particles on the filter. For the present measurements of rBC and OM (see Fig. 9 and 14), the ratio of OM to rBC (the only significant absorbing carbon) is estimated to be  
355 between 15 and 20, consistent with a strong impact of OM on our results. However, Lack et al. (2008) and Cappa et al. (2008a; 2008b) found that for relatively low (OM), the PSAP-based  $\sigma_{abs}$  was only 12% higher than the photoacoustic-based  $\sigma_{abs}$ . At  $1 \mu\text{g m}^{-3}$  or less (Fig. 12), our flight OM concentrations fall well into the low OM concentration range of Cappa et al. (2008a; 2008b) and Lack et al. (2008). Also, consistent with the lowest level OM in Fig. 12, three years of OM measurements at Alert found OM  
360 always less than  $0.5 \mu\text{g m}^{-3}$  (Leitch et al., 2018), suggesting that the average impact of this factor on our  $\sigma_{abs}$  may be in the area of 12%. Considering the ambient temperatures ( $-40^\circ\text{C}$  to  $-15^\circ\text{C}$ ), it is also possible that our non-absorbing OM was in solid forms (e.g. Zobrist et al., 2008), which might render its behavior on the filter more similar to ammonium sulphate. Sinha et al. (2017) studied the effect of volatile material (removed at  $300^\circ\text{C}$ ) on absorption by particles at two Arctic sites. On average and for  
365 particles less than  $1 \mu\text{m}$ , they found that the absorption given by the PSAP was reduced by 22% with the removal of volatile material, which suggests that the present correction based on  $\sigma_{sp}$  from the nephelometer may be insufficient.

3) The BC is underestimated here using rBC. Sharma et al. (2017) found that filter-based thermo-optical measurements of elemental carbon were an average of 1.9 times higher than rBC measured at Alert. This result will explain the high MAC value of  $18.4 \text{ m}^2\text{g}^{-1}$ , but there are many uncertainties associated with the measurement of BC by the many techniques, and the rBC measurement has been recommended for use (Bond et al., 2013). The size limitation of the rBC measurement was taken into account by Sharma et al. (2017), but not in the POLAR 6 dataset. Schulz et al. (2019) estimate the deficiency in the POLAR 6 rBC mass concentrations due to sizing limitations at 7.5%.

370

375

If we assume that our observed  $\sigma_{\text{abs}}$  are overestimated by 22% and our rBC are underestimated by 7.5%, the MAC value is reduced from  $18.4 \text{ m}^2\text{g}^{-1}$  to about  $13.4 \text{ m}^2\text{g}^{-1}$ , which is about halfway between our measured value and the more commonly accepted value in the area of  $9 \text{ m}^2\text{g}^{-1}$ . However, because there are a number of potential factors influencing the measurements of both  $\sigma_{\text{abs}}$  and BC, we cannot attribute one value with the necessary certainty. For that reason, we employ a range for  $\sigma_{\text{abs}}$  by adding to our discussion the POLAR 6 plus Alert (Apr. 1-14) observations (Fig. 5) adjusted to MAC values of  $9.2 \text{ m}^2\text{g}^{-1}$  and  $7.0 \text{ m}^2\text{g}^{-1}$ , respectively, through division of the  $\sigma_{\text{ap}}$  by factors of 2 and 2.62. The  $9.2 \text{ m}^2\text{g}^{-1}$  is derived from an average of the three MAC values referenced above and the two modelled grids for the Allcore assumption, while the  $7.0 \text{ m}^2\text{g}^{-1}$  is an average of the two modelled grids for the Rshell assumption.

380

385

### 3.3 BC, Absorption, Scattering, SSA Vertical Distributions

We expand our dataset of  $\sigma_{\text{ap}}$  values by employing the linear regression between  $\sigma_{\text{ap}}$  and rBC shown in Fig. 5.

This results in 956  $\sigma_{\text{ap}}$  points instead of 220, but because the linear relationship does not fully account for variations of individual points, we restrict the profiles of  $\sigma_{\text{ap}}$  and SSA to median values calculated over approximately equal 50 hPa pressure intervals. Based on the linear fit, the uncertainty in the median  $\sigma_{\text{ap}}$  is 6%

390

at a 99% confidence level, but that excludes possible biases in either the measurement of  $\sigma_{\text{ap}}$  or BC (as represented by rBC) discussed in Section 3.2. The corresponding uncertainty in the median SSA is  $\pm 0.01$  or less for the range of SSA discussed here, again excluding possible biases. As discussed in Section 2.1, the profile data, including  $\sigma_{\text{ap}}$  and  $\sigma_{\text{ap}}$  (SSA is dimensionless), have been adjusted to a standard temperature and pressure (20oC and 1013.25 hPa) for purposes of comparisons. We note that in-situ values of  $\sigma_{\text{ap}}$  and  $\sigma_{\text{ap}}$  are appropriate for calculating radiative effects.

395

Figure 9 shows vertical profiles of the rBC measurements of Schulz et al. (2018) and the modelled BC.

Modelled BC for April 1-14 and rBC overlap well for pressures >900 hPa. In the pressure region of 600-900 hPa, the modelled BC is about a factor of two lower than the POLAR 6 rBC: median rBC is  $0.043 \mu\text{g m}^{-3}$ ; median modelled BC for April 1-14 is  $0.022 \mu\text{g m}^{-3}$ . Using version 10.01 of GEOS-Chem (without TOMAS) and its adjoint,

400 Xu et al. (2017) found that BC corresponding to these observations was dominated by sources from eastern and southern Asia. Xi et al. (2017) also found better agreement of modelled BC with these observations; although the same version of GEOS-Chem is used here, all biomass burning emissions were injected only within the boundary layer, potentially accounting for some of the lower modelled BC relative to rBC. The relative increase in rBC in the 600-900 hPa region is consistent with the mean profiles of McNaughton et al. (2011), and the  
405 median observed rBC concentration in the 600-900 hPa range is similar to the mode concentration of normally distributed values for “free tropospheric background haze” of  $0.06 \mu\text{g m}^{-3}$  estimated by Brock et al. (2011). Also, the lower part of the profile concentration data (<1 km) is similar to the springtime low-level profile BC concentrations from Ny-Ålesund measured by Ferrero et al. (2016). The present result is considerably lower than the medians of  $0.1\text{-}0.5 \mu\text{g m}^{-3}$  measured in the Arctic in 1983 (Hansen and Rosen, 1984), 1986 (Hansen and  
410 Novakov, 1989) and 1992 (Hansen et al., 1997). A decrease in BC at Alert, Nunavut, during the 1990s of more than 50% was associated with a reduction in Eurasian emissions (e.g. Sharma et al., 2019), and the present lower rBC concentrations near the surface may be connected to that reduction. Since east Asian emissions increased during the same time (e.g. van Donkelaar et al., 2008), it is difficult to assess the reason for the present lower concentrations in the 600-900 hPa range. It appears that the present observations mostly  
415 represent particles that spent a considerable length of time in the Arctic atmosphere.

Modelled profiles of  $\sigma_{\text{ap}}$  for April 1-14 and both the Axel and NW Alert grids are shown in Fig. 10a and 10c for the Allcore assumption and in Fig. 10b and 10d for the Rshell assumption; major dust influence is removed, as shown as in Fig. 6. The modelled results in Fig. 10a and 10b are for a BC refractive index of  $1.75+0.45i$ . Results for a BC refractive index of  $1.9+0.95i$  are shown in Fig. 10c and 10d. The median values of  
420 the POLAR 6 rBC-based  $\sigma_{\text{ap}}$ , covering the period of April 7-13, inclusive, are also shown in Fig. 10 along with the hourly  $\sigma_{\text{ap}}$  from Alert for April 1-14. The black POLAR 6 curves and Alert points represent the data associated with the higher MAC, and the red POLAR 6 curves and Alert points ( $\sigma_{\text{ap}}/2$  and  $\sigma_{\text{ap}}/2.62$ ) represent the assumption of absorption overestimation. The median  $\sigma_{\text{ap}}$  from the POLAR 6 agree reasonably with the corresponding Alert values. As expected, the modelled  $\sigma_{\text{ap}}$  based on the Allcore assumption are overall higher

425 than those based on the Rshell assumption. The modelled  $\sigma_{ap}$  for the index with the higher imaginary part are slightly lower, particularly in the 400-600 hPa region. Overall, differences in the modelled  $\sigma_{ap}$  for the two refractive indices are small compared with the estimated range of the observations, and for that reason only the results for the refractive index of  $1.75+0.45i$  are used in the SSA analysis that follows. The modelled values mostly fall in between the two sets of POLAR 6  $\sigma_{ap}$  median values, except for pressures less than 600 hPa, 430 where the modelled values tend to be higher than either POLAR 6 curve. For pressures less than 600 hPa, the model and POLAR 6 both suggest an increasing tendency of  $\sigma_{ap}$  with decreasing pressure, consistent with Samset et al. (2013). At the surface, the modelled  $\sigma_{ap}$  overlap only with the Alert  $\sigma_{ap}$  based on the higher MAC.

Vertical profiles of SSA are shown in Fig. 11. The modelled results are for February, 2015, March, 2015, and April 1-14, 2015 with the upper (a-c) and lower panels (d-f) representing the Allcore and Rshell 435 assumptions, respectively. Only results for the index of refraction of  $1.75+0.45i$  are shown, as discussed above. Median values of the Alert SSA, calculated from the measured  $\sigma_{sp}$  and  $\sigma_{ap}$ , are shown for the corresponding model time period. The POLAR 6 SSA overlap well with the corresponding Alert SSA for April 1-14. The April 1-14 Alert SSA, based on the higher MAC, are the same as reported by Schmeisser et al. (2017) for Alert in April. The Alert SSA increase from February to April, which suggests a combination of the near-surface air cleansing 440 itself of BC faster than the air in the lower troposphere during the high Arctic spring-to-summer transition and long-range transport aloft maintaining its influence longer than transport near the surface. Near the surface (>900 hPa), the modelled SSA are lower than the Alert and POLAR 6 results, in contrast to the modelled near-surface  $\sigma_{ap}$  that fall between the POLAR 6 and April 1-14 Alert results (Fig. 9). This difference is due to the modelled  $\sigma_{sp}$  being overall lower than the observed  $\sigma_{sp}$ , as discussed below in connection with Fig. 12. In Fig. 445 11, the modelled and observation-based SSA have opposite tendencies from the surface to about 600 hPa. In the 700-800 hPa region, there is good agreement between the modelled SSA for the Rshell mixing state and the observation-based SSA for  $\sigma_{ap}/2$ . However, given that the modelled and measured  $\sigma_{sp}$  are in reasonable agreement in that region (Fig. 12), the SSA agreement seems inconsistent with the underestimation by the model of BC in that pressure region (Fig. 9). The POLAR 6 near-surface (>950 hPa) and Alert SSA for early April 450 range between 0.94 and 0.98, roughly consistent with the boundary layer SSA estimates of 0.97(+/-0.02) by Brock et al. (2011) and 0.95-0.96 by McNaughton et al. (2011). In the region of 600-900 hPa, the POLAR 6 ( $\sigma_{ap}/2$ ) SSA range between 0.94 and 0.96, which is also consistent with McNaughton et al. (2011) in this region and the “free tropospheric background haze” estimate of Brock et al. (2011). However, the POLAR 6 SSA



(higher MAC) range between 0.88 and 0.92 for the 600-900 hPa region, indicating substantially more absorption relative to scattering in this altitude range than found by Brock et al. (2011) and McNaughton et al. (2011). As mentioned above, the ejection of biomass burning particles only into the boundary layer at the source may contribute to the higher modelled SSA in the 600-900 hPa layer.

Modelled  $\sigma_{sp}$  for the April 1-14 period and observed  $\sigma_{sp}$  are shown in Fig. 12a; only the model results for the Axel grid are shown, as those for the NW Alert grid are similar. The modelled and observed  $\sigma_{sp}$  are most similar in the 600-900 hPa region, and therefore the differences between modelled and observation-based SSA (Fig. 11) in this pressure region are primarily the result of the variations in  $\sigma_{ap}$  (Fig. 10). Close to the surface (>900 hPa), the April 1-14 modelled  $\sigma_{sp}$  are lower than the POLAR 6  $\sigma_{sp}$  and at the low end of the Alert  $\sigma_{sp}$ , which contributes to the lower modelled SSA relative to the observation-based SSA in Fig. 10. Near the surface, the modelled  $\sigma_{sp}$  for February agree best with the observed  $\sigma_{sp}$ , in part due to the higher submicron particle volumes simulated for February (Fig. 12b). The observed submicron volume concentrations are based on the UHSAS, and the modelled submicron volumes are calculated from the modelled mass concentrations of sulphate, organics and BC:  $(SO_4/1.8 + Org/1.2 + BC/1.8)$ , where 1.8, 1.2 and 1.8 are the respective densities of these components. The modelled scattering efficiency (scattering coefficient divided by volume concentration) is significantly lower than the efficiency based on the observations. Near the surface (>900 hPa), the median of  $\sigma_{sp}/Volume$  from the observations is  $12.1 \mu m^{-1}$  (range: 7.2-26.3) compared with  $7.3 \mu m^{-1}$  (range: 5.6-10.3) from the model. For the pressure interval of 600-900 hPa, the median of  $\sigma_{sp}/Volume$  from the observations is  $10.8 \mu m^{-1}$  (range: 6.2-36.4) versus  $7.3 \mu m^{-1}$  (range: 5.6-12.2) from the model. Model underestimation of submicron particle sizes may contribute to the lower modelled volume scattering efficiencies. In Fig. 13, the modelled particle size distributions for the Axel grid, April 1-14 period and averaged over the indicated pressure intervals are compared with measured distributions from the two flights (April 11 and 13) conducted in that grid. The modelled distributions for the 800-900 hPa and 900-1019 hPa intervals are shifted to slightly lower sizes relative to the average of the observations. The modelled distributions for 600-800 hPa are a closer match to the measurements, and for 400-600 hPa, the average of the modelled sizes is a reasonable match to the April 11 measurements, but exceeds the April 13 observations. The overall pattern is generally consistent with the variation of the modelled  $\sigma_{sp}$  for the Axel grid and April 1-14 period, relative to the observed  $\sigma_{sp}$ , shown in Fig. 12a. In addition, the lower modelled volume scattering efficiencies may result from underestimation of the observed volumes that are based on the UHSAS.

The vertical distribution of the median concentrations of organic mass (OM) and sulphate mass (from Willis et al., 2018) are shown in Fig. 14 along with modelled organic and sulphate mass concentrations for the Axel grid; results for the NW Alert grid are not appreciably different. The increase in rBC from the surface to about 650 hPa (Fig. 9) is paralleled by the increase in OM and its increase relative to sulphate. In Fig. 14, modelled OM near the surface (>900 hPa) is relatively close to the observed OM, but above that the observed OM is higher, by as much as a factor of two near 750 hPa. Near the surface, sulphate is underestimated by the model, which contributes to the model's lower volume and  $\sigma_{sp}$  (Fig. 12) and hence SSA near the surface (Fig. 11).

#### 4. Discussion

Models tend to underestimate Arctic BC (Bond et al., 2013; Eckhardt et al., 2015) and sulphate (Eckhardt et al., 2015). Emissions are an important aspect of such comparisons, and a recent study using the ECHAM model showed that improved emissions inventories increase the annual BC burden over the Arctic by up to 30% (Schacht et al., 2019). The vertical distributions in Fig. 9 and 14 suggest that the BC emissions are coupled with other carbonaceous emissions producing OM. In this case, the tendency of rBC/OM is to increase with altitude, from approximately 5.8% near the surface to 7.3% at 650 hPa; 5.8% is equivalent to the slope of the modelled result in Fig. 7d. Variations in source emissions and secondary organic aerosol formation will explain some of the increase in rBC/OM throughout this altitude range, but precipitation scavenging preferentially removing more organic material than BC is also a possibility. In the high Arctic, the number fractions of particles larger than 100 nm that contain detectable rBC are <30% (e.g. Sharma et al., 2017; Kodros et al., 2018; references therein), which means there is significant potential for nucleation scavenging to separate OM (and sulphate) from BC as part of the precipitation process. As discussed by Targino et al. (2006) and Andrews et al. (2011), this may contribute to the general reduction in SSA at lower  $\sigma_{sp}$ .

The microphysics of the rBC components also supports the importance of wet deposition to the present observations. Schulz et al. (2019) showed that the mass mean diameter of the rBC components decreases with decreasing pressure from the surface, with an increase in the number of rBC-containing particles explaining the higher rBC mass concentrations aloft. Wet deposition by convective clouds near the source regions will lead to smaller BC components of particles aloft, if the larger-sized BC components are scavenged on average more than smaller BC components nearer the emissions sources. The stronger emissions from more distant and

southerly regions reaching higher altitudes (Stohl, 2006; Xu et al., 2017) could explain the general increase in rBC number concentrations with altitude. An important role for particle microphysics, and in particular the representation of BC microphysics, in the modelling of BC transport to the Arctic is indicated. Despite the  
515 detailed microphysical model used here, the simulated BC vertical profiles suggest that the model transports more BC close to the surface. Excessive convective wet deposition that removes BC aloft (e.g. Mahmood et al., 2016) or re-distributes it to the near-surface region as well as the present modelled ejection of biomass burning particles only into the boundary layer are two potential reasons for the lower modelled BC in in this case. These results are consistent with Matsui et al. (2018), who point out that for models to properly simulate BC and its  
520 radiative impact on the Arctic atmosphere, they must have good representations of the microphysics and scavenging aspects of BC and other components of the aerosol. The results of Targino et al. (2005) and Andrews et al. (2011), suggesting lower SSA associated with lower  $\sigma_{sp}$ , as well as the present results are examples that wet scavenging may enhance the relative absorption by the Arctic aerosol. However, despite our results suggesting dust is not a factor at the lower BC in this case (Fig. 7), contributions to absorption from dust are, in general, important.

Many studies have shown that BC may be of great importance to Arctic warming (e.g. Bond et al., 2013), and models compare better with ground-based observations in the Arctic than aloft (e.g. Eckhardt et al., 2015; present results). As discussed by Matsui et al. (2018) and indicated from the present work, representation of particle microphysics, including BC microphysics and scavenging of BC relative to all particles,  
530 is critical. There remains a need for focussed airborne studies conducting high quality and comprehensive measurements of BC microphysics and optical properties in the Arctic. Also, more knowledge of the BC mass absorption coefficient across the particle size spectrum is necessary if BC and its effects on absorption are to be properly assessed.

## 535 5. Conclusions

Relationships among vertically-distributed aerosol optical properties ( $\sigma_{abs}$ ,  $\sigma_{sp}$ , and SSA), microphysics and particle chemistry were examined for a region of the Canadian archipelago between 79.9°N and 83.4°N from near the surface to 500 hPa. The airborne data were collected during April, 2015, and combined with ground-based observations from the observatory at Alert, Nunavut and simulations from the GEOS-Chem-TOMAS  
540 model (Kodros et al., 2018) in an effort to increase our knowledge of the effects of BC on absorption. The

results were constrained for  $\sigma_{sp}$  less than  $15 \text{ Mm}^{-1}$ , which represents 98% of the observed  $\sigma_{sp}$ , or the longer-lived Arctic haze in this case.

Large uncertainties in estimating  $\sigma_{ap}$  and SSA are associated with this combined dataset. Based on observations from the POLAR 6 and Alert Observatory, the average mass absorption coefficient (MAC) for BC of

18.4  $\text{m}^2\text{g}^{-1}$  was about twice the averaged modelled values of  $9.2 \text{ m}^2 \text{ g}^{-1}$  and  $7.0 \text{ m}^2 \text{ g}^{-1}$  (similar to some previous Arctic measurements) that represent different internal mixing assumptions discussed by Kodros et al. (2018). The higher MAC value may be due to a number of factors, including underestimation of BC, morphological arrangements of BC components within particles that are inconsistent with the often-used core-shell concept, and overestimation of  $\sigma_{ap}$  by our observations. Due to the uncertainties, we evaluated  $\sigma_{ap}$  and SSA assuming an overestimation of absorption in addition to the higher MAC value.

Measured  $\sigma_{abs}$ , rBC and organic material in the particles all increased by close to a factor of two going from the surface to 750 hPa. Modelled BC, organics, and  $\sigma_{ap}$  did not reflect the higher measured values in the 600-900 hPa region. The main sources of the differences between modelled and observed optical properties are the differences between the simulated and observed vertical distributions of rBC, organics and sulphate. As discussed, this highlights not only the importance of improving model emissions, deposition, and transport processes, but also microphysical details.

Assuming absorption overestimation, the SSA based on the POLAR 6 measurements ranged between 0.94 and 0.98, consistent with the results of Brock et al. (2011) and McNaughton et al. (2011). Assuming the measured MAC is reasonable, SSA ranged between 0.88 and 0.94, consistent with Schmeisser et al. (2017). The lower  $\sigma_{sp}$  ( $<15 \text{ Mm}^{-1}$ ) constraint has been shown to yield lower SSA values on average (Targino et al., 2005; Andrews et al., 2011; this work). The relatively high frequency of occurrence of  $\sigma_{sp} < 15 \text{ Mm}^{-1}$  in this region of the Arctic during April makes it important to understand the size distributions and morphology of BC in particles at these lower  $\sigma_{sp}$ , and to be able to simulate size-distributed wet scavenging of BC. Also, despite ruling out a dust influence at lower BC concentrations here, our result will not be universal and dust at low BC concentrations should be considered.

This work typifies the large uncertainty that exists in our knowledge of the contribution from BC to direct warming of the Arctic atmosphere. It suggests a lower level of confidence in assessing direct absorption by BC, and the need for more detailed efforts if the impact of BC on Arctic climate is to be properly established.

570 Those efforts include improved measurements of BC and absorption, and more vertical profiles of aerosol chemistry, microphysics and optical properties.

### Acknowledgements

NETCARE was funded by the Natural Sciences and Engineering Research Council (NSERC) of Canada under its Climate Change and Atmospheric Research program, with additional financial and in-kind support from  
575 Environment and Climate Change Canada, Fisheries and Oceans Canada, the Alfred Wegener Institute, the Major Research Project Management Fund at the University of Toronto, and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 268020496 – TRR 172, within the Transregional Collaborative Research Center “Arctic Amplification: Climate Relevant Atmospheric and Surface Processes, and Feedback Mechanisms (AC3)”. Colorado State University researchers were supported  
580 by the US Department of Energy’s Atmospheric System Research, an Office of Science, Office of Biological and Environmental Research program, under grant no. DE-SC0011780, the US National Science Foundation, Atmospheric Chemistry program, under grant no. AGS-1559607, and by the US National Oceanic and Atmospheric Administration, an Office of Science, Office of Atmospheric Chemistry, Carbon Cycle, and Climate Program, under the cooperative agreement award no. NA17OAR430001. We gratefully acknowledge Kenn  
585 Borek Air Ltd, in particular our pilots and crew Garry Murtsell, Neil Traverse and Doug Mackenzie, for their support of our measurements. Logistical and technical support before and during the campaign was provided by a number of contributors, in particular by Desiree Toom (ECCC), Andrew Elford (ECCC), Dan Veber (ECCC), Julia Binder (AWI), Lukas Kandora (AWI), Jens Herrmann (AWI) and Manuel Sellmann (AWI). Extensive logistical and technical support was provided by Andrew Platt (ECCC), Mike Harwood (ECCC) and Martin Gerhmann (AWI).  
590 We are grateful to CFS Alert and Eureka Weather Station for supporting the measurements presented in this work.

### References

- 595 • Adams, P. J. and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, *J. Geophys. Res.*, 107, 4370, <https://doi.org/10.1029/2001JD001010>, 2002.
- Alvarado, M.J., Lonsdale, C.R., Macintyre, H.L., Bian, H., Chin, M., Ridley, D.A., Heald, C. L., Thornhill, K.L., Anderson, B.E., Cubison, M.J., Jimenez, J.L, Kondo, Y., Sahu, L.K., Dibb, J.E., and Wang, C.: Evaluating model parameterizations of submicron aerosol scattering and absorption with in situ data from ARCTAS 2008 *Atmos. Chem. Phys.*, 16, 9435–9455, doi:10.5194/acp-16-9435-2016, 2016.

- 600 • Anderson, T. L. and Ogren, J. A.: Determining aerosol radiative properties using TSI 3563 integrating nephelometer, *Aerosol Sci. Tech.*, 29, 57–69, 1998.
- Andrews, E., J.A. Ogren, P. Bonasoni, A. Marinoni, E. Cuevas, S. Rodriguez, J.Y. Sun, D. Jaffe, E. Fischer, U. Baltensperger, E. Weingartner, M. Collaud Coen, S. Sharma, A. Macdonald, W.R. Leitch, N.-H. Lin, P. Laj, J. Stamenov, I. Kalapov, A. Jefferson, P. Sheridan, *Climatology of Aerosol Radiative Properties in the Free*
- 605 *Troposphere. Atmospheric Research*, 102, 365-393, doi:10.1016/j.atmosres.2011.08.017, 2011.
- Aryal, R. P., Voss, K. J., Terman, P. A., Keene, W. C., Moody, J. L., Welton, E. J., and Holben, B. N.: Comparison of surface and column measurements of aerosol scattering properties over the western North Atlantic Ocean at Bermuda, *Atmos. Chem. Phys.*,
- 14, 7617–7629, doi:10.5194/acp-14-7617-2014, 2014.
- 610 • Barrie, L. A.: Arctic air pollution: An overview of current knowledge, *Atmos. Environ.*, 20, 643–663, doi:10.1016/0004-6981(86)90180-0, 1986.
- Barrie, L. A. and Hoff, R. M.: Five years of air chemistry observations in the Canadian Arctic. *Atmos. Environ.*, 19, 1995-2010, 1985.
- Bigg E. K.: Comparison of aerosol at four baseline monitoring stations. *J. appl. Met.* 19, 521-523, 1980.
- 615 • Blanchet J.-P. and R. List: Estimation of optical properties of Arctic Haze using a numerical model, *Atmos.-Ocean*, 21, 444 – 464, 1987.
- Bohren, C. F. and Huffman, D. R.: *Absorption and scattering of light by small particles*, Wiley Interscience, New York, USA, 1983.
- Bond, T. C., Anderson, T. L., and Campbell, D.: Calibration and intercomparison of filter-based measurements of visible light absorption by aerosols, *Aerosol Sci. Tech.*, 30, 582–600, 1999.
- 620 • Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, *Aerosol Sci. Tech.*, 40, 27–67, 2006.
- Bond, T. C., et al.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res. Atmos.*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.
- 625 • Brock, C. A., Cozic, J., Bahreini, R., Froyd, K. D., Middlebrook, A. M., McComiskey, A., Brioude, J., Cooper, O. R., Stohl, A., Aikin, K. C., de Gouw, J. A., Fahey, D. W., Ferrare, R. A., Gao, R.-S., Gore, W., Holloway, J. S., Hubler, G., Jefferson, A., Lack, D. A., Lance, S., Moore, R. H., Murphy, D. M., Nenes, A., Novelli, P. C., Nowak, J. B., Ogren, J. A., Peischl, J., Pierce, R. B., Pilewskie, P., Quinn, P. K., Ryerson, T. B., Schmidt, K. S., Schwarz, J. P., Sodemann, H., Spackman, J. R., Stark, H., Thomson, D. S., Thornberry, T., Veres, P., Watts, L. A., Warneke, C., and Wollny, A. G.: Characteristics, sources, and transport of aerosols measured in spring 2008 during the aerosol, radiation, and cloud processes affecting Arctic climate (ARCPAC) project, *Atmos. Chem. Phys.*, 11, 2423–2453, doi:10.5194/acp-11-2423-2011, 2011.
- 630 • Clarke, A., McNaughton, C., Kapustin, V., Shinozuka, Y., Howell, S., Dibb, J., Zhou, J., Anderson, B., Brekhovskikh, V., Turner, H., and Pinkerton, M.: Biomass burning and pollution aerosol over North America: Organic components and their influence on spectral optical properties and humidification response, *J. Geophys. Res.*, 112, D12S18, doi:10.1029/2006JD007777, 2007.
- 635 • Croft, B., Pierce, J.R., Martin, R.V., Hoose, C., Lohmann, U.: Uncertainty associated with convective wet removal of entrained aerosols in a global climate model, *Atmospheric Chemistry and Physics*, 12, 10725-10748, doi:10.5194/acp-12-10725-2012, 2012.
- 640 • Delene, D. and Ogren, J.A.: Variability of Aerosol Optical Properties at Four North American Surface Monitoring Sites. *J. Atmos. Sci.*, 59, 1135-1150, 2002.

- Düsing, S., Wehner, B., Müller, T., Stöcker, A., and Wiedensohler, A.: The effect of rapid relative humidity changes on fast filter-based aerosol-particle light-absorption measurements: uncertainties and correction schemes, *Atmos. Meas. Tech.*, 12, 5879–5895, doi.org/10.5194/amt-12-5879-2019, 2019.
- 645 • Dymarska, M., Murray, B. J., Sun, L., Eastwood, M. L., Knopf, D. A., and Bertram, A. K.: Deposition ice nucleation on soot at temperatures relevant for the lower troposphere, *J. Geophys. Res.*, 111, D04204, doi:10.1029/2005JD006627, 2006.
- Eckhardt, S., Quennehen, B., Olivie, D.J.L., Berntsen, T.K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., 650 Klimont, Z., Langner, J., Law, K.S., Lund, M.T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I.E., Nøjgaard, J.K., Quaas, J., Quinn, P.K., Raut, J.-C., Rumbold, S.T., Schulz, M., Sharma, S., Skeie, R.B., Skov, H., Uttal, T., von Salzen, K., and Stohl, A.: Current model capabilities for simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive measurement data set, *Atmos. Chem. Phys.*, 15, 9413–9433, doi:10.5194/acp-15-9413-2015, 2015.
- 655 • Ferrero, L., Cappelletti, D., Busetto, M., Mazzola, M., Lupi, A., Lanconelli, C., Becagli, S., Traversi, R., Caiazzo, L., Giardi, F., Moroni, B., Crocchianti, S., Fierz, M., Močnik, G., Sangiorgi, G., Perrone, M. G., Maturilli, M., Vitale, V., Udisti, R., and Bolzacchini, E.: Vertical profiles of aerosol and black carbon in the Arctic: a seasonal phenomenology along 2 years (2011–2012) of field campaigns, *Atmos. Chem. Phys.*, 16, 12601–12629, https://doi.org/10.5194/acp-16-12601-2016, 2016.
- 660 • Fisher, J. A., Jacob, D. J., Wang, Q., Bahreini, R., Carouge, C. C., Cubison, M. J., Dibb, J. E., Diehl, T., Jimenez, J. L., Leibensperger, E. M., Lu, Z., Meinders, M. B. J., Pye, H. O. T., Quinn, P. K., Sharma, S., Streets, D. G., van Donkelaar, A., and Yantosca, R. M.: Sources, distribution, and acidity of sulfate-ammonium aerosol in the Arctic in winter-spring, *Atmos. Environ.*, 45, 7301–7318, doi:10.1016/j.atmosenv.2011.08.030, 2011.
- Flanner, M., Zender, C., Randerson, J. and Rasch, P.: Present-day climate forcing and response from black carbon in snow, *J. Geophys. Res.*, 112, D11202, doi:10.1029/2006JD008003, 2007.
- 665 • Gysel, M., Laborde, M., Olfert, J. S., Subramanian, R., and Gröhn, A. J.: Effective density of Aquadag and fullerene soot black carbon reference materials used for SP2 calibration, *Atmos. Meas. Tech.*, 4, 2851–2858, https://doi.org/10.5194/amt-4-2851-2011, 2011.
- Hallar, A.G., R. Petersen, E. Andrews, J. Michalsky, I. B. McCubbin, and J. A. Ogren: Contributions of dust and biomass burning to aerosols at a Colorado mountain-top site, *Atmos. Chem. Phys.*, 15, 13665–13679, doi:10.5194/acp-15-13665-2015, 2015.
- 670 • Hansen, A.D.A. and Rosen, H. : Vertical distribution of particulate carbon, sulfur, and bromine in the Arctic haze and comparison with ground-level measurements at Barrow, Alaska, *Geophys. Res. Lett.*, 11, 381–384, 1984.
- 675 • Hansen, A.D.A. and Rosen, H. : Horizontal inhomogeneities in the particulate carbon component of the Arctic haze, *Atmos. Environ.*, 19, 2175–2180, 1985.
- Hansen, A.D.A. and Novakov, T. : Aerosol black carbon measurements in the Arctic haze during AGASP-II. *J. Atmos. Chem.*, 9, 347–361, 1989.
- Hansen, A.D.A., Polissar, A.V., Schnell, R.C.: Airborne aerosol and black carbon measurements over the East 680 Siberian Sea, spring 1992, *Atmos. Res.*, 44, 1534–166, 1997.
- Hansen, J., and L. Nazarenko: Soot climate forcing via snow and ice albedos, *Proc. Natl. Acad. Sci. U. S. A.*, 101, 423 – 428, doi:10.1073/pnas.2237157100, 2004.
- Heintzenberg J.: Particle size distribution and optical properties-of Arctic haze. *Tellus* 32, 251–260, 1980.

- 685 • Heidam, N. Z., Wählin, P., and Christensen, J.H.: Tropospheric gases and aerosols in northeast Greenland, *J. Atmos. Sci.*, 56, 261-278, 1999.
- Hess, M., Koepke, P. and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *Bulletin American Meteor. Soc.*, 79(5), 831–844, doi:[https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2), 1998.
- 690 • Hirdman, D., Burkhardt, J. F., Sodemann, H., Eckhardt, S., Jefferson, A., Quinn, P.K., Sharma, S., Ström, J., and Stohl, A.: Long-term trends of black carbon and sulphate aerosol in the Arctic: changes in atmospheric transport and source region emissions, *Atmos. Chem. Phys.*, 10, 9351–9368, doi:10.5194/acp-10-9351-2010, 2010.
- Holmgren B., Shaw G. E. and Weller G.: Turbidity in the Arctic atmosphere, *AIDJEX Bull.*, 27, 135-148, 1974.
- 695 • Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, *Atmos. Chem. Phys.*, 11, 3137–3157, <https://doi.org/10.5194/acp-11-3137-2011>, 2011.
- Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M.: HTAP\_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmos. Chem. Phys.*, 15, 11411–11432, <https://doi.org/10.5194/acp-15-11411-2015>, 2015.
- 700 • Kahnert, M.: On the Discrepancy between Modeled and Measured Mass Absorption Cross Sections of Light Absorbing Carbon Aerosols, *Aerosol Science and Technology*, 44:6, 453-460, doi: 10.1080/02786821003733834, 2010.
- 705 • Kim, J., Bauer, H., Dobovičnik, T., Hitzenberger, R., Lottin, D., Ferry, D. and Petzold, A.: Assessing Optical Properties and Refractive Index of Combustion Aerosol Particles Through Combined Experimental and Modeling Studies, *Aerosol Science and Technology*, 49, 340-350, doi: 10.1080/02786826.2015.1020996, 2015.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmos. Chem. Phys.*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.
- 710 • Kodros, J. K., Hanna, S., Bertram, A., Leaitch, W. R., Schulz, H., Herber, A., Zanatta, M., Burkart, J., Willis, M., Abbatt, J. 20 P. D. and Pierce, J. R.: Size-resolved mixing state of black carbon in the Canadian high Arctic and implications for simulated direct radiative effect, *Atmospheric Chemistry and Physics Discussions*, 1–32, doi:<https://doi.org/10.5194/acp-2018-171>, 2018.
- 715 • Kopke, P., Hess, M., Schult, I., and Shettle, E. P.: **Global Aerosol Data Set, Max Planck Inst. für Meteorol., Hamburg, Germany, 1997.**
- Kopp, R. E. and Mauzerall, D. L.: Assessing the climatic benefits of black carbon mitigation, *P. Natl. Acad. Sci. USA*, 107, 11703–11708, doi:10.1073/pnas.0909605107, 2010.
- 720 • Laborde, M., Mertes, P., Zieger, P., Dommen, J., Baltensperger, U., and Gysel, M.: Sensitivity of the Single Particle Soot Photometer to different black carbon types, *Atmos. Meas. Tech.*, 5, 1031–1043, <https://doi.org/10.5194/amt-5-1031-2012>, 2012.
- Lack, D. A., C. D. Cappa, D. S. Covert, T. Baynard, P. Massoli, B. Sierau, T. S. Bates, P. K. Quinn, E. R. Lovejoy, and A. R. Ravishankara (2008), Bias in filter-based aerosol light absorption measurements due to organic aerosol loading: Evidence from ambient measurements, *Aerosol Sci. Technol.*, 42(12), 1033–1041.
- 725



- Law, K. S. and Stohl, A.: Arctic air pollution: origins and impacts. *Science*, 315, 1537–1540, 2007.
- Leaitch, W.R., Hoff, R.M., and MacPherson, J.I.: Airborne and lidar measurements of aerosol and cloud particles in the troposphere over Alert Canada in April 1986. *J. Atmos. Chem.*, 9, 187-211, 1989.
- 730 • Leaitch, W.R., S. Sharma, L. Huang, A. M. Macdonald, D. Toom-Sauntry, A. Chivulescu, K. von Salzen, J.R. Pierce, N.C. Shantz, A. Bertram, J. Schroder, A.-L. Norman, R.Y.-W. Chang: Dimethyl Sulphide Control of the Clean Summertime Arctic Aerosol and Cloud, *Elementa: Science of the Anthropocene* 1: 000017, doi: 10.12952/journal.elementa.000017, 2013.
- 735 • Leaitch, W. R., L. M. Russell, J. Liu, F. Kolonjari, D. Toom, L. Huang, S. Sharma, A. Chivulescu, D. Veber, and W. Zhang: Organic functional groups in the submicron aerosol at 82.5 N, 62.5W from 2012 to 2014, *Atmos. Chem. Phys.*, 18, 1–19, doi.org/10.5194/acp-18-1-2018, 2018.
- Lee, Y. H. and Adams, P. J.: A Fast and Efficient Version of the Two-Moment Aerosol Sectional (TOMAS) Global Aerosol Microphysics Model, *Aerosol Sci. Technol.*, 46, 678–689, <https://doi.org/10.1080/02786826.2011.643259>, 2012.
- 740 • Lee, Y. H., Pierce, J. R., and Adams, P. J.: Representation of nucleation mode microphysics in a global aerosol model with sectional microphysics, *Geosci. Model Dev.*, 6, 1221–1232, <https://doi.org/10.5194/gmd-6-1221-2013>, 2013.
- Leighton, H.: Influence of the Arctic Haze on the solar radiation budget. *Atmos. Environ.*, 17, 2065-2068, 1983.
- 745 • Liu, F., Yon, J., Fuentes, A., Lobo, P., Smallwood, G.J. and Corbin, J.C.: Review of recent literature on the light absorption properties of black carbon: Refractive index, mass absorption cross section, and absorption function, *Aerosol Science and Technology*, 54, doi.org/10.1080/02786826.2019.1676878, 2020.
- Mahmood, R., Salzen, K., Flanner, M., Sand, M., Langner, J., Wang, H., and Huang, L.: Seasonality of global and Arctic black carbon processes in the Arctic Monitoring and Assessment Programme models, *J. Geophys. Res.-Atmos.*, 121, 7100–7116, 2016.
- 750 • Mason, B., Wagner, N.L., Adler, G., Andrews, E., Brock, C.A., Gordon, T.D., Lack, D.A., Perring, A.E., Richardson, M.S., Schwarz, J.P., Shook, M.A., Thornhill, K.L., Ziembad, L.D., and Murphy, D.M.: An intercomparison of aerosol absorption measurements conducted during the SEAC4RS campaign, *Aerosol Sci. & Technol.*, 52, 1012–1027, doi.org/10.1080/02786826.2018.1500012, 2018.
- 755 • Matsui, H., Hamilton, D.S., and Mahowald, N.M.: Black carbon radiative effects highly sensitive to emitted particle size when resolving mixing-state diversity, *Nat. Commun.*, doi: 10.1038/s41467-018-05635-1, 2018.
- McNaughton, C. S., A. D. Clarke, S. Freitag, V. N. Kapustin, Y. Kondo, N. Moteki, L. Sahu, N. Takegawa, J. P. Schwarz, J. R. Spackman, L. Watts, G. Diskin, J. Podolske, J. S. Holloway, A. Wisthaler, T. Mikoviny, J. de Gouw, C. Warneke, J. Jimenez, M. Cubison, S. G. Howell, A. Middlebrook, R. Bahreini, B. E. Anderson, E. Winstead, K. L. Thornhill, D. Lack, J. Cozic, and C. A. Brock (2011), Absorbing aerosol in the troposphere of 760 the Western Arctic during the 2008 ARCTAS/ARCPAC airborne field campaigns, *Atmospheric Chemistry and Physics*, 11(15), 7561–7582, doi: 10.5194/acp-11-7561-2011.
- McConnell, J. R., Edwards, E., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J. R., Pasteris, D. R., Carter, M. M., and Kahl, J. D.: 20th-century industrial black carbon emissions altered Arctic climate, *Science*, 317, 1381–1384, 2007.
- 765 • Markowicz, K.M., Ritter, C., Lisoka, J., Makuch, P., Stachlewska, S., Cappelletti, D., Mazzola, M., Chilinski, M. T.: Vertical variability of aerosol single-scattering albedo and equivalent black carbon concentration based

on in-situ and remote sensing techniques during the iAREA campaigns in Ny-Ålesund, *Atmos. Environ.*, 164, 431-447, <https://doi.org/10.1016/j.atmosenv.2017.06.014>, 2017.

- 770 • Mitchell M.: Visual range in the polar regions with particular reference to the Alaskan Arctic. *J. Atmos. Terr. Phys.*, Special Supplement, 195-211, 1956.
- 775 • Moteki, N. and Kondo, Y.: Effects of mixing state on black carbon measurements by laser-induced incandescence, *Aerosol Sci. Technol.*, 41, 398–417, <https://doi.org/10.1080/02786820701199728>, 2007.
- Moteki, N. and Kondo, Y.: Dependence of laser-induced incandescence on physical properties of black carbon aerosols: measurements and theoretical interpretation, *Aerosol Sci. Tech.*, 44, 663–675, <https://doi.org/10.1080/02786826.2010.484450>, 2010.
- Najafi, M. R., Zwiers, F. W. and Gillett, N. P.: Attribution of Arctic temperature change to greenhouse-gas and aerosol 25 influences, *Nature Clim. Change*, 5(3), 246–249, doi:10.1038/nclimate2524, 2015.
- 780 • Navarro, J. C. A., Varma, V., Riipinen, I., Seland, Ø., Kirkevåg, A., Struthers, H., Iversen, T., Hansson, H.-C. and Ekman, A. M. L.: Amplification of Arctic warming by past air pollution reductions in Europe, *Nature Geoscience*, 9(4), 277–281, 30 doi:10.1038/ngeo2673, 2016.
- Ogren, J. A.: Comment on “Calibration and intercomparison of filter-based measurements of visible light absorption by aerosols”, *Aerosol Sci. Tech.*, 44, 589–591, 1608, doi.org/10.1080/02786826.2010.482111, 2010.
- 785 • Ogren, J.A., Wendell, J., Andrews, E., et al.: "Continuous Light Absorption Photometer for Long-Term Studies," *Atmos Meas. Tech.*, 10, 4805-4818, doi:10.5194/amt-10-4805-2017, 2017.
- Pluchino A. B., Goldberg S. S., Dowling J. M. and Randall C. M: Refractive index measurements of single micron-sized carbon particles. *Appl. Opt.* 19, 3370-3372, 1980.
- Pueschel, R.F. and S.A. Kinne: Physical and radiative properties of Arctic atmospheric aerosols, *Sci. Tot. Environ.*, 161, 811-824, 1995.
- 790 • Qi, L., Q. Li, D. K. Henze, H.-L. Tseng, and C. He: Sources of springtime surface black carbon in the Arctic: an adjoint analysis for April 2008, *Atmos. Chem. Phys.*, 17, 9697–9716, doi:10.5194/acp-17-9697-2017, 2017.
- Quinn, P. K. and Coauthors, 2008: Short-lived pollutants in the Arctic: their climate impact and possible mitigation strategies. *Atmos. Chem. Phys.*, 8(6), doi:10.5194/acp-8-1723-2008.
- 795 • Quinn, P.K., T. S. Bates, K. Schulz, and G. E. Shaw: Decadal trends in aerosol chemical composition at Barrow, Alaska: 1976–2008, *Atmos. Chem. Phys.*, 9, 8883–8888, 2009.
- Radke L. F., Lyons J. H., Hegg D. A., Hobbs P. V. and Bailey I. H.: Airborne observations of Arctic aerosols-I. Characteristics of Arctic haze, *Geophys. Res. Lett.* 11, 393-396, 1984.
- Rahn K. A., Boyrs R. and Shaw G. E.: The Asian source of Arctic haze bands, *Nature*, 268, 713-715, 1977.
- Rahn K. A.: The Arctic air-sampling network in 1980, *Atmos. Environ.*, 15, 1349-1352, 1981.
- 800 • Russell, P. B., Bergstrom, R. W., Shinozuka, Y., Clarke, A. D., De-Carlo, P. F., Jimenez, J. L., Livingston, J. M., Redemann, J., Dubovik, O., and Strawa, A.: Absorption Angstrom Exponent in AERONET and related data as an indicator of aerosol composition, *Atmos. Chem. Phys.*, 10, 1155–1169, doi:10.5194/acp-10-1155-2010, 2010.
- 805 • Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty, *Atmos. Chem. Phys.*, 13, 2423-2434, <https://doi.org/10.5194/acp-13-2423-2013>, 2013.

- 810 • Sand, M., Berntsen, T. K., von Salzen, K., Flanner, M. G., Langner, J., and Victor, D. G: Response of the Arctic temperature to changes in emissions of short-lived climate forcers, *Nat. Clim. Chang.*, 6, 286–289, <https://doi.org/10.1038/nclimate2880>, 2016.
- 815 • Schacht, J., Heinold, B., Quaas, J., Backman, J., Cherian, R., Ehrlich, A., Herber, A., Huang, W. T. K., Kondo, Y., Massling, A., Sinha, P. R., Weinzierl, B., Zanutta, M., and Tegen, I.: The importance of the representation of air pollution emissions for the modeled distribution and radiative effects of black carbon in the Arctic, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-71>, in review, 2019.
- Schmeisser, L., Backman, J., Ogren, J. A., Andrews, E., Asmi, E., Starkweather, S., Uttal, T., Fiebig, M., Sharma, S., Eleftheriadis, K., Vratolis, S., Bergin, M., Tunved, P., and Jefferson, A.: Seasonality of aerosol optical properties in the Arctic, *Atmos. Chem. Phys.* 18, 11599–11622, doi:10.5194/acp-18-11599-2018, 2018.
- 820 • Schnell R. C. and Raatz W. E.: Vertical and horizontal characteristics of Arctic haze during AGASP; Alaskan Arctic, *Geophys. Res. Lett.* 11, 369–376, 1984.
- Schulz, H., Bozem, H., Zanutta, M., Leaitch, W. R., Herber, A. B., Burkart, J., Willis, M. D., Hoor, P. M., Abbatt, J. P. D. and Gerdes, R.: High–Arctic aircraft measurements characterising black carbon vertical variability in spring and summer, 35 *Atmospheric Chemistry and Physics Discussions*, 1–34, doi:10.5194/acp-2018-587, 2018.
- 825 • Schwarz, J. P., Spackman, J. R., Gao, R. S., Watts, L. A., Stier, P., Schulz, M., Davis, S. M., Wofsy, S. C. and Fahey, D. W.: Global-scale black carbon profiles observed in the remote atmosphere and compared to models, *Geophys. Res. Lett.*, 37, L18812, doi:10.1029/2010GL044372, 2010.
- 830 • Sirois, A., and L. A. Barrie: Arctic lower tropospheric aerosol trends and composition at Alert, Canada: 1980–1995, *J. Geophys. Res.*, 104, 11,599–11,618, doi:10.1029/1999JD900077, 1999.
- Sharma, S., D. Lavoue, H. Cachier, L. A. Barrie, and S. L. Gong: Long-term trends of the black carbon concentrations in the Canadian Arctic, *J. Geophys. Res.*, 109, D15203, doi:10.1029/2003JD004331, 2004.
- 835 • Sharma, S., E. Andrews, L. A. Barrie, J. A. Ogren, and D. Lavoue: Variations and sources of the equivalent black carbon in the high Arctic revealed by long-term observations at Alert and Barrow: 1989–2003, *J. Geophys. Res.*, 111, D14208, doi:10.1029/2005JD006581, 2006.
- Sharma, S., M. Ishizawa, D. Chan, D. Lavoue, E. Andrews, K. Eleftheriadis, and S. Maksyutov: 16-Year Simulation of Arctic Black Carbon: Transport, Source Contribution, and Sensitivity Analysis on Deposition, *J. Geophys. Res.*, 118, doi:10.1029/2012JD017774, 2013.
- 840 • Sharma, S., W. R. Leaitch, L. Huang, D. Veber, F. Kolonjari, W. Zhang, S. J. Hanna, A. K. Bertram, and J. A. Ogren: An evaluation of three methods for measuring black carbon in Alert, Canada, *Atmos. Chem. Phys.*, 17, 15225–15243, doi.org/10.5194/acp-17-15225-2017, 2017.
- 845 • Sharma, S., Barrie, L. A., Magnusson, E., Brattström, G., Leaitch, W. R., Steffen, A., & Landsberger, S. (2019). A factor and trends analysis of multidecadal lower tropospheric observations of Arctic aerosol composition, black carbon, ozone, and mercury at Alert, Canada, *J. Geophys. Res.: Atmos.*, 124, <https://doi.org/10.1029/2019JD030844>, 2019.
- Shaw, G. E.: Evidence for a central Eurasian source area of Arctic haze in Alaska, *Nature*, 299, 815–818, 1983.
- Sherman, J. P., Sheridan, P. J., Ogren, J. A., Andrews, E., Hageman, D., Schmeisser, L., Jefferson, A., and Sharma, S.: A multi-year study of lower tropospheric aerosol variability and systematic relationships from

- 850 four North American regions, *Atmos. Chem. Phys.*, 15, 12487–12517, <https://doi.org/10.5194/acp-15-12487-2015>, 2015.
- Shindell, D. and G. Faluvegi: Climate response to regional radiative forcing in the twentieth century. *Nature Geosci.*, 2(4), 294–300, doi: 10.1038/NGEO473, 2009.
  - Sinha, P. R., Kondo, Y., Koike, M., Ogren, J. A., Jefferson, A., Barrett, T. E., Sheesley, R. J., Ohata, S., Moteki, N., Coe, H., Liu, D., Irwin, M., Tunved, P., Quinn, P. K., and Zhao, Y.: Evaluation of ground-based black carbon measurements by filter-based photometers at two Arctic sites, *J. Geophys. Res. Atmos.*, 122, 3544–3572, doi:10.1002/2016JD025843, 2017.
  - Slowik, J. G., Cross, E. S., Han, J.-H., Davidovits, P., Onasch, T. B., Jayne, J. T., Williams, L. R., Canagaratna, M. R., Worsnop, D. R., Chakrabarty, R. K., Moosmüller, H., Arnott, W. P., Schwarz, J. P., Gao, R.-S., Fahey, D. W., Kok, G. L., and Petzold, A.: An Inter-Comparison of Instruments Measuring Black Carbon Content of Soot Particles, *Aerosol Sci. Technol.*, 41, 295–314, <https://doi.org/10.1080/02786820701197078>, 2007.
  - Sinha, P. R., Y. Kondo, M. Koike, J. A. Ogren, A. Jefferson, T. E. Barrett, R. J. Sheesley, S. Ohata, N. Moteki, H. Coe, D. Liu, M. Irwin, P. Tunved, P. K. Quinn, and Y. Zhao: Evaluation of ground-based black carbon measurements by filter-based photometers at two Arctic sites, *J. Geophys. Res. Atmos.*, 122, 3544–3572, doi:10.1002/2016JD025843, 2017.
  - Sobhani, N., Kulkarni, S., and Carmichael, G.R.: Source sector and region contributions to black carbon and PM<sub>2.5</sub> in the Arctic, *Atmos. Chem. Phys.*, 18, 18123–18148, doi.org/10.5194/acp-18-18123-2018, 2018.
  - Spackman, J. R., Gao, R. S., Neff, W. D., Schwarz, J. P., Watts, L. A., Fahey, D. W., Holloway, J. S., Ryerson, T. B., Peischl, J., and Brock, C. A.: Aircraft observations of enhancement and depletion of black carbon mass in the springtime Arctic, *Atmos. Chem. Phys.*, 10, 9667–9680, <https://doi.org/10.5194/acp-10-9667-2010>, 2010.
  - Staebler, R.M., G. den Hartog, B. Georgi, and Thorsten Dösterdiek: Aerosol size distributions in Arctic haze during the POLAR Sunrise Experiment 1992, *J. Geophys. Res.*, 99, 25,429–25,437, 1994.
  - Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Berntsen, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinska, M., Eckhardt, S., Fuglestedt, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U., Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Olivie, D., Quaas, J., Quennehen, B., Raut, J.-C., Rumbold, S. T., Samset, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E., and Zhu, T.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmos. Chem. Phys.*, 15, 10529–10566, <https://doi.org/10.5194/acp-15-10529-2015>, 2015.
  - Targino, A.C., Noone, K.J., Ostrom, E: Airborne in-situ characterization of dry aerosol optical properties in a multisource influenced marine region, *Tellus* 57B, 247–260, 2005.
  - Valero F. P. J., Ackerman T. P. and Gore J. Y.: The absorption of solar radiation by the Arctic atmosphere during the haze season and its effects on the radiation balance. *Geophys. Res. Lett.* 11, 465–468, 1984.
  - Valero, F.P.J., T.P. Ackerman, and W.J.R. Gore: The effects of the arctic haze as determined from airborne radiometric measurements during AGASP II, *J. Atmos. Chem.*, 9, 225–244, 1989.
  - van Donkelaar, A., Martin, R. V., Leaitch, W. R., Macdonald, A. M., Walker, T. W., Streets, D.G., Zang, Q., Dunlea, E., Jimenez, J. L., Dibb, J. E., Huley, G., Weber, R., and Andreae, M.O. 2008: Analysis of Aircraft and Satellite Measurements from the Intercontinental Chemical Transport Experiment (INTEX-B) to Quantify Long-Range Transport of Asian Sulfur to North America, *Atmos. Chem. Phys.*, 8, 2999–3014, [www.atmos-chem-phys.net/8/2999/2008/](http://www.atmos-chem-phys.net/8/2999/2008/), 2008.
- 860
- 865
- 870
- 875
- 880
- 885
- 890

- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire Inventory from NCAR (FINN): a high-resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.
- 895 • Willis, M.D., Bozem, H., Kunkel, D., Lee, A.K.Y., Schulz, H., Burkart, J., Aliabadi, A.A., Herber, A.B., Leaitch, W.R., and Abbatt, J.P.D.: Aircraft-based measurements of High Arctic springtime aerosol show evidence for vertically varying sources, transport and composition, *Atmos. Chem. Phys.*, 19, 57–76, <https://doi.org/10.5194/acp-19-57-2019>, 2019.
- 900 • Wofsy, S. C., The HIPPO Science Team and Cooperating Modellers and Satellite Teams: HIAPER Pole-to-Pole Observations (HIPPO): fine-grained, global-scale measurements of climatically important atmospheric gases and aerosols, *Phil. Trans. R. Soc. A* (2011) 369, 2073–2086, doi:10.1098/rsta.2010.0313, 2011.
- Xu, J.-W., Martin, R. V., Morrow, A., Sharma, S., Huang, L., Leaitch, W. R., Burkart, J., Schulz, H., Zanutta, M., Willis, M. D., Henze, D. K., Lee, C. J., Herber, A. B. and Abbatt, J. P. D.: Source attribution of Arctic black carbon constrained by aircraft and surface measurements, *Atmospheric Chemistry and Physics*, 17, 11971–11989, doi:10.5194/acp-17-11971-2017, 2017.
- 905 • Yu, H., Li, W., Zhang, Y., Tunved, P., Dall’Osto, M., Shen, X., Sun, J., Zhang, X., Zhang, J., and Shi, Z.: Organic coating on sulfate and soot particles during late summer in the Svalbard Archipelago, *Atmos. Chem. Phys.*, 19, 10433–10446, <https://doi.org/10.5194/acp-19-10433-2019>, 2019.
- Zanutta, M., Paolo Laj, Martin Gysel, Urs Baltensperger, Stergios Vratolis, Konstantinos Eleftheriadis, Yutaka Kondo, Philippe Dubuisson, Victor Winiarek, Stelios Kazadzis, Peter Tunved and Hand-Werner Jacobi: Effects of mixing state on optical and radiative properties of black carbon in the European Arctic, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2018-455>, 2018.
- 910 • Zender, C. S.: Mineral Dust Entrainment and Deposition (DEAD) model: Description and 1990s dust climatology, *J. Geophys. Res.*, 108, 4416, <https://doi.org/10.1029/2002JD002775>, 2003.
- 915 • Zobrist, B., Marcolli, C., Pedernera, D. A., and Koop, T.: Do atmospheric aerosols form glasses?, *Atmos. Chem. Phys.*, 8, 5221-5244, <https://doi.org/10.5194/acp-8-5221-2008>, 2008.




Figure 1. Map of Arctic centred on North Pole. Stars show centre of model grids used in this study: green is Northwest Alert (NW Alert) grid; orange is Axel grid. Grid sizes are approximately 440 km (N-S) x 72 km (E-W).

920







Figure 2. Particle light absorption coefficient ( $\sigma_{\text{ap}}$ ) at three wavelengths (450 nm, 550 nm and 700 nm) plotted versus refractory black carbon (rBC) mass concentrations for all above-detection-limit data collected during POLAR 6 flights from Alert, Nunavut, Eureka, Nunavut and Inuvik, NWT. Regression lines shown for reference.

925





930

Figure 3. Light-scattering Angstrom exponent between green and red wavelengths plotted versus the number concentrations of coarse particles (> 2 μm diameter). Data are separated between those collected during flights from Alert and Eureka (red boxes) and from Inuvik (black crosses).

935






Figure 4. Single scattering albedo (SSA) at the green wavelength (550 nm) plotted versus the particle light scattering coefficient ( $\sigma_{sp}$ ) at 550 nm. Data are separated between those collected at the Alert Observatory (blue crosses) and those collected from the POLAR 6 aircraft (black dots). In addition, points influenced most strongly by dust are circled in orange. Line of equal particle light absorption ( $\sigma_{ap}$ ) are for reference.

940

945





Figure 5. Particle light absorption coefficient ( $\sigma_{ap}$ ) at 550 nm plotted versus refractory black carbon (rBC) mass concentrations for all above-detection-limit data collected during POLAR 6 flights from Alert, Eureka and Inuvik, NWT, with major dust influence removed (orange dots), for data collected during POLAR 6 flights from Alert and Eureka only (green crosses) and for data collected at the Alert Observatory. The Alert Observatory data are separated between those collected during February, during March and during April 1-14, where the latter corresponds mostly closely with the POLAR 6 flights out of Alert and Eureka. Regressions are shown for all POLAR 6 data (major dust removed; orange curve), for all Alert Observatory data (purple curve) and for the POLAR 6 data from Alert and Eureka combined with the Alert Observatory data for April 1-14 (black curve).






960

Figure 6. Modelled and observed particle light absorption coefficients ( $\sigma_{ap}$ ) at 550 nm plotted versus black carbon and refractory black carbon (rBC) mass concentrations, respectively: a) modelled data for Axel grid showing all data based on the 'Allcore' assumption (green), data based on the 'Allcore' assumption and with dust limited to less than  $1.5 \mu g m^{-3}$  (purple), data based on the 'Rshell' assumption and with dust limited to less than  $1.5 \mu g m^{-3}$  (orange).

965




970

975

980


Figure 7. a) Modelled BC Mass Absorption Coefficient (MAC) plotted against modelled BC for the 'Allcore' (black crosses) and the 'Rshell' (orange crosses) 855 assumptions; modelled dust mass concentrations constrained to dust less than  $1.5 \mu\text{g m}^{-3}$  versus modelled BC mass concentrations (red triangles); modelled organic aerosol (OA) mass concentrations versus BC mass concentrations (green crosses); all modelled values are for April 1-14, 2015. b) MAC values from POLAR 6 flights and Alert Observatory (April 1-14) plotted versus measured refractory black carbon (rBC) mass concentrations (black dots); dust mass concentrations (red crosses), estimated from particle size distributions onboard the POLAR 6, plotted versus rBC mass concentrations; MAC values associated with zero dust points identified (red circles). The confidence level in the negative slope of the black points is greater than 99%. The confidence in the negative slope of the 14 red-circled points is only 80%.



985

Figure 8. Mean mass-weighted diameters of rBC measurements versus rBC mass concentrations, assuming spherical rBC components. Points are averages for mass concentration intervals of 0-0.01, 0.01-0.02, 0.02-0.03, 0.03-0.05, 0.05-0.1 and 0.1 to the maximum observed. The power-law fit ( $y=0.237[x^{0.129}]$ ) is through all points with an  $R^2$  of 0.89 and confidence level of better than 99%. The power law is appropriate based on the relationship between diameter and mass of a sphere, which includes the diameter going to zero as the mass goes to zero.

990



995

**Figure 9.** Vertical profile plot of BC mass concentrations from the model for the Axel grid (green crosses for Feb.-Apr. 14, inclusive; red crosses for April 1-14 only) and rBC mass concentrations from the POLAR 6 observations plotted with pressure.

1000






Figure 10. Vertical profile plots of  $\sigma_{ap}$  with atmospheric pressure showing median values based on POLAR 6 observations (black points). Red points in (a) and (c) indicate the assumption of absorption overestimation of the POLAR 6 observations ( $\sigma_{ap}/2$ ). Red points in (b) and (d) indicate the assumption of absorption overestimation of the POLAR 6 observations ( $\sigma_{ap}/2.62$ ). Data from the Alert Observatory for April 1-14 are shown for the absorption overestimation (red squares) and BC underestimation assumptions (black squares). The model results are for April 1-14 with the green dots representing the Axel grid and blue crosses representing the NW Alert grid. (a) gives the model results for the 'Allcore' assumption and the BC index of refraction of  $1.75-i0.45$ ; (b) gives the model results for the 'Rshell' assumption and the BC index of refraction of  $1.75-i0.45$ ; (c) gives the model results for the 'Allcore' assumption and the BC index of refraction of  $1.9-i0.79$ ; (d) gives the model results for the 'Rshell' assumption and the BC index of refraction of  $1.9-i0.79$ .




Figure 11. Vertical profile plot of SSA with atmospheric pressure showing modelled results for February (a), March (b) and April 1-14 (c) based on the 'Allcore' assumption and results for February (d), March (e) and April 1-14 (f) based on the 'Rshell' assumption. Median values of SSA from the POLAR 6 observations and the Alert Observatory shown in a, b and c are based on the measured absorption (black points) and the absorption overestimation assumption ( $\sigma_{ap}/2$ ; red points), corresponding to the 'Allcore' assumption. Median values of SSA from the POLAR 6 observations and the Alert Observatory shown in a, b and c are based on the measured absorption (black points) and the absorption overestimation assumption ( $\sigma_{ap}/2.62$ ; red points), corresponding to the 'Rshell' assumption.






Figure 12. a) Vertical profile plot of  $\sigma_{sp}$  from the model for the Axel grid (green crosses for Feb.-Apr. 14, inclusive; red crosses for April 1-14 only) and  $\sigma_{sp}$  from the POLAR 6 observations plotted with pressure. Also shown are the Alert Observatory  $\sigma_{sp}$  for April 1-14. b) As in a), but for submicron volume concentrations from the model simulations and POLAR 6 observations. Volume concentrations from the Alert Observatory are not shown.

1035





Figure 13. Comparison of modelled size distributions for the Axel grid and period of April 1-14 with the measured distributions from the POLAR 6 (P6) flights on April 11 and 13. All distributions are averaged for the indicated pressure intervals. a) Modelled and measured distributions for pressure intervals of 400-600 hPa and 600-800 hPa. b) Modelled and measured distributions for pressure intervals of 800-900 hPa and 900-1000 hPa.



1050 Figure 14. Vertical profile plot of mass concentrations of sulphate and of organics with pressure from the model for the Axel grid for April 1-14 (red crosses are sulphate and green crosses are organics) and median values of the POLAR 6 sulphate (open circles) and organic (open squares) mass concentrations.