

Sources of volatile organic compounds and policy implications for regional ozone pollution control in an urban location of Nanjing, East China

Qiuyue Zhao^{1,2}, Jun Bi^{1*}, Qian Liu², Zhenghao Ling^{3*}, Guofeng Shen⁴, Feng Chen², Yuezhen Qiao², Chunyan Li², Zongwei Ma¹

¹State Key Laboratory of Pollution Control and Resource Reuse, School of the Environment, Nanjing University, Nanjing 210023, China

²Jiangsu Key Laboratory of Environmental Engineering, Jiangsu Academy of Environmental Sciences, Nanjing 210036, China

³School of Atmospheric Sciences, Sun Yat-sen University, Guangzhou 510275, China

⁴College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

Correspondence to: Jun Bi (jbi@nju.edu.cn) and Zhenhao Ling (lingzh3@mail.sysu.edu.cn)

Abstract. Understanding the composition, temporal variability, and source apportionment of volatile organic compounds (VOCs) is necessary for determining effective control measures to minimize VOCs and its related photochemical pollution. To provide a comprehensive analysis of VOC sources and their contributions to ozone (O₃) formation in the Yangtze River Delta (YRD) - a region experiencing highest rates of industrial and economic development in China, we conducted a one-year sampling exercise using a thermal desorption-GC (gas chromatography) system for the first time at an urban site in Nanjing (JAES site). Alkanes were the dominant group at the JAES site, contributing ~53% to the observed total VOCs, followed by aromatics (~17%), acetylene (~17%), and alkenes (~13%). We identified seasonal variability in TVOCs with maximum and minimum concentrations in winter and summer, respectively. A morning and evening peak and a daytime trough were identified in the diurnal VOCs patterns. We identified VOC sources using positive matrix factorization and assessed their contributions to photochemical O₃ formation through calculating the O₃ formation potential (OFP) based on the mass concentrations and maximum incremental reactivities of VOCs. The PMF model identified five dominant VOC sources, with highest contributions from diesel vehicular exhausts (34 ± 5%), followed by gasoline vehicular exhausts (27 ± 3%), industrial emissions (19 ± 2%), fuel evaporation (15 ± 2%) and biogenic emissions (4 ± 1%). The results of the OFP calculation inferred that VOCs from industrial and vehicular emissions were found to be the dominance precursors for OFP, particularly the VOC species of xylenes, toluene and propene, which top priorities should be given to the alleviation of photochemical smog. Our results therefore highlight that priority should be given to limited VOC sources and species for effective control of O₃ formation in Nanjing.

1. Introduction

Volatile organic compounds (VOCs) are key precursors of O₃ and secondary organic aerosols (SOA) - a major component of fine particulate matter (PM_{2.5}). VOCs significantly contribute to the formation of photochemical

36 smog, atmospheric oxidative capacity, visibility degradation, and global climate (Jenkin and Clemitshaw, 2000;
37 Seinfeld and Pandis, 2006), and some VOCs are also known to be toxic to human health. Therefore, in recent
38 years, much research has focused on the impacts of VOCs due to their influence on atmospheric chemistry and
39 impacts on human health (Shao et al., 2009 and references therein).

40

41 The Yangtze River Delta (YRD) region (Shanghai-Jiangsu-Zhenjiang region) is one of the fastest growing
42 regions in China, having recently undergone rapid urbanization and industrialization. Rapid economic growth
43 has led to increased photochemical smog and elevated concentrations of ground-level O₃ and fine particulate
44 matter (PM_{2.5}). These conditions have been listed as the most important sources of pollution affecting the
45 population in the YRD region, and are likely caused by increasing concentrations of VOCs. Therefore, it has
46 been suggested that controlling VOC emissions is necessary for the effective alleviation of photochemical smog
47 (Wang et al., 2009; Zhang et al., 2009; Cai et al., 2010; Kurokawa et al., 2013; Ding et al., 2016).

48

49 To further understand VOC characteristics and to develop effective policies towards lowering VOC emissions,
50 a number of sampling campaigns have been conducted to investigate the components, mixing ratios,
51 photochemical reactivity and emissions of VOCs over the YRD region (Cai et al., 2010; An et al., 2014; Mo et
52 al., 2015; Pan et al., 2015; Shao et al., 2016; Xu et al., 2017). For example, based on continuous observation data
53 collected from March, 2011 to February, 2012, An et al. (2014) identified clear seasonal VOC variability in an
54 industrial area of Nanjing, with maximum and minimum levels observed in summer and winter, respectively.
55 VOC variability was also found to be strongly influenced by industrial emissions. In contrast, Mo et al. (2017)
56 found no difference in VOC chemical compositions between residential, industrial and suburban areas of the
57 coastal industrial city, Ningbo. By comparing the emission-based profiles and those extracted from the positive
58 matrix factorization (PMF) model, the petrochemical industry was identified as the highest contributor of
59 ambient VOCs due to the unique industrial structure of Ningbo, which is a coastal city located on the southern
60 wing of the Yangtze River Delta with petrochemical industry as its lead industry (Mo et al., 2015, 2016). Pan et
61 al. (2015) conducted emissions measurements of open biomass burning in the rural area of the YRD region and
62 examined the major contributors to O₃ pollution using a box model together with the Regional Atmospheric
63 Chemical Mechanism. Overall, these studies were conducted in industrialized and/or rural areas of the YRD
64 region and demonstrate the contribution of industrial emissions and biomass burning towards ambient VOC
65 levels and their contributions to O₃ formation. However, VOC studies in urban areas of the YRD region are
66 limited and could help to improve our understanding of the spatial variability of VOCs and their environmental
67 impact, particularly as stricter policies on VOCs and/or photochemical smog have been implemented since 2013
68 (Fu et al., 2016). Furthermore, the sampling resolution and sampling duration of these studies were relatively
69 low as the samples were collected using canisters. High-resolution VOC datasets can provide more detailed
70 information on the temporal and spatial variability, source apportionments, and impact factors of VOCs.

71


72 In this study, we collected continuous one-year observational VOC data at an urban site in Nanjing in the YRD
73 region. The seasonal and diurnal characteristics of VOCs were investigated, and their sources were identified
74 and quantified using the PMF model. Furthermore, we used a box model together with a Master Chemical
75 Mechanism (MCM) (version 3.2) to identify the O₃-precursor relationships and the contributions of VOC sources
76 to photochemical O₃ formation. Our results were compared with VOCs data from other Chinese megacities.
77 Based on these findings, we summarize and propose control strategies to minimize VOCs pollution and assess
78 their implications for Nanjing and the wider YRD region. The results provide useful information towards
79 lowering photochemical pollution in the YRD region as well as other regions in China.

80 **2. Methodology**

81 **2.1. Sampling campaign**

82 We continuously measured VOC concentrations from January to December, 2016, at an observation station on
83 the rooftop of an office building (~80 m above the ground level) of the Jiangsu Academy of Environmental
84 Science (JAES). There is a waterproof layer on the rooftop of the building but there was no guarantee that it was
85 made of asphalt. Furthermore, despite this waterproof layer on the rooftop of the building, the interferences of
86 emissions from this layer were believed to be insignificant because: 1) The waterproof layer was covered by the
87 layer of concrete, which was further covered with a layer of ceramic tile; 2) The building had been built for three
88 years before the sampling campaign was started; 3) It was documented that the VOC emitted from asphalt mainly
89 included benzene, toluene, ethylbenzene and xylene (Gardiner and Lange, 2005). However, the levels of benzene,
90 toluene, ethylbenzene, *m/p*-xylene and *o*-xylene were lower than those observed in other urban, industrial and
91 rural environments in different regions (section 3.1, Zhang et al., 2012; An et al., 2014 and 2015; Mo et al., 2015,
92 2017; He et al., 2019). 4) The sampling inlet was about 2-3 m above the rooftop of the building. It should be
93 noted that there is a waterproof layer on the rooftop of the building. However, it is not sure that the waterproof
94 layer was made of asphalt. Furthermore, though there is a waterproof layer on the rooftop of the building, the
95 interferences of emissions from the layer were believed to be insignificant because: 1) The waterproof layer was
96 covered by the layer of concrete, which was further covered with a layer of ceramic tile; 2) The building has
97 been built for at least three years when the sampling campaign was started; 3) It was documented that the VOC
98 emitted from asphalt mainly included benzene, toluene, ethylbenzene and xylene (Gardiner and Lange, 2005).
99 However, the levels of benzene, toluene, ethylbenzene, *m/p*-xylene and *o*-xylene were lower than those observed
100 in other urban and industrial and rural environments in different regions (Zhang et al., 2012; An et al., 2014 and
101 2015; Mo et al., 2015, 2017; He et al., 2019) (details in section 3.1). 4) The sampling inlet were about 2-3 m
102 above the rooftop of the building. The station is located in an urban area of Nanjing, and is surrounded by heavy
103 road traffic, residential buildings, a plant and flower market, and several auto repair shops (Figure 1). Nanjing,
104 located in the western part of the YRD region, is one of the most urbanized and industrialized areas in the world
105 and consequently experiences severe air pollution. The site is located downwind of both Nanjing city center and

106 the wider YRD region (Zhao et al., 2017; Zhou et al., 2017), and is therefore ideally placed to determine the
107 combined impacts of VOCs from both local and regional atmospheric pollution.



108
109 **Figure 1. (a) Maps of the study location showing VOCs emission at a resolution of 0.25 degrees (MG/a) (The data were from**
110 **MEIC emission inventory (www.meicmodel.org, last access date:15 September 2019). (b) The location of the JAES sampling**
111 **site is indicated by a red circle (The base map was from © Baidu Maps). The blue circles indicate vehicle repair shops, the**
112 **yellow circle indicates chemical industry park and the black solid line indicates a heavy traffic road**

113 Fifty-six VOC species including alkanes, alkenes, aromatics, and acetylene were measured at 1-h intervals using
114 a PerkinElmer Online Ozone Precursor Analyzer based on a thermal desorption-GC (gas chromatography)
115 system. First, the dried air samples were collected by a thermal desorption instrument and subsequently pre-
116 concentrated onto a cold trap. The sampling flow was 15 mL/min. After 600 mL of air was sampled, the cold
117 trap was heated to resolve the compounds adsorbed on to it. By applying the Dean's Switch technology whereby
118 the technology that transfers the effluent from one column to another column with a different stationary phase,
119 the low- and high-volatile components were injected into the $\text{Al}_2\text{O}_3/\text{Na}_2\text{SO}_4$ PLOT column ($50 \text{ m} \times 0.22 \text{ mm} \times$
120 $1 \text{ }\mu\text{m}$) and the dimethyl siloxane column ($50 \text{ m} \times 0.32 \text{ mm} \times 1 \text{ }\mu\text{m}$), respectively, and analyzed using a flame
121 ionization detector (FID). The temperature increased from 46 °C for 15 min to 170 °C at a rate of 5 °C/min, and
122 then to 200 °C at a rate of 15 °C/min. The samples were finally held at 200 °C for 6 min.

123
124 A calibration was performed daily for quality control. The calibration curves showed good linearity with a
125 correlation coefficient of 0.99. Seven analyses were performed repeatedly to test the precision of the 56 species.
126 Calibrant concentrations in the gas standard mixture (56 $\text{C}_2\text{-C}_{12}$ NMHCs, Linde Spectra Environment Gases, Inc,
127 USA) ranged from 20 to 49 ppbC. The relative standard deviations of most of the 56 species were <5%,
128 representing an error of <0.5 ppb.

129 On the other hand, trace gases including CO, NO-NO₂-NO_x, SO₂, and O₃ were measured at 1-min resolution
130 using the commercial instruments of TEI 48i, 42i, 43i and 49i (Thermo Electron Corporation). All these
131 instruments were zero checked daily, span calibrated weekly and multi-point calibrated monthly. Furthermore,

132 meteorological conditions, including the temperature, relative humidity, pressure, wind speed and direction were
133 monitored at 1-min resolution by a weather station (Vantage Pro TM & Vantage Pro 2 plus TM Weather Stations,
134 Davis Instruments).

135 **2.2. The PMF model for VOC source identification**

136 In this study, the US EPA PMF (version 4.1) model, which has been widely used to conduct source apportionment
137 of VOCs (Zhang et al., 2013; Mo et al., 2017; He et al., 2019 and references therein), was applied to the observed
138 VOC data to identify potential VOC sources. A detailed description of the PMF model is provided by Yuan et al.
139 (2009) and Ling et al. (2011). In brief, the PMF model is a receptor model, which can identify the sources and
140 contributions of given species without prior input of their source profiles. In this study, a total of 25 species were
141 selected as the input for the PMF model including species with high abundances as well as typical tracers of
142 emission sources. Species with high percentages of missing values (> 25%) were excluded (i.e., 1,3-butadiene,
143 cis/trans-2-pentene, dimethylpentane, and trimethylpentane). The total concentration of the 25 selected species
144 accounted for ~92% of the total measured VOC composition. Furthermore, we calculated the total reactivity of
145 the selected 25 species to be ~90% of the total measured VOCs through the analysis of maximum incremental
146 reactivity (MIR) (Shao et al., 2009a). The high abundance and total reactivity contributions suggests that the
147 selected 25 species were appropriate for the PMF model simulation.

148
149 The PMF model was tested using a variety of factor numbers, and the optimum source profiles and contributions
150 were determined based on the correlation between modelled and observed data, the comparison of modelled
151 profiles with the results from emission-based measurements, and previous studies involving PMF/other receptor
152 model simulations (i.e., HKEPD, 2015; Wang et al., 2014; An et al., 2014; Liu et al., 2008a). For example,
153 different solution with different factor numbers was explored and the source apportionment results from a five-
154 factor resolution that could sufficiently explain the observed levels of VOCs were selected (details in Section
155 3.3). Compared with five-factor solution, the four-factor solution derived two profiles that attributable to gasoline
156 and diesel vehicular exhaust, while most of the aromatic species in these sources and certain amounts of C₃-C₄
157 species from fuel evaporation were categorized under industrial emission. On the other hand, the six-factor
158 solution has split a factor with high presence of ethyne and certain amounts of ethane (30% in species total), C₃
159 species and benzene (~20% in species total), while some alkenes (18-80% in species total) were incorporated
160 into fuel evaporation. Furthermore, the performance of the five-factor solution was evaluated using various
161 checks and sensitivity tests. Suitable correlations between the observed concentrations and those of each species
162 predicted by the model were observed, with the correlation coefficients (R²) ranging from 0.60 - 0.91, indicating
163 that the solution adequately reproduced the observed variations of each species. All the scale residuals were
164 within ± 3σ with normal distributions for all species (Baudic et al., 2016). Moreover, different numbers of start
165 seeds were tested during the simulation and no-multiple solutions were found. The ratio of Q(robust)/Q(true)
166 obtained was ~0.93, close to 1 as suggested by previous studies and the user guide manual (Paatero, 2000; Lau

167 et al., 2010; Ling et al., 2016). In addition, the results from bootstrapping analysis for the five-factor solution
 168 with bootstrap random seed found that all the factors were mapped to a basic factor in all the 20 bootstrap runs,
 169 while the uncertainties of each species from bootstrapping analysis were within the range of 1~20%. In this study,
 170 different F_{peak} values ranging from -5 to 5 was tested in the 5-factor solution for a more realistic profile (Lau et
 171 al., 2010; Baudic et al., 2016). The profiles with the nonzero F_{peak} values were consistent with those with zero
 172 F_{peak} value, reflecting that there was little rotation for the selected solution, confirming that the profiles were
 173 reasonably explained by the five-factor solution (Baudic et al., 2016). The results of F_{peak} value = 0.5 (the base
 174 run) was selected for analysis in this study. Overall, the above features demonstrated that the five-factor solution
 175 from PMF could provide reasonable and stable apportionment results for the observed VOCs at the JAES site.

176 3. Results and discussion

177 3.1 VOC observation statistics

178 Table 1 shows the average concentration and standard deviation of fifty-six VOC species concentrations
 179 measured at the JAES site, while Figure S1 in the supplementary presented the time series of all pollution data
 180 collected at the JAES site. The annual average total VOC (TVOC, sum of the measured VOCs) concentrations
 181 in 2016 was 25.7 ± 19.1 ppbv, with highest contributions from alkanes (13.6 ± 10.5 ppbv, ~53%), followed by
 182 aromatics (4.4 ± 4.0 ppbv, ~17%), acetylene (4.5 ± 5.5 ppbv, ~17%) and alkenes (3.2 ± 3.3 ppbv, ~13%).
 183 Annually, the most abundant 10 species were acetylene, propane, ethane, ethylene, butane, toluene, *i*-pentane, *i*-
 184 butane, propylene and benzene, with a combined contribution of ~77% of the TVOC. This observed VOC
 185 composition suggests that VOCs at the JAES site are predominantly sourced from combustion emissions (i.e.,
 186 vehicular emissions). Alkenes are mainly associated with vehicular emissions and are more photochemically
 187 reactive relative to alkanes and aromatics. The alkenes were found to have higher mixing ratios during weekdays
 188 relative to the weekends (3.5 ± 0.2 vs 2.9 ± 0.1 ppbv for weekdays and weekend, respectively, $p < 0.05$), further
 189 confirming the dominant contribution of vehicular emissions to VOC levels at the JAES site.

190
 191 **Table 1. The average mixing ratios and standard deviation of VOC species concentrations measured at the JAES site from**
 192 **January to December 2016.**


Species	Average \pm Standard deviation (ppbv)	Species	Average \pm Standard deviation (ppbv)
Alkanes	13.64 ± 10.53	Alkenes	3.24 ± 3.28
ethane	3.63 ± 2.68	ethene	1.72 ± 2.00
propane	3.70 ± 3.01	propylene	0.92 ± 1.16
<i>i</i> -butane	1.03 ± 0.87	1-butene	0.12 ± 0.16
<i>n</i> -butane	1.55 ± 1.26	<i>cis</i> -2-butene	0.06 ± 0.09
cyclopentane	0.08 ± 0.10	<i>trans</i> -2-butene	0.16 ± 0.11
<i>i</i> -pentane	1.15 ± 1.24	1-pentene	0.03 ± 0.03
<i>n</i> -pentane	0.61 ± 0.60	<i>cis</i> -1-pentene	0.02 ± 0.03

2,2-dimethylbutane	0.02 ± 0.02	<i>trans</i> -2-pentene	0.02 ± 0.03
2,3-dimethylbutane	0.05 ± 0.07	isoprene	0.14 ± 0.20
2-methylpentane	0.26 ± 0.29	<i>n</i> -hexene	0.05 ± 0.03
3-methylpentane	0.16 ± 0.21	Aromatics	4.40 ± 4.01
<i>n</i> -hexane	0.40 ± 0.45	benzene	0.80 ± 0.70
methylcyclopentane	0.26 ± 0.27	toluene	1.40 ± 1.35
cyclohexane	0.10 ± 0.16	ethylbenzene	0.50 ± 0.62
2,4-dimethylpentane	0.03 ± 0.01	<i>m/p</i> -xylene	0.70 ± 0.71
2,3-dimethylpentane	0.03 ± 0.02	<i>o</i> -xylene	0.25 ± 0.24
2-methylhexane	0.06 ± 0.09	styrene	0.12 ± 0.17
3-methylhexane	0.07 ± 0.10	<i>n</i> -propylbenzene	0.03 ± 0.03
heptane	0.09 ± 0.11	<i>i</i> -propylbenzene	0.03 ± 0.04
methylcyclohexane	0.07 ± 0.09	<i>m</i> -ethyltoluene	0.11 ± 0.14
2,2,4-trimethylpentane	0.02 ± 0.03	<i>p</i> -ethyltoluene	0.05 ± 0.07
2,3,4-trimethylpentane	0.02 ± 0.01	<i>o</i> -ethyltoluene	0.04 ± 0.05
2-methylheptane	0.02 ± 0.02	1,3,5-trimethylbenzene	0.04 ± 0.06
3-methylheptane	0.02 ± 0.02	1,2,4-trimethylbenzene	0.15 ± 0.21
octane	0.04 ± 0.06	1,2,3-trimethylpentane	0.10 ± 0.14
nonane	0.02 ± 0.02	<i>m</i> -diethylbenzene	0.03 ± 0.06
decane	0.04 ± 0.04	<i>p</i> -diethylbenzene	0.04 ± 0.08
undecane	0.04 ± 0.07	Acetylene	4.47 ± 5.49
dodecane	0.09 ± 0.20	--	--

193
194 The TVOC level in this study was lower than previous measurements from an industrial site in Nanjing, in which
195 43.5 ppbv TVOC was reported (An et al., 2014). However, the high TVOC levels are likely due to the proximity
196 of the observation site (~3 km northeast) to the Nanjing chemical industry area, as well as several iron, steel, and
197 cogeneration power plants (within 2 km) (An et al., 2014). The variability in land-use between these two studies
198 have also resulted in distinct VOC component profiles. In the industrial area, the relative contributions of alkenes
199 and aromatics were as high as 25% and 22%, while the contribution of alkynes was only 7% (An et al., 2014).
200 The alkane, alkene, and aromatic concentrations from the industrial site were 1.4, 3.4, and 2.2 times higher than
201 the concentrations of this study, respectively, while alkyne concentrations were ~30% lower. Given the large
202 variability observed between the two sites, it is crucial to assess the spatial variability of ambient VOCs across
203 the city through a collaboration of multiple research groups using available real-time and online VOC monitoring
204 systems.

205
206 Table S1 compares reported ambient VOCs from continuous measurements of ≥1 year in several megacities in
207 a number of countries, including China. Continuous online measurements of ambient VOCs have only been
208 available in China since 2010, unlike many developed countries whereby online VOC measurements have been
209 available for multiple decades. In China, such measurements are only concentrated in a few megacities, including
210 Beijing, Guangzhou, and Shanghai. The TVOC level reported in Nanjing was close to levels measured in

211 Shanghai (another megacity in the YRD, East China, 27.8 ppbv) (Wang et al., 2013), Tianjin (a megacity in
212 North China, 28.7 ppbv) (Liu et al., 2016), and Wuhan (a megacity located in central China, 24.3 ppbv) (Lyu et
213 al., 2016), but was considerably lower than Beijing (north China, 35.2 ppbv) (Zhang et al., 2017) and Guangzhou
214 (south China, 42.7 ppbv) (Zou et al., 2015). Alkanes were the dominant hydrocarbon group in all the cities;
215 however, some differences in relative contributions of the four classes were observed. The contribution from
216 aromatics was highest in Shanghai (31%) relative to the other cities, which is likely explained by the large
217 petrochemical and steel industry in Shanghai (Huang et al., 2011; Wang et al., 2013). In comparison, the
218 contribution of aromatics in Guangzhou (Zou et al., 2015) and the industrial area in Nanjing (An et al., 2014)
219 were 24% and 22%, respectively, while in other cities the contribution ranged between 17-19%. The current
220 ambient VOC concentrations in Chinese megacities are generally comparable to the urban VOC levels in
221 developed countries during the year 2000. However, in developed countries, the mixing ratios of VOCs were
222 observed to decrease in the recent decades following the implementation and formulation of VOC strategies
223 (Warneke et al., 2012). For example, the mixing ratios of VOCs in Los Angeles have decreased significantly
224 from 1960-2002 at an average annual rate of ~7.5%, while the mixing ratios of VOCs in London presented a
225 higher and faster decreased since 1998 when there were higher VOC mixing ratios than those in Los Angeles,
226 confirming that the earlier implementation of VOC reduction strategies in California had clearly led to the earlier
227 improvement of air quality compared to London (Warneke et al., 2012; von Schneidemesser et al., 2010).
228 Chinese megacities are therefore experiencing significantly higher ambient VOCs contamination, given the
229 remarkable decrease in VOC emissions in developed countries over the last two decades (Pan et al., 2015;
230 European Environment Agency, 2016; U.S. EPA, 2017;). High VOC levels in Chinese megacities are known to
231 impact ambient ozone and secondary particle pollution, as well as cause adverse impacts on human health.
232 However, as China has a solid foundation for VOCs monitoring and control, numerous strict, appropriate and
233 targeted reduction strategies for VOCs have been/are being formulated and implemented in Chinese megacities
234 (Guo et al., 2017). It is expected these measures could help China to reduce VOC emissions/mixing ratios and
235 improve air quality in the future.




236
237 **Figure 2. Comparison of annual average concentrations of ambient VOC in different cities based on real-time online**
238 **continuous measurements of at least one year.**

239 **3.2 Temporal variability**

240 In this study, ambient VOCs showed significant seasonal variability, with relatively high monthly average
241 concentrations in winter (40.2 ± 24.0 ppbv) and spring (23.8 ± 15.0 ppbv), and low concentrations in summer
242 (18.5 ± 14.6 ppbv) and autumn (20.1 ± 12.2 ppbv). As shown in Figure S2, the highest monthly average
243 concentration was observed in December, followed by January. High pollution levels during the winter period
244 are usually expected and is explained by atmospheric temperature inversions caused by cooler weather, which
245 inhibits particle dispersion. Lower concentrations during the summer period are due to both favorable diffusion
246 conditions and photochemical degradation of VOCs.

247
248 High wintertime VOCs pollution were also reported in Shanghai (Wang et al., 2013), Guangzhou (Zou et al.,
249 2015), and Tianjin (Liu et al., 2016), though some differences in the monthly VOC variability were also observed.
250 Except for the winter months, similar (and relatively stable) ambient VOC levels in the remaining months were
251 observed for Guangdong (Figure 3). In Shanghai, relative high levels of VOCs were observed from October to
252 January of the following year and from June to July based on the two-year measurement conducted from 2009
253 to 2010 (Wang et al., 2013). The inversion layer, the effect of cold front or uniform pressure in winter resulted
254 in high levels of VOCs from October to January of the following year, while the frontal inverted trough or
255 frequently observed stagnant high pressure system with southwest flow that could lead to poor diffusion were
256 unfavorable meteorological conditions for high VOC levels from June to July. In addition, air masses transported
257 from upwind chemical and petrochemical industrial factories located in the southwest and south of the
258 monitoring site was another factor for the high VOC levels in summer (*i.e.*, June and July) and winter. VOCs
259 concentrations in Tianjin showed significant monthly variability. Highest concentrations were reported in
260 autumn and lowest concentrations were reported in summer. The observed monthly variability is affected by

261 several factors including the type and level of emissions and local meteorological conditions.



262
263 **Figure 3. Monthly variability of ambient VOCs at the JAES site and three other Chinese cities, Shanghai (Wang et al., 2013),**
264 **Guangzhou (Zou et al., 2015), and Tianjin (Liu et al., 2016).**

265 Figure S3 shows the diurnal trends in ambient VOCs for each month. The diurnal patterns were generally similar
266 for all the months. The observed peak at approximately 8-9 am (local time) corresponds with the city's morning
267 traffic rush. The concentration begins to decrease after 9 am, with lowest concentrations observed at
268 approximately 3 pm. The observed decline was likely due to reduced vehicle emissions, growth of the inversion
269 top, and enhanced photochemical VOC degradation. After 3 pm, the concentrations begin to increase gradually
270 as a result of increased vehicle emissions during the evening rush hour, as well as a reduction in the atmospheric
271 mixing height under evening meteorological conditions. The second evening VOC peak was less prominent than
272 the morning peak. Evening concentrations were generally higher than the daytime concentrations, and the
273 amplitudes of diurnal variability were larger in autumn and summer compared to winter and spring.

274

275 3.3 Source apportionment of VOCs

276 In this study, we applied the PMF model to apportion the sources of VOCs at the sampling site. Figure 4
277 illustrates the source profiles of the VOCs produced by the PMF model. Five VOC sources were resolved by
278 PMF, including biogenic emissions (Source 1), fuel evaporation (Source 2), gasoline vehicular exhausts (Source
279 3), diesel vehicular exhausts (Source 4), and industrial emissions (Source 5).


280

281 Source 1 was identified as biogenic emissions due to the high loading of isoprene – a typical tracer of biogenic
282 emissions (Lau et al., 2010; Yuan et al., 2012). Source 2 was represented by high proportions of 2-methylpentane,
283 3-methylpentane, *i*-pentane, and cyclopentane. Pentanes are mainly associated with profiles from gasoline-
284 related emissions (Barletta et al., 2005; Tsai et al., 2006). However, the low contributions of incomplete
285 combustion tracers in this profile suggested that the VOCs were sourced from fuel evaporation. The high

286 presence of pentanes in this profile was consistent with the source profile of gasoline volatilization extracted
287 from principal component analysis/absolute principal component scores (PCA/APCs) based on the observed
288 VOC data collected in an industrial area of Nanjing (An et al., 2014), the source profile of gasoline evaporation
289 from PMF at the suburban site and urban site in Beijing and Hong Kong (Yuan et al., 2009; Lau et al., 2010).
290 Particularly, based on the emission-based measurement, Liu et al. (2008b) conducted source apportionments of
291 VOCs in the Pearl River Delta region by the chemical mass balance (CMB) receptor model, which attributed the
292 source with high loadings of *n/i*-pentanes, cyclopentane and 2/3-methylpentane as gasoline evaporation.
293 Therefore, Source 2 here was identified as fuel evaporation.

294
295 Source 3 and Source 4 were identified as vehicular exhausts due to their high loadings of incomplete combustion
296 tracers, i.e., C₂-C₄ alkanes and alkenes (Guo et al., 2011a, b; Zhang et al., 2018). Zhang et al. (2018) compared
297 the VOC composition of vehicular emissions from Zhujiang Tunnel in 2014 and 2004 in the Pearl River Delta
298 region with those from other tunnel measurements. C₂-C₄ alkanes and alkenes were found to made the greatest
299 contributions to the loading of VOCs emitted from vehicles in 2014. The higher proportions of *n/i*-pentane, *n*-
300 hexane, and methylcyclopentane in Source 3 relative to Source 4 indicated VOCs sourced from gasoline
301 vehicular exhausts (Liu et al., 2008b; Guo et al., 2011b; Zhang et al., 2018). Source 4 was identified as diesel
302 vehicular exhausts due to the high percentages of ethyne, ethane, and propene, as well as C₂-C₄ alkenes (Ho et
303 al., 2009; Cai et al., 2010; Ou et al., 2015; Liu et al., 2008c). Source 5 was characterized by high concentrations
304 of aromatics. In addition to gasoline vehicle emissions, industrial emission could be another important
305 contributor to ambient aromatic hydrocarbons in the Yangtze River Delta, Pearl River Delta and North China
306 Plain (Yuan et al., 2009; Zhang et al., 2013, 2014; An et al., 2014; Mo et al., 2015, 2017; He et al., 2019). The
307 tunnel studies and emission-based measurement results found that aromatic hydrocarbons from gasoline vehicle
308 exhaust were coherently emitted with pentanes, butenes, *n*-hexane, and cyclopentane, which were more
309 consistent with the profile in source 3 mentioned above (Liu et al., 2008; Ho et al., 2009; Yuan et al., 2009;
310 Zhang et al., 2018). Therefore, the absence of above species in source 5 indicated that this source could be related
311 to industrial emission (Zhang et al., 2014). Particularly, the high presence of toluene, ethylbenzene, xylenes,
312 ethyltoluene and trimethylbenzene was consistent with the emission-base measurement results conducted in
313 paint and printing industries (Yuan et al., 2010) and manufacturing facilities (Zheng et al., 2013). On the other
314 hand, the profile of high presence of aromatic hydrocarbons (C₇-C₉ aromatics) and the certain amount of ethene,
315 was also agree with the profiles measured in the areas dominated by industrial emissions in the Yangtze River
316 Delta region (An et al., 2014; Shao et al., 2016; Mo et al., 2017). For example, An et al. (2014) reported that
317 toluene, ethylbenzene, xylenes, and trimethylbenzenes could be emitted from different industrial processes, and
318 identified that the factors with high loadings of these species as industrial production, solvent usage and
319 industrial production volatilization sources by PAC/APCS at the industrial area in Nanjing. On the other hand,
320 Mo et al. (2017) identified the factors with high concentrations of C₇-C₉ aromatics and ethene as residential
321 solvent usage, chemical and paint industries and petrochemical industry with the PMF model applied to the data

322 collected in an industrialized coastal city of Yangtze River Delta region. To further identify source 3 and source
 323 5, the ratio of toluene/benzene (T/B, ppbv/ppbv) in each profile was compared with those obtained from
 324 emission-based measurements and tunnel study results (Zhang et al., 2018 and references therein). The ratios of
 325 T/B were ~8.2 and ~1.2 for sources 5 and 3, respectively, and were consistent with those of “industrial processes
 326 and solvent application”, and “roadside and tunnel study”, respectively (Zhang et al., 2018 and references
 327 therein). This further confirmed that source 3 was related to gasoline vehicular exhaust, while source 5 was
 328 associated with industrial emission.



329
 330 **Figure 4. Source profiles of VOCs identified using the PMF model and the relative contributions of the individual VOC**
 331 **species.**


332
 333 Vehicular exhausts were found to be the most significant contributor to the TVOCs at the JAES site, with average
 334 contributions of ~34% and ~27% for diesel and gasoline exhausts, respectively, followed by industrial emissions
 335 (19%), fuel evaporation (~15%), and biogenic emissions (~4%). Our results are inconsistent with previous results
 336 observed at industrial sites in Nanjing (An et al., 2014; Xia et al., 2014a). An et al. (2014) found that industrial
 337 activities were the most significant source of VOCs, contributing 45%-63% (mainly aromatic VOCs), followed
 338 by vehicle emission at 34%-50%. Similarly, Xia et al. (2014a) reported solvent usage and other industrial sources
 339 to account for most (31%) of the VOCs in a suburban site in southwestern Nanjing, close in proximity to
 340 Nanjing’s industrial zone. Fossil fuel/biomass/biofuel combustion were the second highest contributors at 28%,
 341 while the average contribution of vehicular emissions was 17%, mainly from the northern center of Nanjing (Xia
 342 et al., 2014a). Combined, these results infer vehicular emissions to be a major component of urban emissions in
 343 Nanjing. The observed spatial variability in the contributions of VOC sources infers the complex emissions
 344 characteristics of VOCs in Nanjing, likely due to the city’s unique industrial structure. For example, the sampling
 345 site (i.e., the JAES site) was located at a more residential and urban area compared to other sites listed in An et

346 al. (2014) and Xia et al. (2014). There are more than 0.22 million people living in the areas surrounding the
347 sampled site (within 3 km of the observation site) which composed of residential communities, schools,
348 government agencies, and business centers. These results also demonstrate that local emissions are dominant
349 contributors to ambient VOCs levels in Nanjing.

350
351 The dominant contribution of vehicular emissions to ambient VOCs in Nanjing is consistent with the
352 urban/central areas of other large cities, including Hong Kong, Guangzhou, Shanghai, and Beijing, as identified
353 and quantified by the PMF model (Yuan et al., 2009; Cai et al., 2010; Guo et al., 2011a; Zhang et al., 2013; Wang
354 et al., 2015). In addition, our results are in agreement with the anthropogenic VOC source emission inventory of
355 Jiangsu Province in 2010 (Xia et al., 2014b), indicating vehicular emissions and industrial emissions (i.e., solvent
356 usage and industrial process source) to be the two dominant sources of VOCs in the region. However, the
357 contributions of vehicle related emissions (i.e., ~25%) and industrial emissions were lower and higher than those
358 quantified by the PMF model in this study, respectively. The observed discrepancy between the two studies may
359 be due to differences in source categories, measured VOC species, and/or sampling locations and methods used
360 in the different models. For example, the VOC sources in Jiangsu province were categorized into vehicular
361 related emission (~26%), industrial solvent usage (~25%), fossil fuel combustion (~24%), industrial processes
362 (~22%) and biomass burning (~3%). Further, vehicle related emissions only included emissions from motor
363 vehicles and ships, and the volatilization of fuel, while solvent usage included organic solvents volatilized from
364 a variety of industries (the industrial produce process of electronic equipment manufacturing, furniture
365 manufacturing, printing, packaging, inks, adhesives, etc. and other dry cleaning, catering, and architectural
366 decoration processes). Higher vehicular emission contribution in this study may also be due to the increasing
367 number of vehicles from 2010-2014 as a result of increased urbanization and industrialization (Statistical
368 yearbook of Nanjing, 2014).

369
370 Figure 5 illustrates the mean diurnal variability of all identified source at the JAES site. These trends were
371 influenced by the variability in emission strength, mixing height, and the concentrations and photochemical
372 reactivity of individual species in each source profile. For example, we observed a typical diurnal pattern with a
373 broad peak between 9 am-6 pm for biogenic emissions, as the emission rate of isoprene from vegetation is largely
374 depended on ambient temperature and sunlight intensity. Higher levels of diesel and gasoline vehicular emissions
375 were observed in the evening and early morning due to a reduced mixing height and increased emissions from
376 the morning and evening rush hour. Lower concentrations observed during daytime hours were likely due to
377 decreased emissions, an increased mixing height and enhanced photochemical loss (Gillman et al., 2009; Yuan
378 et al., 2009; Wang et al., 2013). A diurnal pattern of fuel evaporation that was similar to that of vehicular
379 emissions. Though the evaporation of fuel is dependent on temperature, the average temperature in the morning
380 and evening (i.e., 0800-1000 and 1700-1900 LT, respectively) when peaks of fuel evaporation were found was
381 only about ~1.2 °C lower than that observed from noon to afternoon (1100-1600 LT), which may not result in

382 much higher fuel evaporation at noon (the difference between maximum and minimum values for fuel
 383 evaporation was found to be $\sim 6 \mu\text{g}/\text{m}^3$). On the other hand, in addition to evaporation from the gas station, fuel
 384 could evaporate from hot engines, fuel tanks and the exhaust system when the car is running. Furthermore, the
 385 engine remains hot for a period of time after the car is turned off, and gasoline evaporation continues when the
 386 car is parked (Technology center, University of Illinois, <https://mste.illinois.edu/tcd/ecology/fuelevap.html>,
 387 access date: 25 December 2019). The similarity of diurnal variations of fuel evaporation to vehicular emissions
 388 suggested that the prominent peak in the morning and evening hours were related to the increased vehicles in
 389 the traffic rush hour and emissions accumulated in the relatively low boundary layer. Moreover, we identified
 390 higher concentrations of industrial emissions at night and in the early morning, with values remaining fairly
 391 stable during daytime hours. This finding is consistent with other observations in urban and rural areas (Yuan et
 392 al., 2009; Leuchner and Rappenglück, 2010).




393
 394 **Figure 5. Diurnal patterns in source concentrations of the five identified sources**

395 3.4 Contributions of VOC sources to O₃ formation

396 As important O₃ precursors, information on the contributions of VOC sources and related species to O₃ formation
 397 is necessary for the formulation and implementation of VOC control measures. To achieve this goal, the
 398 Maximum Incremental Reactivity (MIR) method, which evaluates the O₃ formation potential (OFP) on the basis
 399 of mass concentrations and maximum incremental reactivities of VOCs with the OH radical, were adopted in
 400 the present study (Shao et al., 2009b, 2011; Mo et al., 2017). Figure 6 presented the relative contributions of
 401 individual VOC sources and related VOC species from PMF to OFP at the JAES site. Industrial emissions were
 402 found to have the largest OFP at JAES due to the high loadings of aromatic VOC species that have relatively
 403 high OH reactivities in this source profile (Atkinson and Arey, 2003), with the OFP value of $\sim 43 \mu\text{g}/\text{m}^3$ and the
 404 contribution percentage of $\sim 32\%$ to the total OFP of all VOC sources, followed by diesel vehicular exhausts

405 (~36 $\mu\text{g}/\text{m}^3$, ~27%), gasoline vehicular exhausts (~32 $\mu\text{g}/\text{m}^3$, ~24%) and fuel evaporation (~13 $\mu\text{g}/\text{m}^3$, ~10%).
 406 Furthermore, though the MIR value of isoprene was much higher than other VOC species, biogenic emissions
 407 only contributed ~7% (~9 $\mu\text{g}/\text{m}^3$) to the total OFP of all VOC sources as the relatively low mixing ratio of
 408 isoprene at the JAES site. Similarly, using the same method to evaluate OFP of different VOC sources, Mo et al.
 409 (2017) found that industrial emissions (including the emissions of petrochemical industry, chemical and paint
 410 industries, solvent usage) and vehicular emissions were the dominant VOC sources of the total OFP in an
 411 industrialized coastal city (i.e., Ningbo) in the YRD region. Therefore, our results further demonstrated the need
 412 to minimize VOC emissions from industrial emissions and vehicle exhausts in order to lower O_3 formation and
 413 photochemical pollution in YRD.



414
 415 **Figure 6. (a) The contribution of individual sources to the total OFP of all sources extracted from PMF and (b) OFP values**
 416 **of the top 10 VOC species in the different source categories.**

417
 418 Based on the mass concentrations of individual species in each source, we found that *m,p*-xylene and toluene in
 419 industrial emissions and gasoline vehicular emissions, propene, ethene, toluene and *m,p*-xylene in diesel
 420 vehicular emissions, and *o*-xylene, 1,2,4-trimethylbenzene and ethene in industrial emissions to be the dominant
 421 species from VOC emissions contributing to photochemical O_3 formation. Thus, only a small number of VOC
 422 species can be monitored for the effective control of O_3 formation.

423
 424 **3.5 Policy summary and implications**

425 To effectively control photochemical pollution, the Prevention and Control of Atmospheric Pollution Act was
 426 passed in 1987 and amended in 2015. As a result, a series of measures to prevent and control VOCs levels have
 427 been and are being implemented by central and local governments, including the implementation of new laws
 428 and regulations, and the advancement of technology. The results of this study suggest that photochemical O_3

429 formation within the urban areas of Nanjing city are VOC-limited, which is consistent with observations in the
430 urban locations of other regions, including the North China Plain, the Yangtze River Delta and the Pearl River
431 Delta. Minimizing VOC emissions and their concentrations should therefore be prioritized in order to alleviate
432 O₃ pollution in urban environments. The prevention and control of VOC pollution has been listed as one of the
433 key tasks of “the Blue Sky” Project initiated in 2012 by the Department of Environmental Protection of Jiangsu
434 Province. Furthermore, the administrative measures on the Prevention and Control of Volatile Organic
435 Compounds Pollution in Jiangsu (Order No. 119 of the Provincial Government) was enacted on March 6, 2018
436 and implemented on May 1, 2018, with the aim of controlling VOC emissions in Jiangsu Province.

437
438 In order to achieve these goals, various measures have been implemented (Table S2), including: 1) investigating
439 the current pollution status and identifying the progress of VOC prevention and control in Jiangsu Province
440 (Provincial Office of the Joint Conference on the prevention and control of air pollution [2012] No. 2); 2)
441 conducting a strict industry access system, under the Advice on Promoting Air Pollution Joint Prevention and
442 Control Work to Improve Regional Air Quality (Office of the State Council [2010] No. 33); 3) strengthening the
443 remediation on existing sources of VOCs and reducing VOC emissions from these sources, under the Guidelines
444 for the Implementation of Leak Detection and Repair (LDAR) in Jiangsu Province (Trial) (Provincial Office of
445 Environmental Protection [2013] No. 318); 4) strengthening the VOC monitoring capacity, under the Guidelines
446 for Control of Volatile Organic Compounds Pollution in Key Industries in Jiangsu Province (Provincial Office
447 of Environmental Protection [2013] No. 128); 5) improving standards regarding VOC emissions for key
448 industries, including standards for surface coating of the automobile manufacturing industry (DB32/2862-2016),
449 the chemical industry (DB32/3151-2016), and furniture manufacturing operations (DB32/3152-2016), which are
450 still effective since their enforcement; 6) implementing the Pilot Measures for Volatile Organic Compounds
451 Discharge Charges (Ministry of Finance [2015] No. 71) on October 1, 2015 to raise awareness pertaining to
452 emissions reduction in factories and to control VOC emissions from industrial sources; 7) encouraging the public
453 to live a low-carbon life and supervise and offer recommendations in accordance with the laws, under the
454 Measures for Public Participation in Environmental Protection in Jiangsu Province (Trial) (Provincial Regulation
455 of Environmental Protection Office [2016] No. 1).

456
457 Based on the VOC source apportionment results in this study, we identified vehicular emissions and industrial
458 emissions as the two major VOC sources contributing to photochemical O₃ formation. Other measures and/or
459 regulations have been conducted in the Jiangsu Province to effectively control VOC emissions from vehicles
460 and industry. For vehicular emissions, the Regulations on Prevention and Control of Vehicle Exhaust Pollution
461 in Nanjing was amended in July 2017, and subsequently in March, 2018
462 (<http://hbt.jiangsu.gov.cn/col/col11590/index.html>). The new regulation not only focusses on vehicle emissions,
463 but also incorporates a number of additional topics, including optimizing the function and distribution of urban
464 areas, limiting the number of vehicles in the region, promoting new green energy vehicles, and improving the

465 quality of fuel. The promotion of intelligent traffic management, implementation of a priority strategy for public
466 transportation, and construction of more efficient traffic systems to promote pedestrian and bicycle use is
467 recommended. Further studies should be conducted to estimate and manage the increasing quantity of vehicles
468 on the road. As of January 1, 2017, regulation stipulate that all new and used vehicles should meet the fifth phase
469 of vehicle emission standards, including vehicle manufacture, sales, registration and importation. For vehicles
470 already in use, an environmental protection examination should be conducted annually, based on the standards
471 of GB 14622-2016, GB 18176-2016, GB 19755-2016, and HJ 689-2014. Penalties are issued if qualified
472 vehicles excessively emit pollutants due to poor maintenance.

473
474 For industrial emissions, various policies have been implemented to reduce VOC emissions, particularly in
475 chemical industries: including, 1) investigations on the VOC emissions of the chemical industry and the
476 establishment of an archive system for VOC pollution control, particularly the inspection of industry information,
477 products and materials, unorganized emission of storage and exhaust gas treatment facilities, under the Plan for
478 Investigation of Volatile Organic Pollutant Emissions in Jiangsu Province, mentioned in the Provincial Office of
479 Environmental Protection [2012] No. 183; 2) exhaust gas remediation in the chemical industry park, under the
480 Technical Specifications for Prevention and Control of Air Pollution in Chemical Industries in Jiangsu Province
481 (Provincial Office of Environmental Protection [2014] No. 3), which requires the establishment of the long-term
482 supervision of exhaust gas remediation in the chemical industry park of Jiangsu Province; 3) a pilot project on
483 the leak detection and repair (LDAR) technology in the chemical industry park, under the notification on carrying
484 out the technical demonstration and pilot work of leak detection and repair (LDAR) in petrochemical and
485 chemical industries (Provincial Office of Environmental Protection [2015] No. 157). The TVOC removal
486 efficiency of organic exhaust vents should be >95%, and higher for areas of excessive environmental pollution
487 at >97% (GB 31571-2015).

488
489 Furthermore, though measures have been adopted to improve standards and control vehicle VOC emissions,
490 most of these policies only focus on total VOC emissions (or the mass of total emissions) and do not consider
491 the impacts of individual VOC species. To accelerate the implementation of existing policies and to strengthen
492 collaborative regional prevention and control, priority should be placed on specific high-impact VOC species
493 (i.e., *m,p*-xylene and toluene in the industrial emission and gasoline vehicular emission) by considering both
494 their reactivity and abundance.

495
496 Last but not the least, O₃ pollution is a regional cross-boundary environmental issue rather than a local pollution
497 problem. Apart from VOCs, NO_x was another important precursor for O₃ formation with its dual roles in O₃
498 production (enhancing O₃ formation in non NO_x-saturated environment and titrating O₃ in NO_x-saturated
499 environment). In other areas (i.e., the rural environment and/or the downwind areas of urban center in the same
500 region) where the concentrations of NO_x are low and/or there is a non NO_x-saturated environment, the situation

501 may be different and controlling VOCs should be conducted cautiously (Zheng et al., 2010; Yuan et al., 2013;
502 Ou, et al., 2016). Therefore, from a regional perspective, the benefits of VOCs control measures could be further
503 evaluated with those of NO_x (i.e., the appropriate ratios of VOC/NO_x for the reduction of O_3 pollution) as well
504 as the associated O_3 -VOCs- NO_x sensitivity. Therefore, one important concern for the policy formulation and
505 implementation system is whether controlling VOCs and NO_x individually or controlling both VOCs and NO_x
506 is more effective and appropriate for alleviating O_3 pollution. It is necessary to consider the reduction ratios of
507 VOC/NO_x when VOCs and NO_x are simultaneously controlled. Finally, long-term monitoring studies are
508 necessary to determine the cost-benefits and performance of each policy.

509 **4. Conclusion**

510 In this study, a one-year field sampling campaign was conducted to investigate the VOC characteristics at an
511 urban site in Nanjing (the JAES site), Jiangsu province. In total, 56 VOCs including 29 alkanes, 10 alkenes, 16
512 aromatics and acetylene were identified and quantified. The composition analysis found that alkanes were the
513 dominant group of VOCs observed at the JAES site (~53%), followed by aromatics, acetylene, and alkenes. This
514 finding is consistent with the VOC measurements in studies conducted in the North China Plain, Pearl River
515 Delta, and Yangtze River Delta. We observed distinct seasonal patterns of TVOCs, with maximum values in
516 winter and minimum values in summer. Similarly, prominent morning and evening peaks were observed in the
517 diurnal variability of TVOCs, influenced by local emissions and meteorology.

518
519 Based on the observed VOC data, we identified five dominant VOC sources at the JAES site using a PMF model.
520 By considering both the abundance and maximum incremental reactivity of individual VOC species in each
521 source, the OFP values identified industrial and vehicular emissions, particularly *m,p*-xylene, toluene and
522 propene, as the main contributors of O_3 pollution. Local governments have strengthened several measures to
523 minimize VOC pollution from vehicle and industrial emissions in the Jiangsu province in recent years, though
524 most of these policies focus particularly on lowering the total emissions of VOCs. Furthermore, from a regional
525 perspective, it is suggested that appropriate ratios of VOC/NO_x , their associated sensitivity to O_3 formation and
526 relative benefits/disbenefits of reducing VOCs/NO_x should be investigated and evaluated when control measures
527 of VOCs and NO_x were both conducted.

528

529 **Author Contributions.** Jun Bi, Zhenhao Ling, and Qiuyue Zhao designed the research and carried them out.
530 Zhenhao Ling performed the data simulation. Qiuyue Zhao and Guofeng Shen performed the observation data
531 analysis. Qiuyue Zhao prepared the manuscript with contributions from all co-authors.

532 **Competing Interests.** The authors declare that they have no conflict of interest.

533 **Acknowledgements.** This work was supported by National Key R&D Program of China (No.2016YFC0207607,
534 2017YFC0210106), National Science Foundation of Jiangsu Province of China (General Program, No.
535 BK20161601), Open Research Fund of Jiangsu Province Key Laboratory of Environmental Engineering
536 (No.ZX2016002) and National Natural Science Foundation of China (No. 41775114). This work was also partly
537 supported by the Pearl River Science and Technology Nova Program of Guangzhou (grant no. 201806010146).

538 Reference

- 539 An, J., Zhu, B., Wang, H., Li, Y., Lin, X., and Yang, H.: Characteristics and source apportionment of VOCs
540 measured in an industrial area of Nanjing, Yangtze River Delta, China, *Atmospheric Environment*, 97, 206-
541 214, 2014
- 542 An, J., Zou, J., Wang, J., Lin, X., and Zhu, B.: Differences in ozone photochemical characteristics between the
543 megacity Nanjing and its suburban surroundings, Yangtze River Delta, China, *Environmental Science and
544 Pollution Research International*, 22(24), 19607, 2015.
- 545 Barletta, B., Meinardi, S., Rowland, F.S., Chan, C.Y., Wang, X.M., Zou, S.C., Chan, L.Y. and Blake, D.R.:
546 Volatile organic compounds in 43 Chinese cities, *Atmos. Environ.*, 39 (32), 5979-5990, 2005.
- 547 Baudic, A., Gros, V., Sauvage, S., Locoge, N., Olivier, S., Sarda-Estève, R., Kalogridis, C., Petit, J.-E., Bonnaire,
548 N., Baisnee, D., Favez, O., Albinet, A., Sciare, J. and Bonsang, B.: Seasonal variability and source
549 apportionment of volatile organic compounds (VOCs) in the Paris megacity (France), *Atmospheric
550 Chemistry and Physics*, 16, 11961-11989, 2016.
- 551 Cai, C. J., Geng, F. H., Tie, X.X., and An, J. L.: Characteristic and source apportionment of VOCs measured in
552 Shanghai, China, *Atmospheric Environment*, 44, 5005-5014, 2010.
- 553 Ding, A.J., Nie, W., Huang, X., Chi, X.G., Xu, Z., et al.: Ozone in the western Yangtze River Delta of China: a
554 synthesis study based on ground, aircraft and sounding measurement 2011-2015, The 6th PEEEX meeting,
555 May 2016, 2016
- 556 Fu, L., Wan, W., Zhang, W., and Cheng, H.: Air pollution prevention and control progress in Chinese cities,
557 *Clean Air Asia*, China, available at: [http://cleanairasia.org/wp-content/uploads/2016/08/China-Air-2016-
558 Report-Full.pdf](http://cleanairasia.org/wp-content/uploads/2016/08/China-Air-2016-Report-Full.pdf), 2016, last access: 16 May 2018.
- 559 Gardiner, M.S. and Lange, C.R.: Comparison of laboratory generated and field-obtained HMA VOCs with odour
560 potential, *The International Journal of Pavement Engineering*, 6, 257-263, 2005.
- 561 Gilman, J.B., Kuster, W.C., Goldan, P.D., Herndon, S.C., Zahniser, M.S., Tucker, S.C., Brewer, W.A., Lerner,
562 B.M., Williams, E.J., Harley, R.A., Fehsenfeld, F.C., Warneke, C. and de Gouw, J.A.: Measurements of
563 volatile organic compounds during the 2006 TexAQs/GoMACCS campaign: industrial influences, regional
564 characteristics, and diurnal dependencies of the OH reactivity, *Journal of Geophysical Research*, 114,
565 D7, doi: 10.1029/2008JD011525, 2009.
- 566 Guo, H., Cheng, H. R., Ling, Z. H., Louie, P. K. K., and Ayoko, G. A.: Which emission sources are responsible
567 for the volatile organic compounds in the atmosphere of Pearl River Delta?, *Journal of Hazardous Materials*,
568 188, 116-124, 2011a.
- 569 Guo, H., Zheng J. Y., Characterisation of VOC sources and integrated photochemical ozone analysis in Hong
570 Kong and the Pearl River Delta region. Report to the Environmental Protection Department of Hong Kong
571 (Tender Reference AS12-02909), Department of Civil and Environmental Engineering, The Hong Kong
572 Polytechnic University, 2016.

573 Guo, H., Zou, S. C., Tsai, W. Y., Chan, L. Y., and Blake, D. R.: Emission characteristics of non-methane
574 hydrocarbons from private cars and taxis at different driving speeds in Hong Kong, *Atmos. Environ.*, 45,
575 2711–2721, 2011b.

576 Guo, J.P., Miao, Y.C., Zhang, Y., Liu, H., Li, Z.Q., Zhang, W.C., He, J., Lou, M.Y., Yan, Y., Bian, L.G. and Zhai,
577 P.M.: The climatology of planetary boundary layer height in China derived from radiosonde and reanalysis
578 data, *Atmospheric Chemistry and Physics*, 16, 13309-13319, 2016.

579 He, Z., Wang, X., Ling, Z., Zhao, J., Guo, H., Shao, M., and Wang, Z.: Contributions of different anthropogenic
580 volatile organic compound sources to ozone formation at a receptor site in the Pearl River Delta region and
581 its policy implications, *Atmos. Chem. Phys.*, 19, 8801-8816, 2019.

582 HKEPD (Hong Kong Environmental Protection Department): Characterisation of VOC sources and integrated
583 photochemical ozone analysis in Hong Kong and the Pearl River Delta region. Final report, , Hong Kong,
584 2015.

585 Ho, K.F., Lee, S.C., Ho, W.K., Blake, D.R., Cheng, Y., Li, Y.S., Ho, S.S.H., Fung, K., Louie, P.K.K. and Park,
586 D.: Vehicular emission of volatile organic compounds (VOCs) from a tunnel study in Hong Kong,
587 *Atmospheric Chemistry and Physics*, 9, 7491-7504, 2009.

588 Huang, C., Chen, C. H., Li, L., Cheng, Z., Wang, H. L., Huang, H. Y., Streets, D. G., Wang, Y. J., Zhang, G. F.,
589 and Chen, Y.R.: Emission inventory of anthropogenic air pollutants and VOC species in the Yangtze River
590 Delta region, China. *Atmos. Chem. Phys.*, 11, 4105-4120, 2011a

591 Jenkin, M. E. and Clemitshaw, K. C.: Ozone and other secondary photochemical pollutants: chemical processes
592 governing their formation in the planetary boundary layer, *Atmospheric Environment*, 34, 2499-2527, 2000.

593 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and
594 Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asia regions during 2000-2008:
595 Regional Emission inventory in Asia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058, 2013.

596 Lau, A. K. H., Yuan, Z., Yu, J. Z., and Louie, P. K.: Source apportionment of ambient volatile organic compounds
597 in Hong Kong, *Science of the Total Environment*, 408, 4138–4149, 2010.

598 Leuchner, M. and Rappenglück, B.: VOC source-receptor relationships in Houston during TexAQS-II,
599 *Atmospheric Environment*, 44, 4056-4067, 2010.

600 Liu, B., Liang, D., Yang, J., Dai, Q., Bi, X., and Feng, Y., Yuan, J., Xiao, Z.M., Zhang, Y.F., Xu, H.:
601 Characterization and source apportionment of volatile organic compounds based on 1-year of observational
602 data in Tianjin, China. *Environmental Pollution*, 218, 757-769 2015.

603 Liu, Y., Shao, M., Fu, L. L., Lu, S. H., Zeng, L. M., and Tang, D. G.: Source profiles of volatile organic
604 compounds (VOCs) measured in China: Part I, *Atmospheric Environment* 42, 6247-6260, 2008a.

605 Liu, Y., Shao, M., Lu, S., Chang, C. C., Wang, J. L. and Fu, L.: Source apportionment of ambient volatile organic
606 compounds in the Pearl River delta, China: part II, *Atmospheric Environment*, 42(25), 6261-6274, 2008b.

607 Liu, Y., Shao, M., Lu, S., Chang, C., Wang, J. L., and Chen, G.: Volatile organic compound (VOC) measurements
608 in the Pearl River delta (PRD) region, China, *Atmos. Chem. Phys.*, 7(5), 1531-1545, 2008c.

609 Lyu, X., Chen, N., Guo, H., Zhang, W., Wang, N., Wang, Y., and Liu, M.: Ambient volatile organic compounds
610 and their effect on ozone production in Wuhan, central China, *Science of the Total Environment*, 548-
611 549(60), 483-483, 2016.

612 Lyu, X.P., Wang, N., Guo, H., Xue, L.K., Jiang, F., Zeren, Y.Z., Cheng, H.R., Cai, Z., Han, L.H. and Zhou, Y.:
613 Causes of a continuous summertime O₃ pollution event in Jinan, a central city in the North China Plain,
614 *Atmospheric Chemistry and Physics* 19, 3025-3042, 2019.

615 Mo, Z., Shao, M., and Lu, S.: Compilation of a source profile database for hydrocarbon and OVOC emissions
616 in China, *Atmospheric Environment*, 143, 209-217, 2016.

617 Mo, Z., Shao, M., Lu, S. H., Niu, H., Zhou, M. Y., Sun, J.: Characterization of non-methane hydrocarbons and
618 their sources in an industrialized coastal city, Yangtze River Delta, China, *Science of the Total Environment*,
619 593-594, 641-653, 2017.

620 Mo, Z., Shao, M., Lu, S., Qu, H., Zhou, M., and Gou, B.: Process-specific emission characteristics of volatile
621 organic compounds (VOCs) from petrochemical facilities in the Yangtze River Delta, China, *Science of the*
622 *Total Environment*, 533, 422-431, 2015.

623 Ou, J., Guo, H., Zheng, J., Cheung, K., Louie, P. K. K., Ling, Z., and Wang D. W.: Concentrations and sources
624 of non-methane hydrocarbons (NMHCs) from 2005 to 2013 in Hong Kong: a multi-year real-time data
625 analysis, *Atmospheric Environment*, 103(103), 196-206, 2015.

626 Ou, J.M., Yuan, Z.B., Zheng, J.Y., Huang, Z.J., Shao, M., Li, Z.K., Huang, X.B., Guo, H. and Louie, P.K.K.:
627 Ambient ozone control in a photochemically active region: short-term despiking or long-term attainment?
628 *Environmental Science and Technology*, 50, 5720-5728, 2016.

629 Pan, X., Kanaya, Y., Tanimoto, H., Inomata, S., Wang, Z., Kudo, S., and Uno, I.: Examining the major
630 contributors of ozone pollution in a rural area of the Yangtze River Delta region during harvest season,
631 *Atmos. Chem. Phys.*, 15, 6101-6111, 2015

632 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: from air pollution to climate change*, 2nd
633 edition. Wiley Publisher, New Jersey, USA, 2006.

634 Shao, M., Lu, S., Liu, Y., Xie, X., Chang, C., Huang, S., and Chen, Z.: Volatile organic compounds measured in
635 summer in Beijing and their role in ground-level ozone formation, *Journal of Geophysical Research*, 114,
636 D00G06, doi: 10.1029/2008JD010863, 2009a.

637 Shao, M., Zhang, Y.H., Zeng, L.M., Tang, X.Y., Zhang, J., Zhong, L.J. and Wang, B.G.: Ground-level ozone in
638 the Pearl River Delta and the roles of VOC and NO_x in its production, *Journal of Environmental*
639 *Management*, 90, 512-518, 2009b.

640 Shao, P., An, J.L., Xin, J.Y., Wu, F.K., Wang, J.X., Ji, D.S. and Wang, Y.S.: Source apportionment of VOCs and
641 the contribution to photochemical ozone formation during summer in the typical industrial area in the
642 Yangtze River Delta, China, *Atmospheric Research*, 176-177, 64-74, 2016.

643 Song, Y., Dai, W., Shao, M., Liu, Y., Lu, S. H., Kuster, W, and Goldan, P.: Comparison of receptor models for
644 source apportionment of volatile organic compounds in Beijing, China, *Environmental Pollution*, 156, 174-
645 183, 2008.

646 Statistical yearbook of Nanjing. Nanjing Statistic Bureau:
647 <http://221.226.86.104/file/nj2004/2014/jiaotongyunshu/index.htm>, last access: 17 May 2018, 2014

648 Tasi, W.Y., Chan, L.Y., Blake, D.R. and Chu, K.W.: Vehicular fuel composition and atmospheric emissions in
649 South China: Hong Kong, Macau, Guangzhou, and Zhuhai, *Atmospheric Chemistry and Physics*, 6, 3281-
650 3288, 2006.

651 von Schneidemesser, E., Monks, P.S. and Plass-Duelmer, C.: Global comparison of VOC and CO observations
652 in urban areas, *Atmospheric Environment*, 44, 5053-5064, 2010.

653 Wang, H. L., Lou, S. R., Huang, C., Qiao, L. P., Tang, X. B., Chen, C. H., Zeng, L. M., Wang, Q., Zhou, M., Lu,
654 S. H., and Yu, X.N.: Source profiles of volatile organic compounds from biomass burning in Yangtze River
655 Delta, China, *Aerosol and Air Quality Research*, 14, 818-828, 2014

656 Wang, H., Chen, C., Wang, Q., Huang, C., Su, L., Huang, H., Lou, S., Zhou, M., Li, L., Qiao, L., and Wang, Y.:
657 Chemical loss of volatile organic compounds and its impact on the source analysis through a two-year
658 continuous measurement, *Atmospheric Environment*, 80, 488-498, 2013.

659 Wang, M., Shao, M., Chen, W., Lu, S., Liu, Y., Yuan, B., and Li, Y.: Trends of non-methane hydrocarbons
660 (NMHCs) emissions in Beijing during 2002-2013, *Atmos. Chem. Phys.*, 15, 1489-1502, 2015.

661 Wang, N., Guo, H., Jiang, F., Ling, Z.H. and Wang, T.: Simulation of ozone formation at different elevations in
662 mountainous area of Hong Kong using WRF-CMAQ model, *Science of the Total Environment*, 505, 939-
663 951, 2015.

664 Wang, T., Wei, X. L., Ding, A. J., Poon, C. N., Lam, K. S., Li, Y. S., Chan, L. Y., and Anson, M.: Increasing
665 surface ozone concentrations in the background atmosphere of Southern China, 1994–2007, *Atmos. Chem.*
666 *Phys.*, 9, 6217-6227, 2009.

667 Warneke, C., de Gouw, J.A., Holloway, J.S., Peischl, J., Ryerson, T.B., Atlas, E., Blake, D., Trainer, M. and
668 Parrish, D.D.: Multiyear trends in volatile organic compounds in Los Angeles, California: Five decades of
669 decreasing emissions, *Journal of Geophysical Research*, 117, D00V17, doi: 10.1029/2012JD017899, 2012.

670 Xia, L., Cai, C., Zhu, B., An, J., Li, Y., and Li, Y.: Source apportionment of VOCs in a suburb of Nanjing, China,
671 in autumn and winter, *Journal of Atmospheric Chemistry*, 71(3), 175-193, 2014a.

672 Xia, S., Zhao, Q., Li, B., and Shen, G.: Anthropogenic source VOCs emission inventory of Jiangsu Province (In
673 Chinese), *Research of Environmental Sciences*, 27(2), 120-126, 2014b.

674 Xu, Z. N., Huang, X., Nie, W., Chi, X. G., Xu, Z., Zheng, L. F., Sun, P., Ding, and A. J.: Influence of synoptic
675 condition and holiday effects on VOCs and ozone production in the Yangtze River Delta region, China,
676 *Atmospheric Environment*, 168, 112-124, 2017.

677 Yuan, B., Chen, W.T., Shao, M., Wang, M., Lu, S.H., Wang, B., Liu, Y., Chang, C.C., Wang, B: Measurements
678 of ambient hydrocarbons and carbonyls in the Pearl River Delta (PRD), China, *Atmos. Res.*, 116, 93-104,

2012.

680 Yuan, Z., Lau, A. K. H., Shao, M., Louie, P. K. K., Liu, S. C., and Zhu, T.: Source analysis of volatile organic
681 compounds by positive matrix factorization in urban and rural environments in Beijing, *Journal of*
682 *Geophysical Research*, 114, D00G15, doi:10.1029/2008JD011190, 2009.

683 Yuan, Z.B., Zhong, L.J., Lau, A.K.H., Yu, J.Z. and Louie, P.K.K.: Volatile organic compounds in the Pearl River
684 Delta: identification of source regions and recommendation for emission-oriented monitoring strategies,
685 *Atmospheric Environment*, 76, 162-172, 2013.

686 Zhang, J., Wang, Y., Wu, F., Lin, H., and Wang, W.: Nonmethane hydrocarbon measurements at a suburban site
687 in Changsha City, China. *Sci. Total Environ.*, 408 (2), 312-317, 2009.

688 Zhang, X., Xue, Z., Li, H., Yan, L., Yang, Y., Wang, Y., Duan, J., Li, L., Chai, F., Cheng, M., and Zhang, W.:
689 Ambient volatile organic compounds pollution in China, *Journal of Environmental Sciences*, 55, 69-75,
690 2017.

691 Zhang, Y. L., Wang, X. M., Barletta, B., Simpson, I.J., Blake, D.R., Fu, X.X., Zhang, Z., He Q.F., Liu, T.Y., Zhao,
692 X.Y., and Ding, X.: Source attributions of hazardous aromatic hydrocarbons in urban, suburban and rural
693 areas in the Pearl River Delta (PRD) region, *Journal of Hazardous Materials*, 250-251, 403-411, 2013.

694 Zhang, Y. X., An, J. L., Wang, J. X., Shi, Y. Z., Liu, J. D., and Liang, J. S.: Source analysis of volatile organic
695 compounds in the Nanjing industrial area and evaluation of their contribution to ozone, *Environmental*
696 *Science*, 39(2), 502-510, 2018.

697 Zhang, Y.L., Wang, X., Barletta, B., Simpson, I.J., Blake, D.R., Fu, X., Zhang, Z., He, Q., Liu, T., Zhao, X. and
698 Ding, X.: Source attributions of hazardous aromatic hydrocarbons in urban, suburban and rural areas in the
699 Pearl River Delta (PRD) region, *Journal of Hazardous Materials*, 250, 403-411, 2013.

700 Zhang, Y.L., Wang, X.M., Blake, D.R., Li, L.F., Zhang, Z., Wang, S.Y., Guo, H., Lee, F.S.C., Gao, B., Chan,
701 L.Y., Wu, D. and Rowland, F.S.: Aromatic hydrocarbons as ozone precursors before and after outbreak of
702 the 2008 financial crisis in the Pearl River Delta region, south China, *Journal of Geophysical Research*, 117,
703 D15306, doi:10.1029/2011JD017356, 2012

704 Zhang, Y.L., Yang, W.Q., Simpson, I.J., Huang, X.Y., Yu, J.Z., Huang, Z.H., Wang, Z.Y., Zhang, Z., Liu, D.,
705 Huang, Z.Z., Wang, Y.J., Pei, C.L., Shao, M., Blake, D.R., Zheng, J.Y. and Huang, Z.J.: Decadal changes
706 in emissions of volatile organic compounds (VOCs) from on-road vehicles with intensified automobile
707 pollution control: case study in a busy urban tunnel in south China, *Environmental Pollution*, 233, 806-819,
708 2018.

709 Zhao, Q, Shen, G., Li, L., Chen, F., Qiao, Y.Z., Li C.Y., Liu Q., and Han, J.Z.: Ambient Particles (PM₁₀, PM_{2.5},
710 and PM_{1.0}) and PM_{2.5} chemical components in Western Yangtze River Delta (YRD): An Overview of Data
711 from 1-year Online Continuous Monitoring at Nanjing, *Aerosol Science and Engineering*, 1, 107-118, 2017.

712 Zheng, J. Y., Shao, M., Che, W. W., Zhang, L. J., Zhong, L. J., Zhang, Y. H., and Streets, D.: Speciated VOC
713 emission inventory and spatial patterns of ozone formation potential in the Pearl River Delta, China,
714 *Environ. Sci. Technol.*, 43, 8580-8586, 2009.

715 Zheng, J.Y., Yu, Y.F., Mo, Z.W., Zhang, Z., Wang, X.M., Yin, S.S., Peng, K., Yang, Y., Feng, X.Q. and Cai, H.H.:
716 Industrial sector-based volatile organic compound (VOC) source profiles measured in manufacturing
717 facilities in the Pearl River Delta, China, *Science of the Total Environment*, 456-457, 127-136, 2016.

718 Zheng, J.Y., Zhong, L.J., Wang, T., Louie, P.K.K. and Li, Z.C.: Ground-level ozone in the Pearl River Delta
719 region: analysis of data from a recently established regional air quality monitoring network, *Atmospheric*
720 *Environment* 44, 814-823, 2010.

721 Zhou, D., Li, B., Huang X., Virkkula, A., Wu, H.S., Zhao, Q.Y., Zhang, J., Liu, Q., Li, L., Li, C.Y., Chen, F.,
722 Yuan, S.Y., Qiao, Y.Z., Shen, G.F., and Ding A.J.: The impacts of emission control and regional transport
723 on PM_{2.5} ions and carbon components in Nanjing during the 2014 Nanjing Youth Olympic Games, *Aerosol*
724 *and Air Quality Research*, 17(3), 730-740, 2017.

725 Zou, Y., Deng, X. J., Zhu, D., Gong, D. C., Wang, H., Li, F., Tan, H. B., Deng, T., Mai, B. R., Liu, X. T., and
726 Wang, B. G.: Characteristics of 1 year of observational data of VOCs, NO_x and O₃ at a suburban site in
727 Guangzhou, China, *Atmos. Chem. Phys.*, 15, 6625–6636, 2015.