

We appreciate the reviewer for carefully reviewing our manuscripts and providing the valuable suggestion to improve quality and readability for our paper. We carefully revised the manuscripts as suggested. The following are our responses to your comments point by point. The red sentences are the reviewer's comments (italics), the black sentences are the author's responses, and the blue sentences and words are the specific revisions.

Anonymous Referee #2

This paper is a case study of a period in November and December 2015 that was characterized by many haze days in the North China plain. The authors describe the synoptic conditions that led to an environment that was conducive to the occurrence of haze and relate this to the larger scale waveguide and quasi-stationary wave environment. Overall, it's concluded that Rossby wave trains propagating from Western Europe along two waveguides toward east Asia were responsible for setting up decent over the North China plain and a weaker than normal winter monsoon - both of which were favorable for haze. Overall, I found this to be a worthwhile study and I think it nicely describes the conditions that have led to this event. In the end, there is also some discussion of the extent to which other events have related to this kind of large-scale environment, which I think is valuable. I have only relatively minor comments to suggest before publications, although I do have quite a number of them that are mostly aimed at improving readability. I think there is some confusion throughout the text of the difference between the waveguide and the wave themselves. The authors often refer to waveguide when I think they should really be referring to the waves. I've pointed out a couple of cases below, but I suggest attention be paid to that during the revisions.

Response: We thanks for your suggestions and we have revised our manuscripts as suggested which is marked in the blue in the new manuscripts.

Minor comments by line number

116: it is stated that "two Rossby waveguides within the westerly jet" are responsible for the haze. But isn't it really the Rossby waves that propagate along these waveguides that are ultimately responsible. If so, the wording could be clearer with something along the lines of "...anomalous Rossby waves that propagated along two waveguides within the westerly jet..."

Response: We thank the reviewer for the suggestion. Indeed, it is the Rossby waves that propagate along two waveguides within the westerly jet. We agree with the reviewer and the sentence has been revised as suggested. In addition, we have revised some other similar statements.

115: In this paper, the combined effect of **the anomalous Rossby waves within two westerly jet waveguides** on...

117: ...that the **anomalous Rossby waves that propagated along two waveguides within the westerly jet** originating from the...

168: In summary, **the Rossby waves within the East Asian upper westerly jet waveguides** has important influences on the East Asian...

172-75: ...circulations in the upper troposphere on haze, especially the combined effect of **the Rossby waves within two westerly jet waveguides**, are rare. Given the above content, the objective of the present study is to determine whether the effects of **the Rossby waves within two westerly jet waveguides** on haze were significant in the NCP in November and December 2015 and, if so, to identify the principal mechanism behind the effects of **the Rossby waves within two westerly jet waveguides** on haze.

178: Section 4 demonstrates the influencing mechanisms of **the Rossby waves within two westerly jet waveguides** on haze events.

1218: 4 Evolution of **the Rossby waves within two westerly jet waveguides** and principal mechanism analysis

1228: To investigate the relationship between the position change in the jet and **the Rossby waves**, the time series of the 200hPa...

1231: Figure 8a and Figure 8b. That is, the position movement of the jet and **the Rossby waves** was closely related in winter, which...

1234: To reveal the influence of [the Rossby waves within jet waveguide](#) on atmospheric circulation in the middle and lower...

1246: To further examine the mechanism behind the propagation of the [Rossby waves](#), two-dimensional wave activity flux was...

1258: flux was obviously enhanced in the Sea of Japan area, which may be due to the concurrent effect of [two waves](#)...

1261: Based on the analysis above, a diagram of [Rossby waves within the waveguide of the westerly jet](#) affecting haze events in...

1275: [The Rossby waves within two westerly jet waveguides](#) originating...

1286: [The Rossby waves within two westerly jet waveguides](#) may be the main mechanism leading to the occurrence and maintenance...

1294-300: The linear correlation coefficient of the PC1 of the leading EOF mode for the 200 hPa meridional wind anomaly and EUI is approximately 0.92, which further shows that the combined effect of [the waves within two westerly jet waveguides](#) may have an important influence on heavy haze in the NCP. In addition to 2015, the [two waves](#) also existed in 1989, 1994, 1996, 2004, 2006, and 2011. The visibility anomaly in the NCP in these years was negative. However, [the waves within two westerly jet waveguides](#) were also strong in 1982, 1986, 1988, 1991, and 1992, and the visibility anomaly in the NCP was positive (Figure 13). In addition, compared to [the waves](#) in 2015, [the Rossby waves](#) were weak or out of phase in 2000, 2002, 2003, 2007, 2013, and 2014, while the visibility anomaly in the NCP was negative.

127: I would recommend simply stating "This study elucidates the formation...". Let the science speak for itself and determine whether it is of "great significance" or not.

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

127: [This study elucidates the formation](#) and maintenance mechanism of large-scale haze in the NCP in late fall and boreal winter.

146: It's not clear what the anomalies here are referring to. I assume it's geopotential height, so suggest stating "In the negative EU, there is a positive anomaly in geopotential height in Europe..."

Response: We are sorry for this confusion. It is the anomalies of geopotential height. The sentence has been revised as suggested.

147-48: [In the negative EU, there is a positive anomaly in geopotential height in Europe](#) and East Asia and, a negative anomaly in Siberia at 500 hPa.

194-95: It doesn't seem like the Cressman paper is actually cited here. Suggest "using the Cressman interpolation method (Cressman 1959)"

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

195-96: Visibility station data were interpolated on the regular grid of $0.5^\circ \times 0.5^\circ$ using [Cressman interpolation method \(Cressman, 1959\)](#).

1103: I don't find it clear what "function of interpolation" means. Would it be clearer to state, "is the interpolated value at point i,j ". Also, since the same symbol is used for latitude below, perhaps it would be better to choose a different symbol for this.

Response: We agree with the reviewer. The sentence has been revised as suggested. Also, we replace $\Phi_{i,j}$ with $S_{i,j}$ to distinguish it from latitude below.

1104: Where $S_{i,j}$ is the interpolated value at point i, j and S_{obs}^k is station data. N is number of stations.

1105: I think it should be "Plumb" not "Plum"

Response: Yes, it is Plumb. As you suggested, the word of “Plum” been corrected into “Plumb”.

1105-106: To analyze the anomalous propagation of Rossby waves, we calculated horizontal stationary wave activity flux to show the propagation of wave energy using the method of [Plumb](#).

1115: I think some more explanation of what standardization means. Does this mean they are anomalies from the mean and normalized to have standard deviation = 1?

Response: We are sorry for this confusion. The standardization means they are anomalies from the mean and normalized to have standard deviation = 1 and mean value = 0. We have added formula for calculating standardized anomalies as follows:

1116:
$$Norm(x) = \frac{x - mean(x)}{\sigma}$$

1117: ... x is a variable. σ is the number of non-missing value.

1126: I'm confused about what 70% is referring to. 70% of the total of what? 22 days isn't 70% of the total days in November and December, so I'm not sure what this is referring to.

Response: We are sorry for this confusion. 22 days isn't the total days in November and December, but in November and December, respectively. So, 70% is the proportion of haze days every month. The sentence has been revised as following.

1129-130: There were 22 haze days, accounting for more than 70% of the total days, in November and December, [respectively](#) (Figure 2).

1133: suggest being more specific since this is only referring to the lower troposphere e.g., "so that the lower troposphere was relatively stable".

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

1136-137: The value of the static stability was small [so that the lower troposphere was relatively stable](#) (Figure 3a).

1153: Are these number of 37% and 25% referring to this particular event or to air pollution in China more generally. I think this could be stated more clearly to distinguish between the two.

Response: We thank the reviewer for the suggestion. According to Yang et al. (2016), data of daily PM_{2.5} concentration and daily meteorological variables in eastern China (105–122.5°E, 20–45°N) in DJF of 1985–2005 were from GEOS-4 meteorology. So, we think number of 37% and 25% referring to air pollution in China more generally. The sentence has been revised as following.

1154-157: In addition, compared with surface temperature, surface specific humidity, cloud fraction, precipitation rate, and upward flux (UP), the contributions of horizontal wind and the boundary layer height to PM_{2.5} concentration in eastern China [in winter 1985-2005](#) were 37% and 25%, respectively (Yang et al., 2016).

1161: There are many anomalies in this figure from west to east. I suggest being more explicit about which you are referring to e.g., "there was a clear northerly anomaly in western China and southerly anomaly in eastern China in the meridional wind..."

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

1164-165: At 200 hPa (Figure 4a), [there was a clear northerly anomaly in western China and southerly anomaly in eastern China in the meridional wind](#) within the mid-high latitude westerly jet.

1164: Similarly, it's not very clear what "the wave train" is referring to. Suggest pointing to Figure 4b here and describe the wave train of relevance.

Response: We are sorry for this confusion. This wave train only refers to the 200hPa meridional wind anomaly (Figure 4a), not the 500hPa EU (Figure 4b). The sentence has been revised as following.

1165-169: Within the subtropical westerly jet (Figure 4a, green contour line), there existed a southerly/northerly wind wave train package which is similar to study of Ding et al. (2017) (Figure 4a). There were northerly wind anomalies in the Mediterranean region, southerly wind anomalies in the Arabian Peninsula, northerly wind anomalies in western China, and southerly wind anomalies in the Yellow and Bohai Sea of China. Hoskins et al. (1993) and Hsu et al. (1992) found that there are Rossby wave in mid latitude upper troposphere of northern hemisphere.

1172: suggest pointing to the figures for the variables described here.

Response: We thank the reviewer for the suggestion. We have pointed to the figures for the variables.

1175-177: The northerly and southerly wind anomalies over the Asian continent resulted in anomalous circulation southern China, with abnormal ascending movement on the eastern side of the cyclonic anomaly according to the value of divergence at 850 hPa (Figure 4c) and 200 hPa (Figure 4a) and the omega value at 500 hPa (Figure 4b).

Figure 5: Why show the anomalies in figure 4 and the actual values here. It makes it difficult to compare them. Suggest that it might be more useful to show the anomalies in Figure 5 as opposed to the actual values.

Response: We thank the reviewer for the suggestion. It will be more clearly to show the anomalies in Figure 5 if only to compare with figure 4. Our aim is not only to compare with Figure 4, but also to look at the background of large-scale circulation as well as the development of trough and ridge when haze occurs. We also show the anomalies (not shown in manuscripts). The anomalies are similar to those in Figure 4. There is anomalous divergence at 200hPa and anomalous convergence at 850hPa in NCP, which is not conducive to descend motion. However, the actual value shows that there is descend motion (Figure 5 in manuscripts). Besides, southerly anomaly occurred at 850hPa (Figure 4c in manuscripts). The two above indicate that the descend movement and the southerly wind anomaly caused haze, which is consistent with the results of the paper. In order to show the background of large-scale circulation when haze occurs, we use weather map of the actual value. The picture below shows the anomalies.




Figure: The same as Figure 4 but a composite of anomalous weather maps during the periods of November 9-15 and 19-21, December 6-10 and 19-25 in 2015: (a) meridional wind (black contours, CI (contour interval) = 1 m s^{-1} , solid (dashed) lines represent southerly (northerly) wind), divergence (shading, 10^{-5} m s^{-1}) and westerly jet (green line) (zonal mean wind, CI > 30 m s^{-1}) at 200 hPa; (b) geopotential height (black contours, CI = 10 gpm) and vertical velocity (shading, $10^{-2} \text{ Pa s}^{-1}$); negative(positive) values represent ascent (descent) at 500hPa; (c) divergence (shading, 10^{-5} m s^{-1}) and wind vector (vector, m s^{-1}) at 850hPa.

l216: it doesn't make much sense that the positive phase of the EOF means that the jet is strong when above it has been stated that the EOF represents a north-south movement of the subtropical jet. Presumably accompanying this north-south movement is an overall change in the jet strength, so I suggest stating that where the north-south movement is mentioned.

Response: We thank the reviewer for the suggestion. And we think suggestion is very well. The sentence has been revised as suggested.

l222-223: From the first modal time series, it can be seen that the exponent is positive in 2015, [meaning that the location of the subtropical jet is south.](#)

l217: Here's another place where I think there is some confusion between a Rossby wave and a Rossby waveguide. The meridional wind is associated with the Rossby wave which is propagating along the Rossby waveguide, so I suggest "waveguide" -> "wave".

Response: We are sorry for this confusion and we thank the reviewer for the suggestion. The word has been revised as suggested.

l217: The first mode of the meridional wind is a Rossby [wave](#) manifested by the north-south anomaly of the 200 hPa meridional wind, following Li et al. (2017).

l222-224: I think the wording here would be clearer as "To investigate the relationship between the position change in the jet and the Rossby waveguide, the time series of the 200hPa vector wind field and meridional wind are regressed onto the principal component time series of the first EOF of meridional wind and the first EOF of zonal wind in Figs 8 a and b, respectively"

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

l222-224: [To investigate the relationship between the position change in the jet and the Rossby waveguide, the time series of the 200hPa vector wind field and meridional wind are regressed onto the principal component time series of the first EOF of meridional wind and the first EOF of zonal wind in Figs 8 a and b, respectively.](#)

l224: To back up this conclusion that the first EOF of meridional and zonal wind are strongly related to each other, why not just state the correlation between the principal component time series of the two EOFs?

Response: We thank the reviewer for the suggestion. Yes, it will be good to find correlation between the principal component time series of EOF. In our article, we not only want to show the correlation between the principal component time series of the two EOFs, but also want to know whether they will really cause the anomalous circulation at 200hPa corresponding to haze days mentioned above. So, we use the method of regression. we can clearly see the anomalous wind field at 200hPa by using regression analysis.

l248: It's not clear to me that the dispersion of Rossby wave energy necessarily should result in heavy rainfall. I'm not sure to what extent these two things are always connected. Perhaps this could be clarified.

Response: First of all, we're sorry for this confusion. We admit that the statement here is not right. The South Branch of Rossby wave propagated eastward and break in South China, meaning that there is convergence of Rossby wave energy here, which may be beneficial to the circulation for precipitation in South China. For example, Li et al. (2015) and Ding et al. (2017) found the convergence of Rossby wave energy lead to heavy rainfall in Southern China in winter. We have revised the original expression.

l254-256: Almost all of its energy was [converged](#) within South China, [resulting in anomalous circulation, conducive](#) to a large

amount of heavy rainfall in South China.

1281: A value of 0.38 quoted here sounds like it's a correlation coefficient that's being quoted as opposed to a regression coefficient. If it's a regression coefficient then there should be some units.

Response: Yes, it's a regression coefficient with units km. We have added units of regression coefficient in manuscript.

1289: ...approximately 0.38 km from 1980 to 2015 (Figure 13).

1288: I don't think this further shows the combined effect of the two waveguides has important influence on Haze. Doesn't it just show that the meridional wind and the geopotential height are related to each other? If so, this is to be expected, so I suggest deleting this statement.

Response: We are sorry for this confusion. The 200hPa meridional wind anomaly according to EOF is the South Branch wave train propagating along the subtropical westerly jet expressed with PC1, while the 500hPa geopotential height anomaly is the North Branch wave train propagating along the polar front jet expressed with EUI. The correlation coefficient between PC1 and EUI is approximately 0.92, meaning that the two waveguides have important influence on haze.

Figure 3 caption: I think what's plotted in (a) is the difference in temperature between 1000hPa and 850hPa, but that's referred to simply as the lapse rate in the caption. I think it would be clearer to state that this is the vertical temperature difference between those levels. Suggest stating also what this is an Average over e.g., "Anomalies averaged over November and December 2015."

Response: We thank the reviewer for the suggestion. Yes, what's plotted in (a) is the vertical temperature difference between 1000hPa and 850hPa. Besides, in order to reflect the real background field, figure 3a, 3b,3c,3d is the actual values rather than the anomalies. But Figure 3e is anomalies. The caption has been revised as suggested.

Figure 3: Spatial distribution of **average over November and December 2015: the vertical temperature difference between 1000hPa and 850hPa** (unit: °C; Figure 3a); relative humidity at 925 hPa (unit: %; Figure 3b); wind vector (unit: m s^{-1}) at 10 m (Figure 3c) and 850hPa (Figure 3d) in NCP, with shading indicating the wind speed in the appropriate level. **Vertical distribution of anomalies averaged over November and December 2015:** air temperature (AT) (blue dashed line), horizontal wind speed (WS) (red solid line) and relative humidity (RH) (red dash-dotted line) in NCP (Figure 3e).

Figure 4b: I'm not sure if it'll make too much difference, but since the intention is to look at the wave motions, it might be clearer to plot eddy geopotential height as opposed to just geopotential height.

Response: We are sorry for this confusion. And we admit it might be clearer to look at the wave motions by using eddy geopotential height as opposed to just geopotential height. The aim of using geopotential height instead of eddy geopotential height here is not only to look at the wave motions, but also to look at the circulation situation at 500hPa conducive to the formation of haze. For example, according to the results in our paper, the anticyclonic anomaly at 500hPa in Sea of Japan and the cyclonic anomaly in South China are conducive to the formation of haze in the North China Plain. Besides, geopotential height anomalies at 500hPa was used to study EU pattern in many previous researches. We think that geopotential height anomalies may show more information that we need.

Figure 6: Best specify in the caption whether these are anomalies or not.

Response: We are sorry for this confusion. These are the actual values, not anomalies in Figure 6. The detail information has been added in Figure 6.

Figure 6: Composed sections of **the actual values** during the periods of November 9-15 and 19-21, December 6-10 and 19-25 in 2015: latitude-height sections with average longitude in 112°E-120°E of vertical velocity (shading, unit: -5 m s^{-1}) and wind vector (u and ω) (a); longitude-height sections with average latitude in 20°N -30°N (b) and in 30°N-40°N (c) of vertical velocity (shading, unit: -5 m s^{-1}) and wind vector (u and ω).

Figure 8 caption: line 2 suggest "(a) meridional wind and zonal wind" -> "(a) meridional wind and (b) zonal wind"

Response: We thank the reviewer for the suggestion. We have noticed this error. We are sorry that it's not updated in the latest manuscript. The sentence has been revised as suggested.

Figure 8: The 200 hPa horizontal wind anomalies (vector, m s^{-1}) and meridional wind anomalies (contours, m s^{-1}) regressed onto the standardized PC1 of the leading EOF mode for 200 hPa (a) meridional and (b) zonal wind anomalies. The values of the contours and arrows are regression coefficients. The solid red (dashed blue) contours indicate the positive (negative) meridional wind anomaly. The thick black line in (a) and (b) delineates the climatological jet axis (the gradient of the zonal wind with the longitude is 0). Only anomalies statistically significant at the 0.1 level based on Student's t test are shown in (a) and (b).

Typo's/wording:

134: "The causes of haze in China, except for pollutant emissions, ..." "Aside from pollutant emissions, ..."

Response: The sentence has been revised as suggested.

134: Aside from pollutant emissions...

170: suggest deleting "is analyzed only"

Response: We thank the reviewer for the suggestion. We have deleted it as following.

169-73: Previous research on the meteorological parameters influencing haze in the NCP has focused mainly on local meteorological conditions in the middle and lower troposphere, or only the correlation between haze in China and North Atlantic SST or other patterns, while studies about the specific mechanisms behind the influence of large atmospheric circulations in the upper troposphere on haze, especially the combined effect of two westerly jet waveguides, are rare.

172: "are significant" -> "were significant"

Response: The sentence has been revised as suggested.

174: ...on haze were significant in the NCP ...

1121: "mean in November" -> "mean over November"

Response: The sentence has been revised as suggested.

1123-124: Here, $hgt_{a_{Nov2015}}$, $hgt_{Nov2015}$ and $hgt_{m_{Nov\ during\ 1981\ to\ 2010}}$ represent the geopotential height anomaly in November 2015, the geopotential height in November 2015 and geopotential height mean over November 1981 to 2010.

1128: "less 10" -> "of less than 10"

Response: The sentence has been revised as suggested.

1131-132: A regional mean visibility of less than 10 km appeared on 4-16, 18-22 and 27-30, with minimal daily values of 5.16 km, 5.86 km and 4.14 km, respectively.

1145: "was a negative" -> "exhibited a negative"

Response: The sentence has been revised as suggested.

1148-149: The horizontal wind speed exhibited a negative anomaly from the lower troposphere to the upper troposphere (Figure 3e).

1155: "anticyclone" -> "anticyclonic"

Response: The word has been revised as suggested.

1158-159: From the 850 hPa wind vector, it can be found that there was an anticyclonic circulation in southern China.

1168: "anomaly" -> "anomalies" (since this is referring to multiple anomalies.

Response: Yes, we agree with the reviewer. The word has been revised as suggested.

1173-175: The centers of the southerly wind **anomalies** were located over the Arabian Peninsula and East Asia, and the centers of the northerly wind anomaly were located over the European continent and Central Asia.

1170: "circulation southern" -> "circulation in southern"

Response: The sentence has been revised as suggested.

1175-176: The northerly and southerly wind anomalies over the Asian continent resulted in anomalous **circulation in southern China**...

1181: "conductive" -> "conducive", but also it may be clearer to state "stabilizes the atmosphere, creating an environment that is conducive to haze"

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

Due to the anticyclonic anomaly in the troposphere over the Sea of Japan, the descending movement over the NCP **stabilizes the atmosphere, creating an environment that is conducive to haze**.

1191: "at the 200hPa" -> "at 200hPa"

Response: The sentence has been revised as suggested.

1197: ...anomalous meridional wind **at 200 hPa**...

1191: "average locations" -> "averages"

Response: The sentence has been revised as suggested.

1197: ...monthly **averages** (Figure 4a, 4c and Figure 5a, 5c)...

1196: "intruding China" -> "intruding into China"

Response: We admit it's missing "into". The sentence has been revised as suggested.

1202: This circulation situation reduced the amount of cold air **intruding into China** and the winter monsoon also weakened.

1197: "provides" -> "provided"

Response: The word has been revised as suggested.

1203: The Southern Branch trough **provided** rising...

1213: I think this needs some rewording. Perhaps "To explore the influences on Rossby wave propagation along the subtropical westerly jet and associated influences on haze..."

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

1219: **To explore the influences on Rossby wave propagation along the subtropical westerly jet and associated influences on haze**...

1228: "regressed by" -> "regressed onto"

Response: The sentence has been revised as suggested.

1235: ...were **regressed onto** the time series...

1230-231: "the adjacent Sea of Japan areas" -> "the Sea of Japan"?

Response: Yes, it is the Sea of Japan. The sentence has been revised as suggested.

1237-238: The two positive centers of the regression coefficients of the geopotential height anomaly are located over the

Mediterranean Sea and [the Sea of Japan](#).

l252: "is very weak" -> "was very weak" and "is still" -> "was still"

Response: We thank the reviewer for the suggestion. The sentence has been revised as suggested.

l258-260: Thus, the energy from the southern branch of the Rossby wave [was very weak](#) over the Sea of Japan, but the anticyclone anomaly over the Sea of Japan [was still](#) very strong.

l275: "of Rossby" -> "of the Rossby"

Response: We are sorry that it's missing "the". The sentence has been revised as suggested.

l282: ...of the southern path [of the Rossby](#) wave...

Reference

Ding, F. and Li, C.: Subtropical westerly jet waveguide and winter persistent heavy rainfall in south China, J. Geophys. Res. Atmos., 122, 7385-7400, doi:10.1002/2017JD026530, 2017.

Li, C. and Sun, J. L.: Role of the subtropical westerly jet waveguide in a southern China heavy rainstorm in December 2013, Adv. Atmos. Sci., 32, 601-612, doi:10.1007/s00376-014-4099-y, 2015.

The combined effect of two westerly jet waveguides on heavy haze in the North China Plain in November and December 2015

Xiadong An¹, Lifang Sheng^{1, 2}, Qian Liu³, Chun Li^{1, 2}, Yang Gao⁴, and Jianping Li⁵

¹Department of Marine Meteorology, College of Oceanic and Atmospheric Sciences, Ocean University of China, Qingdao, 266100, China

²Ocean-Atmosphere Interaction and Climate Laboratory, Key Laboratory of Physical Oceanography, Ocean University of China, Qingdao, 266100, China

³School of Atmospheric Sciences, and Guangdong Province Key Laboratory for Climate Change and Natural Disaster Studies, Sun Yat-sen University, Guangzhou, 510275, China

⁴College of environmental Science and Engineering, Ocean University of China, Qingdao, 266100, China

⁵Key Laboratory of Physical Oceanography–Institute for Advanced Ocean Studies, Ocean University of China and Qingdao National Laboratory for Marine Science and Technology, Qingdao, 266003, China

Correspondence to: Lifang Sheng (shenglf@ouc.edu.cn)

Abstract. Severe haze occurred in the North China Plain (NCP) from November to December 2015, with a wide spatial range and long duration. In this paper, the combined effect of [the anomalous Rossby waves within two westerly jet waveguides](#) on haze in the NCP was investigated based on visibility observational data and NCEP/NCAR reanalysis data. The results showed that [the anomalous Rossby waves that propagated along two waveguides within the westerly jet](#) originating from the Mediterranean were responsible for haze formation in the NCP. The Rossby wave propagated eastward along the subtropical westerly jet and the polar front jet, causing an anomalous anticyclone over the Sea of Japan and anticyclonic wind speed shear at 850 hPa over the NCP, which enhanced the anomalous descending air motion in the middle and lower troposphere and subsequently resulted in a stable atmosphere. Furthermore, the Rossby wave weakened the East Asia trough and Ural ridge, and strengthened the anomalous southerly wind at 850 hPa over the coastal areas of east China, decelerating the East Asia winter monsoon. The above meteorological conditions modulated haze accumulation in November and December 2015. Meanwhile, continuous rainfall related to ascending motion due to Rossby wave propagation along the subtropical westerly jet occurred in a large area of southern China. The latent heat released by rainfall acted as a heat source, inducing convection over South China. This further strengthened the ascending motion over South China so that the descending motion over the NCP was maintained, favoring the maintenance of severe haze. [This study elucidates the formation](#) and maintenance mechanism of large-scale haze in the NCP in late fall and boreal winter.

1 Introduction

30 Haze is the phenomenon of reduced visibility caused by the increase in aerosols or the hygroscopic growth of aerosols at a high relative humidity (Ma et al., 2014). In December 2015, the air quality was poor in the North China Plain (NCP), especially in the Beijing-Tianjin-Hebei region (Chang et al., 2016; Zhang et al., 2016; Zhang et al., 2019). Based on the emissions level in December 2015 and the Weather Research and Forecasting-Community Multiscale Air Quality (WRF-CMAQ) regional model, Zhang et al. (2019) found higher monthly mean PM_{2.5} concentrations. [Aside from pollutant emissions](#), the causes of

35 haze in China, the weather conditions and climate change also play a significant role in modulating haze formation, distribution, maintenance and change (Ding et al., 2009; Tai et al., 2012; Zhang et al., 2013). Yang et al. (2016) found that changes in meteorological parameters contributed 17 (± 14)% to the increasing trend in PM_{2.5} concentration from 1985 to 2005. Dang and Liao (2019) found that severe winter haze days showed large interannual variations in frequency and intensity, which were mainly driven by changes in meteorology.

40 Winter haze- fog in northern China in recent years are related to weak winter circulation (Wu et al., 2014). Some studies have noted that the circulation in mid- and high latitude regions plays an important role in wind variations in northern China (Wang et al., 2016; Li et al., 2017; He et al., 2019). Zhao et al. (2013) and Zhang et al. (2013) showed that the weak East Asian winter monsoon (EAWM) is particularly unfavorable to the outward transport of aerosols in northern China, which results in continuous strong haze in northern China. In addition, the stable weather conditions associated with the Siberian high pressure

45 are also conducive to the accumulation and maintenance of haze (Zhang et al., 2016; Liu et al., 2017). Li et al. (2019) reported that the Eurasian teleconnection (EU) at 500 hPa is the most important pattern affecting haze- fog in northern China, and its contribution rate is 45%. In the negative EU, there is a positive anomaly [in geopotential height](#) in Europe and East Asia and, a negative anomaly in Siberia at 500 hPa. As a result, the cold air mass from the arctic is too weak to reach northern China (Ren et al., 2010; Zhai et al., 2016), which is not conducive to the dissipation of haze.

50 According to many studies, the North African-Asian (NAA) jet has an important impact on the Asian climate in winter (Syed et al., 2006; Feldstein and Dayan, 2008; Wen et al., 2009; Ni et al., 2010; Li et al., 2013). On the decadal timescale, the strength of the jet is highly consistent with the frequency of cold air temperature extremes in China (Chen et al., 2013). The divergence in the upper troposphere induced by the subtropical westerly jet waveguide is important for spatially large and persistent winter rainfall in South China (Li and Sun, 2015; Ding and Li, 2017). Based on previous theoretical work, Xu et al. (2019) noted that

55 the British-Baikal Corridor teleconnection on the polar front jet waveguide in Eurasia has an important impact on the climate of East Asia in summer. Zhang et al. (2019) found that the winter concurrent meridional shift in the East Asian jet streams is related to the east-ward propagating Rossby wave. Huang et al. (2019) further found that the concurrent change in the location of the subtropical and polar front jet in East Asia in winter has an important impact on the climate of East Asia. In addition, previous studies showed that as the teleconnection wave source location, the anomaly in atmosphere circulation related to sea

60 surface temperature (SST) in the North Atlantic can significantly influence winter weather in the Northern Hemisphere (Wallace and Gutzler, 1981; Peng et al., 1995; Czaja and Frankignoul, 2002; Li and Betas, 2007), which can influence haze in China (Xiao et al., 2014; Gao and Chen, 2017). Chen et al. (2013) noted that the cold air activity was possibly associated with the development of the Siberian high pressure and a wave train from the North Atlantic. Yang et al. (2019) stated that the East Asian wind and temperature anomalies during boreal winter are determined by the combination of the two wave trains
65 propagating along the subtropical jet and the polar front jet. The SST anomalies in the North Atlantic can induce a downstream Rossby wave train, resulting in anomalous circulations over the NCP, which are favorable for haze over central and south China (Feng et al., 2019; Wang et al., 2019).

In summary, [the Rossby waves within the East Asian upper westerly jet waveguides](#) has important influences on the East Asian climate. Previous research on the meteorological parameters influencing haze in the NCP has focused mainly on local
70 meteorological conditions in the middle and lower troposphere, or only the correlation between haze in China and North Atlantic SST or other patterns, while studies about the specific mechanisms behind the influence of large atmospheric circulations in the upper troposphere on haze, especially the combined effect of [the Rossby waves within two westerly jet waveguides](#), are rare. Given the above content, the objective of the present study is to determine whether the effects of [the Rossby waves within two westerly jet waveguides](#) on haze were significant in the NCP in November and December 2015 and,
75 if so, to identify the principal mechanism behind the effects of [the Rossby waves within two westerly jet waveguides](#) on haze. The structure of this paper is as follows: data and methodology used in this paper are described briefly in Section 2. Section 3 presents the haze event and describes major meteorological parameters and atmosphere circulation patterns during haze events. Section 4 demonstrates the influencing mechanisms of [the Rossby waves within two westerly jet waveguides](#) on haze events. Further discussions and conclusions of our findings are provided in Section 5.

80 **2 Data and Methodology**

2.1 Data

Visibility station data from the stations in the region (15-55°N, 105-135°E) in November and December 2015 were from the China Meteorological Administration (CMA). These data were recorded four times daily (02:00, 08:00, 14:00, and 20:00, local time (LCT)). The data were averaged monthly in this study.

85 The monthly mean and daily mean reanalysis data are from the National Centers for Environmental Prediction-National Center for Atmospheric Research (NCEP/NCAR) NCEP-DOE AMIP-II Reanalysis (R-2) dataset (Kanamitsu et al., 2002). The dataset covers a 39-yr period from 1979 to 2017, with a latitude-longitude spatial resolution of $2.5^\circ \times 2.5^\circ$, including the geopotential height, wind vector, relative humidity, air temperature, and omega data at each standard level from 10 hPa to 1000 hPa. The daily precipitation data from November to December 2015 are provided by the NASA/Goddard Space Flight Center (Huffman

2.2 Methodology

Haze was defined as a day when the daily mean visibility and relative humidity were less than 10 km and 80%, respectively, and no rain, snow, sand and dust storms occurred in accordance with the standards set by the CMA (China Meteorological Administration, 2010).

95 Visibility station data were interpolated on the regular grid of $0.5^\circ \times 0.5^\circ$ using Cressman interpolation method (Cressman, 1959). Cressman objective analysis adopts the method of successive corrections, which has been widely used in various climate diagnosis analysis and numerical simulation studies. The most important Cressman objective analysis is the determination of the weight function $W_{i,j}$ (Feng et al., 2004) (1-2):

$$W_{i,j} = \frac{R^2 - r_{i,j}^2}{R^2 + r_{i,j}^2} \quad r_{i,j}^2 < R, \quad (1)$$

$$100 \quad W_{i,j} = 0 \quad r_{i,j}^2 > R, \quad (2)$$

Where R is the influence radius, $r_{i,j}^2$ is the distance between two interpolation points, i and j are each interpolation point. R are 5, 4, 3 in this paper. Unit of R is degree.

$$S_{i,j} = \frac{\sum_{k=1}^N W_{i,j}^k S_{obs}^k}{\sum_{k=1}^N W_{i,j}^k} \quad (3)$$

Where $S_{i,j}$ is the interpolated value at point i, j and S_{obs}^k is station data. N is number of stations.

105 To analyze the anomalous propagation of Rossby waves, we calculated horizontal stationary wave activity flux to show the propagation of wave energy using the method of Plumb (1985) (4):

$$F = p_0 \cos \phi \left(\frac{1}{2a^2 \cos^2 \phi} \left[\left(\frac{\partial \psi'}{\partial \lambda} \right)^2 - \psi' \frac{\partial^2 \psi'}{\partial \lambda^2} \right] - \frac{1}{2a^2 \cos^2 \phi} \left(\frac{\partial \psi'}{\partial \lambda} \frac{\partial \psi'}{\partial \phi} \right) - \psi' \frac{\partial^2 \psi'}{\partial \lambda \partial \phi} \right), \quad (4)$$

Here, F is the horizontal stationary wave activity flux, p_0 is pressure / (1000 hPa), ψ' is the perturbation stream function, a is Earth's radius, ϕ is the latitude, and λ is the longitude.

110 The East Asian winter monsoon composite index defined by He and Zhou (2012) was used to characterize the East Asia winter monsoon. The formula is as follows (5-8):

$$I_1 = Norm[\bar{p}_s(40^\circ \sim 60^\circ N, 80^\circ \sim 125^\circ E)] , \quad (5)$$

$$I_4^* = -1 \times I_4 = -1 \times Norm[\bar{h}_{500}(25^\circ \sim 45^\circ N, 110^\circ \sim 145^\circ E)] , \quad (6)$$

$$I_5 = Norm[\bar{u}_{300}(25^\circ \sim 40^\circ N, 80^\circ \sim 180^\circ E) - \bar{u}_{300}(45^\circ \sim 60^\circ N, 60^\circ \sim 160^\circ E)] , \quad (7)$$

$$115 \quad EAWMII = \frac{I_1 + I_4^* + I_5}{3} , \quad (8)$$

$$Norm(x) = \frac{x - mean(x)}{\sigma} , \quad (9)$$

Here, Norm represents standardization, x is a variable. σ is the number of non-missing value, and \bar{p}_s , \bar{h}_{500} , and \bar{u}_{300} are

the mean sea level pressure, 500 hPa mean geopotential height and 300 hPa mean zonal wind over the region defined above, respectively.

120 Please note that all the anomalies calculated in this paper were based on the climatological mean during the thirty-year period of 1981–2010. For example, the calculation procedure can be written as (10)

$$hgt_a_{Nov2015} = hgt_{Nov2015} - hgt_m_{Nov \text{ during } 1981 \text{ to } 2010} \quad (10)$$

Here, $hgt_a_{Nov2015}$, $hgt_{Nov2015}$ and $hgt_m_{Nov \text{ during } 1981 \text{ to } 2010}$ represent the geopotential height anomaly in November 2015, the geopotential height in November 2015 and geopotential height mean over November 1981 to 2010.

125 3 Persistent haze events and associated weather patterns

Heavy haze occurred over the NCP in November and December 2015. The locations where the average monthly visibility was less than 10 km covered northern China and eastern China (Figure 1). The haze events were characterized by wide range, strong intensity and long duration, which was consistent with the analysis of Chang et al. (2016), Zhang et al. (2016) and Zhang et al. (2019). There were 22 haze days, accounting for more than 70% of the total days, in November and December 2015, respectively (Figure 2). In November, the mean monthly visibility over the NCP was 8.33 km with a minimal value of 4.14 km. A regional mean visibility of less than 10 km appeared on 4-16, 18-22 and 27-30, with minimal daily values of 5.16 km, 5.86 km and 4.14 km, respectively. Similarly, the mean monthly visibility in December over the NCP was 8.63 km with a minimal daily value of 2.62 km. The dates when the regional mean visibility was less than 10 km were 6-14, 19-24 and 28-31 December, with minimal daily values of 3.87 km, 2.62 km and 6.00 km, respectively.

130 Static stability is expressed by the vertical difference in the temperature between 1000 hPa and 850 hPa (Liu et al., 2017), which is on average 4.5°C-7.5°C in the NCP. The value of the static stability was small so that the lower troposphere was relatively stable (Figure 3a). In addition, the vertical profiles of the temperature anomalies between 1000 hPa and 700 hPa show warm anomalies in the middle and lower troposphere and cold anomalies near the surface (Figure 3e), which further indicates that the atmosphere is relatively stable. When haze occurred, the mean monthly relative humidity (RH) in the NCP was less than 70% (Figure 3b), and the regional daily relative humidity was also relatively low (not shown). According to the haze identification conditions issued by the China Meteorological Administration (2010), haze rather than haze- fog can be considered the main pollution event in November and December 2015. An RH of approximately 55-70% may promote the liquid phase reaction of aerosol particles. RH has an important influence on the hygroscopic growth of aerosol particles (Ma et al., 2014), which can promote the photochemical reaction of aerosol particles. Wu et al. (2019) found that the air over the NCP during the haze episode was humid, with an average simulated RH of approximately 71%.

145 The ground was mainly controlled by an anticyclone in November and December 2015 according to the ground wind field at 10 m. The NCP was in the center of the anticyclone, and the wind speed was very weak (Figure 3c). However, in the periphery

of the NCP, the wind speed was relatively strong (Figure 3c). The horizontal wind speed exhibited a negative anomaly from the lower troposphere to the upper troposphere (Figure 3e). The 850 hPa wind field showed a northeast-southwest wind speed shear (Figure 3d), which corresponded to negative vorticity anomalies. The atmosphere was mainly composed of descending movement, so the stratification was relatively stable. According to Liu et al. (2017), the frequency of haze occurrence was 24.5% under low wind speed conditions at the ground and low stability over 1303 noncold wave days, accounting for the highest proportion of the four conditions (including low wind speed and low stability (20.4%), high wind speed and low stability (14.4%), and high wind speed and high stability (14.6%)). In addition, compared with surface temperature, surface specific humidity, cloud fraction, precipitation rate, and upward flux (UP), the contributions of horizontal wind and the boundary layer height to $PM_{2.5}$ concentration in eastern China in winter 1985-2005 were 37% and 25%, respectively (Yang et al., 2016). Therefore, wind speed and atmospheric stability play vital roles in haze formation during November and December in the NCP. From the 850 hPa wind vector, it can be found that there was an anticyclonic circulation in southern China. The southerly wind on the northwest side of the anticyclone circulation prevented the southward dispersion of pollutants over the NCP. At the same time, the anticyclone circulation weakened the EAWM in the NCP. According to the winter monsoon composite index defined by He et al. (2012), the EAWM index in November and December 2015 was -0.29 (not shown). The weak EAWM was beneficial to the occurrence and maintenance of haze.

Figure 4 shows the mean monthly circulation anomalies and the average jet positions in November and December 2015. At 200 hPa (Figure 4a), there was a clear northerly anomaly in western China and southerly anomaly in eastern China in the meridional wind within the mid-high latitude westerly jet. Within the subtropical westerly jet (Figure 4a, green contour line), there existed a southerly/northerly wind wave train package which is similar to study of Ding et al. (2017) (Figure 4a). There were northerly wind anomalies in the Mediterranean region, southerly wind anomalies in the Arabian Peninsula, northerly wind anomalies in western China, and southerly wind anomalies in the Yellow and Bohai Sea of China. Hoskins et al. (1993) and Hsu et al. (1992) found that there are wave train in mid latitude upper troposphere of northern hemisphere. This was also similar to the teleconnection of the Northern Hemisphere Silk Road in summer proposed by Lu et al. (2002). There was a strong northerly anomaly over the European continent. The strong northerly wind formed a convergence of cold air in the Middle East and the Mediterranean Sea, which intruded into the entrance of the westerly jet stream and propagated eastward in the form of a wave train along the westerly jet stream. The centers of the southerly wind anomalies were located over the Arabian Peninsula and East Asia, and the centers of the northerly wind anomaly were located over the European continent and Central Asia. The northerly and southerly wind anomalies over the Asian continent resulted in anomalous circulation in southern China, with abnormal ascending movement on the eastern side of the cyclonic anomaly according to the value of divergence at 850 hPa (Figure 4c) and 200 hPa (Figure 4a) and the omega value at 500 hPa (Figure 4b). The NCP experienced an anticyclonic anomaly with abnormal descent. Atmospheric motion ascended over southern China and descended over the

NCP to form a north-south circulation system from 20°N to 50°N at 112°E-120°E (Figure 4a, Figure 6). In addition, the NCP
180 was located in the middle of the cyclonic and anticyclonic anomalies, with strong southerly winds at 850 hPa (Figure 4b,
Figure 4c), weakening the EAWM and favoring haze accumulation. There was a negative phase of the EU at 500 hPa (Figure
4b): the negative geopotential height anomaly over Siberia weakened the Ural ridge, and the positive geopotential height
anomaly over the Sea of Japan weakened the East Asian trough. According to Li et al. (2019), the direct effect of the EU
negative phase is that the arctic cold air does not easily flow to East Asia, and the EAWM weakens. Thus, a negative EU may
185 have an important impact on haze. Due to the anticyclonic anomaly in the troposphere over the Sea of Japan, the descending
movement over the NCP [stabilizes the atmosphere, creating an environment that is conducive to haze](#). In southern China, there
was mainly ascending motion, with an anomalous southwesterly wind at 850 hPa (Figure 4c). The convergence of
southwesterly airflow was conducive to the occurrence of large-scale continuous precipitation over the southern China, as
reported by Li and Sun (2015) and Ding and Li (2017). As shown in Figure 2, strong precipitation occurred in November and
190 December 2015 in southern China (20°N-30°N, 100°E-120°E), i.e., during November 6-13, 14-17, and 18-21 and December
2-6, 7-10 and 21-23. The north-south circulation systems were consistent, which was implicative of a barotropic stable
circulation structure triggering long-lasting of the haze events.

To more clearly depict the circulation in detail during the persistent haze events, the circulation field was composited by
selecting haze events lasting at least three days with a mean daily visibility of less than 8 km (Figure 5). The periods are
195 November 9-15, 19-21 and December 6-10, 19-25. The results during the severe haze events are in generally consistent with
those of the monthly average. The locations of the centers of convergence and divergence at 850 hPa and 200 hPa and the
anomalous meridional wind [at 200 hPa](#) are similar to the monthly [averages](#) (Figure 4a, 4c and Figure 5a, 5c), but the intensity
of the anomalous meridional wind over the Mediterranean and East Asia seems to be stronger, and the strength of the
divergence is enhanced as well. There is a relatively strong trough over the North Atlantic, a weak trough over the Ural
200 Mountains and a trough over the North Pacific at 500 hPa. In addition, there was a weak Southern Branch trough over the
southern Qinghai-Tibet Plateau and southern China. The subtropical high over the western Pacific was stronger and located in
the west (Figure 5b). This circulation situation reduced the amount of cold air [intruding into China](#) and the winter monsoon
also weakened. The Southern Branch trough [provided](#) rising conditions for precipitation over southern China. The southerly
airflow on the west side of the subtropical high (Figure 5c; near 115°E) carried a large amount of water vapor originating from
205 the South China Sea into southern China. In contrast, convergence of the southerly wind occurred over the NCP, leading to
haze formation.

Through the analysis above, we found that the main meteorological factors modulating the large-scale persistent haze in the
NCP are wind vector and atmospheric stability. The anomalous southerly wind over the eastern coast of China weakens the
winter monsoon, and the anticyclonic shear of the wind at 850 hPa in the NCP yields an increase in negative vorticity and the

210 subsequent enhancement of atmospheric descent. In addition, the anomalous anticyclone over the Sea of Japan strengthens the descending air motion and increases atmospheric stability. From the geopotential height anomaly at 200 hPa (not shown), 500 hPa (Figure 4b), 850 hPa (Figure 4c) and the tropospheric vertical motion (Figure 6), it can be seen that the upper and lower levels exhibit barotropic structures. The north-south anomaly of the meridional wind from west to east appears in the upper tropospheric subtropical westerly jet, which is similar to the Silk Road teleconnection proposed by Lu et al. (2002). The anticyclone anomaly over the NCP strengthens the descending motion, and the ascending motion related to upper divergence and lower convergence over southern China is strong. The north-south circulation system leads to the maintenance of haze in the NCP, whereas large-scale precipitation occurred over southern China.

4 Evolution of the Rossby waves within two westerly jet waveguides and principal mechanism analysis

To explore the influence on Rossby wave propagation along the subtropical westerly jet and associated influences on haze, the first leading EOF mode of the 200 hPa zonal and meridional winds from November 1979 to December 2017 was calculated. The first mode of zonal wind represents the north-south movement of the subtropical jet (Figure 7a), following Hong et al. (2016). This mode's variance is 25.3%. From the first modal time series, it can be seen that the exponent is positive in 2015, meaning that the location of the subtropical jet is south. The first mode of the meridional wind is a Rossby wave manifested by the north-south anomaly of the 200 hPa meridional wind, following Li et al. (2017). The variance is 23.4%. The first modal time series shows that the exponent has a negative index in 2015, with strong intensity (Figure 7c). The northerly wind anomaly appears in the Mediterranean region and the Indian Peninsula region, and the southerly wind anomaly appears in the Arabian Peninsula region, North China and the Yellow Sea.

To investigate the relationship between the position change in the jet and the Rossby waves, the time series of the 200hPa vector wind field and meridional wind are regressed onto the principal component time series of the first EOF of meridional wind and the first EOF of zonal wind in Figs 8 a and b, respectively. The results of the regression are very similar as shown in Figure 8a and Figure 8b. That is, the position movement of the jet and the Rossby waves was closely related in winter, which is basically consistent with the results of Hong et al. (2016) on the relationship between the Silk Road teleconnection (summer) and the change in jet stream location.

To reveal the influence of the Rossby waves within jet waveguide on atmospheric circulation in the middle and lower troposphere, 500 hPa geopotential height anomalies and 850 hPa wind field anomalies were regressed onto the time series of the first EOF mode of meridional wind (Figure 9). The EU negative phase appears in the regressed geopotential height field. The two positive centers of the regression coefficients of the geopotential height anomaly are located over the Mediterranean Sea and the Sea of Japan. The positive geopotential height anomaly near the Sea of Japan weakens the East Asian trough, weakening the EAWM. In addition, the anticyclone anomaly also causes the southerly wind anomaly at 850 hPa over the

240 eastern coast of China, further weakening the winter monsoon. The center of the negative geopotential height anomaly is located in Siberia, weakening the Ural ridge and weakening the Lower Siberian High and favoring haze formation. The regression coefficient of the wind field anomaly shows that there is a southerly anomaly over the NCP, and a southwesterly anomaly in South China. The southerly anomaly over the NCP tends to weaken the winter wind, while the southwesterly anomaly over South China carries water vapor from the Bay of Bengal and South China Sea to South China, providing
245 favorable moisture conditions for precipitation in South China.

To further examine the mechanism behind the propagation of the Rossby waves, two-dimensional wave activity flux was calculated and generated according to the formula for the wave action flux defined by Plum (1985). Figure 10 shows that the European continent is the source of the Rossby wave. Rossby waves propagate into two pathways: a subtropical westerly jet and polar front jet. The Rossby wave in the polar front jet was stronger than that in the subtropical westerly jet. Rossby waves
250 in the polar front jet, named the EU, resulted in energy dispersion over the Siberian area, leading to a negative geopotential height anomaly in this location, weakening the Ural ridge and preventing cold air advection towards China. The remaining energy from the easterly propagation caused the positive geopotential height anomaly over the Sea of Japan, weakening the East Asian trough and inducing the southerly wind anomaly over the eastern coast of China. The southern branch of the Rossby wave propagated within the subtropical westerly jet that passed over the Arabian Peninsula, India and other places. Almost all
255 of its energy was converged within South China, resulting in anomalous circulation, conducive to a large amount of heavy rainfall in South China. The latent heat released by precipitation as a heat source strengthened the abnormally ascending motion in South China, which further strengthened the anticyclone anomaly in the NCP. The wave action flux was obviously enhanced in the Sea of Japan area, which may be due to the concurrent effect of two waves superposing in this area. Thus, the energy from the southern branch of the Rossby wave was very weak over the Sea of Japan, but the anticyclone anomaly over the Sea
260 of Japan was still very strong.

Based on the analysis above, a diagram of the Rossby waves within the waveguide of the westerly jet affecting haze events in the NCP is drawn (Figure 11, Figure 12). The Rossby wave originated from the Mediterranean region (Li and Sun, 2015) and propagated eastward along the westerly jet, resulting in an anomalous anticyclone over the Sea of Japan and strengthening descending air motion over the NCP. In addition, there was a southerly wind anomaly at 850 hPa over the eastern coastal areas
265 of China, which weakened the winter monsoon (Figure 12). Both of these features are conducive to haze accumulation. In addition, the Rossby waves also caused a strong abnormal upward movement over South China, which combined with the rich water vapor carried by the southwesterly airflow resulted in large-scale continuous rainfall (Figure 11, Figure 12). The latent heat released by precipitation as a heat source strengthened the abnormal ascending motion over South China, which further helped to maintain the sinking motion in the NCP. A local circulation in the southern and northern parts of China was formed
270 (Figure 12): the air rises in the south and subsides in the north, facilitating the intensification of the descending air motion in

the NCP and the persistence of haze.

5 Discussions and conclusions

A wide range of haze pollution occurred in the NCP in November and December 2015. The meteorological factors, climatic system and mechanisms affecting haze were elucidated in this study. Haze was mainly modulated by the weakening of the EAWM and the strengthening of descending air in the NCP. [The Rossby waves within two westerly jet waveguides](#) originating from the Mediterranean region was the main mode fostering the atmospheric circulation patterns tightly associated with haze formation in the NCP.

The Rossby wave propagated easterly primarily along two pathways. The northern path seemed to yield positive geopotential height anomalies in the troposphere over the Sea of Japan and the NCP, strengthening the descending air movement over the NCP. In addition, it weakened the East Asia trough and Ural ridge, resulting in the southerly wind anomaly at 850 hPa over the eastern coast of China, weakening the winter monsoon, and leading to subsequent haze accumulation. Due to the influence of the southern path [of the Rossby wave](#), there was abnormal upward air movement over South China, which combined with a large amount of water vapor carried by the 850 hPa anticyclone circulation over South China, provided favorable conditions for continuous rainfall in South China. As a heat source, the latent heat of condensation released by precipitation, further strengthened abnormal upward movement in South China, favorable for the subsidence and haze formation in the NCP.

[The Rossby waves within two westerly jet waveguides](#) may be the main mechanism leading to the occurrence and maintenance of large-scale haze in the NCP in November and December 2015. The linear regression coefficient of PC1 (bar) of the leading EOF mode for the 200 hPa meridional wind anomaly onto the mean visibility anomaly in November and December is approximately [0.38 km](#) from 1980 to 2015 (Figure 13). The location of the EU mentioned above is slightly different from that of the EU defined by predecessors. To describe the EU more accurately in this study, a new EU index (EUI) was defined according to the definition of the EUI of Wallace and Gutzler (1981).

$$EUI = -1/4 \times \text{hgt500_ano} (40^\circ\text{N}, 0^\circ) + 1/2 \times \text{hgt500_ano} (50^\circ \text{N}, 50^\circ \text{E}) - 1/4 \times \text{hgt500_ano} (40^\circ \text{N}, 140^\circ \text{E})$$

Here, hgt500_ano is the mean geopotential height anomaly at 500 hPa in November and December.

The linear correlation coefficient of the PC1 of the leading EOF mode for the 200 hPa meridional wind anomaly and EUI is approximately 0.92, which further shows that the combined effect of [the Rossby waves within two westerly jet waveguides](#) may have an important influence on heavy haze in the NCP. In addition to 2015, [the two waves](#) also existed in 1989, 1994, 1996, 2004, 2006, and 2011. The visibility anomaly in the NCP in these years was negative. However, [the Rossby waves within two westerly jet waveguides](#) were also strong in 1982, 1986, 1988, 1991, and 1992, and the visibility anomaly in the NCP was positive (Figure 13). In addition, compared to [the waves](#) in 2015, [the Rossby waves](#) were weak or out of phase in 2000, 2002, 2003, 2007, 2013, and 2014, while the visibility anomaly in the NCP was negative. This means that under similar pollutant

emission conditions, the large-scale circulation conditions that cause air pollution may take on diversified modes. This paper mainly focused the haze in November and December 2015; therefore, the robustness of the mechanism may be further verified if other haze events can be evaluated in future studies. We can understand the complexity behind the climatic causes of severe air pollution from some existing studies. For example, the combined effects of the negative North Atlantic Oscillation (NAO) and El Niño worsened air conditions over central and south China in the winters of 2000 and 2003 (Feng et al., 2019). The autumnal increased number of haze days in 2013 and 2014 was closely associated with simultaneous sea surface warming in the western North Pacific sector and the North Atlantic subtropical sector (Wang et al., 2019).

Acknowledgements

This study was supported by the National Natural Science Foundation of China (Grant No. 41675146) and Fundamental Research Funds for the Central Universities (Grant No. 201941006). All authors would like to express their great thanks to the China Meteorological Administration (<http://data.cma.cn/>), NCEP/NCAR (<https://www.esrl.noaa.gov/psd/data/gridded/>), and NASA (<https://pmm.nasa.gov/data-access/downloads/gpm>) for supplying the research data. They are also very grateful for anonymous reviewers for their kind efforts that improved the quality of this manuscript significantly.

Data availability

The visibility observational data is available at the China Meteorological Administration (<http://data.cma.cn/>). The reanalysis dataset is available at NCEP/NCAR (<https://www.esrl.noaa.gov/psd/data/gridded/>). The precipitation data is available at NASA (<https://pmm.nasa.gov/data-access/downloads/gpm>).

Author contribution

XA, LS, CL and JL designed the study. LS and YG acquisitioned the funding. LS obtained observation data. XA and QL dealt with the visibility observational data. XA downloaded, analyzed the reanalysis data and prepared all figures. XA led the writing with the help of LS, QL and YG. All the authors discussed the results and commented on the paper.

Competing interests

The authors declare no conflicts of interest.

Financial support

This study was supported by the National Natural Science Foundation of China (Grant No. 41675146) and Fundamental Research Funds for the Central Universities (Grant No. 201941006).


References

- 330 Chang, L. Y., Xu, J. M., Tie, X. X., and Wu, J. B.: Impact of the 2015 El Niño event on winter air quality in China, *Sci. Rep.*, 6, 34275, doi:10.1038/srep34275, 2016.
- Chen, H. S., Liu, L., and Zhu, Y. J.: Possible linkage between winter extreme low temperature events over China and synoptic-scale transient wave activity, *Sci. China Ser.*, 56, 1266-1280, doi:10.1007/s11430-012-4442-z, 2013.
- China Meteorological Administration.: QXT 113-2010 Observation and forecasting levels of haze (in Chinese), 1-1, China Meteorol. Press, Beijing, China, 2013.
- 335 Cressman, G. P.: An operational objective analysis system, *Mon. Wea. Rev.*, 87, 367–374, 1959.
- Czaja, A. and Frankignoul, C.: Observed Impact of Atlantic SST Anomalies on the North Atlantic Oscillation, *J. Clim.*, 15, 606-623, doi:10.1175/1520-0442(2002)0152.0.CO;2, 2002.
- Ding, F. and Li, C.: Subtropical westerly jet waveguide and winter persistent heavy rainfall in south China, *J. Geophys. Res. Atmos.*, 122, 7385-7400, doi:10.1002/2017JD026530, 2017.
- 340 Ding, Y. H., Li, Q. P., Liu, Y. J., Zhang, L., Song, Y. F., and Zhang, J.: Atmospheric Aerosols, Air Pollution and Climate Change, *Meteorological Monthly*, 35, 3-15 (in Chinese), 2009.
- Feldstein, S. B. and Dayan, U.: Circumglobal teleconnections and wave packets associated with Israeli winter precipitation, *Quart. J. Roy. Meteor. Soc.*, 134, 455- 467, doi:10.1002/qj.225, 2008.
- Feng, J., Li, J. P., Liao, H., and Zhu, J. L.: Simulated coordinated impacts of the previous autumn North Atlantic Oscillation (NAO) and winter El Niño on winter aerosol concentrations over eastern China, *Atmos. Chem. Phys.*, 19, 10787–10800, doi:10.5194/acp-19-10787-2019, 2019.
- 345 Feng, J. M., Zhao, T. B., and Zhang, Y. J.: Intercomparison of Spatial Interpolation Based on Observed Precipitation Data, *Climatic Environ. Res.*, 02, 261-277 (in Chinese), 2014.
- Gao, Y. and Chen, D.: A dark October in Beijing 2016, *Atmospheric and Oceanic Science Letters*, 10:3, 206-213, doi:10.1080/16742834.2017.1293473, 2017.
- 350 Hoskins, B. J. and Ambrizzi, T.: Rossby wave propagation on a realistic longitudinally varying flow, *J. Atmos. Sci.* 50, 1661–1671, doi :10.1175/1520-0469(1993)050<1661:RWPOAR>2.0.CO;2, 1993.
- Hsu, H. H. and Lin, S. H.: Global teleconnections in the 250-mb streamfunction field during the in the geopotential height field during the Northern Hemisphere winter, *Mon. Wea. Rev.*, 120, 1169–1190, doi:10.1175/1520-0493(1992)120<1169:GTITMS>2.0.CO;2, 1992.
- 355


- He, C., Liu, R., Wang, X. M., Liu, S. C., Zhou, T. J., and Liao, W. H.: How does El Niño-Southern Oscillation modulate the interannual variability of winter haze days over eastern China? *Sci. Total Environ.*, 651, 1892-1902, doi:10.1016/j.scitotenv.2018.10.100, 2019.
- 360 He, S. P. and Wang, H. J.: An integrated East Asian winter monsoon index and its inter-annual variability, *Chinese Journal of Atmospheric Sciences*, 36(3), 523-538 (in Chinese), 2012.
- Hong, X. and Lu, R.: The meridional displacement of the summer Asian jet, Silk Road Pattern, and tropical SST anomalies, *J. Clim.*, 29, 3753-3766, doi:10.1175/JCLI-D-15-0541.1, in press, 2016.
- Huang, D., Dai, A., Yang, B., Yan, P., Zhu, J., and Zhang, Y.: Contributions of Different Combinations of the IPO and AMO to Recent Changes in Winter East Asian Jets, *J. Clim.*, 32, 1607-626, doi:10.1175/JCLI-D-18-0218.1, 2019.
- 365 Huffman, G., Bolvin, D., Braithwaite, D., Hsu, K., Joyce, R., and Xie, P.: Integrated Multi-satellite Retrievals for GPM (IMERG), version 4.4. NASA's Precipitation Processing Center, accessed 09 June, 2019, 2014.
- Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S. -K., Hnilo, J.J., Fiorino, M., and Potter, G. L.: NCEP-DOE AMIP-II Reanalysis (R-2), *Bulletin of the American Meteorological Society*, Nov 2002, 1631-1643, 2002.
- Li, C. and Sun, J. L.: Role of the subtropical westerly jet waveguide in a southern China heavy rainstorm in December 2013, 370 *Adv. Atmos. Sci.*, 32, 601-612, doi:10.1007/s00376-014-4099-y, 2015.
- Li, C. F., Chen, W., Hong, X. W., and Lu, R. Y.: Why was the strengthening of rainfall in summer over the Yangtze River valley in 2016 less pronounced than that in 1998 under similar preceding El Nino events? Role of midlatitude circulation in August, *Adv. Atmos. Sci.*, 34, 1290-1300, doi:10.1007/s00376-017-7003-8, 2017.
- Li, J., Sun, C., and Jin, F. F.: NAO implicated as a predictor of Northern Hemisphere mean temperature multi-decadal 375 variability, *Geophys. Res. Lett.*, 40, 5497- 5502, doi:10.1002/2013GL057877, 2013.
- Li, S., Han, Z., and Chen. H.: A comparison of the effects of inter-annual Arctic sea ice loss and ENSO on winter haze days: observational analyses and AGCM simulations, *J. Meteor. Res.*, 31, 820-833, doi:10.1007/s13351-017-7017-2, 2017.
- Li, S. L. and Bates, G.: Influence of the Atlantic Multidecadal Oscillation (AMO) on the winter climate of East China, *Adv. Atmos. Sci.*, 24, 126–135, doi:10.1007/s00376-007-0126-6, 2007.
- 380 Li, Y., Sheng, L. F., Li, C., and Wang, Y. H.: Impact of the Eurasian Teleconnection on the Interannual Variability of Haze-Fog in Northern China in January, *Atmosphere*, 10, 113, doi:10.3390/atmos10030113, 2019.
- Liu, Q., Sheng, L., Cao, Z., Diao, Y., Wang, W., and Zhou, Y.: Dual effects of the winter monsoon on haze-fog variations in eastern China, *J. Geophys. Res. Atmos.*, 122, 5857-5869, doi:10.1002/2016JD026296, 2017.
- Lu, R. Y., Jai, Ho. Oh, and Baek, Jo. Kim.: A teleconnection pattern in upperlevel meridional wind over the North African and 385 Eurasian continent in summer, *Tellus A: Dynamic Meteorology and Oceanography*, 54, 44-55, doi:10.3402/tellusa.v54i1.12122, 2002.

- Ma, N., Zhao, C. S., Chen, J., Xu, W. Y., Yan, P., and Zhou, X. J.: A novel method for distinguishing fog and haze based on PM_{2.5}, visibility, and relative humidity, *Science China: Earth Sciences*, 57, 2156-2164, doi:10.1007/s11430-014-4885-5, 2014.
- 390 Ni, D.-H., Sun, Z.-B., Li, Z.-X., Zeng, G., and Deng, W.-T.: Relation of Middle East jet stream and China climate anomaly in winter, (in Chinese) *Scientia Meteor. Sinica*, 30, 301-307, 2010.
- Peng, S., Mysak, L. A., Derome, J., Ritchie, H., and Dugas, B.: The Differences between Early and Midwinter Atmospheric Responses to Sea Surface Temperature Anomalies in the Northwest Atlantic, *J. Clim.*, 8, 137-157, 1995.
- Plumb, R.A.: On the three-dimensional propagation of stationary waves, *J. Atmos. Sci.*, 42, 217-229, doi:10.1175/1520-395 0469(1985)042<0217:OTTDPO>2.0.CO:2, 1985.
- Ren, X. J., Yang, X. Q., Zhou, T. J., and Fang, J. B.: Diagnostic comparison of the East Asian subtropical jet and polar-front jet: Large-scale characteristics and transient eddy activities, *Acta Meteor. Sinica*, 68, 1-11, doi:10.1007/s13351-011-0002-2, 2010.
- Syed, F. S., Giorgi, F., Pal, J. S., and King, M. P.: Effect of remote forcings on the winter precipitation of central southwest 400 Asia Part 1: Observations, *Theor. Appl. Climatol.*, 86, 147-160, doi:10.1007/s00704-005-0217-1, 2006.
- Tai, A. P. K., Mickley, L. J., Jacob, D. J., Leibensperger, E. M., Zhang, L., Fisher, J. A., and Pye, H. O. T.: Meteorological modes of variability for fine particulate matter (PM_{2.5}) air quality in the United States: implications for PM_{2.5} sensitivity to climate change, *Atmos. Chem. Phys.*, 12, 3131-3145, doi:10.5194/acp-12-3131-2012, 2012.
- Wallace, J. M. and Gutzler, D. S.: Teleconnections in the geopotential height field during the Northern Hemisphere winter, 405 *Mon. Wea. Rev.*, 109, 784–812, doi:10.1175/1520-0493(1981)109<0784:TITGHF>2.0.CO;2, 1981.
- Wang, H. J. and Chen, H. P.: Understanding the recent trend of haze pollution in eastern China: roles of climate change, *Atmos. Chem. Phys.*, 16, 4205-4211, doi:10.5194/acp-16-4205-2016, 2016.
- Wang, J., Zhu, Z., Qi, L., Zhao, Q., He, J., and Wang, J. X. L.: Two pathways of how remote SST anomalies drive the interannual variability of autumnal haze days in the Beijing–Tianjin–Hebei region, China, *Atmos. Chem. Phys.*, 19, 1521–410 1535, doi:10.5194/acp-19-1521-2019, 2019.
- Wang, L., Chen, W., and Huang, R.: Interdecadal modulation of PDO on the impact of ENSO on the East Asian winter monsoon, *Geophys. Res. Lett.*, 35(20), L20702, doi:10.1029/2008GL035287, 2008b.
- Wen, M., Yang, S., Kumar, A., and Zhang, P. Q.: An analysis of the Large-Scale climate anomalies associated with the snowstorms affecting China in January 2008, *Mon. Wea. Rev.*, 137, 1111-1131, doi:10.1175/2008MWR2638.1, 2009.
- 415 Wu, R. G. and Wang, B.: Interannual variability of summer monsoon onset over the western North Pacific and the underlying processes, *J. Clim.*, 13, 2483-2501, doi:10.1175/15200442(2000)013<2483:IVOSMO>2.0.CO;2, 2001.
- Xiao, D., Li, Y., Fan, S. J., Zhang, R. H., Sun, J. R., and Wang, Y.: Plausible influence of Atlantic Ocean SST anomalies on

- winter haze in China, *Theor. Appl. Climatol.*, 122, 249-257, doi:10.1007/s00704-014-1297-6, 2014.
- Xu, P., Wang, L., and Chen, W.: The British Baikal Corridor: A Teleconnection Pattern along the Summer
420 time Polar Front Jet over Eurasia, *J. Clim.*, 32, 877-896, doi:10.1175/JCLI-D-18-0343, 2019.
- Yang, J., Wu, R., and Song, L.: Individual and combined impacts of two Eurasian wave trains on intra-seasonal East Asian
winter monsoon variability, *J. Geophys. Res. Atmos.*, 124, 4530-4548, doi:10.1029/2018JD029953, 2019.
- Yang, Y., Liao, H., and Lou, S.: Increase in winter haze over eastern China in recent decades: Roles of variations in
meteorological parameters and anthropogenic emissions, *J. Geophys. Res. Atmos.*, 121, 13050-13065, doi:
425 10.1002/2016JD025136, 2016.
- Zhai, P. M., Yu, R., Guo, Y. J., Li, Q. X., Ren, X. J., Wang, Y. Q., Xu, W. H., Liu, Y. J., and Ding, Y. H.: The strong El Niño of
2015/16 and its dominant impacts on global and China's climate, *J. Meteor. Res.*, 30, 283-297, doi: 10.1007/s13351-016-6101-
3, 2016.
- Zhang, G., Gao, Y., Cai, W. J., , Ruby Leung, L., Wang, S. X., Zhang, B., Wang, M. H., Shan, H. Y., Yao, X. H., and Gao, H.
430 W.: Seesaw haze pollution in North China modulated by the sub-seasonal variability of atmospheric circulation, *Atmos. Chem.
Phys.*, 19, 565-576, doi:10.5194/acp-19-565-2019, 2019.
- Zhang, L. Y., Lu, X. B., Yang, L. L., Ding, F., Zhu, Z. F., and Rui, D. M.: Study on the Effects of the Strong El Nino Event on
Air Quality of Eastern China in Winter, *The Administration and Technique of Environmental Monitoring*, 28, 23-27, 2016.
- Zhang, X. Y., Sun, J. Y., Wang, Y. Q., Li, W. J., Zhang, Q., Wang, W. G., Quan, J. N., Cao, G. L., Wang, J. Z., Yang, Y. Q., and
435 Zhang, Y. M.: Factors contributing to haze and fog in China, *Chin. Sci. Bull (Chin Ver)*, 58, 1178-1187 (in Chinese), 2013.
- Zhang, Y., Yan, P., Liao, Z., Huang, D., and Zhang, Y.: The Winter Concurrent Meridional Shift of the East Asian Jet Streams
and the Associated Thermal Conditions, *J. Clim.*, 32, 2075-2088, doi:10.1175/JCLI-D-18-0085.1, 2019.
- Zhang, Z., Zhang, X., Gong, D., Kim, S.-J., Mao, R., and Zhao, X.: Possible influence of atmospheric circulations on winter
haze pollution in the Beijing–Tianjin–Hebei region, northern China, *Atmos. Chem. Phys.*, 16, 561–571, doi:10.5194/acp-16-
440 561-2016, 2016.
- Zhao, X. J., Zhao, P. S., Xu, J., Meng, W., Pu, W. W., Dong, F., He, D., and Shi, Q. F.: Analysis of a winter regional haze event
and its formation mechanism in the North China Plain, *Atmos. Chem. Phys.*, 13, 5685-5696, doi:10.5194/acp-13-5685-2013,
2013.



445 **Figure 1: Spatial distribution of monthly mean visibility (unit: km, shading) in (a) November and (b) December 2015. The black box indicates the NCP (30°N-40.5°N, 112°E-121.5°E). Shading indicates the value of visibility.**



450 **Figure 2: Variations in the regional mean daily visibility (bars; unit: km) is obtained by calculating the average value of all stations data in the NCP (30°N-40.5°N, 112°E-121.5°E) and the regional mean daily precipitation (lines; unit: mm day⁻¹) in southern China (20°N-30°N, 100°E-120°E) (a) in November and (b) in December 2015.**




Figure 3: Spatial distribution of average over November and December 2015: the vertical temperature difference between 1000hPa and 850hPa (unit: °C; Figure 3a); relative humidity at 925 hPa (unit: %; Figure 3b); wind vector (unit: m s^{-1}) at 10 m (Figure 3c) and 850hPa (Figure 3d) in NCP, with shading indicating the wind speed in the appropriate level. Vertical distribution of anomalies averaged over November and December 2015: air temperature (AT) (blue dashed line), horizontal wind speed (WS) (red solid line) and relative humidity (RH) (red dash-dotted line) in NCP (Figure 3e).

455




Figure 4: Anomalous weather and westerly jet maps in November and December 2015: (a) meridional wind (black contours, CI (contour interval) = 1 m s⁻¹, solid (dashed) lines represent southerly (northerly) wind anomaly), divergence (shading, 10⁻⁵ m s⁻¹) and westerly jet (green line) (zonal mean wind, CI > 30 m s⁻¹) at 200 hPa; (b) geopotential height (black contours, CI = 10 gpm (geopotential meters), solid (dashed) lines represent positive (negative) geopotential height anomaly) and vertical velocity (shading, 10⁻² Pa s⁻¹); negative(positive) values represent ascent (descent) at 500 hPa; (c) geopotential height (red contours, CI = 10 gpm), divergence (shading, 10⁻⁵ m s⁻¹) and wind vector (vector, m s⁻¹) at 850 hPa.




Figure 5: The same as Figure 4 but a composite of non-anomalous weather maps during the periods of November 9-15 and 19-21, December 6-10 and 19-25 in 2015: (a) meridional wind (black contours, CI (contour interval) = 1 m s^{-1} , solid (dashed) lines represent southerly (northerly) wind), divergence (shading, 10^{-5} m s^{-1}) and westerly jet (green line) (zonal mean wind, CI $> 30 \text{ m s}^{-1}$) at 200 hPa; (b) geopotential height (black contours, CI = 50 gpm) and vertical velocity (shading, $10^{-2} \text{ Pa s}^{-1}$); negative(positive) values represent ascent (descent) at 500 hPa; (c) divergence (shading, 10^{-5} m s^{-1}) and wind vector (vector, m s^{-1}) at 850 hPa.

470




Figure 6: Composed sections of the actual values during the periods of November 9-15 and 19-21, December 6-10 and 19-25 in 2015: latitude-height sections with average longitude in 112°E-120°E of vertical velocity (shading, unit: -5 m s^{-1}) and wind vector (u and ω) (a); longitude-height sections with average latitude in 20°N -30°N (b) and in 30°N-40°N (c) of vertical velocity (shading, unit: -5 m s^{-1}) and wind vector (u and ω).





Figure 7: The (a) spatial distribution and (b) its principal component (PC) of the leading EOF mode for 200 hPa zonal wind anomaly within the domain 0-75°N, 0-160°E in November and December 1979-2017, where the solid (dashed) lines represent westerly (easterly) winds; (c) and (d) same as (a) and (b) but for meridional wind within the domain 0-75°N, 0-160°E, where solid (dashed) lines represents southerly (northerly) winds. The percentage values in the upper-right corners of (a) and (c) show the percentage variance explained by this mode.



485 **Figure 8: The 200 hPa horizontal wind anomalies (vector, m s^{-1}) and meridional wind anomalies (contours, m s^{-1})**
regressed onto the standardized PC1 of the leading EOF mode for 200 hPa (a) meridional and (b) zonal wind anomalies.
The values of the contours and arrows are regression coefficients. The solid red (dashed blue) contours indicate the
positive (negative) meridional wind anomaly. The thick black line in (a) and (b) delineates the climatological jet axis
(the gradient of the zonal wind with the longitude is 0). Only anomalies statistically significant at the 0.1 level based on
490 **Student's t test are shown in (a) and (b).**




Figure 9: Regression of 500 hPa geopotential height anomalies and 850 hPa wind vector onto the PC1 of the leading EOF mode for the 200 hPa meridional wind anomaly. The values of shading and arrows represent regression coefficients. Regions of dotted areas indicate anomalies exceeding the 0.05 confidence level. Only anomalies statistically significant at the 0.05 level based on Student's t test are given for the wind vector.

495






Figure 10: Anomalous geopotential height (shading, 10 gpm) at 200 hPa in November and December 2015 and its stationary wave activity flux (vector, $\text{m}^{-2} \text{s}^{-2}$).



500 Figure 11: A schematic diagram of the negative EU in the 200 hPa geopotential height field and meridional wind anomaly at 200 hPa in November and December 2015. H (L) denotes positive (negative) geopotential height anomalies; the plus and minus represent positive (southerly) and negative (northerly) meridional wind anomalies, respectively; and the shaded belt of arrow represents the subtropical westerly jet.



505 Figure 12: Schematic illustration showing circulation system affecting the haze events in November and December 2015. The red (blue) circular of arrow represents convergence (divergence); the red (blue) translucent arrow represents ascending (descending) air; the brown translucent arrow represents the southerly wind anomaly; the purple solid arrow represents the water vapor transported by southwesterly airflow from the Bay of Bengal and South China Sea; the thick (thin) black arrow represents northward (southward) movement in the atmosphere; H denotes positive

510 geopotential height anomaly; the white weather symbol represents precipitation.




Figure 13: The mean visibility anomaly (dashed line, purple) in November and December in the NCP, the EUI (solid line, green), and PC1 (bars, red and blue) of the leading EOF mode for the 200 hPa meridional wind anomaly from 1980 to 2015. The linear regression coefficient of the mean visibility onto PC1 is approximately 0.38 km, which can pass the significance test at the 95% level.

515