

The isotopic composition of atmospheric nitrous oxide observed at the high-altitude research station Jungfraujoch, Switzerland

Longfei Yu^{1*}, Eliza Harris^{1†}, Stephan Henne¹, Sarah Eggleston¹, Martin Steinbacher¹, Lukas
5 Emmenegger¹, Christoph Zellweger¹ and Joachim Mohn¹

¹Laboratory for Air Pollution & Environmental Technology, Empa, Swiss Federal Laboratories
for Materials Science and Technology, Ueberlandstr. 129, CH-8600 Duebendorf, Switzerland.

[†]Current address: Institute of Ecology, University of Innsbruck, Sternwartestrasse 15, A-6020
Innsbruck, Austria

10 * Correspondence: L. Yu (longfei.yu@empa.ch)

Abstract

Atmospheric nitrous oxide (N₂O) levels have been continuously growing since preindustrial times. Mitigation requires information about sources and sinks on the regional and global scales. Isotopic composition of N₂O in the atmosphere could contribute valuable constraints. However, isotopic records of N₂O in the unpolluted atmosphere remain too scarce for large-scale N₂O models. Here, we report the results of discrete air samples collected weekly to bi-weekly over a five-year period at the high-altitude research station Jungfraujoch, located in central Switzerland. High-precision N₂O isotopic measurements were made using a recently developed preconcentration-laser spectroscopy technique. The measurements of discrete samples were accompanied by *in situ* continuous measurements of N₂O mixing ratios. Our results indicate a pronounced seasonal pattern with minimum N₂O mixing ratios in late summer, associated with a maximum in $\delta^{15}\text{N}^{\text{bulk}}$ and a minimum in intramolecular ¹⁵N site preference ($\delta^{15}\text{N}^{\text{SP}}$). This pattern is most likely due to stratosphere-troposphere exchange (STE), which delivers N₂O-depleted but ¹⁵N-enriched air from the stratosphere into the troposphere. Variability in $\delta^{15}\text{N}^{\text{SP}}$ induced by changes in STE may be masked by biogeochemical N₂O production processes in late summer, which are possibly dominated by a low- $\delta^{15}\text{N}^{\text{SP}}$ pathway of N₂O production (denitrification), providing an explanation for the observed seasonality of $\delta^{15}\text{N}^{\text{SP}}$. Footprint analyses and atmospheric transport simulations of N₂O for Jungfraujoch suggest that regional emissions from the planetary boundary layer contribute to seasonal variations of atmospheric N₂O isotopic composition at Jungfraujoch, albeit more clearly for $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ than for $\delta^{15}\text{N}^{\text{bulk}}$. With the time-series of five years, we obtained a significant interannual trend for $\delta^{15}\text{N}^{\text{bulk}}$ after deseasonalization ($-0.052 \pm 0.012\text{‰ a}^{-1}$), indicating that the atmospheric N₂O increase is due to isotopically depleted N₂O sources. We estimated the average isotopic signature of anthropogenic N₂O sources with a two-box model to be $-8.6 \pm 0.6\text{‰}$

35 for $\delta^{15}\text{N}^{\text{bulk}}$, $34.8\pm 3\text{‰}$ for $\delta^{18}\text{O}$ and $10.7\pm 4\text{‰}$ for $\delta^{15}\text{N}^{\text{SP}}$. Our study demonstrates that seasonal
variation of N_2O isotopic composition in the background atmosphere is important when
determining interannual trends. More frequent, high-precision and inter-laboratory compatible
measurements of atmospheric N_2O isotopocules, especially for $\delta^{15}\text{N}^{\text{SP}}$, are needed to better
constrain anthropogenic N_2O sources, and thus the contribution of biogeochemical processes to
40 N_2O growth on the global scale.

1 Introduction

Nitrous oxide (N₂O) is a potent greenhouse gas (Fowler et al., 2015) and a strong stratospheric ozone-depleting substance (Ravishankara et al., 2009). For several decades, near-surface atmospheric N₂O mixing ratios have been continuously measured at a series of remote sites, within
45 the networks of the Global Atmosphere Watch Programme (JMA and WMO, 2018), the Advanced Global Atmospheric Gases Experiment (AGAGE) (Prinn et al., 2018), and the National Oceanic and Atmospheric Administration (NOAA) Earth System Research Laboratory (ESRL) Global Monitoring Division (GMD) (Nevison et al., 2011). These measurements have shown a significant increase in atmospheric N₂O mixing ratio, at a current growth rate of about 0.93 nmol mol⁻¹ a⁻¹
50 (WMO, 2018). On the global scale, given excessive nitrogen (N) fertilizer application, agriculture is known to be the largest and most important anthropogenic source of N₂O (Reay et al., 2012; Tian et al., 2019). However, long-term observations of N₂O in the unpolluted atmosphere have shown seasonal and interannual variabilities as well as interhemispheric differences in N₂O mixing ratios (Nevison et al., 2011; Thompson et al., 2014a, 2014b), which cannot yet be resolved by
55 atmospheric transport models and existing emission inventories. Moreover, regional contributions of N₂O emissions and the strengths of individual N₂O production pathways remain difficult to quantify.

Isotopic signatures of atmospheric N₂O can provide important constraints on N₂O sources (Denk et al., 2017) and trends (Kim and Craig, 1993). The ratios of ¹⁵N/¹⁴N and ¹⁸O/¹⁶O in N₂O are often
60 reported in δ notation as $\delta(^{15}\text{N}/^{14}\text{N})$ and $\delta(^{18}\text{O}/^{16}\text{O})$, abbreviated as $\delta^{15}\text{N}^{\text{bulk}}$ (average for ¹⁴N¹⁵N¹⁶O and ¹⁵N¹⁴N¹⁶O) and $\delta^{18}\text{O}$, respectively. A large fraction of N₂O emitted to the atmosphere originates from soil bacterial processes, which usually emit N₂O that is more enriched in light (¹⁴N, ¹⁶O) isotopes than the tropospheric background (Pérez et al., 2001; Snider et al., 2015a; Toyoda et

al., 2017). By contrast, N₂O produced in the oceans (Bourbonnais et al., 2017; Fujii et al., 2013) and emitted from fossil fuel combustion (Ogawa and Yoshida, 2005; Toyoda et al., 2008) has higher $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$ values which are comparable to the tropospheric background. A recent study has summarized isotopic signatures of anthropogenic N₂O sources divided into the EDGAR (Emissions Database for Global Atmospheric Research) emission categories (Janssens-Maenhout et al., 2019), showing differences in isotopic signatures between agricultural ($\delta^{15}\text{N}^{\text{bulk}} = -17.8$ to -1.0‰ and $\delta^{18}\text{O} = 23.9$ to 29‰) and industrial sources ($\delta^{15}\text{N}^{\text{bulk}} = -28.7$ to 5.5‰ and $\delta^{18}\text{O} = 28.6$ to 40.3‰) (Harris et al., 2017). These empirical ranges, together with isotopic mixing models, provide a valuable approach to interpret variability in atmospheric N₂O mixing ratios.

A number of studies have analyzed temporal trends in N₂O isotopic composition in the modern atmosphere (Kaiser et al., 2003; Park et al., 2012; Röckmann and Levin, 2005; Toyoda et al., 2013) and in the past from firn and ice cores (Bernard et al., 2006; Ishijima et al., 2007; Prokopiou et al., 2018; Röckmann et al., 2003; Sowers et al., 2002). These isotopic measurements have shown a decrease in both $\delta^{15}\text{N}^{\text{bulk}}$ - and $\delta^{18}\text{O}$ -N₂O associated with an increasing trend in atmospheric N₂O mixing ratios since preindustrial times, indicating that the recent increase of atmospheric N₂O may be due to agricultural emissions (¹⁵N and ¹⁸O depleted). The reported trend since the 1960s seems rather steady ($-0.034 \pm 0.005 \text{‰ a}^{-1}$ for $\delta^{15}\text{N}^{\text{bulk}}$ and $-0.016 \text{‰} \pm 0.006 \text{‰ a}^{-1}$ for $\delta^{18}\text{O}$) (Bernard et al., 2006; Ishijima et al., 2007; Park et al., 2012; Prokopiou et al., 2017; Röckmann et al., 2003; Röckmann and Levin, 2005). However, a more recent (1999-2010) study reported a smaller decreasing trend in $\delta^{15}\text{N}^{\text{bulk}}$ and only an insignificant trend in $\delta^{18}\text{O}$ for the Northern Hemisphere (Toyoda et al., 2013). Several hypotheses were proposed to explain the differences in the observed trends: 1) the interhemispheric difference in N₂O emission sources results in inconsistent isotopic signatures among different studies (Thompson et al., 2014b); 2) uncertainties in isotopic

measurements and variable sampling schemes (air type, sampling frequency and time) mask the small secular trend of N₂O isotopic composition in the background atmosphere (Toyoda et al., 2013); and/or 3) N₂O source isotopic signatures have changed in recent years, possibly due to shifts in N fertilizer type and climatic forcing (Tian et al., 2019). Hence, further investigation into the global N₂O source inventory and its evolution over time requires more frequent, precise measurements of N₂O isotopocules in the unpolluted atmosphere, particularly in the Northern Hemisphere.

Recently, site-specific composition of N₂O isotopomers (site preference: $\delta^{15}\text{N}^{\text{SP}}$), which denotes the difference of ¹⁵N between the central (¹⁴N¹⁵N¹⁶O, α position) and terminal (¹⁵N¹⁴N¹⁶O, β position) N atoms, has been applied to constrain sources contributing to atmospheric N₂O (Toyoda et al., 2013; Yoshida and Toyoda, 2000). $\delta^{15}\text{N}^{\text{SP}}$ of N₂O is particularly effective for distinguishing between the major N₂O production processes, i.e. nitrification and denitrification, generally referred to as aerobic and anaerobic N₂O production, with high and low $\delta^{15}\text{N}^{\text{SP}}$, respectively (Sutka et al., 2006). However, despite the advantages of $\delta^{15}\text{N}^{\text{SP}}$ measurements, existing long-term studies have not yet been able to reach a definitive understanding of the $\delta^{15}\text{N}^{\text{SP}}$ -N₂O trend, showing both positive (Bernard et al., 2006; Park et al., 2012; Röckmann and Levin, 2005) and negative tendencies (Röckmann et al., 2003) over the last four decades. This is probably due to an insufficient analytical precision and poor inter-laboratory agreement, in particular as the aforementioned studies are all based on isotope ratio mass spectrometry (IRMS). To retrieve site-specific isotopic information by IRMS, the N₂O⁺ molecular ions and the NO⁺ fragment ions are analyzed and raw data have to be corrected for rearrangements of central and terminal N and ¹⁷O content (Toyoda et al., 2001). Inappropriate correction algorithms and the limited availability of

reference materials (Ostrom et al., 2018) further enlarge the analytical uncertainty (Mohn et al.,
110 2014).

Seasonal variability in atmospheric N₂O isotopic composition, which could affect the longer-term trends, is still rarely reported in the literature (Park et al., 2012; Toyoda et al., 2013). Moreover, studies of seasonality of N₂O isotopic composition are limited to the recent past since the air samples derived from firn and ice cores suffer from coarse temporal resolution (< 2 samples per
115 year). Park et al. (2012) studied seasonality of atmospheric N₂O isotopic composition by analyzing a set of archived air samples collected from Cape Grim (Australia) using a sophisticated mathematical modeling approach. They found consistent seasonal patterns in $\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{18}\text{O}$ and $\delta^{15}\text{N}^{\text{SP}}$ of atmospheric N₂O, showing highest ¹⁵N/¹⁸O enrichment in June and lowest in December. This pattern was negatively correlated with the seasonality of the N₂O mixing ratios (lowest in
120 April-May and highest in December), which is in agreement with a previous study by Nevison et al. (2011). The negative correlation between isotopic composition and mixing ratios has been explained by stratosphere-troposphere exchange (STE), which transports N₂O-depleted but isotopically enriched stratospheric air (prevailing reduction process) into the lower atmosphere (Yung and Miller, 1997). However, in a more recent study from Hateruma Island (Japan), Toyoda
125 et al. (2013) reported insignificant seasonal patterns in atmospheric N₂O isotopocules (smaller variability than measurement precision), despite their finding of a somewhat similar seasonal pattern in N₂O mixing ratio (minimum in July). Although there are interhemispheric differences in N₂O sources and distinct sampling frequencies in the two studies discussed above (2-3 times per year versus monthly), it is noteworthy that both studies observed significantly larger variability
130 in $\delta^{15}\text{N}^{\text{SP}}$ than in $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$. Whether the fluctuations in $\delta^{15}\text{N}^{\text{SP}}$ are mainly caused by the

limited repeatability of the chosen analytical techniques or interplay of processes or mechanisms regulating atmospheric N₂O remains to be tested (Park et al., 2012).

With inherent selectiveness, in particular for site-specific isotopic composition, laser spectroscopy provides a new analytical approach for direct, precise measurements of all four N₂O isotopocules (Harris et al., 2014; Mohn et al., 2012). The recent development of quantum cascade laser absorption spectroscopy (QCLAS) coupled with an automated preconcentration unit has been applied to measure N₂O isotopocules in ambient air, with comparable precision for $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$ and superior precision for $\delta^{15}\text{N}^{\text{SP}}$ relative to IRMS systems (Harris et al., 2017; Mohn et al., 2014). Here, we present results from the application of a preconcentration unit coupled to QCLAS to measure atmospheric N₂O isotopocules in background air collected at the high altitude research station Jungfraujoch, Switzerland. Between April 2014 and December 2018, we collected weekly to bi-weekly air samples for N₂O isotopic analyses, in parallel with online measurement of N₂O mixing ratios. To our knowledge, this work reports the first time-series of background atmospheric N₂O isotopic composition using laser spectroscopy. With this unique dataset, we aim to 1) constrain seasonal patterns of three N₂O isotopic signatures at the Jungfraujoch observatory; 2) determine interannual trends in N₂O isotopocules, especially $\delta^{15}\text{N}^{\text{SP}}$; and 3) interpret the observed patterns in N₂O mixing ratios using temporal trends in N₂O isotopic composition and reported isotopic signatures of anthropogenic sources.

2 Materials and Method

150 2.1 Site description

The high altitude research station Jungfraujoch (3580 m above sea level), located on the northern ridge of the Swiss Alps, is a well-established site for studying unpolluted atmosphere over Central Europe (e.g. Buchmann et al., 2016). Although the station is located in the free troposphere most of the time, it is occasionally affected by air recently lifted from the planetary boundary layer (Herrmann et al., 2015; Zellweger et al., 2003). Henne et al. (2010) investigated the representativeness of 35 European monitoring stations and categorized Jungfraujoch as “mostly remote”. The Jungfraujoch station is part of several national and international networks, like the meteorological SwissMetNet network operated by MeteoSwiss, the Swiss National Air Pollution Monitoring Network (NABEL), the Global Atmospheric Watch Programme (GAW) of the World Meteorological Organization (WMO) and the Integrated Carbon Observation Systems (ICOS) Research Infrastructure. This results in an extended set of long-term and continuously available parameters such as meteorological variables (Appenzeller et al., 2008), greenhouse gases (Schibig et al., 2015; Sepúlveda et al., 2014; Yuan et al., 2018), CO₂ isotopic composition (Sturm et al., 2013; Tuzson et al., 2011), ozone-depleting substances and their replacement products (Reimann et al., 2008), atmospheric pollutants (Logan et al., 2012; Pandey Deolal et al., 2012; Zellweger et al., 2009) and aerosol parameters (Bukowiecki et al., 2016).

2.2 *In situ* measurements and discrete air sampling (flasks)

In situ observations of N₂O mixing ratios commenced at Jungfraujoch in December 2004. Initially, measurements were made with gas chromatograph (GC) (Agilent 6890N, USA) followed by electron capture detection (ECD). The time resolution of these measurements was 24 to 30 minutes.

In late 2014, we implemented a cavity-enhanced off-axis integrated cavity out-put spectroscopy analyzer (OA-ICOS, Los Gatos Research Inc., Mountain View, CA, USA), which measures the atmospheric N₂O mixing ratio continuously. Measurements of N₂O mixing ratios at Jungfraujoch were calibrated with three standard gases (319, 327 and 342 nmol mol⁻¹) and accompanied with measurement of a working standard (331 nmol mol⁻¹) every 160 minutes to account for instrumental drift. In addition, short- (two times every 40 hours) and long-term (every 40 hours) target measurements were included to monitor the data quality. Due to the superior measurement precision compared to the GC-ECD method (Lebague et al., 2016), the OA-ICOS record has become the primary time-series since January 2015. The GC-ECD observations continued until summer 2016 for comparison and quality control.

Additional parameters, recorded within the NABEL network and the ICOS infrastructure, were included in the analysis below. These data were carbon monoxide (CO) (measured by cavity ring-down spectroscopy; Model G2401, Picarro Inc., USA), the sum of oxidized nitrogen species (NO_y) (measured by chemiluminescence detection after conversion of NO_y to NO on a heated gold catalyst; CLD 89p, Eco Physics, Switzerland) and O₃ (measured by UV absorption; TEI 49i, Thermo Scientific, USA). Details on measurement methods and calibration strategies can be found in Zellweger et al. (2009) for CO, Pandey Deolal et al. (2012) for NO_y and Logan et al. (2012) for O₃.

In conjunction with the online measurements, we deployed an automated sampling system (Fig. S1) to collect pressurized air samples in aluminum cylinders from the same air inlet at the Sphinx observatory of the Junfraujoch station, for subsequent N₂O mixing ratio and isotopic analyses. The sample collection was conducted weekly from April 2014 to February 2016. After a sampling gap of five months due to a technical failure, we reinitiated a bi-weekly sampling, which continued

from August 2016 to December 2018. The sampling system, automated by a customized LabVIEW
195 program (National Instruments Corp., USA), consisted of a Nafion drier (PD-100T-48MSS, Perma
Pure LLC, USA), a membrane gas compressor (KNF Neuberger, USA; Type N286 series), a 16-
port selector valve (EMT2CSD16MWEPPH, VICI AG, Switzerland), and a rack to accommodate
nine 2-L aluminum flasks (Luxfer, Messer Schweiz AG, Switzerland). During sample filling, pre-
evacuated flasks were first purged with ambient air five times (1 hour), and then filled to 12000
200 hPa within 40 min, resulting in approximately 24 L (298 K and 1000 hPa) of air per flask for
isotopic analysis. Air sample filling generally took place between 2:00 and 3:00 pm local time at
each sampling day. Sample flasks were sent back to the laboratory at Empa for analyses every few
months. For this study, 142 air samples were collected in flasks and analyzed for N₂O isotopocules.

2.3 Analyses of discrete air samples

205 Discrete air samples were regularly analyzed in batches but not in chronological order to prevent
the imprint of analytical drifts on temporal trends of the samples. N₂O mole fractions were
analyzed by QCLAS (CW-QC-TILDAS-76-CS, Aerodyne Research Inc., USA) against NOAA
standards on the WMO-X2006A calibration scale (Hall et al., 2007), at a precision around 0.1
nmol mol⁻¹ (determined with the average of 1-min data).

210 The four most abundant N₂O isotopocules (¹⁴N¹⁴N¹⁶O, 99.03%; ¹⁴N¹⁵N¹⁶O, 0.36%; ¹⁵N¹⁴N¹⁶O,
0.36%; ¹⁴N¹⁴N¹⁸O, 0.20%) were analyzed using a customized QCLAS system (Aerodyne Research,
Inc., USA) (Heil et al., 2014) coupled with an automated preconcentration device (Mohn et al.,
2010). Before entering the pre-concentration unit, sample air is passed through a Sofnocat 423 trap
(Molecular Products Limited, GB) to remove CO, and subsequently through an Ascarite trap
215 (Ascarite: 6 g, 10–35 mesh, Sigma Aldrich, Switzerland, bracketed by Mg(ClO₄)₂, 2 × 1.5 g, Alfa
Aesar, Germany) to remove CO₂ and water. Approximately 5.5 L of air with a flow of 250 ml min⁻¹

¹ (at 295 K and 3500 hPa) is then passed through a HayeSep D trap cooled to -145 °C to collect N₂O (Mohn et al., 2010). For N₂O release to the multipath cell of the QCLAS, the HayeSep D trap is quickly heated to 10 °C and flushed with high-purity synthetic air (20.5% of O₂ in N₂) carrier gas at a flow rate of 25 ml min⁻¹ (at 295 K and 3500 hPa). A final cell pressure around 16 hPa is achieved, which results in an N₂O mixing ratio of about 45 μmol mol⁻¹. More instrumental details can be found in previous studies (Harris et al., 2017; Mohn et al., 2010, 2012). Sample tanks were each analyzed twice to yield duplicates for N₂O isotopic results, which left sufficient air for amount fraction analysis as described in the previous paragraph.

2.4 Data analyses

We used 10-minute averages of the continuous *in situ* measurements from the Jungfraujoch station across this study. For a point-to-point comparison of continuous and discrete measurements of N₂O mixing ratio, we aggregated 10-minute averages of *in situ* data for the same period when the discrete sample was filled into the cylinder (40 min).

In this study, we report abundances of N₂O isotopocules using δ notation (‰) as below:

$$\delta X = \frac{(R_{sample} - R_{standard})}{R_{standard}} \quad (1)$$

where X refers to ¹⁵N^α (¹⁴N¹⁵N¹⁶O), ¹⁵N^β (¹⁵N¹⁴N¹⁶O) and ¹⁸O (¹⁴N¹⁴N¹⁸O); R refers to the ratio between the amount fractions of the rare isotopocules as mentioned above and the amount fraction of ¹⁴N¹⁴N¹⁶O; isotope standards refer to atmospheric N₂ for ¹⁵N and Vienna Standard Mean Ocean Water (VSMOW) for ¹⁸O.

Hence, the total ¹⁵N content of N₂O and site-specific composition of N₂O isotopomers could be further illustrated as $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{15}\text{N}^{\text{SP}}$, respectively, according to the equations below:

$$\delta^{15}N^{bulk} = (\delta^{15}N^{\alpha} + \delta^{15}N^{\beta})/2 \quad (2)$$

$$\delta^{15}N^{SP} = \delta^{15}N^{\alpha} - \delta^{15}N^{\beta} \quad (3)$$

240 Two standards (CG1 and CG2; in 79.5% N₂ and 20.5% O₂) with distinct isotopic signatures ($\delta^{15}N^{\alpha}$
 = $16.29 \pm 0.07\text{‰}$ (CG1) and $-51.09 \pm 0.07\text{‰}$ (CG2); $\delta^{15}N^{\beta} = -2.59 \pm 0.06\text{‰}$ and $-48.12 \pm 0.04\text{‰}$;
 $\delta^{18}O = 39.37 \pm 0.04\text{‰}$ and $30.81 \pm 0.03\text{‰}$) were used for calibrating isotopic composition. The
 calibration gases CG1 and CG2 were calibrated on the Tokyo Institute of Technology (TIT) scale,
 based on cross-calibration with primary standards assigned by TIT (Mohn et al., 2012, 2014). In
 245 addition, CG1 was measured repeatedly between samples and target gases to account for
 instrumental drift. Both CG1 and CG2 have N₂O mixing ratios of 45 $\mu\text{mol mol}^{-1}$, similar to the
 N₂O amount fraction of the samples after preconcentration. However, to correct for possible
 instrumental dependence on N₂O mixing ratio, CG1 was diluted to N₂O mole fractions of 35-40
 $\mu\text{mol mol}^{-1}$ within each measurement batch. In general, duplicated isotopic measurements of flask
 250 samples yielded values of repeatability of 0.10-0.20‰ for $\delta^{15}N^{bulk}$ and $\delta^{18}O$, and 0.15-0.25‰ for
 $\delta^{15}N^{SP}$.

At the beginning of the project, a batch of three cylinders (50 L water volume, Luxfer, Italy) were
 filled with pressurized ambient air in Dübendorf with an oil-free, three stage compressor (SA-3,
 Rix Industries, USA) and used as long-term target gases. The pressurized ambient air target gas
 255 was analyzed with identical treatment as Jungfrauoch air samples during every analysis batch, to
 monitor long-term analytical drift. Standard deviations for repeated target gas measurements
 throughout the period of Jungfrauoch sample measurements, were 0.13‰ for $\delta^{15}N^{bulk}$, 0.21‰ for
 $\delta^{15}N^{SP}$, and 0.11‰ for $\delta^{18}O$ (Fig. S2).

2.5 Surface air footprint analysis and simulated regional N₂O enhancement

260 We analyzed the air mass origin at Jungfraujoch by applying the Lagrangian particle dispersion
model (LPDM) FLEX-PART in the backward mode (Stohl et al., 2005). The model was driven by
meteorological fields taken from the ECMWF-IFS operational analysis cycle, extracted at a
resolution of $1^\circ \times 1^\circ$, 90/137 levels globally, and at higher horizontal resolution of $0.2^\circ \times 0.2^\circ$ for
central Europe. We released 50000 virtual air parcels every 3 hours at 3000 m a.s.l. from
265 Jungfraujoch to perform backward dispersion simulations over 10 days, which allowed us to
calculate surface source sensitivities (concentration footprints). A release height of 3000 m a.s.l.
was previously determined to be an optimum for simulating concentration footprints at
Jungfraujoch, given the stated horizontal resolution which results in a considerable smoothing of
the complex, alpine orography (Keller et al., 2012). The 3-hourly surface footprints for the whole
270 observation period were used to categorize different transport regimes using the clustering
approach outlined in Sturm et al. (2013). This allowed us to distinguish among six different source
regions: Free Troposphere (FT), Southwest (SW), East (E), Local (L), West (W) and Northwest
(NW).

Similar to Henne et al. (2016) for CH_4 and based on spatially resolved N_2O emission inventories
275 (Meteotest for Switzerland; EDGAR for Europe), we used the FLEXPART concentration
footprints to calculate time-series of atmospheric mole fraction increases at Jungfraujoch resolved
by emission sectors (Henne et al., 2016). The emission inventory by Meteotest consists of 12
emission sectors, among which all sectors except “organic soils” are comparable to sectors in the
EDGAR inventory (See Table S1) (Janssens-Maenhout et al., 2019). To improve seasonal
280 representation of the emissions in our model, we used a monthly resolved, optimized version of
the emission inventory, which was obtained through inverse modeling using the N_2O atmospheric
mole fractions observed between March 2017 and September 2018 at the tall tower site

Beromuenster on the Swiss plateau (Henne et al., 2019). Therefore, in this study, source contributions to Jungfraujoch were estimated specifically for the period mentioned above.

285 **2.6 Evaluation of seasonal pattern and interannual trend for time-series**

To explore seasonality and interannual trends, we fit the time-series of *in situ* measurements of N₂O and O₃ mixing ratios, NO_y-to-CO ratios and isotopic measurements of N₂O with polynomial functions and Fourier series (four harmonics for *in situ* measurements and two harmonics for discrete measurements) (Thoning et al., 1989). Time-series were then decomposed into a linear
290 trend, seasonal variability (per 12 months) and residuals. This fit was conducted with a nonlinear least-squares (NLS) model with R-3.5.3 (R Core Team, 2016). The detrended seasonality was examined by comparing peak-to-peak amplitudes with our analytical precisions and the uncertainty given by the one standard deviation of monthly residuals. To determine interannual trends, a linear regression was applied to both the raw and the deseasonalized datasets. The
295 significance level is set to $p < 0.01$. The interannual trends for N₂O mixing ratios were found to be little affected by seasonality, so growth rates were determined only based on the raw datasets.

Although Jungfraujoch is a remote site, episodic influence from the planetary boundary layer can be observed at the station (Pandey Deolal et al., 2012; Zellweger et al., 2003). For evaluating trends of N₂O mixing ratio measurements, we filtered out *in situ* data with significant influence of plenary
300 boundary layer, in order to represent a major air mass footprint from the free troposphere (FT). In addition to the air transport regimes, an alternative filtering criterion for the free troposphere was based on the published mean ranges of NO_y mixing ratios (501-748 pmol mol⁻¹ depending on the season) and NO_y to CO ratios (0.003-0.005 depending on the season) at Jungfraujoch (Zellweger et al., 2003). This criterion is less strict than that given by footprint analyses (Herrmann et al.,

305 2015). After applying this criterion to the isotopic time-series (which led to the exclusion of 32 measurement points), we re-evaluated the seasonal and interannual trends in the N₂O isotopic composition. In addition, because of the strong variability observed for isotopic data during the first 1.5 years (until February 2016), we performed an independent evaluation for the time-series starting from August 2016.

310 **2.7 Two-box model simulation**

A two-box model representing a well-mixed troposphere and stratosphere was used to estimate the anthropogenic N₂O source strength and isotopic composition from the trends measured at Jungfraujoch, similar to the approaches used by several previous studies (Ishijima et al., 2007; Röckmann et al., 2003; Schilt et al., 2014; Sowers et al., 2002). The input variables used to run the
315 model are given in Table S2. 200 iterations of the model were run using a Monte Carlo-style approach to approximate the uncertainty considering the uncertainty distribution for each input variable as given in Table S2. All variables were set independently within the Monte Carlo approximation except for preindustrial N₂O life time (τ_{PI}), which was fixed to 106% of the present-day N₂O life time τ_{PD} (Prather et al., 2015).

320 Within each iteration of the model, the preindustrial N₂O burden was first described, assuming steady state in the preindustrial era. The preindustrial stratospheric N₂O mixing ratio ($c_{S,PI}$) (270 ± 7.5 nmol mol⁻¹) was taken from Sowers et al. (2002):

$$0 = F_{ex} (c_{PI} - c_{S,PI}) - (M_{PI} + M_{S,PI})/\tau_{PI} \quad (4)$$

where F_{ex} refers to the troposphere-stratosphere exchange rate; c_{PI} refers to the preindustrial
325 tropospheric N₂O mixing ratio; and M_{PI} and $M_{S,PI}$ are the masses of N₂O in the troposphere and stratosphere respectively. The preindustrial terrestrial flux in Sowers et al. (2002) (equation 2) was

used here assuming no anthropogenic emissions. The delta values for the preindustrial stratosphere and the fractionation factor for the stratospheric sink were taken from equations 6 and 7 from Sowers et al. (2002) assuming steady state and no anthropogenic emissions. The model was run
330 with a yearly time step starting from the preindustrial assuming that anthropogenic emissions began in 1845 (Sowers et al., 2002). For each year of the model run, the anthropogenic flux was calculated according to the exponential increase described by Sowers et al. (2002):

$$F_{\text{anth},t} = e^{\alpha(t-t_0)} - 1 \quad (5)$$

where t is the current year, $t_0 = 1845$ and α is the growth rate (assumed to be constant). The rates
335 of change for tropospheric and stratospheric N₂O mixing ratios were then retrieved from equations 2 and 3 in Sowers et al. (2002), and for the isotopic composition of stratospheric and tropospheric N₂O from equations 6 and 7 in Sowers et al. (2002).

The values of the parameters describing the anthropogenic flux were optimized to fit both the trend and the absolute values for the five years of Jungfraujoch isotope data, and the mixing ratio data
340 from the Jungfraujoch flasks and *in situ* data since 2005 (GAW data source). The uncertainties in α and in the anthropogenic source isotopic signatures were approximated by one standard deviation of values derived from repeated model runs.

2.8 Evaluation of the combined effects from STE and soil emission on $\delta^{15}\text{N}^{\text{SP}}$

To evaluate the combined effects of STE and soil emission on the seasonal variability of $\delta^{15}\text{N}^{\text{SP}}$
345 (i.e. August minima), we made a mixing calculation as below:

Soil emission: Based on the determined seasonality of N₂O mole fraction at Jungfraujoch, the maximum N₂O mole fraction enhancement was approximately 0.2 nmol mol⁻¹ above baseline (Fig.

1b). Hence, we assumed N₂O enhancement from soil emission to be close to 0.15 to 0.20 nmol mol⁻¹, which is close to the maximum N₂O enhancement in our observation. The isotopic effect from soil emission can be derived from the difference between soil emission (7.2‰; Table S1) and tropospheric air (18‰, Fig. 2) in δ¹⁵N^{SP}, i.e. -10.8‰.

Mixing with stratospheric air: The minimum of N₂O mole fraction in August (-0.20 nmol mol⁻¹) is likely to be the result of both N₂O mole fraction enhancement from soil emission and N₂O mole fraction depletion due to STE. Given the assumed N₂O enhancement from soil emission, we estimated the N₂O depletion due to STE as -0.35 to -0.40 nmol mol⁻¹. The isotopic effect due to mixing with stratospheric air can be approximated using the apparent isotopic fractionation ε_{app} (Kaiser et al., 2006), which was derived from the slope of Rayleigh plot with normalized N₂O mole and isotope ratios. For ¹⁵N^{SP}, ε_{app} is calculated from the difference between ¹⁵N/¹⁴N isotope fractionations at the central and terminal N atoms, i.e. α_{ε_{app}} - β_{ε_{app}}. Therefore, for the lower stratosphere, ε_{app}(¹⁵N^{SP}) was calculated to be about -15‰ (see more details in Kaiser et al., 2006).

Overall effect: Combing the isotope effects and contributions to the change of N₂O mole fraction by the two processes, the net effect is [(-0.35 to -0.40 nmol mol⁻¹) (-15‰) + (0.15 to 0.20 nmol mol⁻¹) (-10.8‰)] / (330 nmol mol⁻¹) ≈ 0.01‰. Such isotope effect is below our analytical precision and too small to be measured in the background atmosphere.

2.9 “Bottom-up” estimates of source isotopic signatures

To gauge the accuracy of the two-box model, we deployed a “bottom-up” approach as an alternative method of estimating the N₂O source signatures. The isotopic signatures of most N₂O source sectors given in the Meteotest/EDGAR emission inventory are available from the literature, except for the “Refinery” (Table S1). As “Refinery” generally contributes only about 0.02% of the

370 N₂O emission at Jungfraujoch, it was excluded for source isotopic signature estimation. The simulated N₂O emissions by variable sources were categorized according to the EDGAR emission types (Janssens-Maenhout et al., 2019). We then calculated isotopic signatures for the overall source and the anthropogenic sources alone (excluding indirect natural emission) as weighted averages.

375 **3 Results**

3.1 Atmospheric N₂O mixing ratios at Jungfraujoch

We observed a linear growth of atmospheric N₂O at Jungfraujoch during the period 2014-2018 (Fig. 1a). A point-to-point comparison of discrete and *in situ* measurements showed good agreement, in particular after the first year (2015-2018), where the data quality of *in situ* measurements was largely improved due to the implementation of the more precise laser spectroscopy method as compared to GC-ECD (Fig. S3). The improvement in analytical precision for N₂O mixing ratio was due to better temporal coverage by the OA-ICOS instrument, in contrast with the GC analyses which conduct one measurement per 24-30 minutes. The annual growth rates from 2014 to 2018 determined with *in situ* measurements were 0.880 ± 0.001 and 0.993 ± 0.001 nmol mol⁻¹ a⁻¹ with and without GC-ECD measurements in 2014, respectively. This difference in N₂O growth rates is probably due to the limited data quality of GC-ECD, although a lower growth rate in 2014 compared to 2015-2018 cannot be excluded. It is noteworthy that the N₂O growth rate determined for 2015 to 2019 at Jungfraujoch is slightly above the global mean growth rate for the recent decade reported by NOAA (0.93 ± 0.03 nmol mol⁻¹ a⁻¹) (WMO, 2018). If we filter the *in situ* dataset to examine only the “free troposphere” periods, we obtain a lower increase (0.858 ± 0.002 nmol mol⁻¹ a⁻¹). By comparison, the absolute growth rate determined from the discrete gas samples was even lower albeit larger uncertainty (0.813 ± 0.027 nmol mol⁻¹ a⁻¹).

A significant seasonal pattern was observed for N₂O mixing ratios measured *in situ*, with a maximum in early summer and a minimum in late summer (Fig. 1b). For discrete N₂O measurements a similar trend was observed, but the detrended seasonality was not significant, which might be due to the much lower number of samples (Fig. S4).

3.2 Interannual trends of N₂O isotopic composition and anthropogenic source signatures

Time-series of $\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ for atmospheric N₂O at Jungfraujoch are shown in Figure 2. The NLS model simulation accounts well for the variabilities of isotopic time-series. Interannual trends of three isotopic deltas were determined for both raw and deseasonalized datasets by linear regression (Table 1). The deseasonalized interannual trends were slightly smaller than the trends determined with the raw datasets. For the whole dataset, the deseasonalized trend indicates a significant decrease in $\delta^{15}\text{N}^{\text{bulk}}$, of $-0.052\pm 0.012\%$ a⁻¹. In contrast, deseasonalized time-series of $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ increased, albeit insignificantly, by $0.065\pm 0.027\%$ a⁻¹ and $0.019\pm 0.011\%$ a⁻¹, respectively. The trends determined for periods with major air mass footprints from the free troposphere were close to those calculated for the whole dataset, except that $\delta^{15}\text{N}^{\text{SP}}$ trends decreased after filtering out the samples with significant impact from plenary boundary layer. This indicates that N₂O interannual trends observed at Jungfraujoch are of regional relevance, despite the fact that a small impact from local sources can be seen. Because of the observed irregular variability and the change in sampling frequency (though no change in daily sampling time) in our dataset, we separated the time-series into two phases: April 2014-February 2016 (first phase; weekly sampling) and August 2016-December 2018 (second phase; bi-weekly sampling). In the first phase, the rates of increase in $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ were almost one order of magnitude larger than over the whole dataset. This is most likely due to the unexpectedly low $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ in summer 2014 followed by a distinct increase in winter 2014-2015, which results in large rates of increase over short periods. Such growth rates were not seen in the second phase, when both $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ showed small and insignificant variations. $\delta^{15}\text{N}^{\text{bulk}}$ displayed a decreasing interannual trend in both phases; however, the rate of decrease was larger in the second phase ($-0.130\pm 0.045\%$ a⁻¹).

We tuned our two-box model to best match the observed N₂O mixing ratios and isotopic composition at Jungfraujoch. An estimate of anthropogenic emissions and source signatures is given in Table 2. For 2018, annual N₂O emissions were estimated to be 8.6±0.6 Tg N₂O-N a⁻¹ equivalents. The average isotopic signatures for anthropogenic sources were -8.6±4‰, 34.8±3‰ and 10.7±4‰ for δ¹⁵N^{bulk}, δ¹⁵N^{SP} and δ¹⁸O, respectively, which are clearly lower than those for preindustrial N₂O in the tropospheric background (Table S2; Toyoda et al., 2013).

3.3 Seasonal variation of N₂O isotopic composition

δ¹⁵N^{SP} of N₂O showed the most pronounced variability among all isotopic time-series (Fig. 2), spanning 2.5‰ for individual flask sample measurements. Seasonal variability was estimated with the NLS model and presented as mean seasonal cycles (Fig. 3). For δ¹⁵N^{SP} a “summer minimum” was found regardless of whether the entire dataset or only the second phase was considered (Fig. 3), although seasonal variability of the second time-series was smaller and showed the minimum occurring earlier. The seasonal pattern of δ¹⁵N^{bulk} determined from the whole dataset indicates a significant summer maximum, but this was not seen when only the data from the second phase was taken, as there was no significant seasonal pattern over this period alone. For δ¹⁸O, we observed only small temporal variability and a lack of seasonal pattern. In addition, seasonal variations of time-series filtered for free troposphere were evaluated; these show temporal patterns similar to the whole dataset (Fig. S5).

3.4 Air mass origin and *in situ* measurements at Jungfraujoch

Back-trajectory simulations indicate six major transport clusters during 2014-2018, as shown in Figure 4a. Four of these transport regimes (SW, E, L and NW) dominate, accounting for about 60-90% coverage of the whole period. By contrast, the free troposphere cluster only represents 10-

20% of the data. Averaged monthly contributions of transport clusters are shown in Figure 4b, with more pronounced impact by the L, E and NW regions in summer and stronger contribution by FT and SW in winter. The source patterns of the air masses at Jungfraujoch were generally consistent across the years in the present study. However, an apparent discrepancy was found for
445 discrete sampling times in the last two years (e.g. particularly low contribution from SW) which is most likely due to the low and variable sampling frequency of the discrete sample collection (Fig. 4b).

The detrended seasonal variability of *in situ* measurements indicates summer maxima for NO_y mixing ratios as well as NO_y-to-CO ratios at Jungfraujoch (Fig. S6). This likely indicates stronger
450 exchange with the polluted planetary boundary layer in summer (Herrmann et al., 2015; Zellweger et al., 2003), which is consistent with the seasonal pattern of air mass footprint derived from back-trajectory simulations. The late spring-to-summer maxima for O₃ mixing ratios may be attributed to air mixing with stratosphere and/or planetary boundary layer, similar to the findings from a previous study at Jungfraujoch (Tarasova et al., 2009). On the other hand, CO shows a maximum
455 in early spring and decreases in summer when its atmospheric lifetime is shortest. Atmospheric O₃, NO_y and CO measurements during our discrete sampling periods also well represented seasonal variability shown for *in situ* measurements, except for 2016-2017 where there was a five-month sampling gap (Fig. S6).

Comparisons of air mass footprints as well as O₃, NO_y and CO mixing ratios between *in situ* and
460 discrete sampling indicate that the discrete sampling covers the main air source regions and variabilities in local pollution/free troposphere fairly well (Fig. 4 and S6). In the second phase (2016-2018), the less frequent sampling impedes evaluation of the seasonal and interannual variabilities.

3.5 Relationship between N₂O isotopic signatures and air mass footprints

465 We categorized N₂O mixing ratio and isotopic signature time-series into subsets based on the six
air mass transport clusters. One-way ANOVA among clusters indicates that N₂O mixing ratios in
air masses originating from cluster L were significantly higher and those from clusters FT and W
were significantly lower than the others (Fig. 5). In accordance with the pattern found for mixing
ratios, $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ were high for cluster FT, and low for cluster L. For $\delta^{15}\text{N}^{\text{bulk}}$, little difference
470 between transport clusters was detected.

4 Discussion

4.1 Quality assurance of isotopic measurements

This study reports the first results of background N₂O isotopic measurements based on a laser spectroscopic technique. Benefiting from the preconcentration process, we achieved measurement
475 repeatability for a target gas of 0.10-0.20‰ for $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$ (Fig. S2), which is comparable to that of IRMS measurements of ambient atmosphere (Park et al., 2012; Prokopiou et al., 2017; Röckmann et al., 2003; Toyoda et al., 2013). The long-term robustness of our technique is adequate for disentangling both seasonal and interannual temporal variability as shown in Figure 2. In particular, our repeatability of target measurements for $\delta^{15}\text{N}^{\text{SP}}$ (0.15-0.25‰) appears to be better
480 than previous studies measuring background atmosphere or firn air (0.8‰, Park et al., 2012; 0.3‰, Prokopiou et al., 2017; 0.3‰, Toyoda et al., 2013).

4.2 Seasonal variabilities of atmospheric N₂O isotopic composition

In situ measurements of N₂O mixing ratios showed a clear early summer maximum and late summer minimum (Fig. 1). Such a seasonal pattern was previously found for a number of NOAA
485 and AGAGE sites analyzing long-term N₂O records in the NH (Jiang et al., 2007; Nevison et al., 2011). One explanation of the late-summer minimum is a strong influence of the STE process in this period, which transports N₂O-depleted but isotopically enriched air downward from the stratosphere into the troposphere (Park et al., 2012; Snider et al., 2015b). During the late summer at Jungfraujoch, we find strong enrichment of ¹⁵N in atmospheric N₂O according to the detrended
490 seasonality for the whole dataset (Fig. 3). This is supported by a FLEXPART model simulation of the contribution of upper tropospheric air to Jungfraujoch station, showing highest contributions in August (Fig. S7; Henne et al., Personal Communication). At Hateruma Island, Japan, Toyoda

et al. (2013) observed a seasonal pattern of atmospheric N₂O mixing ratios which is comparable with our study, but found insignificant variations of isotopic composition. On the other hand, N₂O seasonal variability could be influenced by oceanic emission sources (Jiang et al., 2007; Nevison et al., 2005), complicating the explanations for the observed temporal patterns. For example, in another study looking at archived air from Cape Grim, Australia, Park et al. (2012) detected an April-May minimum and a November-December maximum for N₂O. This is expected for the SH, as STE is most prevalent in April (Nevison et al., 2011). They observed negative correlations of $\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\alpha}$ and $\delta^{18}\text{O}$ with N₂O mixing ratios, appearing to support the idea that the STE process is responsible for seasonal variabilities in N₂O mixing ratios and isotopic composition at Cape Grim. However, the seasonal cycle for $\delta^{15}\text{N}^{\alpha}$ was much larger than $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$, which could not be explained by STE alone. They suggested that the seasonal patterns of N₂O isotopes at Cape Grim may be due to mixing between oceanic sources (high N₂O with low ¹⁵N and ¹⁸O) and STE (low N₂O with high ¹⁵N and ¹⁸O) (Nevison et al., 2011; Park et al., 2012). However, because we observed a concurrent minimum of $\delta^{15}\text{N}^{\text{SP}}$ and maximum of $\delta^{15}\text{N}^{\text{bulk}}$ in July-August with low N₂O at Jungfraujoch (Fig. 3), additional mechanisms must be considered here.

Regional model simulations based on Swiss N₂O emissions derived from the inverse method were used to explore contributions from different sources to the variability in N₂O enhancements at Jungfraujoch. As shown in Figure 6a&6b, soil emissions, including direct and indirect emissions from agricultural lands and emissions from (semi-)natural areas, account for more than 70% of the total N₂O enhancements, while manure and waste management contribute another 20%. Total N₂O enhancements appeared to be highest in May to July (Fig. 6c), in accordance with the highest contribution by soil emissions. The early-to-middle summer maximum in the simulated N₂O enhancements is comparable with maximum of N₂O mixing ratios in early summer as observed at

Jungfraujoch (Fig. 1b). This underlines the importance of soil emission in accounting for atmospheric N₂O variability (Saikawa et al., 2014). In late summer, the minimum of $\delta^{15}\text{N}^{\text{SP}}$ (Fig. 3) may be then attributed to the influence of soil emitted N₂O, which has lower $\delta^{15}\text{N}^{\text{SP}}$ ($7.2\pm 3.8\text{‰}$; Table S1) than the troposphere (Fig. 2). However, the STE process, which resulted in the minimum of N₂O mixing ratio, likely contributes a positive isotope effect in the meanwhile (Kaiser et al., 2006). In order to evaluate the combined effect of STE and soil emission on $\delta^{15}\text{N}^{\text{SP}}$ in late summer, we applied a mixing calculation. Such estimate was made based on the approximated N₂O enhancement/depletion contributed by the two processes and the assumed isotope effects (see more details in M&M). The mixing calculation indicated an overall isotope effect of about 0.01‰, which is extremely small and below our analytic precision. This practice suggests that it is still challenging to build a direct link of N₂O sources/processes with the observed isotope signature in the background atmosphere. It is also noteworthy that the $\delta^{15}\text{N}^{\text{SP}}$ used in the calculation ($7.2\pm 3.8\text{‰}$) may underestimate the isotope effects of soil emission, given that denitrification, as a major N₂O process in soils, produces N₂O with $\delta^{15}\text{N}^{\text{SP}}$ close to 0‰ (Sutka et al., 2006). Previous field studies have demonstrated that low- $\delta^{15}\text{N}^{\text{SP}}$ N₂O emissions ($\sim 0\text{‰}$), i.e. following the denitrification pathway, predominates during summer periods at Swiss (Wolf et al., 2015) and German (Ibrahim et al., 2019) grasslands. By contrast, the influence of biogeochemical processes (nitrification and denitrification) on $\delta^{15}\text{N}^{\text{bulk}}$ is generally smaller than that on $\delta^{15}\text{N}^{\text{SP}}$ (Toyoda et al., 2011), and such effect on $\delta^{15}\text{N}^{\text{bulk}}$ are usually overwritten by the wide range of isotopic signatures in soil N substrates (Sutka et al., 2006). Hence, given the distinct $\delta^{15}\text{N}^{\text{bulk}}$ maximum and N₂O minimum in late summer during our observation (Figs. 1 and 3), we suggest that the STE process is mainly responsible for the seasonal variability in $\delta^{15}\text{N}^{\text{bulk}}$.

The footprint analyses based on air mass residence time revealed a seasonal pattern, with a higher

contribution of background air from the FT and SW regions in winter and more pronounced
540 contribution of local planetary boundary layer air from the L, E and NW regions in summer (Fig.
4b). The higher frequency of air mass footprints recently in contact with the surface in summer is
consistent with inverse modeling results, indicating a larger contribution of soil N₂O emissions in
June/July (Fig. 6). For the air mass regime representing the free troposphere, N₂O mixing ratios
observed at Jungfraujoch were significantly below the average, while $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ were higher
545 (Fig. 5). By contrast, the local cluster (L) representing a strong impact from the planetary boundary
layer had higher N₂O mixing ratios and lower isotopic signatures (except $\delta^{15}\text{N}^{\text{bulk}}$) than the other
source regions. In addition, the ratio of NO_y to CO, which is a more straightforward indicator of
the free troposphere (Zellweger et al., 2003), show significant negative correlations with $\delta^{15}\text{N}^{\text{SP}}$
and $\delta^{18}\text{O}$, but not with $\delta^{15}\text{N}^{\text{bulk}}$ (Fig. S8). This further suggests that the seasonal variability of
550 $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ observed at Jungfraujoch is most likely influenced by ground-derived emissions,
while fluctuations in N₂O mixing ratios and $\delta^{15}\text{N}^{\text{bulk}}$ are possibly driven by STE.

Considering the complexity in mechanisms responsible for N₂O isotopic variations, we strongly
recommend more field measurements of N₂O isotopic signatures at higher frequency and at
different background sites, in order to cover spatial and temporal variability in N₂O sources. For
555 example, in the second phase, we only detected a significant seasonality of $\delta^{15}\text{N}^{\text{SP}}$, with a minimum
in July, which is one month earlier than the summer minimum found for the whole dataset (Fig.
3). This may be attributed to a difference in source regions, as Northwest regions appeared to be
significantly more important during 2017 (second phase). However, due to low sampling
frequency, it is challenging to overcome the large uncertainty in seasonality analysis for a two-
560 year period such as the second phase.

Based on our bottom-up approach, we simulated isotopic signatures for the overall N₂O sources

responsible for the N₂O mixing ratio increase in the atmosphere (Fig. S9). However, the interpretation of simulated versus observed variability in N₂O isotopic composition was difficult, except for the somewhat similar patterns in $\delta^{18}\text{O}$. Our results suggest a limitation in the current
565 knowledge and literature values on isotopic signatures of most N₂O sources. In addition, most N₂O sources may not exhibit a well-defined isotopic signature but a range of values regulated under a number of processes/environmental factors. For example, isotopic signatures of soil-derived N₂O are often determined by an interaction of several soil and climatic factors. It might be possible in the future to model these changes implementing isotopes in ecosystem models, as recently
570 demonstrated by Denk et al. (2019).

4.3 Interannual trends of atmospheric N₂O isotopic composition

Over a period of almost five years, our observations show an interannual increase in N₂O mixing ratio and decrease in $\delta^{15}\text{N}^{\text{bulk}}$ (Fig. 7). This is to be expected, assuming that the atmospheric N₂O increase is primarily attributed to anthropogenic sources, which emit isotopically lighter N₂O
575 relative to the tropospheric background (Table S1) (Rahn and Wahlen, 2000). Compared to several studies on firn air (Ishijima et al., 2007; Röckmann et al., 2003) and surface air (Park et al., 2012; Röckmann and Levin, 2005; Toyoda et al., 2013), the rate of decrease for $\delta^{15}\text{N}^{\text{bulk}}$ at Jungfraujoch is relatively high (-0.05 to -0.06 ‰ a⁻¹, Table 1). Such a discrepancy in the $\delta^{15}\text{N}^{\text{bulk}}$ trend could be due to a large contribution of terrestrial N₂O emission from the European continent to Jungfraujoch
580 (Figs. 1 and 5), as N₂O originating from soil emissions is significantly more isotopically depleted than that of oceanic sources (Snider et al., 2015b). Nevertheless, our observation period is shorter than that of other studies, so the interannual trends determined here are more likely affected by year-to-year variability. Among all reported records, the decrease of $\delta^{15}\text{N}^{\text{bulk}}$ observed at Hateruma Island was the most up-to-date and smallest (-0.020-0.026‰ a⁻¹) (Toyoda et al., 2013). The authors

585 argued that the smaller declining trend for $\delta^{15}\text{N}^{\text{bulk}}$ may be explained by the recent increase in anthropogenic isotopic ratios particularly for agricultural N_2O emissions, although Ishijima et al. (2007) suggested a decline in both $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$ in anthropogenic N_2O from 1952-1970 to 1970-2001 based on inverse modeling.

For the interannual trends observed at Jungfraujoch, it is noteworthy to point out that our 590 observations covering a rather short period may lead to large uncertainties despite statistical significance. The discrepancy found in the trends between the first and second phases indicates that variability of N_2O isotopic composition is likely to obscure interannual trends over shorter periods (Toyoda et al., 2013). Hence, extended time-series of isotopic measurements are needed to reevaluate, for example, the observed tendency of increase in $\delta^{18}\text{O}$ and $\delta^{15}\text{N}^{\text{SP}}$ at Jungfraujoch 595 (Table 1; only significant during the first phase). For $\delta^{18}\text{O}$ of atmospheric N_2O , a generally declining trend smaller than that of $\delta^{15}\text{N}^{\text{bulk}}$ has been indicated by a number of observations (Bernard et al., 2006; Ishijima et al., 2007; Park et al., 2012; Röckmann et al., 2003; Röckmann and Levin, 2005). This is expected as $\delta^{18}\text{O}$ of anthropogenic N_2O is not much different from that of the natural background, assuming that the oxygen atom in N_2O is largely derived from soil water 600 and ambient oxygen during production (Rahn and Wahlen, 2000).

It is still a challenging task to disentangle interannual trends of $\delta^{15}\text{N}^{\text{SP}}\text{-N}_2\text{O}$ in the background atmosphere, due to limitations in analytical repeatability and precision (Harris et al., 2017; Mohn et al., 2014). Past results have reached inconsistent conclusions, showing positive (Bernard et al., 2006; Park et al., 2012; Prokopiou et al., 2017; Röckmann and Levin, 2005) or negative 605 (Röckmann et al., 2003; Toyoda et al., 2013) trends of similar magnitude (Fig. 7). On the one hand, the negative trend in $\delta^{15}\text{N}^{\text{SP}}$ could be explained by the significantly lower $\delta^{15}\text{N}^{\text{SP}}$ from anthropogenic sources (e.g. agricultural sources; Table S1) than of the tropospheric background

(near 18‰; Fig. 7). On the other hand, Park et al. (2012) suggested that the increase of $\delta^{15}\text{N}^{\text{SP}}$ in the atmospheric N_2O may reflect a global increase in importance of the contribution by nitrification (high- $\delta^{15}\text{N}^{\text{SP}}$ process) to agricultural N_2O emissions. This is based on the assumption that the growth of N_2O emissions is largely due to enhanced fertilizer application which promotes nitrification activity (Pérez et al., 2001; Tian et al., 2019). The observed mean increase rate of 0.02‰ a^{-1} for $\delta^{15}\text{N}^{\text{SP}}$ by Park et al. (2012) could then be translated into an increase of 13-23% for the relative amount of nitrification-derived N_2O between 1750 and 2005. However, this should be further evaluated with more frequent sampling (Park et al. (2012) only sampled 1-6 times per year) and tested with isotopic measurements across the NH, where agricultural N_2O emissions are more dominant than in the SH. In addition, the strong seasonal pattern of $\delta^{15}\text{N}^{\text{SP}}$ at Jungfraujoch suggests that seasonal variations of $\delta^{15}\text{N}^{\text{SP}}$ in response to climatic or source factors are crucial and must be taken into consideration for evaluating interannual $\delta^{15}\text{N}^{\text{SP}}$ trends.

620 **4.4 Simulated anthropogenic N_2O sources with the two-box model and comparison with other studies**

To further evaluate anthropogenic source signatures of N_2O isotopic composition, we applied a two-box model representing a well-mixed troposphere and stratosphere (Röckmann et al., 2003; Schilt et al., 2014; Sowers et al., 2002). The model runs with the whole dataset and the dataset filtered for free-troposphere only (Table 2) exhibit statistically identical results, supporting that our model estimates, with observations at Jungfraujoch, are representative of the background atmosphere. The simulated trends of the N_2O mixing ratios and isotopic composition show a gradual increase in N_2O and decrease in the isotopic signatures (see Fig. 7), which agree with existing observations within the model uncertainty. However, this does not hold for individual studies considered separately. For example, the N_2O mixing ratios observed by Röckmann et al.

(2003) and Prokopiou et al. (2017) would lead to a higher preindustrial N₂O compared to our model simulation, which is likely due to the uncertainty in the firm air records (Prokopiou et al., 2017).

We compared the anthropogenic isotopic signatures determined by our two-box model with other
635 similar studies in Table 2. Our estimates generally lie within the ranges given in the earlier studies (Ishijima et al., 2007; Park et al., 2012; Prokopiou et al., 2017; Röckmann et al., 2003; Sowers et al., 2002; Toyoda et al., 2013). However, isotopic signatures of N₂O sources estimated for 2018 in this study are higher in $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$ (by 4-8‰), and lower in $\delta^{15}\text{N}^{\text{SP}}$ (by 2-7‰) than model estimates for the early 2000s from two other studies from SH (Park et al., 2012; Prokopiou et al.,
640 2017). Such differences in $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{18}\text{O}$ could be related to interhemispheric differences, as the relative contributions of N₂O sources vary between the two hemispheres (Toyoda et al., 2013). Also, more interestingly, this could suggest a shift in the N₂O source isotopic signatures over the last few decades. For example, an increase of $\delta^{15}\text{N}^{\text{bulk}}$ in anthropogenic N₂O sources over time may be attributed to growing contributions of other industrial/waste sources with high $\delta^{15}\text{N}^{\text{bulk}}$
645 (Prokopiou et al., 2017). In addition, if the assumption of increasing $\delta^{15}\text{N}^{\text{bulk}}$ and decreasing $\delta^{15}\text{N}^{\text{SP}}$ in anthropogenic N₂O sources over time holds, it points to a recently growing contribution of denitrification relative to nitrification, to the global atmospheric N₂O increase (Sutka et al., 2006; Toyoda et al., 2013). By contrast, Park et al. (2012) and Prokopiou et al. (2017) proposed an increasing importance of nitrification for anthropogenic N₂O emissions based on the increasing
650 $\delta^{15}\text{N}^{\text{SP}}$ trend since 1940. This may suggest that a strong climate change feedback has recently resulted in significant shifts in N₂O source process, hence twisting the isotopic signatures of anthropogenic sources (Griffis et al., 2017; Xu-Ri et al., 2012). Alternatively, the uncertainty in determining N₂O isotopic signatures in the background atmosphere and inter-laboratory

comparability may play a role in the observed discrepancy.

655 Given the strong heterogeneity in source contributions to N₂O emissions around the globe (Saikawa et al., 2014), current two- and four-box model estimates based on observations at individual sites or regions are likely to reflect latitudinal or even interhemispheric differences in anthropogenic isotopic signatures. On the other hand, previous discussions of the model sensitivities by Röckmann et al. (2003) and Toyoda et al. (2013) have suggested that anthropogenic isotopic values are most sensitive to the trends in tropospheric isotopic values and the relative
660 difference in tropospheric isotopic values between present and preindustrial times. For example, given the similar parameters used for preindustrial times as our study, Park et al. (2012) observed much lower $\delta^{15}\text{N}^{\text{bulk}}$ in the recent troposphere than in our case, hence resulting in significantly lower $\delta^{15}\text{N}^{\text{bulk}}$ for the anthropogenic source. Furthermore, Park et al. (2012) and Prokopiou et al.
665 (2017) simulated a positive trend in $\delta^{15}\text{N}^{\text{SP}}$ relative to preindustrial times, which in return resulted in a much higher $\delta^{15}\text{N}^{\text{SP}}$ for the anthropogenic sources.

Using an alternative bottom-up approach, we estimated the anthropogenic source isotopic signatures based on the N₂O emission inventory simulated for Jungfraujoeh and published source isotopic signatures as summarized by Harris et al. (2017) (Table S1). The retrieved anthropogenic isotopic signatures (Table 3) were largely in agreement with the isotopic signature of agricultural soil emissions (Snider et al., 2015b; Wolf et al., 2015), indicating that this source could explain more than 60% of the total N₂O emissions. However, the anthropogenic isotopic signatures estimated by this approach were lower than the results from our two-box model (Table 2). In contrast, another similar bottom-up estimate based on the global N₂O emission inventory (Toyoda et al., 2013) reported anthropogenic isotopic values that agree well with our box-model results.
675 This may be explained by the different isotopic signatures used to describe agricultural N₂O

emissions, as those values used for the bottom-up estimates by Toyoda et al. (2013) were significantly lower (Toyoda et al., 2011) than those used in this study (Snider et al., 2015b; Wolf et al., 2015). Such bottom-up estimation suggests that more isotopic measurements of the background atmosphere from different regions, and better constraints on individual anthropogenic (especially agricultural) N₂O isotopic signatures, are necessary for a better representation of N₂O isotopic composition in atmospheric modeling studies.

5 Conclusions

With the recently developed laser spectroscopic technique coupled with a preconcentration device, we achieved good repeatability in measurements of N₂O isotopic composition from the background atmosphere at Jungfraujoch, Switzerland. This time-series covered a period of five years and showed a distinct seasonality, with $\delta^{15}\text{N}^{\text{bulk}}$ maxima and $\delta^{15}\text{N}^{\text{SP}}$ minima in late summer, associated with the lowest N₂O mixing ratios over the year. The seasonal fluctuation of $\delta^{15}\text{N}^{\text{bulk}}$ was associated with the stratosphere-troposphere exchange process, in agreement with other monitoring networks (Nevison et al., 2011), while the contrasting depletion of $\delta^{15}\text{N}^{\text{SP}}$ in later summer is possibly a combined result of STE and agricultural emissions, with the latter being more important. The analyses of air mass transport regimes together with the simulation of N₂O enhancements for Jungfraujoch supported our explanations and highlighted that the fluctuation between the free troposphere and local contributions dominated by soil emission drives the seasonality of $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ as observed at Jungfraujoch.

We found statistically significant interannual trends for $\delta^{15}\text{N}^{\text{bulk}}$, which is expected as anthropogenic N₂O sources are characterized by low ¹⁵N abundance. For $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$, interannual trends were highly uncertain and possibly masked by their large temporal variabilities. Using a two-box model approach, we simulated the evolution of N₂O isotopic composition from preindustrial times to the present. This model suggests an overall decreasing trend for all isotopic deltas in conjunction with the atmospheric N₂O increase. The anthropogenic source signatures given by the model generally agreed with previous studies. However, these model results are still sensitive to the ranges and trends of the observed N₂O isotopic signatures in the present troposphere. In the future, more extended records of high-precision N₂O isotopic measurements

705 and application of multiple-box modeling approaches (Rigby et al., 2013) are necessary to account for the global N₂O budget and evolution of anthropogenic sources.

Data availability

Data for this study have been deposited in a general data repository (<https://figshare.com/s/077562ab408dd1bd0880>; doi:10.6084/m9.figshare.12032760.v1, 2020).

710 **Author contribution:**

LY, EH and JM led and designed this study. LY, EH, SE conducted sample collection at Jungfrauoch; LY and EH analyzed discrete samples at Empa; MS and CZ contributed *in situ* measurements of N₂O, NO_y, CO and O₃ at Jungfrauoch; LY, EH and SH performed data analyses for the time-series and conducted model simulations. LY wrote the main manuscript; EH, SH and
715 JM were involved in the revisions of the manuscript and commenting. SE, MS, LE and CZ were also involved in scientific discussion and commenting on the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

720 We are thankful to the research infrastructure provided by the High Altitude Research Stations Jungfrauoch and Gornergrat. We are grateful to the help from the custodians (Mr. and Mrs. Fischer and Mr. and Mrs. Käser) at the research station of Jungfrauoch. We would like to thank Simon Wyss, Kerstin Zeyer, Patrik Zanchetta and Flurin Dietz for their support with the sample collection as well as laboratory assistance. The NABEL network is operated by Empa in collaboration with
725 the Swiss Federal Office for the Environment. Prof. Sakae Toyoda and Prof. Naohiro Yoshida from Tokyo Institute of Technology are acknowledged for their analyses of the applied reference standards. This study was financially supported by the Swiss National Science Foundation (grant number 200021_163075) and the Swiss contribution to the Integrated Carbon Observation System

(ICOS) Research Infrastructure (ICOS-CH). ICOS-CH is funded by the Swiss National Science
730 Foundation and in-house contributions. Longfei Yu was additionally supported by the
EMPAPOSTDOCS-II program, which receives funding from the European Union's Horizon 2020
research and innovation program under the Marie Skłodowska-Curie grant agreement number
754364.

735 **References**

- Appenzeller, C., Begert, M., Zenklusen, E. and Scherrer, S. C.: Monitoring climate at Jungfrauoch in the high Swiss Alpine region, *Sci. Total Environ.*, 391(2–3), 262–268, doi:10.1016/j.scitotenv.2007.10.005, 2008.
- 740 Bernard, S., Röckmann, T., Kaiser, J., Barnola, J.-M., Fischer, H., Blunier, T. and Chappellaz, J.: Constraints on N₂O budget changes since pre-industrial time from new firn air and ice core isotope measurements, *Atmos. Chem. Phys. Discuss.*, 5(4), 7547–7575, doi:10.5194/acpd-5-7547-2005, 2006.
- 745 Bourbonnais, A., Letscher, R. T., Bange, H. W., Échevin, V., Larkum, J., Mohn, J., Yoshida, N. and Altabet, M. A.: N₂O production and consumption from stable isotopic and concentration data in the Peruvian coastal upwelling system, *Global Biogeochem. Cycles*, 31(4), 678–698, doi:10.1002/2016GB005567, 2017.
- Buchmann, B., Hueglin, C., Reimann, S., Vollmer, M. K., Steinbacher, M. and Emmenegger, L.: Reactive gases, ozone depleting substances and greenhouse gases, in *From weather observations to atmospheric and climate sciences in Switzerland*, edited by S. Willemse and M. Furger, vdf Hochschulverlag AG., 2016.
- 750 Bukowiecki, N., Weingartner, E., Gysel, M., Coen, M. C., Zieger, P., Herrmann, E., Steinbacher, M., Gäggeler, H. W. and Baltensperger, U.: A review of more than 20 years of aerosol observation at the high altitude research station Jungfrauoch, Switzerland (3580 m asl), *Aerosol Air Qual. Res.*, 16(3), 764–788, doi:10.4209/aaqr.2015.05.0305, 2016.
- 755 Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R. and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 368, 20130122, doi:10.1098/rstb.2013.0122, 2013.
- Decock, C. and Six, J.: How reliable is the intramolecular distribution of ¹⁵N in N₂O to source partition N₂O emitted from soil?, *Soil Biol. Biochem.*, 65(2), 114–127, doi:10.1016/j.soilbio.2013.05.012, 2013.
- 760 Denk, T. R. A., Mohn, J., Decock, C., Lewicka-Szczepak, D., Harris, E., Butterbach-Bahl, K., Kiese, R. and Wolf, B.: The nitrogen cycle: A review of isotope effects and isotope modeling approaches, *Soil Biol. Biochem.*, 105, 121–137, doi:10.1016/j.soilbio.2016.11.015, 2017.
- Denk, T. R. A., Kraus, D., Kiese, R., Butterbach-Bahl, K. and Wolf, B.: Constraining N cycling in the ecosystem model LandscapeDNDC with the stable isotope model SIMONE, *Ecology*, 100(5), c02675, doi:10.1002/ecy.2675, 2019.
- 765 Fowler, D., Steadman, C. E., Stevenson, D., Coyle, M., Rees, R. M., Skiba, U. M., Sutton, M. a., Cape, J. N., Dore, a. J., Veno, M., Simpson, D., Zaehle, S., Stocker, B. D., Rinaldi, M., Facchini, M. C., Flechard, C. R., Nemitz, E., Twigg, M., Erisman, J. W. and Galloway, J. N.: Effects of global change during the 21st century on the nitrogen cycle, *Atmos. Chem. Phys. Discuss.*, 15(2), 1747–1868, doi:10.5194/acpd-15-1747-2015, 2015.
- 770 Fujii, A., Toyoda, S., Yoshida, O., Watanabe, S., Sasaki, K. and Yoshida, N.: Distribution of nitrous oxide dissolved in water masses in the eastern subtropical North Pacific and its origin inferred from isotopomer analysis, *J. Oceanogr.*, 69(2), 147–157, doi:10.1007/s10872-012-0162-4, 2013.
- Griffis, T. J., Chen, Z., Baker, J. M., Wood, J. D., Millet, D. B., Lee, X., Venterea, R. T. and Turner, P. A.: Nitrous oxide emissions are enhanced in a warmer and wetter world, *Proc. Natl. Acad. Sci.*, 201704552, doi:10.1073/pnas.1704552114, 2017.
- 775 Hall, B. D., Dutton, G. S. and Elkins, J. W.: The NOAA nitrous oxide standard scale for atmospheric observations, *J. Geophys. Res. Atmos.*, 112(9), 1–9, doi:10.1029/2006JD007954, 2007.

- 780 Harris, E., Ibraim, E., Henne, S., Hüglin, C., Zellweger, C., Tuzson, B., Emmenegger, L. and Mohn, J.: Tracking nitrous oxide emission processes at a suburban site with semicontinuous, in situ measurements of isotopic composition, *J. Geophys. Res. Atmos.*, 122, 1850–1870, doi:10.1002/2016JD025906, 2017.
- Harris, E. J., Nelson, D. D., Olsewski, W., Zahniser, M., Potter, E., Mcmanus, B. J., Whitehill, A., Prinn, R. G., Ono, S. and Harris, E.: Development of a spectroscopic technique for continuous online monitoring of oxygen and site-specific nitrogen isotopic com, *Anal. Chem.*, 86(3), 1726–1734, 2014.
- 785 Heil, J., Wolf, B., Brüggemann, N., Emmenegger, L., Tuzson, B., Vereecken, H. and Mohn, J.: Site-specific ^{15}N isotopic signatures of abiotically produced N_2O , *Geochim. Cosmochim. Acta*, 139, 72–82, doi:10.1016/j.gca.2014.04.037, 2014.
- Henne, S., Brunner, D., Folini, D., Solberg, S., Klausen, J. and Buchmann, B.: Assessment of parameters describing representativeness of air quality in-situ measurement sites, *Atmos. Chem. Phys.*, 10(8), 3561–3581, doi:10.5194/acp-10-3561-2010, 2010.
- 790 Henne, S., Brunner, D., Oney, B., Leuenberger, M., Eugster, W., Bamberger, I., Meinhardt, F., Steinbacher, M. and Emmenegger, L.: Validation of the Swiss methane emission inventory by atmospheric observations and inverse modelling, *Atmos. Chem. Phys.*, 16(6), 3683–3710, doi:10.5194/acp-16-3683-2016, 2016.
- 795 Henne, S., Mohn, J. and Brunner, D.: Quantification of Swiss nitrous oxide emissions through atmospheric observations and inverse modelling, Final Report, Project of FOEN, 2019.
- Herrmann, E., Weingartner, E., Henne, S., Vuilleumier, L., Bukowiecki, N., Steinbacher, M., Conen, F., Coen, M. C., Hammer, E., Juranyi, Z., Baltensperger, U. and Gysel, M.: Analysis of long-term aerosol size distribution data from Jungfraujoch with emphasis on free tropospheric conditions, cloud influence, and air mass transport, *J. Geophys. Res. Atmos.*, 120, 1751–1762, doi:10.1002/2015JD023660, 2015.
- 800 Ibraim, E., Wolf, B., Harris, E., Gasche, R., Wei, J., Yu, L., Kiese, R., Eggleston, S., Butterbach-Bahl, K., Zeeman, M., Tuzson, B., Emmenegger, L., Six, J., Henne, S. and Mohn, J.: Attribution of N_2O sources in a grassland soil with laser spectroscopy based isotopocule analysis, *Biogeosciences*, 16, 3247–3266, doi.org/10.5194/bg-16-3247-2019, 2019.
- 805 Ishijima, K., Sugawara, S., Kawamura, K., Hashida, G., Morimoto, S., Murayama, S., Aoki, S. and Nakazawa, T.: Temporal variations of the atmospheric nitrous oxide concentration and its $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ for the latter half of the 20th century reconstructed from firn air analyses, *J. Geophys. Res. Atmos.*, 112(3), doi:10.1029/2006JD007208, 2007.
- 810 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi, P., Pagliari, V., Olivier, J., Peters, J., van Aardenne, J., Monni, S., Doering, U., Petrescu, R., Solazzo, E. and Oreggioni, G.: EDGAR v4.3.2 Global Atlas of the three major Greenhouse Gas Emissions for the period 1970-2012, *Earth Syst. Sci. Data Discuss.*, 2010, 1–52, doi:10.5194/essd-2018-164, 2019.
- Jiang, X., Ku, W. L., Shia, R. L., Li, Q., Elkins, J. W., Prinn, R. G. and Yung, Y. L.: Seasonal cycle of N_2O : Analysis of data, *Global Biogeochem. Cycles*, 21, GB1006, doi:10.1029/2006GB002691, 2007.
- 815 JMA and WMO: World Meteorological Organization - Global Atmosphere Watch - World Data Centre for Greenhouse Gases, Data Summary, No. 42, 101 p., [online] Available from: <https://gaw.kishou.go.jp/static/publications/summary/sum42/sum42.pdf>, 2018.
- Kaiser, J., Röckmann, T. and Brenninkmeijer, C. A. M.: Complete and accurate mass spectrometric isotope analysis of tropospheric nitrous oxide, *J. Geophys. Res. Atmos.*, 108, 4476, doi:10.1029/2003JD003613, D15, 2003.
- 820 Kaiser, J., Engel, A., Borchers, R. and Röckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N_2O isotope distribution, *Atmos. Chem. Phys.*, 6(11), 3535–3556, doi:10.5194/acp-6-3535-2006, 2006.

- 825 Keller, C. A., Hill, M., Vollmer, M. K., Henne, S., Brunner, D., Reimann, S., O'Doherty, S., Arduini, J.,
Maione, M., Ferenczi, Z., Haszpra, L., Manning, A. J. and Peter, T.: European emissions of halogenated
greenhouse gases inferred from atmospheric measurements, *Environ. Sci. Technol.*, 46(1), 217–225,
doi:10.1021/es202453j, 2012.
- Kim, K.-R. and Craig, H.: Nitrogen-15 and Oxygen-18 Characteristics of Nitrous Oxide: A Global
Perspective, *Science*, 262, 1855-1857, 1993.
- 830 Lebegue, B., Schmidt, M., Ramonet, M., Wastine, B., Yver Kwok, C., Laurent, O., Belviso, S., Guemri,
A., Philippon, C., Smith, J. and Conil, S.: Comparison of nitrous oxide (N₂O) analyzers for high-precision
measurements of atmospheric mole fractions, *Atmos. Meas. Tech.*, 9(3), 1221–1238, doi:10.5194/amt-9-
1221-2016, 2016.
- 835 Logan, J. A., Staehelin, J., Megretskaia, I. A., Cammas, J. P., Thouret, V., Claude, H., De Backer, H.,
Steinbacher, M., Scheel, H. E., Stbi, R., Frhlich, M. and Derwent, R.: Changes in ozone over Europe:
Analysis of ozone measurements from sondes, regular aircraft (MOZAIC) and alpine surface sites, *J.
Geophys. Res. Atmos.*, 117(9), 1–23, doi:10.1029/2011JD016952, 2012.
- Mohn, J., Guggenheim, C., Tuzson, B., Vollmer, M. K., Toyoda, S., Yoshida, N. and Emmenegger, L.: A
liquid nitrogen-free preconcentration unit for measurements of ambient N₂O isotopomers by QCLAS,
Atmos. Meas. Tech., 3(3), 609–618, doi:10.5194/amt-3-609-2010, 2010.
- 840 Mohn, J., Tuzson, B., Manninen, A., Yoshida, N., Toyoda, S., Brand, W. A. and Emmenegger, L.: Site
selective real-time measurements of atmospheric N₂O isotopomers by laser spectroscopy, *Atmos. Meas.
Tech.*, 5(7), 1601–1609, doi:10.5194/amt-5-1601-2012, 2012.
- 845 Mohn, J., Wolf, B., Toyoda, S., Lin, C. T., Liang, M. C., Brüggemann, N., Wissel, H., Steiker, A. E.,
Dyckmans, J., Szwec, L., Ostrom, N. E., Casciotti, K. L., Forbes, M., Giesemann, A., Well, R., Doucett,
R. R., Yarnes, C. T., Ridley, A. R., Kaiser, J. and Yoshida, N.: Interlaboratory assessment of nitrous
oxide isotopomer analysis by isotope ratio mass spectrometry and laser spectroscopy: Current status and
perspectives, *Rapid Commun. Mass Spectrom.*, 28(18), 1995–2007, doi:10.1002/rcm.6982, 2014.
- 850 Nevison, C. D., Keeling, R. F., Weiss, R. F., Popp, B. N., Jin, X., Fraser, P. J., Porter, L. W. and Hess, P.
G.: Southern Ocean ventilation inferred from seasonal cycles of atmospheric N₂O and O₂/N₂ at Cape
Grim, Tasmania, *Tellus, Ser. B Chem. Phys. Meteorol.*, 57(3), 218–229, doi:10.1111/j.1600-
0889.2005.00143.x, 2005.
- 855 Nevison, C. D., Dlugokencky, E., Dutton, G., Elkins, J. W., Fraser, P., Hall, B., Krummel, P. B.,
Langenfelds, R. L., O'Doherty, S., Prinn, R. G., Steele, L. P. and Weiss, R. F.: Exploring causes of
interannual variability in the seasonal cycles of tropospheric nitrous oxide, *Atmos. Chem. Phys.*, 11(8),
3713–3730, doi:10.5194/acp-11-3713-2011, 2011.
- Ogawa, M. and Yoshida, N.: Intramolecular distribution of stable nitrogen and oxygen isotopes of nitrous
oxide emitted during coal combustion, *Chemosphere*, 61(6), 877–887,
doi:10.1016/j.chemosphere.2005.04.096, 2005.
- 860 Ostrom, N. E., Gandhi, H., Coplen, T. B., Toyoda, S., Böhlke, J. K., Brand, W. A., Casciotti, K. L.,
Dyckmans, J., Giesemann, A., Mohn, J., Well, R., Yu, L. and Yoshida, N.: Preliminary assessment of
stable nitrogen and oxygen isotopic composition of USGS51 and USGS52 nitrous oxide reference gases
and perspectives on calibration needs, *Rapid Commun. Mass Spectrom.*, 32(15), 1207–1214,
doi:10.1002/rcm.8157, 2018.
- 865 Pandey Deolal, S., Brunner, D., Steinbacher, M., Weers, U. and Staehelin, J.: Long-term in situ
measurements of NO_x and NO_y at Jungfraujoch 1998-2009: Time series analysis and evaluation, *Atmos.
Chem. Phys.*, 12(5), 2551–2566, doi:10.5194/acp-12-2551-2012, 2012.
- Park, S., Croteau, P., Boering, K. A., Etheridge, D. M., Ferretti, D., Fraser, P. J., Kim, K.-R., Krummel, P.

- B., Langenfelds, R. L., van Ommen, T. D., Steele, L. P. and Trudinger, C. M.: Trends and seasonal cycles in the isotopic composition of nitrous oxide since 1940, *Nat. Geosci.*, 5(4), 261–265, doi:10.1038/ngeo1421, 2012.
- 870 Pérez, T., Trumbore, S. E., Tyler, S. C., Matson, P. A., I., O.-M., Rahn, T. and Griffiths, D. W. T.: Identifying the agricultural imprint on the global N₂O budget using stable isotopes, *J. Geophys. Res.*, 106, 9869–9878, doi:10.1179/1607845413y.0000000087, 2001.
- 875 Prather, M. J., Hsu, J., Deluca, N. M., Jackman, C. H., Oman, L. D., Douglass, A. R., Fleming, E. L., Strahan, S. E., Steenrod, S. D., Søvdé, O. A., Isaksen, I. S. A., Froidevaux, L. and Funke, B.: Measuring and modeling the lifetime of nitrous oxide including its variability Michael, *J. Geophys. Res. Atmos.*, 120, 5693–5705, doi:10.1002/2015JD023267. Received, 2015.
- 880 Prinn, R. G., Weiss, R. F., Arduini, J., Arnold, T., Dewitt, H. L., Fraser, P. J., Ganesan, A. L., Gasore, J., Harth, C. M., Hermansen, O., Kim, J., Krummel, P. B., Li, S., Loh, Z. M., Lunder, C. R. and Maione, M.: History of chemically and radiatively important atmospheric gases from the Advanced Global Atmospheric Gases Experiment (AGAGE), *Earth Syst. Sci. Data*, 10, 985–1018, 2018.
- 885 Prokopiou, M., Martinerie, P., Sapart, C. J., Witrant, E., Monteil, G. A., Ishijima, K., Bernard, S., Kaiser, J., Levin, I., Sowers, T., Blunier, T., Etheridge, D., Dlugokencky, E., van de Wal, R. S. W. and Röckmann, T.: Constraining N₂O emissions since 1940 using firn air isotope measurements in both hemispheres, *Atmos. Chem. Phys.*, 2011(June), 1–50, doi:10.5194/acp-2016-487, 2017.
- Prokopiou, M., Sapart, C. J., Rosen, J., Sperlich, P., Blunier, T., Brook, E., van de Wal, R. S. W. and Röckmann, T.: Changes in the Isotopic Signature of Atmospheric Nitrous Oxide and Its Global Average Source During the Last Three Millennia, *J. Geophys. Res. Atmos.*, 1–17, doi:10.1029/2018JD029008, 2018.
- 890 Rahn, T. and Wahlen, M.: A reassessment of the global isotopic budget of atmospheric nitrous oxide, *Global Biogeochem. Cycles*, 14(2), 537–543, doi:10.1029/1999GB900070, 2000.
- Ravishankara, A. R., Daniel, J. S. and Portmann, R. W.: Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st century., *Science*, 326(5949), 123–5, doi:10.1126/science.1176985, 2009.
- 895 Reay, D. S., Davidson, E. a., Smith, K. a., Smith, P., Melillo, J. M., Dentener, F. and Crutzen, P. J.: Global agriculture and nitrous oxide emissions, *Nat. Clim. Chang.*, 2(6), 410–416, doi:10.1038/nclimate1458, 2012.
- 900 Reimann, S., Vollmer, M. K., Folini, D., Steinbacher, M., Hill, M., Buchmann, B., Zander, R. and Mahieu, E.: Observations of long-lived anthropogenic halocarbons at the high-Alpine site of Jungfraujoch (Switzerland) for assessment of trends and European sources, *Sci. Total Environ.*, 391(2–3), 224–231, doi:10.1016/j.scitotenv.2007.10.022, 2008.
- 905 Rigby, M., Prinn, R. G., O’Doherty, S., Montzka, S. A., McCulloch, A., Harth, C. M., Mühle, J., Salameh, P. K., Weiss, R. F., Young, D., Simmonds, P. G., Hall, B. D., Dutton, G. S., Nance, D., Mondeel, D. J., Elkins, J. W., Krummel, P. B., Steele, L. P. and Fraser, P. J.: Re-evaluation of the lifetimes of the major CFCs and CH₃CCl₃ using atmospheric trends, *Atmos. Chem. Phys.*, 13(5), 2691–2702, doi:10.5194/acp-13-2691-2013, 2013.
- Röckmann, T., Kaiser, J. and Brenninkmeijer, C. A. M.: The isotopic fingerprint of the pre-industrial and the anthropogenic N₂O source, *Atmos. Chem. Phys.*, (3), 315–323, 2003.
- 910 Röckmann, T. and Levin, I.: High-precision determination of the changing isotopic composition of atmospheric N₂O from 1990 to 2002, *J. Geophys. Res. Atmos.*, 110(21), 1–8, doi:10.1029/2005JD006066, 2005.
- Saikawa, E., Prinn, R. G., Dlugokencky, E., Ishijima, K., Dutton, G. S., Hall, B. D., Langenfelds, R.,

- 915 Tohjima, Y., Machida, T., Manizza, M., Rigby, M., O'Doherty, S., Patra, P. K., Harth, C. M., Weiss, R. F., Krummel, P. B., Van Der Schoot, M., Fraser, P. J., Steele, L. P., Aoki, S., Nakazawa, T. and Elkins, J. W.: Global and regional emissions estimates for N₂O, *Atmos. Chem. Phys.*, 14(9), 4617–4641, doi:10.5194/acp-14-4617-2014, 2014.
- Schibig, M. F., Steinbacher, M., Buchmann, B., Van Der Laan-Luijkx, I. T., Van Der Laan, S., Ranjan, S. and Leuenberger, M. C.: Comparison of continuous in situ CO₂ observations at Jungfraujoch using two different measurement techniques, *Atmos. Meas. Tech.*, 8(1), 57–68, doi:10.5194/amt-8-57-2015, 2015.
- 920 Schilt, A., Brook, E. J., Bauska, T. K., Bagginstos, D., Fischer, H., Joos, F., Petrenko, V. V., Schaefer, H., Schmitt, J., Severinghaus, J. P., Spahni, R. and Stocker, T. F.: Isotopic constraints on marine and terrestrial N₂O emissions during the last deglaciation, *Nature*, 516(7530), 234–237, doi:10.1038/nature13971, 2014.
- 925 Sepúlveda, E., Schneider, M., Hase, F., Barthlott, S., Dubravica, D., García, O. E., Gomez-Pelaez, A., González, Y., Guerra, J.C., Gisi, M., Kohlhepp, R., Dohe, S., Blumenstock, T., Strong, K., Weaver, D., Palm, N., Sadeghi, A., Deutscher, N. M., Warneke, T., Notholt, J., Jones, N., Griffith, D. W. T., Smale, D., Brailsford, G. W., Robinson, J., Meinhardt, F., Steinbacher, M., Aalto, T. and Worthy, D.: Tropospheric CH₄ signals as observed by NDACC FTIR at globally distributed sites and comparison to GAW surface in situ measurements, *Atmos. Meas. Tech.*, 7(7), 2337–2360, doi:10.5194/amt-7-2337-2014, 2014.
- 930 Snider, D., Thompson, K., Wagner-Riddle, C., Spoelstra, J. and Dunfield, K.: Molecular techniques and stable isotope ratios at natural abundance give complementary inferences about N₂O production pathways in an agricultural soil following a rainfall event, *Soil Biol. Biochem.*, 88, 197–213, doi:10.1016/j.soilbio.2015.05.021, 2015a.
- 935 Snider, D. M., Venkiteswaran, J. J., Schiff, S. L. and Spoelstra, J.: From the ground up: Global nitrous oxide sources are constrained by stable isotope values, *PLoS One*, 10(3), 1–19, doi:10.1371/journal.pone.0118954, 2015b.
- 940 Sowers, T., Rodebaugh, A., Yoshida, N. and Toyoda, S.: Extending records of the isotopic composition of atmospheric N₂O back to 1800 A.D. from air trapped in snow at the South Pole and the Greenland Ice Sheet Project II ice core, *Global Biogeochem. Cycles*, 16(4), 1–10, doi:10.1029/2002GB001911, 2002.
- Stohl, A., Forster, C., Frank, A., Seibert, P. and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, doi:10.3390/atmos9020076, 2005.
- 945 Sturm, P., Tuzson, B., Henne, S. and Emmenegger, L.: Tracking isotopic signatures of CO₂ at the high altitude site Jungfraujoch with laser spectroscopy: Analytical improvements and representative results, *Atmos. Meas. Tech.*, 6(7), 1659–1671, doi:10.5194/amt-6-1659-2013, 2013.
- 950 Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Breznak, J. a, Pitt, a J., Li, F. and Gandhi, H.: Distinguishing Nitrous Oxide Production from Nitrification and Denitrification on the Basis of Isotopomer Abundances, *Appl. Environ. Microbiol.*, 72(1), 638–644, doi:10.1128/AEM.72.1.638, 2006.
- Tarasova, O. A., Senik, I. A., Sosonkin, M. G., Cui, J., Staehelin, J. and Péevot, A. S. H.: Surface ozone at the Caucasian site Kislovodsk High Mountain Station and the Swiss Alpine site Jungfraujoch: Data analysis and trends (1990-2006), *Atmos. Chem. Phys.*, 9(12), 4157–4175, doi:10.5194/acp-9-4157-2009, 2009.
- 955 Team, R. C.: A language and environment for statistical computing. R Foundation for statistical computing, 2015; Vienna, Austria, 2016.
- Thompson, R. L., Patra, P. K., Ishijima, K., Saikawa, E., Corazza, M., Karstens, U., Wilson, C.,

- 960 Bergamaschi, P., Dlugokencky, E., Sweeney, C., Prinn, R. G., Weiss, R. F., O'Doherty, S., Fraser, P. J., Steele, L. P., Krummel, P. B., Saunio, M., Chipperfield, M. and Bousquet, P.: TransCom N₂O model inter-comparison-Part 1: Assessing the influence of transport and surface fluxes on tropospheric N₂O variability, *Atmos. Chem. Phys.*, 14(8), 4349–4368, doi:10.5194/acp-14-4349-2014, 2014a.
- 965 Thompson, R. L., Ishijima, K., Saikawa, E., Corazza, M., Karstens, U., Patra, P. K., Bergamaschi, P., Chevallier, F., Dlugokencky, E., Prinn, R. G., Weiss, R. F., O'Doherty, S., Fraser, P. J., Steele, L. P., Krummel, P. B., Vermeulen, A., Tohjima, Y., Jordan, A., Haszpra, L., Steinbacher, M., Van Der Laan, S., Aalto, T., Meinhardt, F., Poppo, M. E., Moncrieff, J. and Bousquet, P.: TransCom N₂O model inter-comparison - Part 2: Atmospheric inversion estimates of N₂O emissions, *Atmos. Chem. Phys.*, 14(12), 6177–6194, doi:10.5194/acp-14-6177-2014, 2014b.
- 970 Thoning, K. W., Tans, P. P. and Komhyr, W. D.: Atmospheric carbon dioxide at Mauna Loa Observatory: 2. Analysis of the NOAA GMCC data, 1974-1985, *J. Geophys. Res.*, 94(D6), 8549–8565, doi:10.1029/jd094id06p08549, 1989.
- 975 Tian, H., Yang, J., Xu, R., Lu, C., Canadell, J., Davidson, E. A., Jackson, R., Arneeth, A., Chang, J., Ciais, P., Gerber, S., Ito, A., Joos, F., Lienert, S., Messina, P., Olin, S., Pan, S., Peng, C., Saikawa, E., Thompson, R., Vuichard, N., Winiwarter, W., Zaehle, S. and Zhang, B.: Global soil nitrous oxide emissions since the preindustrial era estimated by an ensemble of terrestrial biosphere models: Magnitude, attribution, and uncertainty, *Glob. Chang. Biol.*, 25(2), 640–659, doi:10.1111/gcb.14514, 2019.
- Toyoda, S., Yoshida, N., Urabe, T., Aoki, S., Nakazawa, T., Sugawara, S. and Honda, H.: Fractionation of N₂O isotopomers in the stratosphere, *J. Geophys. Res.*, 106(D7), 7515, doi:10.1029/2000JD900680, 2001.
- 980 Toyoda, S., Yamamoto, S., Arai, S., Nara, H., Yoshida, N., Kashiwakura, K. and Akiyama, K.: Isotopomeric characterization of N₂O produced, consumed, and emitted by automobiles, *rapid comm*, 22, 603–612, doi:10.1002/rcm, 2008.
- 985 Toyoda, S., Yano, M., Nishimura, S., Akiyama, H., Hayakawa, A., Koba, K., Sudo, S., Yagi, K., Makabe, A., Tobari, Y., Ogawa, N. O., Ohkouchi, N., Yamada, K. and Yoshida, N.: Characterization and production and consumption processes of N₂O emitted from temperate agricultural soils determined via isotopomer ratio analysis, *Global Biogeochem. Cycles*, 25(2), 1–17, doi:10.1029/2009GB003769, 2011.
- 990 Toyoda, S., Kuroki, N., Yoshida, N., Ishijima, K., Tohjima, Y. and Machida, T.: Decadal time series of tropospheric abundance of N₂O isotopomers and isotopologues in the Northern Hemisphere obtained by the long-term observation at Hateruma Island, Japan, *J. Geophys. Res. Atmos.*, 118(8), 3369–3381, doi:10.1002/jgrd.50221, 2013.
- Toyoda, S., Yoshida, N. and Koba, K.: Isotopocule analysis of biologically produced nitrous oxide in various environments, *Mass Spectrom. Rev.*, (36), 135–160, doi:doi:10.1002/mas.21459, 2017.
- 995 Toyoda, S., Yoshida, N., Morimoto, S., Aoki, S., Nakazawa, T., Sugawara, S., Ishidoya, S., Uematsu, M., Inai, Y., Hasebe, F., Ikeda, C., Honda, H. and Ishijima, K.: Vertical distributions of N₂O isotopocules in the equatorial stratosphere, *Atmos. Chem. Phys.*, (18), 833–844, 2018.
- Tuzson, B., Henne, S., Brunner, D., Steinbacher, M., Mohn, J., Buchmann, B. and Emmenegger, L.: Continuous isotopic composition measurements of tropospheric CO₂ at Jungfraujoch (3580 m a.s.l.), Switzerland: Real-time observation of regional pollution events, *Atmos. Chem. Phys.*, 11(4), 1685–1696, doi:10.5194/acp-11-1685-2011, 2011.
- 1000 WMO: WMO Greenhouse Gas Bulletin., 2018.
- Wolf, B., Merbold, L., Decock, C., Tuzson, B., Harris, E., Six, J., Emmenegger, L. and Mohn, J.: First on-line isotopic characterization of N₂O above intensively managed grassland, *Biogeosciences*, 12(8),

2517–2531, doi:10.5194/bg-12-2517-2015, 2015.

1005 Xu-Ri, Prentice, I. C., Spahni, R. and Niu, H. S.: Modelling terrestrial nitrous oxide emissions and implications for climate feedback, *New Phytol.*, 196(2), 472–488, doi:10.1111/j.1469-8137.2012.04269.x, 2012.

Yoshida, N. and Toyoda, S.: Constraining the atmospheric N₂O budget from intramolecular site preference in N₂O isotopomers, *Nature*, 405(6784), 330–4, doi:10.1038/35012558, 2000.

1010 Yuan, Y., Ries, L., Petermeier, H., Steinbacher, M., Gómez-Pelaéz, A. J., Leuenberger, M. C., Schumacher, M., Trickl, T., Couret, C., Meinhardt, F. and Menzel, A.: Adaptive selection of diurnal minimum variation: A statistical strategy to obtain representative atmospheric CO₂ data and its application to European elevated mountain stations, *Atmos. Meas. Tech.*, 11(3), 1501–1514, doi:10.5194/amt-11-1501-2018, 2018.

1015 Yung, Y. L. and Miller, C. E.: Isotopic fractionation of stratospheric nitrous oxide, *Science*, 278(5344), 1778–1780, doi:10.1126/science.278.5344.1778, 1997.

Zellweger, C., Forrer, J., Hofer, P., Nyeki, S., Schwarzenbach, B., Weingartner, E., Ammann, M. and Baltensperger, U.: Partitioning of reactive nitrogen (NO_y) and dependence on meteorological conditions in the lower free troposphere, *Atmos. Chem. Phys.*, 3(3), 779–796, doi:10.5194/acp-3-779-2003, 2003.

1020 Zellweger, C., Hüglin, C., Klausen, J., Steinbacher, M., Vollmer, M. and Buchmann, B.: Inter-comparison of four different carbon monoxide measurement techniques and evaluation of the long-term carbon monoxide time series of Jungfraujoch, *Atmos. Chem. Phys.*, 9(11), 3491–3503, doi:10.5194/acp-9-3491-2009, 2009.

1025

Table 1 Trends of atmospheric $\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ at Jungfraujoch determined using discrete measurements between April 2014 and December 2018. The trends are determined for the whole dataset, the dataset filtered for free troposphere (removing data points with significant influence from plenary boundary layer) and the second-phase dataset with bi-weekly measurements (August 2016 to December 2018).

	$\delta^{15}\text{N}^{\text{bulk}}$ (‰ a^{-1})		$\delta^{15}\text{N}^{\text{SP}}$ (‰ a^{-1})		$\delta^{18}\text{O}$ (‰ a^{-1})	
	Raw	Deseasonalized	Raw	Deseasonalized	Raw	Deseasonalized
Whole dataset	$-0.059 \pm 0.012^*$	$-0.052 \pm 0.012^*$	0.069 ± 0.029	0.065 ± 0.027	0.020 ± 0.011	0.019 ± 0.011
Free troposphere	$-0.060 \pm 0.014^*$	$-0.054 \pm 0.013^*$	0.054 ± 0.034	0.036 ± 0.030	0.024 ± 0.013	0.019 ± 0.011
First phase (Apr. 2014-Feb. 2016)	-0.036 ± 0.038	-0.041 ± 0.035	$0.449 \pm 0.100^*$	$0.314 \pm 0.082^*$	$0.238 \pm 0.029^*$	$0.207 \pm 0.026^*$
Second phase (Aug. 2016-Dec. 2018)	-0.105 ± 0.049	$-0.130 \pm 0.045^*$	0.028 ± 0.067	-0.007 ± 0.066	-0.007 ± 0.042	-0.001 ± 0.040

* Indicate significance of linear regression.

Table 2 Results of the two-box model simulations and selected literature values for comparison.

	This study		RMSE ^o	Sowers et al. (2002) ^a	Röckmann et al. (2003) ^b	Ishijima et al. (2007) ^c	Toyoda et al. (2013) ^d	Park et al. (2012) ^e	Prokopiou et al. (2017) ^f
Air sample age	NH [†] 2014-2018	NH(FT ⁿ) 2014-2018		FA [†] , IC [†] 1785-1995	FA NA	FA 1960-2001	NH 1999-2010	SH [†] , FA 1940-2005	FA 1940-2008
α^*	0.0154±0.004	0.0154±0.004	0.65 nmol mol ⁻¹	0.0111 to 0.0128	NA	NA	NA	NA	NA
$F_{\text{anth},2018}$ (Tg N a ⁻¹)	8.6±0.6	8.5±0.6	NA [†]	4.2 to 5.7	6.9	NA	5.5	6.6	5.4±1.7
$\delta^{15}\text{N}^{\text{bulk-}}_{\text{anth}}$ (‰)	-8.6±4	-8.5±4	0.23	-7 to -13	-11.4	-11.6	-9.84	-15.6±1.2	-18.2±2.6
$\delta^{18}\text{O-}^{\text{anth}}$ (‰)	34.8±3	34.3±3	0.22	17 to 26	31.7	NA	35.95	32.0±1.3	27.2±2.6
$\delta^{15}\text{N}^{\text{SP-}}_{\text{anth}}$ (‰)	10.7±4	10.7±4	0.50	NA	11.3	NA	8.52	13.1±9.4	18.0±8.6

1030 [†] NH and SH: surface atmosphere from the Northern and Southern Hemisphere, respectively; FA: firm air; IC: ice core air; NA: not available.

ⁿFT: Jungfraujoch dataset filtered for free troposphere (based on NO_y:CO).

* “Value” is the dimensionless constant α describing the exponential increase in the anthropogenic flux

1035 ^o RMSE refers to root mean square error. It is in nmol mol⁻¹ for α , referring to the present day tropospheric mixing ratio for N₂O. For source isotopic values, RMSE is in the unit of ‰. Simulations with the whole dataset and the dataset filtered for free troposphere yielded the same RMSE.

^a Estimates are for 1995.

^b Estimates are for 1998; isotopic signatures of anthropogenic sources were calculated assuming modern tropospheric values to be the same as this study.

1040 ^c Estimate is for 2000, for $\delta^{18}\text{O}$ calibration is not comparable.

^d Estimates are for 2012 using the “Base” scenario.

^e Estimates are for 2005.


^f δ_{anth} values are averaged values for the period of 1940-2008.

Table 3 Isotopic signatures for the overall, anthropogenic and major N₂O sources contributing to N₂O variations at Jungfrauoch. Source signatures were estimated based on a “bottom-up” approach, with literature-derived isotopic signatures and fluxes for variable sources under the Swiss Meteotest emission inventory.

	Emission inventory (%)	$\delta^{15}\text{N}^{\text{bulk}}$ (‰)	$\delta^{15}\text{N}^{\text{SP}}$ (‰)	$\delta^{18}\text{O}$ (‰)	Reference
Overall source	100	-15.8 (6.2)	7.3 (3.9)	29.4 (5.5)	-
Anthropogenic source	89.4	-15.6 (6.3)	7.4 (4.0)	29.5 (5.7)	-
Agricultural emission	61.5	-17.8 (5.7)	7.2 (3.8)	29.0 (3.7)	Snider et al. (2015) Wolf et al. (2015)
Manure management	7.4	-17.5 (6.2)	6.5 (4.1)	23.9 (3.8)	Maeda et al. (2010)
Waste *	7.2	-11.5 (12.6)	10.4 (5.7)	31.3 (14.0)	Ogawa and Yoshida (2005) Snider et al. (2015)
Natural emission	10.9	-17.8 (5.7)	7.2 (3.8)	29.0 (3.7)	Snider et al. (2015) Wolf et al. (2015)

* “Waste” sources consist of both wastewater treatment and agricultural waste burning (biomass burning).

Figures



1050

Figure 1a *In situ* (10-min averages) and discrete measurements of N₂O mixing ratios from April 2014 to December 2018 at Jungfraujoch. *In situ* N₂O mixing ratio measurements were performed with GC-ECD method between April and December 2014. After that, OA-ICOS became the major analytical method for *in situ* measurements. Discrete sample points are presented as averages with error bars (one standard deviation). Annual N₂O growth rates determined by linear regression are given in the figure (uncertainty shown as one standard deviation). A sampling gap exists for discrete samples between February and August 2016.

1055

1b Seasonality of N₂O mixing ratios at Jungfraujoch derived from *in situ* measurements. Datasets with/without GC-ECD measurements are compared for seasonality evaluation. The NLS model simulation for time-series gives the detrended seasonality, with error bars indicating one standard deviation of monthly residuals.

1060




Figure 2 Time-series of isotopic composition of atmospheric N₂O observed at Jungfraujoch from April 2014 to December 2018. Error bars indicate one standard deviation of repeated measurements. Blue lines indicate the simulated trends by the NLS model.

1065






Figure 3 Seasonality of isotopic signatures of atmospheric N₂O observed at Jungfraujoch. Top panels: seasonality obtained using the whole dataset from April 2014 to December 2018; lower panels: seasonality obtained using bi-weekly data collected between August 2016 and December 2018. Red dashed lines refer to zero variability. The NLS model simulation for time-series gives the detrended seasonality, with error bars indicating one standard deviation of monthly residuals.

1070






1075 **Figure 4a** Clusters of air mass transport regimes for Jungfraujoch shown as normalized surface source sensitivities over our sampling period. Cluster abbreviations refer to Free Troposphere (FT), Southwest (SW), East (E), Local (L), West (W) and Northwest (NW). The normalization was done by calculating the difference between cluster average source sensitivity and whole period average source sensitivities, divided by the period average. Orange colors indicate the main source regimes in each cluster, whereas blue colors indicate little to no influence on Jungfraujoch observations. The free tropospheric cluster showed lower than average surface sensitivity everywhere.

1080 **4b** Cluster frequency of air mass transport regimes (%) shown as a monthly pattern (left) and interannual patterns for the whole periods (middle) and for the periods of discrete sampling (right). Numbers above the right figure indicate the total number of discrete samples per year.



1085 Figure 5 Comparison of N_2O mixing ratios and isotopic signatures (with linear trends removed) for the six air mass footprint clusters used in the present study. Green and blue stars indicate significantly larger and smaller values than the others, respectively; red dashed lines indicate mean levels; grey points indicate outliers.




1090

Figure 6a Mean monthly stacked-bar plots of source contributions (%) to atmospheric N₂O at Jungfraujoch derived from inversion modeling.

6b Overall contributions of N₂O sources responsible for emission to Jungfraujoch.

1095

6c Simulated 3-hourly N₂O mixing ratios, N₂O mixing ratio baseline and N₂O enhancements in nmol mol⁻¹.



1100 **Figure 7** Two-box model results showing the influence of anthropogenic emissions on N₂O mixing ratio and isotopic composition in the troposphere. Left: full time range from the start of the anthropogenic period (1845) to present day; Right: zoom to the last two decades. Isotopic measurements at Jungfraujoch were used as the only constraint of current tropospheric N₂O isotopic composition for the model. See the materials and method as well as the SI for more details and other input parameters. Atmospheric as well as firm air measurements of $\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\text{SP}}$ and $\delta^{18}\text{O}$ from the literature are presented for comparison. Blue shaded areas indicate one standard deviation of the model iterations.