

Interactive comment on “Protein aggregates nucleate ice: the example of apoferritin” by María Cascajo-Castresana et al.

Janine Froehlich

j.frohlich@mpic.de

Received and published: 4 December 2019

The statement in the introduction (page 2, line 39) "In screening experiments, fungi and lichen failed to show IN activity above -25°C , with the exception of *Fusarium acuminatum* and *Fusarium avenaceum* (Pouleur et al., 1992; Pummer et al., 2013)." is not supported by the literature since several studies found IN activity in other fungal species as well as lichen (e.g., Fröhlich-Nowoisky et al., 2015; Haga et al., 2013; 2014; Huffmann et al., 2013; Kieft 1988; Kieft and Ahmadjian, 1989; Kieft and Ruscetti, 1990; Moffet et al., 2015; Morris et al., 2013).

Moreover, the authors cite Fröhlich-Nowoisky et al., 2015 (page 3, line 1) to support their statement "Surveys of the IN ability of pollen showed that only few types were

Printer-friendly version

Discussion paper



active, the most active ones stemming from birch and conifer trees, yet, only at temperatures below -9°C ".

This should be corrected as Fröhlich-Nowoisky et al., 2015 performed a screening of soil fungi and found ice nucleation activity in the widespread soil fungus *Mortierella alpina*.

References:

Fröhlich-Nowoisky, J., Hill, T. C. J., Pummer, B. G., Yordanova, P., Franc, G. D., and Pöschl, U.: Ice nucleation activity in the widespread soil fungus *Mortierella alpina*, *Biogeosciences*, 12, 1057–1071, doi:10.5194/bg-12-1057-2015, 2015.

Fröhlich-Nowoisky, J., Hill, T. C. J., Pummer, B. G., Yordanova, P., Franc, G. D., and Pöschl, U.: Ice nucleation activity in the widespread soil fungus *Mortierella alpina*, *Biogeosciences*, 12, 1057–1071, doi:10.5194/bg-12-1057-2015, 2015.

Haga, D.I., Iannone, R., Wheeler, M.J., Mason, R., Polishchuk, E.A., Fetch, T., van der Kamp, B.J., McKendry, I.G., Bertram, A.K.: Ice nucleation properties of rust and bunt fungal spores and their transport to high altitudes, where they can cause heterogeneous freezing. *J. Geophys. Res. Atmos.* 118, 7260–7272. <http://dx.doi.org/10.1002/jgrd.50556>, 2013.

Haga, D. I., Burrows, S. M., Iannone, R., Wheeler, M. J., Mason, R. H., Chen, J., Polishchuk, E. A., Pöschl, U., and Bertram, A. K.: Ice nucleation by fungal spores from the classes Agaricomycetes, Ustilaginomycetes, and Eurotiomycetes, and the effect on the atmospheric transport of these spores, *Atmos. Chem. Phys.*, 14, 8611–8630, doi:10.5194/acp-14-8611-2014, 2014.

Huffman, J. A., Prenni, A. J., DeMott, P. J., Pöhlker, C., Mason, R. H., Robinson, N. H., Fröhlich-Nowoisky, J., Tobo, Y., Després, V. R., Garcia, E., Gochis, D. J., Harris, E., Müller-Germann, I., Ruzene, C., Schmer, B., Sinha, B., Day, D. A., Andreae, M. O., Jimenez, J. L., Gallagher, M., Kreidenweis, S. M., Bertram, A. K., and Pöschl, U.:

High concentrations of biological aerosol particles and ice nuclei during and after rain, *Atmos. Chem. Phys.*, 13, 6151–6164, doi:10.5194/acp13-6151-2013, 2013.

Kieft, T. L.: Ice nucleation activity in lichens, *Appl. Environ. Microbiol.*, 54, 1678–1681, 1988.

Kieft, T. L. and Ahmadjian, V.: Biological ice nucleation activity in lichen mycobionts and photobionts, *Lichenol.*, 21, 355–362, 1989.

Kieft, T. L. and Ruscetti, T.: Characterization of biological ice nuclei from a lichen, *J. Bacteriol.*, 172, 3519–3523, 1990.

Moffett, B.F., Getti, G., Hill, T.C.J.: Ubiquity of ice nucleation in lichen – possible atmospheric implications. *Lindbergia* 38, 39–43, 2015.

Morris, C. E., D. C. Sands, C. Glaux, J. Samsatly, S. Asaad, A. R. Moukahel, F. L. T. Gonçalves, and E. K. Bigg: Urediospores of rust fungi are ice nucleation active at > -10 °C and harbor ice nucleation active bacteria. *Atmos. Chem. Phys.* 13:4223–4233. <http://www.atmos-chem-phys.net/13/4223/2013/acp-13-4223-2013>, 2013.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-861>, 2019.

Printer-friendly version

Discussion paper

