

Responses to the Reviewer's Comments

We appreciate the reviewer for the thorough reading and thoughtful comments and suggestions, which greatly improve the quality of the manuscript. We revised the MS accordingly. The point-to-point responses to all the comments are given below in blue.

Suggestions/comments:

1. Title: the title could be modified a bit. What does “molecular distribution” mean? “over the South China Sea and East Indian Ocean” could be changed to “from the South China Sea to the East Indian Ocean” .

Response: Molecular distribution means the distribution characteristics of the concentrations of various organic acids with different carbon numbers at different time and places. And, following the reviewer's suggestion, we have changed the title to “Molecular and spatial distributions of dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls in marine aerosols from the South China Sea to East Indian Ocean”.

2. Table 3 could be moved to the supporting information.

Response: In the revision, we have moved Table 3 to the supporting information (SI).

3. The concentrations in the SLDP should be taken for further consideration. As mentioned in the sampling part, these samples mainly show the severe influence of ship exhaust. They are not ambient aerosol samples if compared to other samples.

Response: The aerosol samples of SLDP were affected by multiple sources, namely local coastal and terrestrial biological sources and anthropogenic emissions (especially fossil fuel combustion). In addition, the 5-day backward trajectory showed that the air mass originated from the land of Sri Lanka and the surrounding sea area, which indicated that the samples were not only polluted by ship exhaust. Thus, the SLDP samples were ambient aerosol samples. The reasons for the analysis of SLDP samples in this paper are as follows: firstly, it can be roughly compared with aerosol samples derived from SCS and EIO; secondly, it is meaningful and interesting to study the samples affected by ship exhaust pollution along the coast. Therefore, it is necessary to study and analyze SLDP samples in this paper.

4. The comparison should be more logical in the section 3.1 (L156-173), 3.2 (L217-232) and other parts. It is not easy to follow these parts. For what reason the authors would like to make the comparison if it is really necessary. E.g. The concentrations of OC and EC in the SCS could

be further compared with many other recent observations in this area. Why compared with those from Okinawa (P160-166)?

Response: In the section 3.1 (L156-173), we have compared the results of this study with those of other sea areas that were more affected by terrestrial long-distance transmission and those that were less affected by terrestrial anthropogenic emissions. We have added comparisons with some other recent observations in this area and its vicinity. The revised content is as follows.

“The mass concentrations of OC and EC in SCS were lower than those in the western South China Sea (OC: $2.04 \mu\text{g m}^{-3}$, EC: $3.0 \mu\text{g m}^{-3}$) which were strongly affected by biomass burning (Song et al., 2018). In addition, the carbon component concentration levels in Malacca were similar to those in the Bay of Bengal (OC: $1.9 \mu\text{g m}^{-3}$, EC: $0.4 \mu\text{g m}^{-3}$) (Kumar et al., 2008).” (see Page 6, Line 165–168).

“The concentration levels of carbon components in EIO-WI and EIO-SL were lower than those in the Indian Ocean (OC: $0.71 \mu\text{g m}^{-3}$, EC: $0.49 \mu\text{g m}^{-3}$) (Fu et al., 2011).” (see Page 7, Line 175–176).

In the section 3.2 (L217–232), we want to compare the results of this study with other marine regions, polar regions, and urban regions. In order to make the comparison more logical, the contents are modified as follows.

“Table 1 shows the total concentrations of diacids in aerosol samples from five sea areas in this study (SCS: $0.27 \pm 0.15 \text{ ng m}^{-3}$; EIO-WI: $0.05 \pm 0.05 \text{ ng m}^{-3}$; EIO-SL: $0.04 \pm 0.18 \text{ ng m}^{-3}$; Malacca: $0.24 \pm 0.18 \text{ ng m}^{-3}$; and SLDP: $0.27 \pm 0.15 \text{ ng m}^{-3}$). Fu et al. (2013) reported that the mean total diacids concentration of aerosol samples in the South China Sea and the Indian Ocean with the sampling period of three days was 489 ng m^{-3} and 301 ng m^{-3} , respectively. Comparing this study with aerosols from other marine areas, it was found that the total diacids concentration of aerosol samples in the EIO was similar to that in the western Pacific Ocean ($60 \pm 39 \text{ ng m}^{-3}$) (Wang et al., 2006). What’s more, the concentration of total diacids in aerosol samples from five sea areas was all lower than that of the East China Sea (850 ng m^{-3}) (Mochida et al., 2003), and higher than that of the Southern Ocean ($4.5 \pm 4.0 \text{ ng m}^{-3}$) (Wang et al., 2006). In addition, the concentration of total diacids in aerosol samples in all sea areas was higher than that in the Alert, Arctic (25 ng m^{-3}) (Kawamura et al., 1996) in the polar region, and lower than that in the urban center of Xi'an, China in different dust events ($932\text{--}2240 \text{ ng m}^{-3}$) (Wang et al., 2015), the megacity Chennai, India (694.5 ± 176.3 and $640.6 \pm 150.6 \text{ ng m}^{-3}$ in early and late winter) (Pavuluri et al., 2010), and the megacity Chennai, India in summer ($502.9 \pm 117.9 \text{ ng m}^{-3}$) (Pavuluri et al., 2010) and other urban areas.” (see Page 8, Line 220–232).

5. L216: what do the similar formation processes mean? The sources of diacids in the coastal and in the remote regions could be very different if you read through these and other related references.

Response: The revised content is as follows.

“The dominance of C₂ followed by C₃ and C₄ diacids is consistent with those coastal marine aerosols (Kundu et al., 2010;Kunwar and Kawamura, 2014) and remote marine aerosols from the western North Pacific (Bikkina et al., 2015), suggesting similar formation processes of dicarboxylic acids in the atmosphere of the coastal or remote regions.” (see Page 8, Line 216–219)

6. L286-287: Again, why compare with the concentrations in urban Tokyo?

Response: The revised content is as follows.

“Concentrations of Ph acid in EIO-WI and EIO-SL were similar to those of the western North Pacific (MBA: 0.6 ng m⁻³; LBA: 0.5 ng m⁻³) (Bikkina et al., 2015), where were marine-derived contributions. In addition, concentrations of Ph acid over the SCS to EIO were much lower than those reported in the urban atmosphere such as Chennai (21 ng m⁻³) (Pavuluri et al., 2010) and Chinese cities (90 ng m⁻³) (Ho et al., 2007).” (see Page 10, Line 287–291)

7. L291-294: How about the influence of ship exhaust on the increase of Ph?

Response: Ship exhaust can increase the atmospheric level of phthalic acid (Ph). Phthalic acid is either formed via photochemical pathways of naphthalene or directly released into air by fossil fuel burning and the incomplete combustion of aromatic hydrocarbons in motor vehicles. Moreover, the abundance of Ph may also be caused by increased phthalates emissions from plastic waste burnings in heavily polluted areas in China (Deshmukh et al., 2016). In this study, Ph was more abundant in SLDP by several folds than other regions. Combined with the geographical location of SLDP and the fact that the samples were collected in the ship's docking port, it is speculated that the organic aerosol samples of SLDP may be affected by local coastal and terrestrial biological sources and anthropogenic emissions (especially fossil fuel combustion). In addition, EC is the product of residential coal, fossil fuel combustion and incomplete combustion of biomass. The concentration of EC in SLDP aerosol (4.24±3.02 µg m⁻³) was also much higher than that in other marine areas (SCS: 0.27±0.15 µg m⁻³, EIO-WI: 0.05±0.05 µg m⁻³, EIO-SL: 0.21±0.18 µg m⁻³, Malacca: 0.24±0.18 µg m⁻³), indicating that the aerosol samples of SLDP were seriously polluted by ship exhaust but not for the other marine samples. EC is the product of residential coal, fossil fuel combustion and incomplete

combustion of biomass. The concentration of EC in SLDP aerosol ($4.24 \pm 3.02 \mu\text{g m}^{-3}$) was also much higher than that in other marine areas (SCS: $0.27 \pm 0.15 \mu\text{g m}^{-3}$, EIO-WI: $0.05 \pm 0.05 \mu\text{g m}^{-3}$, EIO-SL: $0.21 \pm 0.18 \mu\text{g m}^{-3}$, Malacca: $0.24 \pm 0.18 \mu\text{g m}^{-3}$), indicating that the aerosol samples of SLDP were seriously polluted by ship exhaust but not for the other marine samples.

8. Section 3.5 could be merged in Section 3.2.

Response: Section 3.5 has been merged into Section 3.2 and becomes Section 3.2.4.

References:

- Bikkina, S., Kawamura, K., and Miyazaki, Y.: Latitudinal distributions of atmospheric dicarboxylic acids, oxocarboxylic acids, and α -dicarbonyls over the western North Pacific: Sources and formation pathways, *Journal of Geophysical Research: Atmospheres*, 120, 5010-5035, 10.1002/2014jd022235, 2015.
- Deshmukh, D. K., Kawamura, K., and Deb, M. K.: Dicarboxylic acids, omega-oxocarboxylic acids, alpha-dicarbonyls, WSOC, OC, EC, and inorganic ions in wintertime size-segregated aerosols from central India: Sources and formation processes, *Chemosphere*, 161, 27-42, 10.1016/j.chemosphere.2016.06.107, 2016.
- Fu, P., Kawamura, K., and Miura, K.: Molecular characterization of marine organic aerosols collected during a round-the-world cruise, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Fu, P., Kawamura, K., Usukura, K., and Miura, K.: Dicarboxylic acids, ketocarboxylic acids and glyoxal in the marine aerosols collected during a round-the-world cruise, *Marine Chemistry*, 148, 22-32, 10.1016/j.marchem.2012.11.002, 2013.
- Ho, K., Cao, J., Lee, S., Kawamura, K., Zhang, R., Chow, J. C., and Watson, J. G.: Dicarboxylic acids, ketocarboxylic acids, and dicarbonyls in the urban atmosphere of China, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Kawamura, K., Kasukabe, H., and Barrie, L. A.: Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in Arctic aerosols: One year of observations, *Atmospheric Environment*, 30, 1709-1722, 1996.
- Kumar, A., Sudheer, A., and Sarin, M.: Chemical characteristics of aerosols in MABL of Bay of Bengal and Arabian Sea during spring inter-monsoon: a comparative study, *Journal of earth system science*, 117, 325-332, 2008.
- Kundu, S., Kawamura, K., and Lee, M.: Seasonal variations of diacids, ketoacids, and α -dicarbonyls in aerosols at Gosan, Jeju Island, South Korea: Implications for sources, formation, and degradation during long-range transport, *Journal of Geophysical Research*, 115, 10.1029/2010jd013973, 2010.
- Kunwar, B., and Kawamura, K.: Seasonal distributions and sources of low molecular weight dicarboxylic acids, ω -oxocarboxylic acids, pyruvic acid, α -dicarbonyls and fatty acids in ambient aerosols from subtropical Okinawa in the western Pacific Rim, *Environmental Chemistry*, 11, 673-689, 2014.
- Mochida, M., Kawamura, K., Umemoto, N., Kobayashi, M., Matsunaga, S., Lim, H. J., Turpin, B. J., Bates, T. S., and Simoneit, B. R.: Spatial distributions of oxygenated organic compounds (dicarboxylic acids, fatty acids, and levoglucosan) in marine aerosols over the western Pacific and off the coast of East Asia: Continental outflow of organic aerosols during the ACE-Asia campaign, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Pavuluri, C. M., Kawamura, K., and Swaminathan, T.: Water-soluble organic carbon, dicarboxylic acids, ketoacids, and β -dicarbonyls in the tropical Indian aerosols, *Journal of Geophysical Research Atmospheres*, 115, 10.1029/2009JD012661, 2010.
- Song, J., Zhao, Y., Zhang, Y., Fu, P., Zheng, L., Yuan, Q., Wang, S., Huang, X., Xu, W., Cao, Z., Gromov, S., and Lai, S.: Influence of biomass burning on atmospheric aerosols over the western South China Sea: Insights from ions, carbonaceous fractions and stable carbon isotope ratios, *Environ Pollut*, 242, 1800-1809, 10.1016/j.envpol.2018.07.088, 2018.
- Wang, G., Cheng, C., Meng, J., Huang, Y., Li, J., and Ren, Y.: Field observation on secondary organic aerosols during Asian dust storm periods: Formation mechanism of oxalic acid and related compounds on dust surface, *Atmospheric Environment*, 113, 169-176, 10.1016/j.atmosenv.2015.05.013, 2015.
- Wang, H., Kawamura, K., and Yamazaki, K.: Water-soluble dicarboxylic acids, ketoacids and dicarbonyls in the atmospheric aerosols over the Southern Ocean and western Pacific Ocean, *Journal of Atmospheric Chemistry*, 53, 43-61, 2006.

A list of all relevant changes made in the manuscript

1. The title of the manuscript is changed from “...over the South China Sea and East Indian Ocean” to “...from the South China Sea to the East Indian Ocean”.
2. Extra description is provided about the spatial distribution of OC, EC and WSOC with those reported in previous studies in Section 3.1 in the revised manuscript.
3. The original Section 3.5 has been merged into Section 3.2 and becomes the new Section 3.2.4.
4. Table 3 is moved to the Supporting Information.

Molecular and spatial distributions of dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls in marine aerosols from the South China Sea to East Indian Ocean

Jing Yang^{1,2}, Wanyu Zhao^{2,4}, Lianfang Wei², Qiang Zhang¹, Yue Zhao¹, Wei Hu¹, Libin Wu¹, Xiaodong Li¹, Chandra Mouli Pavuluri¹, Xiaole Pan², Yele Sun², Zifa Wang², Cong-Qiang Liu¹, Kimitaka Kawamura³, and Pingqing Fu¹

¹Institute of Surface-Earth System Science, Tianjin University, Tianjin 300072, China

²State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

³Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan

⁴College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China

Correspondence to: Pingqing Fu (fupingqing@tju.edu.cn)

Abstract. Marine aerosol samples collected from the South China Sea (SCS) to the East Indian Ocean (EIO) during a cruise from March 10 to April 26, 2015 were studied for diacids and related compounds. In view of the air masses backward trajectories and source regions of geographical features, the cruise area was categorized into the South China Sea (SCS), the East Indian Ocean off the coast of western Indonesia (EIO-WI), the EIO off the coast of Sri Lanka (EIO-SL), Malacca and Sri Lanka docking point (SLDP). Total concentrations of diacids, oxoacids and α -dicarbonyls were much higher at SLDP followed by the SCS, Malacca, and the lowest at the EIO-WI. In this study, oxalic acid (C₂) was the dominant diacid during the cruise, followed by malonic acid (C₃) in the SCS, EIO-WI, EIO-SL and Malacca, whereas succinic acid (C₄) diacid was relatively more abundant than C₃ diacid in SLDP. Except for SLDP, C₃/C₄ mass ratios were always greater than 1, and no significant difference was observed among the cruise. The C₂/C₄ and C₂/total diacids ratios also showed the similar trends. Average mass ratios of adipic acid (C₆) to azelaic acid (C₉) were less than unity except for at EIO-WI; the mass ratios of phthalic acid (Ph) to azelaic acid (C₉) were less than 2 except for at SCS. The concentrations of diacids were higher when the air masses originated from the terrestrial regions than those from the remote oceanic regions. Based on the molecular distributions of organic acids, the mass ratios and linear correlations of selected compounds in each area, we found that the oxidation of biogenic volatile organic compounds (BVOCs) released from the ocean surface and subsequent *in-situ* photochemical oxidations was the main contributor to diacids, oxocarboxylic acids and α -dicarbonyls over the SCS to EIO. In addition, the continental outflow that enriched with the anthropogenic VOCs and their aging influenced the organic aerosol loading, particularly over the SCS. The emissions from Sri Lanka terrestrial

vegetations as well as fossil fuel combustion and subsequent photochemical oxidation also played a prominent role in controlling the organic aerosols loading and molecular distributions of diacids and related compounds at SLDP.

1 Introduction

35 Land-sea-air interaction is one of the most important issues in earth system science. Atmospheric aerosol is the major component in the Earth's atmosphere and is one of the key carriers of the global biogeochemical cycle of nutrients (Zhuang et al., 1992; Li et al., 2017; Tan et al., 2011). Atmospheric aerosol containing inorganic and organic materials plays a vital role in air quality, atmospheric chemistry, health of human and ecosystems, and has a critical impact on the global climate system (Ramanathan et al., 2001). Organic aerosols in tropospheric regions
40 account for up to 90% (Kanakidou et al., 2005), and most of them are water-soluble (Kanakidou et al., 2005). The abundant diacids, oxoacids and α -dicarbonyls are reported to be the major fraction of water-soluble organic aerosols (Sorooshian et al., 2010; Ervens et al., 2011; Cong et al., 2015), which enhance the capacity of aerosol particles to act as cloud condensation nuclei and ice nuclei in the atmosphere (Ervens et al., 2011; Zhao et al., 2016a; Vergara-
Temprado et al., 2017; Huang et al., 2018), whereas the water-insoluble substances such as lipid class compounds
45 can reduce their hygroscopic activity (Kawamura et al., 2017).

It has been well established that the aerosols in the marine atmosphere are influenced by the emissions from the ocean surface, such as plankton activities (Cavalli et al., 2004; O'Dowd and De Leeuw, 2007; Facchini et al., 2008). Marine emissions account for more than half of the global natural aerosol burden (1000–3000 Tg yr⁻¹) (Iii and Duce, 1988), and thus significantly influence the Earth's climate system (Haywood and M., 1999). Previous studies
50 on marine aerosols have focused mainly on the measurements of non-sea-salt/sea-salt sulfates (Charlson et al., 1987; Charlson et al., 1992) and mineral dust (Schulz et al., 2012). However, diacids and related compounds in marine aerosols have been paid little attention to, despite the fact that they account for >10% of total carbon (TC) in the remote oceanic regions (Kawamura and Sakaguchi, 1999; Wang and Kawamura, 2006), which highlights their importance in the marine atmosphere.

55 Generally, molecular distributions of dicarboxylic acids in atmospheric aerosols are characterized by the dominance of oxalic (C₂) acid, followed by malonic (C₃) and succinic (C₄) acids (Fu et al., 2013a; Kawamura and Bikkina, 2016). They can be directly emitted from anthropogenic emissions such as fossil fuel combustion (Kawamura and Kaplan, 1987) and biomass burning (Narukawa et al., 1999; Kawamura et al., 2013), and formed by atmospheric oxidation of various volatile organic compounds (VOCs) emitted from primary sources (Kawamura

60 et al., 1996a). In addition, unsaturated fatty acids emitted from the oceanic biota can be further photo-oxidized to form the corresponding diacids in the atmosphere (Kawamura and Sakaguchi, 1999; Rinaldi et al., 2011). Isoprene, an abundant VOC from terrestrial higher plants as well as oceanic biota, is a major precursor of secondary organic aerosols (Lim et al., 2005; Carlton et al., 2009), including low molecular weight diacids (C₂–C₄) (Nguyen et al., 2010).

65 Chemical characterization of dicarboxylic acids, oxoacids and α -dicarbonyls in atmospheric aerosols provides useful knowledge on the relative contribution of possible primary sources, the long-range atmospheric transport and the photo-oxidation routes of organic compounds (Kawamura and Bikkina, 2016). Many studies have investigated the impacts of the continental air outflows on the concentrations, compositions and distributions of organic aerosols in the marine areas (Fu et al., 2013a). For example, a long-term observation of aerosols at the
70 Chichi-jima Island in the western North Pacific found a remarkable increase in the concentrations of dicarboxylic acids in winter and spring due to the influence of the East Asian air outflow by the westerlies (Mochida et al., 2003a). Moreover, organic compounds from marine sources, such as unsaturated fatty acids, are main contributors to the secondary formation of diacids and related compounds in the ocean atmosphere (Miyazaki et al., 2010; Kawamura and Gagosian, 1987). Bikkina et al. (2014) found that the abundance of C₂–C₆ and C₉ diacids
75 continued to increase with the increase in phytoplankton activities in the North Pacific Ocean. Meanwhile, the oxidation products (e.g. pyruvic and glyoxylic acids) of isoprene observed in the marine boundary layer had a similar concentration trend with that of C₂, suggesting that the secondary oxidation of biogenic VOCs from marine sources significantly contribute to the atmospheric loading of diacids and related compounds.

The oceans account for more than 70% of the earth's surface, and marine aerosols are important components of
80 the global aerosol system of natural sources. However, current knowledge about the biogeochemical cycles of organic matters in the tropical marine atmosphere is very limited. The South China Sea (SCS) is a large semi-closed marginal basin and one of the largest marginal seas in the world (Liu et al., 2002). The seasonal division of prevailing winds in the South China Sea is mainly influenced by the northeast monsoon roughly from mid-October to mid-March of the following year and by the southwest monsoon from mid-May to mid-September; while from
85 mid-March to mid-May is the spring transition period, during which the wind direction is variable. The climate of the South China Sea is part of the East Asian monsoon system (Lau et al., 1998). The Indian Ocean is the third largest ocean in the world, with distinct tropical maritime and monsoon climate characteristics. The prevailing wind over the Indian Ocean in summer is the southwest monsoon, while the prevailing wind in winter is the northeast monsoon (Fu et al., 2016; Ramanathan et al., 2005). The Indian Ocean is warmer than the Pacific and Atlantic at

90 the same latitude, so it is called the tropical ocean. The tropical East Indian Ocean (10°S–15°N, 65°E–100°E), including the southern bay of Bengal, southeastern Arabian Sea, eastern equatorial Indian Ocean and parts of the southern Indian Ocean, is one of the key regions affecting climate change such as drought and flood in China (Yao et al., 2015).

95 Although great progress has been made in the research field of atmospheric organic aerosols, the studies on diacids, oxoacids and α -dicarbonyls in the marine atmosphere are scarce and limited to the Mediterranean, North Pacific, South Pacific, Caribbean and Atlantic Ocean (Kawamura and Bikkina, 2016), while little is known about the marine aerosols over the Indian Ocean. Studies on the molecular composition and distribution of dicarboxylic acids and related compounds can provide useful information for source analysis, secondary formation and photochemical transformation processes of atmospheric organic aerosols. In this study, total suspended particle 100 (TSP) samples collected on day- and night-time basis over the South China Sea and the East Indian Ocean were studied for diacids, oxoacids, and α -dicarbonyls, which provided an ideal opportunity to investigate the spatial distributions of marine organic aerosols which influenced by organics from both marine emissions and long-range transported continental aerosols. In addition, carbonaceous components, i.e. organic (OC) and elemental carbon (EC) and water-soluble OC (WSOC) in the TSP samples were measured. Based on the molecular and spatial 105 distributions, and mass ratios of organic compounds and relations with bulk components together with the backward air mass trajectories, we discuss their sources and possible formation pathways in the studied region.

2 Materials and methods

2.1 Marine aerosol sampling

110 Marine aerosol sampling was performed on the *R/V* “Shiyan 1” during the cruise NORC2015-10 in the South China Sea and the East Indian Ocean (Figure 1) from 10 March to 26 April 2015. TSP samples were collected on daytime ($n = 44$) and nighttime ($n = 43$) using a high-volume air sampler (Kimoto AS 810A, KIMOTO, Japan) placed on the front upper deck of the vessel’s navigation room. The exhaust outlet of the ship was located at the stern, and the aerosol samples were collected underway to avoid the potential pollution from the ship emissions. The three 115 samples collected at the Sri Lanka docking point (SLDP) when the ship was docked in the port, which were seriously polluted by ship exhaust emission. Therefore, these three samples were discussed separately to compare with other samples. The air sampler was operated at a flow rate of $1.0 \text{ m}^3 \text{ min}^{-1}$ with pre-combusted (450°C, 6h) quartz filters (25 cm \times 20 cm, PALLFLEX®TM, 2500 QAT-UP). We also collected field blanks ($n = 4$) during the

cruise. After sampling, the filter samples were wrapped in aluminum foil and packed in zip-lock bag and stored in
120 dark at -20°C until the chemical analysis.

Five-day air mass backward trajectories arriving at 100 m above the sea level on each day were computed using
the HYSPLIT model (<https://www.ready.noaa.gov/HYSPLIT.php>) (Figure 1). The air masses arriving over the
South China Sea (SCS) originated from East Asia and surrounding oceanic regions. Those arrived over the East
Indian Ocean off the coast of western Indonesia (EIO-WI) originated from the remote EIO, while those arriving
125 over the EIO off the coast of Sri Lanka (EIO-SL) were mixed from the remote EIO and Bay of Bengal. The air
masses arrived over Malacca from the Southeast Asian subcontinent and the surrounding EIO. In addition, the
vessel was docked for 3 days at Sri Lanka docking point (SLDP), during which the air masses were originated from
the surrounding oceanic regions. Thus, the TSP samples collected in these different locales during the cruise may
have been affected by the long-distance atmospheric transport, and hence we segregated the TSP samples into five
130 categories: SCS, EIO-WI, EIO-SL, SLDP and Malacca (Figure 1).

2.2 Determination of diacids and related compounds

Aerosol samples were determined for water-soluble organic acids using a previously reported method (Kawamura,
1993). Briefly, small pieces (47 mm in diameter) of TSP filter samples were extracted with organic-free Milli-Q
water (10 mL) under ultrasonication for three times. Then, the extracts were concentrated to dryness and reacted
135 with 14% BF_3/n -butanol at 100°C for 1 hour. The derivatized acids and carbonyls were extracted with *n*-hexane
and injected into a split/splitless gas chromatography (GC-FID, Agilent 6980) equipped with an HP-5 column for
the determination of diacids and related compounds. The field blank filters were also analyzed using the same
experiment procedure. Recoveries of major organic acids were better than 85%. The analytical errors in duplicate
analysis were within 10% for major species. Concentrations of diacids and related compounds were corrected
140 according to the field blanks.

2.3 Measurements of WSOC, OC and EC

For WSOC measurement, 17.3 cm^2 or 34.7 cm^2 of each filter was extracted with Milli-Q water (20 mL) under
ultrasonication for 20 min. The extracts were analyzed using a total organic carbon (TOC) analyzer (TOC-L, 5000A,
Shimadzu, Japan) for WSOC. OC and EC were determined using a carbon analyzer (Sunset Laboratory Inc., USA)
145 using a piece (1.5 cm^2) of each filter following the Interagency Monitoring of Protected Visual Environments
(IMPROVE) protocol (Pavuluri et al., 2011). The limits of detection (LODs) for both OC and EC were $0.1\text{ }\mu\text{gC}$

cm⁻² with a precision of >10%. Concentrations of all the carbonaceous components reported here were corrected using the field blanks.

Table 1 shows the concentration levels of OC and EC in TSP samples collected from different marine regions. The concentrations of EC (4.24±3.02 µg m⁻³) in SLDP aerosols were much higher than those in other regions (SCS: 0.27±0.15 µg m⁻³; EIO-WI: 0.05±0.05 µg m⁻³; EIO-SL: 0.04±0.18 µg m⁻³; and Malacca: 0.24±0.18 µg m⁻³), which proved that the samples of SLDP were seriously polluted by ship exhaust, but other sea areas should be minor.

3 Results and discussion

3.1 OC, EC and WSOC in marine aerosols

The concentrations of OC, EC and WSOC in each marine region (SCS, EIO-WI, EIO-SL, SLDP and Malacca) are summarized in Table 1. In the TSP samples collected in the South China Sea to the East Indian Ocean from March to April 2015, the mass concentrations of OC and EC varied from 0.02 to 20.9 µg m⁻³ (average 1.29±2.72 µg m⁻³) and from 0.00 to 7.57 µg m⁻³ (average 0.33±0.90 µg m⁻³), respectively. Generally, the mass concentrations of OC and EC in SLDP were the highest, followed by Malacca and SCS, and the lowest in EIO-WI and EIO-SL. In this study, the mean mass concentrations of OC and EC in SCS were 1.5 times (OC: 0.82 µg m⁻³) and 1.7 times (EC: 0.16 µg m⁻³) higher than those reported by Fu et al. (2013a) in the South China Sea in late February 1990. Because of the same sampling season, the above situation indicated that with the economic development of the surrounding areas of the South China Sea, the input of anthropogenic emissions to the South China Sea carbon aerosols were also increasing. And, the mass concentrations of OC and EC in SCS were lower than those in the western South China Sea (OC: 2.04 µg m⁻³, EC: 3.0 µg m⁻³) which were strongly affected by biomass burning (Song et al., 2018). In addition, the carbonaceous aerosol levels in Malacca were similar to those in the Bay of Bengal (OC: 1.9 µg m⁻³, EC: 0.4 µg m⁻³) (Kumar et al., 2008). The concentrations of carbon components in SLDP were relatively close to that of Jeju Island (OC: 11.3 µg m⁻³, EC: 3.0 µg m⁻³) (Jung and Kawamura, 2011), and were higher than that of Sapporo Island (OC: 6.1 µg m⁻³, EC: 1.9 µg m⁻³) (Simoneit et al., 2004), these two sites were greatly affected by terrestrial long-distance transmission. Meanwhile, the concentrations of carbon components in EIO-WI and EIO-SL were lower than those in Northern Atlantic (OC: 0.93 µg m⁻³, EC: 0.27 µg m⁻³) (Zhang et al., 2010), Pacific (OC: 0.82 µg m⁻³, EC: 0.22 µg m⁻³) (Zhang et al., 2010), Amsterdam Island (OC: 0.24 µg m⁻³, EC: n.d.) (Claeys et al., 2010), Arctic ocean (OC:0.56 µg m⁻³) (Fu et al., 2013b) and so on. These places were all less affected by

175 terrestrial anthropogenic emissions. The concentration levels of carbonaceous components in EIO-WI and EIO-SL were lower than those in the Indian Ocean (OC: $0.71 \mu\text{g m}^{-3}$, EC: $0.49 \mu\text{g m}^{-3}$) (Fu et al., 2011).

EC is the product of residential coal, fossil fuel combustion and incomplete combustion of biomass, while OC contains both primary organic carbon and secondary organic aerosol generated by photochemical reactions of organic gases in the atmosphere. The concentration of EC in SLDP aerosol ($4.24 \pm 3.02 \mu\text{g m}^{-3}$) was much higher than that in other marine areas (SCS: $0.27 \pm 0.15 \mu\text{g m}^{-3}$, EIO-WI: $0.05 \pm 0.05 \mu\text{g m}^{-3}$, EIO-SL: $0.21 \pm 0.18 \mu\text{g m}^{-3}$, Malacca: $0.24 \pm 0.18 \mu\text{g m}^{-3}$), indicating that the aerosol samples of SLDP were seriously polluted by ship exhaust but not for the other marine samples. OC and EC can roughly determine the source of the aerosols. There was a weak correlation between the concentration of OC and EC in aerosols from the South China Sea to the East Indian Ocean ($R^2=0.34$, $P<0.001$), indicating that the origins of organic aerosols were different from that of EC and were affected by multiple sources (Turpin and Huntzicker, 1995). In addition, the OC/EC ratio can be used to indicate the contribution of different sources (Chow et al., 1996). The contribution of different emission sources, the influence of secondary organic aerosol formation, and the removal path of OC and EC can all affect the OC/EC ratio. When the OC/EC ratio was greater than 2, it indicates the presence of secondary organic aerosols. The OC/EC ratio was 5.52 ± 5.33 (range 1.02–29.7) in SCS, 11.84 ± 17.44 (range 1.55–51.0) in EIO-WI, 0.96 ± 2.33 (range 0.96–9.28) in EIO-SL, 8.76 ± 6.84 (range 2.71–24.4) in Malacca, and 2.98 ± 0.97 (range 2.14–4.04) in SLDP. Among them, the ratio of OC/EC in SCS was very similar to that measured by Fu et al. (2013a) in TSP samples collected in the South China Sea (OC/EC: 5.1). The above results indicated that the contribution of secondary organic aerosols was widespread in aerosol samples from the South China Sea to the East Indian Ocean, except for some samples in EIO-WI and EIO-SL.

195 The concentrations of WSOC were higher (range 1.12 – 7.44 ng m^{-3} , average $4.43 \pm 3.17 \text{ ng m}^{-3}$) in the SLDP followed by Malacca (0.09 – 2.72 ng m^{-3} , $1.15 \pm 0.92 \text{ ng m}^{-3}$), SCS (0.22 – 2.31 ng m^{-3} , $1.05 \pm 0.65 \text{ ng m}^{-3}$), EIO-SL (0.06 – 1.00 ng m^{-3} , $0.35 \pm 0.25 \text{ ng m}^{-3}$) and EIO-WI (0.05 – 0.58 ng m^{-3} , $0.15 \pm 0.12 \text{ ng m}^{-3}$). The WSOC concentrations over the SCS to EIO are much lower than those in urban aerosols from Tokyo (average $13 \mu\text{g m}^{-3}$) (Sempéré and Kawamura, 1994) and in the biomass burning aerosols from Amazonia (18 – $51 \mu\text{g m}^{-3}$) (Kundu et al., 2010a). However, WSOC concentrations in the EIO-SL are comparable to those reported in marine aerosols from the central Pacific (average $0.30 \mu\text{g m}^{-3}$) (Hoque and Kawamura, 2016), the western Pacific (average $0.33 \mu\text{g m}^{-3}$) (Sempéré and Kawamura, 2003), and Hawaii (average $0.39 \mu\text{g m}^{-3}$) (Hoffman and Duce, 2013). Kawamura et al. (2010) determined lower WSOC concentrations (0.04 – $0.30 \mu\text{g m}^{-3}$, average $0.18 \mu\text{g m}^{-3}$) in the Arctic aerosols, which are comparable to those observed in the EIO-WI.

205 3.2 Concentrations and molecular distributions of dicarboxylic acids and related compounds

The concentrations of dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls in each marine region (SCS, EIO-WI, EIO-SL, SLDP and Malacca) are summarized in Table 1. Their chemical structures are shown in Fig. S1. Most of the organic species were found to be more abundant at SLDP followed by the SCS, Malacca, EIO-WI and EIO-SL, respectively (Table 1). For the aerosol samples of SCS, the 5-day backward trajectories (Fig. 1b) showed that the air masses were originated from East Asia, whereas the air masses of Malacca were delivered from Southeast Asia. The concentrations of diacids and related compounds were the highest in SCS followed by Malacca, due to increased anthropogenic activities through long-range atmospheric transport in East Asia (Zhao et al., 2016b) and Southeast Asia, respectively. C_2 diacid showed the highest abundance followed by $C_3 > C_4 > \text{MeGly} > \omega C_2 > C_5 > \text{Ph} \approx \text{Gly} > C_9$ (see Table 1 for full form of abbreviation) among the measured major species over the SCS (Fig. 2a), while $C_3 > \text{MeGly} > C_4 > F > \text{Ph} > C_6 \approx C_9$ over the EIO-WI (Fig. 2b), $C_3 > C_4 > C_9 \approx \text{MeGly} > \omega C_2 > \text{Gly} > \text{Ph}$ at EIO-SL. The order in Malacca was $C_2 > C_3 > C_4 > \text{kC}_7 > C_5 > \omega C_2 > C_9 \approx \text{MeGly}$. The dominance of C_2 followed by C_3 and C_4 diacids is consistent with those coastal marine aerosols (Kundu et al., 2010b; Kunwar and Kawamura, 2014) and remote marine aerosols from the western North Pacific (Bikkina et al., 2015), suggesting similar formation processes of dicarboxylic acids in the atmosphere of the coastal or remote regions.

220 Table 1 shows the total concentrations of diacids in marine aerosols from five regions in this study (SCS: $0.27 \pm 0.15 \text{ ng m}^{-3}$; EIO-WI: $0.05 \pm 0.05 \text{ ng m}^{-3}$; EIO-SL: $0.04 \pm 0.18 \text{ ng m}^{-3}$; Malacca: $0.24 \pm 0.18 \text{ ng m}^{-3}$; and SLDP: $0.27 \pm 0.15 \text{ ng m}^{-3}$). Fu et al. (2013a) reported that the mean concentration of total diacids in marine aerosol samples in the South China Sea and the Indian Ocean was 489 ng m^{-3} and 301 ng m^{-3} , respectively. By comparison this study with aerosols from other marine areas, it was found that total diacids in the EIO was similar to that in the western Pacific Ocean ($60 \pm 39 \text{ ng m}^{-3}$) (Wang et al., 2006). Furthermore, the concentrations of total diacids from five sea areas were all lower than that of the East China Sea (850 ng m^{-3}) (Mochida et al., 2003b), and higher than that of the Southern Ocean ($4.5 \pm 4.0 \text{ ng m}^{-3}$) (Wang et al., 2006). In addition, the concentrations of total diacids in aerosol samples in all sea areas were higher than that in the Alert, Canadian high Arctic (25 ng m^{-3}) (Kawamura et al., 1996a), and lower than that in the urban aerosol from Xi'an, China in different dust events ($932\text{--}2240 \text{ ng m}^{-3}$) (Wang et al., 2015), and the megacity Chennai, India (694.5 ± 176.3 and $640.6 \pm 150.6 \text{ ng m}^{-3}$ in early and late winter) and in summer ($502.9 \pm 117.9 \text{ ng m}^{-3}$) (Pavuluri et al., 2010). Combined with the geographical location of SLDP and the fact that the samples were collected in the ship's docking port, it is speculated that the organic aerosol samples of SLDP may be affected by local coastal and terrestrial biological sources and anthropogenic emissions

(especially fossil fuel combustion). The pattern of diacids abundance order ($C_2 > C_9 > C_4 > C_3 > Ph > C_5$) observed at SLDP was similar to that in continental aerosols (Kawamura and Bikkina, 2016).

3.2.1 Dicarboxylic acids

C_2 – C_4 , C_6 , C_9 and Ph are important species of dicarboxylic acids; this section briefly summarizes the main sources and/or formation processes of these organic acids. C_2 diacid concentrations ranged from 4.43 to 70.1 ng m^{-3} with an average of $18.2 \pm 18.4 \text{ ng m}^{-3}$ in the EIO-WI, 5.21–232 ng m^{-3} (average $58.2 \pm 55.3 \text{ ng m}^{-3}$) in the EIO-SL, 14.5–373 ng m^{-3} ($145 \pm 112 \text{ ng m}^{-3}$) in Malacca, 30.3–626 ng m^{-3} ($233 \pm 143 \text{ ng m}^{-3}$) in the SCS and 179–454 ng m^{-3} ($303 \pm 140 \text{ ng m}^{-3}$) in SLDP. It can be clearly seen that the concentrations of C_2 in SCS and Malacca were higher than those in EIO-SL and EIO-WI (Table 1). The large amount of C_2 can be generated from the following sources: fossil fuel combustion (Kawamura and Kaplan, 1987;Donnelly et al., 2010), biomass combustion (Schauer et al., 2001;Narukawa et al., 1999), cooking emissions (Kawamura and Kaplan, 1987;Rogge et al., 1993), photooxidation of VOCs and other precursors (Kawamura and Yasui, 2005;Kundu et al., 2010b). The diurnal variation trend of C_2 was similar to that of C_3 and C_4 , indicating that these compounds may have similar photochemical oxidation pathways or emission sources in the atmosphere.

C_9 diacid is the most abundant diacid in the range of C_7 – C_{11} diacids, which is similar to that in marine aerosols from the southern and western Pacific Ocean (Wang et al., 2006). Such high abundance of C_9 diacid indicates that marine biogenic emissions and subsequent photochemical formation of organic aerosols were likely significant over the SCS and EIO, because C_9 diacid is a photochemical oxidation product of unsaturated fatty acids emitted from the productive marine regions (Bikkina et al., 2014;Hoque et al., 2017). In addition, the photochemical oxidation of C_9 diacid leads to the generation of its lower homologues including C_4 – C_6 diacids (Yang et al., 2008;Kawamura et al., 1996a). Thus, the similar abundance of diacids (C_2 to C_6 and C_9) in the SCS and Malacca (Fig. 3) suggest that the marine biogenic emission was a major source, although the influence of the continental air masses transported from East and Southeast Asia was also significant. However, the concentrations of diacids over Malacca were relatively low, which indicated that the influence of polluted air masses transported from Southeast Asia was lower than those from East Asia. In general, the abundance of diacids in the EIO-SL and EIO-WI regions was lower than that in the SCS and Malacca. It was likely because the air masses arrived over EIO-SL and EIO-WI regions mostly originated from the remote oceanic regions and thus, the marine biogenic emission should be the major potential source of organic aerosols with no significant influence from the Asian outflow. Interestingly, the concentrations of C_9 diacid (average $68.9 \pm 65.1 \text{ ng m}^{-3}$) in SLDP samples were several times higher than that observed in other regions over the SCS and EIO. As noted above, C_9 diacid is mostly derived from unsaturated

fatty acids of biogenic origin and has been considered as a marker for biogenic organic aerosols (Kawamura and Gagosian, 1987). Therefore, the high concentration of C₉ diacid in SLDP is likely derived by the oxidation of unsaturated fatty acids from mainland biota (Zhao et al., 2018). In fact, the terrestrial air masses that enriched with higher plant emissions and SOA produced upon subsequent oxidation of BVOCs during daytime should be transported onshore during nighttime by land breezes (Kang et al., 2017). In addition, the photochemical breakdown of C₉ diacid should be insignificant during nighttime.

Figure 3 shows the temporal variations in selected diacids over the SCS and EIO during the sampling periods. The spatial patterns of C₆ and C₉ diacids were similar (Fig. 3). C₆ diacid could be produced via the oxidation of cyclic olefins emitted from anthropogenic sources (e.g. fossil-fuel combustion) (Kawamura et al., 1996a), and could also be derived from photochemical breakdown of biogenic C₉ diacid (Kawamura et al., 1996b). Bikkina et al. (2015) concluded that the high mass concentrations of C₉ in marine aerosols were consistent with the high biological activity at the ocean surface. Such similar temporal variations in C₆ and C₉ diacids together with the results of backward trajectories (Fig. 1b) indicate that the formation of C₆ diacid is mostly from the photochemical breakdown of C₉ diacid which should have been derived from marine biogenic emissions. Based on ultraviolet (UV) irradiation of oleic acid and ozone system, Matsunaga et al. (1999) found a series of C₂ to C₉ diacids, with high abundance of C₉ diacid. Temporal variations in C₉, C₆, and C₂–C₄ diacids were also similar (Fig. 3), suggesting that photochemical breakdown of C₉ might be the major formation pathway of short-chain diacids such as C₆, C₅, and C₄ diacids. Further aging processes eventually led to the formation of C₂ diacid in the EIO. Thus, the molecular distributions and temporal variations of diacids inferred that the organic aerosols collected over the EIO should be mainly derived from marine biogenic emissions. However, the contribution of diacids to WSOC was reduced in the EIO-WI and EIO-SL due to low biological activity in those oceanic regions as reflected by low C₉ diacid loading.

Phthalic acid (Ph acid) is a tracer for anthropogenic organic aerosols. Concentrations of Ph acid in EIO-WI and EIO-SL were similar to those in the western North Pacific aerosol (MBA: 0.6 ng m⁻³; LBA: 0.5 ng m⁻³) (Bikkina et al., 2015). In addition, concentrations of Ph acid over the SCS to EIO were much lower than those reported in the urban atmosphere such as Chennai (21 ng m⁻³) (Pavuluri et al., 2010) and Chinese cities (90 ng m⁻³) (Ho et al., 2007). Ph acid can be released directly from fossil fuel combustion (Kawamura and Kaplan, 1987). Secondary formation through atmospheric oxidation of aromatic hydrocarbons such as naphthalene is also important (Fu et al., 2009; Ho et al., 2010). Such low levels of Ph acid suggested that the marine aerosols of EIO were not so seriously impacted by continental air masses from East and Southeast Asia. It was worth noting that Ph was more abundant

in SLDP by several folds than other regions (Table 1), so we do not preclude the influence of anthropogenic emissions, particularly fossil fuel combustion, associated with the terrestrial air masses arrived in SLDP.

295 3.2.2 Oxocarboxylic acids

Oxocarboxylic acids (ω C₂– ω C₉ and pyruvic acid), the intermediate products of the oxidation of monocarboxylic acids, can be further oxidized to form diacids (Warneck, 2003;Carlton et al., 2007). Being similar to diacids, oxocarboxylic acids are mainly derived from combustion sources, but can also be produced by photooxidation of various organic precursors in the atmosphere from anthropogenic and biological sources (Kawamura and Bikkina, 2016). The concentrations of total oxoacids ranged from 1.48 to 13.2 ng m⁻³ (average 6.51 ± 3.99 ng m⁻³) in the SCS, 0.16–2.01 ng m⁻³ (0.96 ± 0.46 ng m⁻³) in the EIO-WI, 0.23–5.06 ng m⁻³ (1.85 ± 1.51 ng m⁻³) in the EIO-SL, 0.29–10.4 ng m⁻³ (4.22 ± 3.75 ng m⁻³) in Malacca and 7.46–40.3 ng m⁻³ (24.0 ± 16.4 ng m⁻³) in SLDP. The concentrations of total oxoacids are lower than those from Gosan, Jeju Island, South Korea (average 53 ng m⁻³) (Kawamura et al., 2004) and urban sites in China (45 ng m⁻³) (Ho et al., 2007). The concentrations of oxoacids were lower in the EIO-WI than the other regions (Table 1). The spatial distributions of ω -oxoacids showed a pattern of SLDP > SCS > Malacca > EIO-SL > EIO-WI, being consistent with those of major diacids (C₂, C₃, and C₄), which indicated that these oxoacids were potential precursors of dicarboxylic acids (Sempère and Kawamura, 1994).

Glyoxylic acid (ω C₂) was the most abundant oxoacid followed by pyruvic acid (Pyr) acid (Table 1). All of them are important intermediates in photo-oxidation processes and are used in the production of low carbon-number diacids such as C₂, C₃ and C₄ diacids (Hatakeyama et al., 1987). Several studies reported that Pyr is produced by in-cloud oxidation of isoprene in the atmosphere (Carlton et al., 2006;Carlton et al., 2009) that can be transported from mainland regions and/or emitted from the ocean surface (Shaw et al., 2010).

3.2.3 α -Dicarbonyls

The concentrations of total α -dicarbonyls (glyoxal and methylglyoxal) varied over a wide range (0.56–20.2 ng m⁻³) with relatively high abundance in the SCS (average 5.25 ± 4.46 ng m⁻³) and SLDP (average 7.35 ± 3.58 ng m⁻³). The average concentration of total α -dicarbonyls during the cruise was higher than those in super-micron and submicron aerosols collected over the North Pacific in summer (0.51 ± 0.22 ng m⁻³ in more biologically influenced aerosols and 0.66 ± 0.20 ng m⁻³ in less biologically influenced aerosols) (Miyazaki et al., 2010). The abundances of α -dicarbonyls were higher in the SCS and Malacca than those in the EIO-SL and EIO-WI, which were similar to diacids and oxoacids (Fig. 2 and Table 1). The concentrations of both glyoxal (Gly) and methylglyoxal (MeGly) were higher in the SCS and their spatial distributions were consistent with those of diacids. Gly and MeGly can be

formed via atmospheric oxidation of isoprene, emitted from the terrestrial vegetation and/or marine phytoplankton (Sorooshian et al., 2009). Besides, Gly is also a photochemical oxidation product of aromatic hydrocarbons from combustion of fossil fuels (Carlton et al., 2007; Volkamer et al., 2007). Both species can be existed in the aerosol phase and further oxidized to form less-volatile organic acids such as Pyr, ω C₂, and C₂ (Sorooshian et al., 2006). Therefore, the higher abundances of Gly and MeGly over the SCS than in other three regions could be due to enhanced continental outflow as well as the high biological activity in the sea surface. The spatial distributions of total oxoacids and α -dicarbonyls were similar to those of total diacids, indicating that they were similar in origin or formation mechanism (Kunwar et al., 2017). Based on the backward trajectory analysis, we identified four source regions of diacids and related compounds: (a) East Asia, (b) Southeast Asia, (c) the Bay of Bengal, and (d) the East Indian Ocean. When air masses originated from East Asia and Southeast Asia and their coastal areas (i.e. SCS and Malacca), the concentrations of these compounds were high; when air masses were mainly derived from the East Indian Ocean (i.e. EIO-WI), their concentrations was the lowest; and when the air masses originated from the Bay of Bengal and the East Indian Ocean (i.e. EIO-SL), the concentration of these compounds was between the former two.

3.2.4 Fractions of diacids and related compounds in WSOC and OC

Spatial/temporal distributions of WSOC showed much higher values in the SCS and Malacca than those in the EIO-WI and EIO-SL regions (Fig. 3j). As discussed above, such high abundance of WSOC in the SCS and Malacca may be caused by enhanced aging of organic compounds from the surface ocean by high biological activities and then subsequent oxidation to produce more water-soluble organics. In this study, the spatial/temporal trends of diacids and related compounds are generally consistent with that of OC, EC and WSOC, which are high in the SCS and low in other regions. The fractions of some major diacids, oxoacids and α -dicarbonyls in WSOC and OC are presented in Table 2. The total diacids account for $12.0 \pm 5.75\%$ (range 2.00–32.4%) of WSOC over the EIO. Whilst, the fractions of oxoacids and α -dicarbonyls in WSOC ranged from 0.03 to 1.22% (average $0.34 \pm 0.20\%$) and 0.02–3.12% ($0.60 \pm 0.68\%$), respectively. In addition, a strong linear correlation was found between WSOC and C₂ diacid (Table 2), which suggests that the WSOC was mainly generated by secondary formation, rather than biomass-burning emissions associated with the Asian outflow.

3.3 Relative abundances of diacids and related compounds

350 The concentrations of diacids and related compounds and their relative abundances in the atmosphere are controlled by the emission of their precursors and subsequent oxidation processes. Pie diagrams of the percentage contributions of individual straight chain diacids to total aliphatic diacids in the different regions over the SCS to EIO (SCS, EIO-WI, EIO-SL, Malacca and SLDP) are depicted in Fig 4. The percentage contributions of LMW-diacids (C_2-C_4) in total mass concentrations ($\Sigma(C_2-C_{12})$) varied from 96.4 to 98.9% ($97.6 \pm 1.25\%$) in the SCS, 85.2
355 to 96.0% ($90.6 \pm 5.38\%$) in the EIO-WI, 90.6 to 99.4% ($95.0 \pm 4.44\%$) in the EIO-SL, 94.2 to 98.0% ($96.1 \pm 1.88\%$) in Malacca, and 72.0 to 95.0% ($83.5 \pm 11.5\%$) in SLDP. Interestingly, the relative abundances of C_2 to total mass concentrations of C_2 to C_{12} diacids (ΣC_2-C_{12}) were similar in four regions (Fig. 4). A significant difference was found in the relative abundance of C_2 diacid between SLDP and other four regions with the lower value in SLDP, whereas that of C_9 diacid was the opposite.

360 Figure 5 presents the relative abundances of the sums of short-chain diacids (C_2-C_4), long-chain diacids (C_5-C_{12}), unsaturated diacids (M, F, mM, Ph, iPh, and tPh), diacids containing additional functional group (hC₄, kC₃, and kC₇), oxoacids ($\omega C_2-\omega C_9$ and Pyr) and α -dicarbonyls (MeGly, Gly) in the measured total diacids and related compounds in different regions from the SCS to EIO. The relative abundances of short-chain diacids were much higher accounting for about 90% in the SCS and about 60% in the EIO-WI. Long-chain diacids were found to be
365 second highest in SLDP and Malacca whereas that of α -dicarbonyls in the EIO-WI and EIO-SL, with high abundance in the SLDP and EIO-SL, respectively (Fig. 5). Furthermore, the abundance of long-chain diacids was relatively high in the EIO-WI and EIO-SL but was the lowest in the SCS. The relative abundances of unsaturated diacids were the highest in the EIO-WI followed by SLDP; while oxoacids were the highest in SLDP, followed by the EIO-WI, with the lowest values in the SCS and Malacca (Fig. 5). The high abundances of short-chain diacids
370 indicate that the aerosols over the SCS to EIO were significantly aged during the long-distance transport, while the high abundances of long-chain diacids and α -dicarbonyls suggest that the contributions from biogenic emissions were much higher over the EIO and at SLDP. The high atmospheric levels of unsaturated fatty acids and oxoacids in SLDP infer that the contributions from anthropogenic sources are also significant at SLDP.

3.4 Implications for the origins and formation pathways of diacids and related compounds

375 Different emission sources and the extent of aging lead to large differences in the concentrations of diacids and related compounds, and hence, mass concentration ratios of selected species can serve as effective markers for the identification of their origins and formation/transformation processes. Temporal variations in mass ratios of several

organic acids as well as the relative abundance of C_2 to total aliphatic diacids are depicted in Fig. 6. The spatial distributions of C_3/C_4 , C_2/C_4 and $C_2/(\sum C_2-C_{12})$ ratios are similar to each other. It has been reported that C_3 diacid can be produced from C_4 diacid by oxidative reaction (Kawamura and Ikushima, 1993). There were robust relationships among C_2 , C_3 , and C_4 diacids in SCS, EIO-WI and EIO-SL regions, and between C_3 and C_4 diacids in Malacca (Table S1), suggesting that they should have been derived from similar sources. The C_3/C_4 ratio can be used as an effective marker to assess the extent of organic aerosol aging (Kawamura and Ikushima, 1993). The C_3/C_4 ratios were reported to be low (0.25–0.44) in vehicular emissions, because the C_3 diacid is thermally unstable to be easily degraded during combustion (Kawamura and Kaplan, 1987). In contrast, the C_3/C_4 ratios reported to be high (range, 1.0–11, average 3.9) in marine aerosols collected from the North Pacific (including tropics), which have been considered to be photochemically aged during the atmospheric transport (Kawamura and Sakaguchi, 1999). The C_3/C_4 ratios in this study ranged from 0.3 to 4.6 with an average value of 2.0. They are higher than those in urban aerosols from Chinese megacities (0.6–1.1, average 0.74) (Ho et al., 2007), Tokyo (0.56–2.9, average 1.6) (Kawamura and Ikushima, 1993) and New Delhi, India (0.40–1.1; average 0.66 in daytime and 0.58 in nighttime) (Miyazaki et al., 2009). It is worth noting that both malonic and succinic acids show a net loss from coastal areas to the open oceans, but the net loss is less for C_3 diacid—potentially owing to photochemical processing (Fu et al., 2013a). Therefore, such higher C_3/C_4 ratios suggest that the organic aerosols over the SCS to EIO are significantly subjected for intensive photochemical aging during the atmospheric long-range transport of the outflows from East and Southeast Asia to remote oceanic regions (Fig. 1). Furthermore, the C_3/C_4 ratios increased with the decrease of latitude over the SCS, suggesting that the photochemical transformation of C_4 to C_3 diacid was enhanced at lower latitudes due to the increase of temperature and solar radiation. However, such a trend did not appear during the return voyage and the C_3/C_4 ratios were much lower than those reported over the North Pacific. In addition, the temporal variations in their concentrations were similar throughout the campaign. Such results indicate that their *in-situ* formation from the BVOCs (e.g. isoprene) emitted from marine biota might also be significant.

The C_2/C_4 mass ratios can also be used to assess the extent of organic aerosols aging (Sorooshian et al., 2007). In the SCS, the C_2/C_4 ratios (7.6–28.9) showed an increasing trend with the decreasing latitude. The C_2/C_4 ratios in the EIO are higher than those reported in Chinese cities (average 7.1) (Ho et al., 2007), Sapporo (average 3.1) (Aggarwal and Kawamura, 2008), and South Korea (average 8.6) (Kundu et al., 2010b). Such a result indicates that C_2 could be largely generated by the photochemical degradation of C_4 diacid. Also, a significant correlation was found between relative abundance of C_2 (C_2 %) and C_2/C_4 mass ratio (Fig. S2c), suggesting the potential formation of C_2 diacid from photochemical breakdown of C_4 diacid. A concurrent trend (Fig. 6c) was also obtained

for the increased contribution of C₂ to total aliphatic diacids (C₂/Σ(C₂–C₁₂)), suggesting more photochemical aging of organic aerosols in low latitudes with stronger solar radiation.

410 The C₂/Σ(C₂–C₁₂) ratio has also been considered as an indicator to the degree of organic aerosols aging during long-distance atmospheric transport (Wang et al., 2006). In general, the more the aerosol particles are aged, the higher the C₂/Σ(C₂–C₁₂) ratios are (Kawamura and Sakaguchi, 1999). The C₂/Σ(C₂–C₁₂) was higher (0.76±0.06) in the SCS than that in the EIO-SL (0.74±0.08) and Malacca (0.71±0.11), followed by the EIO-WI (0.65±0.11) and SLDP (0.60±0.11). It is lower than that (0.8±0.04) reported in wintertime and higher than in summertime (0.5±0.01)
415 Himalayan aerosols (Hegde and Kawamura, 2012). These results indicate that the organic aerosols in the SCS should have been more aged and/or influenced by anthropogenic emissions from East Asia, whereas in the EIO-WI and Malacca the aging of organic aerosols might be less intensive and influenced by only the marine biogenic emissions. Whilst in SLDP, the low C₂/Σ(C₂–C₁₂) should have been driven by the local coastal biota and the terrestrial biogenic and anthropogenic emissions as well as the less aging, particularly in the nighttime.

420 Maleic acid (M, *cis*-form) is transformed to fumaric acid (F, *trans*-form) with enhanced photochemical activity and hence, the mass ratio of M/F has been considered as a proxy to assess the degree of aerosol aging (Kawamura and Sakaguchi, 1999). In general, low M/F values reflect secondary oxidation as an important source. M/F ratios (0.07–0.73) were found to be lower in the EIO-WI than other (0.9–2.3) regions (Table 3). The mean value of M/F was 1.33 in samples collected in the SCS, 0.29 in the EIO-WI, 1.30 in the EIO-SL, 2.08 in Malacca, and 1.77 in
425 SLDP. They were all lower than those in Xi'an aerosols (summer: 2.22, winter: 2.38), and were about the same or less than those reported at the Korean Gosan site (spring: 1.38, summer: 0.76, autumn: 1.62, winter: 2.21), which have been considered to be aged, indicating that the aging of organic aerosols was more intensive in the EIO-WI, whereas in Malacca and SLDP, the organic aerosols were less aged and/or influenced by local emissions, and thus the *cis-trans* transformation is weak.

430 According to the description of the origins of C₆, Ph and C₉ in the above sections, the C₆/C₉ and Ph/C₉ ratios could provide insights into the relative strengths of biogenic and anthropogenic sources in the given area. C₆/C₉ ratios (Fig. 6f) showed lower values in the EIO (except sample No. 2), which can be attributed to large influence of the marine air masses containing fatty acids like oleic acid, a precursor of C₉ diacid (Kawamura and Gagosian, 1987). Temporal variations in Ph/C₉ ratios over the SCS to EIO are shown in Fig. 6g, with the averages of 2.02 in
435 the SCS, 1.56 in the EIO-WI, 1.22 in the EIO-SL, and 0.90 in Malacca. They were lower than that (5.71) reported in wintertime Gosan aerosols from South Korea (Kundu et al., 2010a), but higher than these in summertime (1.7) and wintertime (1.78) aerosols in Xi'an, China (Wang et al., 2012) and that in the western Pacific (1.41) (Sempéré

and Kawamura, 2003). These comparisons suggest significant secondary formation of organic aerosols from anthropogenic precursors (e.g. naphthalene) transported from East and Southeast Asia (Fig. 1). However, C₆ and C₉ diacids showed a positive correlation ($R = 0.51$, $p < 0.0001$) among all samples, indicating that C₆ should also be significantly derived from photo-oxidation of BVOCs such as cyclic olefins. The higher Ph/C₉ ratios found over the SCS than in other regions (Fig. 6g) again suggest that the contribution from fossil-fuel combustion is also significant, associated with the continental outflow mainly over the SCS and slightly over the EIO.

The origins of MeGly, Gly, Pyr and isoprene have been discussed in the above sections. The mass ratios such as Pyr/C₂, ω C₂/C₂, Gly/C₂, and MeGly/C₂ can be used to explore the C₂ formation pathway and/or its origin. Here, all the ratios were found to peak in the EIO-WI and/or EIO-SL regions (Fig. 6), where the clean oceanic air masses arrived at the sampling points (Fig. 1). Whereas in the SCS, Malacca and SLDP, those ratios were insignificant except ω C₂/C₂ (Fig. 6). Such spatial distributions of Pyr/C₂, ω C₂/C₂, Gly/C₂, and MeGly/C₂ ratios indicate that the loading of organic aerosols in the EIO-WI and EIO-SL regions should have been influenced by the marine biogenic emissions and their subsequent oxidation processes. In addition, there were good correlations among Pyr, ω C₂ and MeGly in the SCS and Malacca, while the correlations were poor in the samples from EIO-WI and EIO-SL (except for Pyr and MeGly in EIO-WI, and ω C₂ and Pyr in EIO-SL). The results showed that organic aerosols produced by BVOCs (e.g. isoprene) emitted from the ocean surface were more aged in the EIO-WI and EIO-SL than the SCS and Malacca where were affected by anthropogenic emissions.

Finally, to better understand the enhanced emission of unsaturated fatty acids and BVOCs from the biologically active ocean surface over the SCS to EIO, we presented the satellite image of Chlorophyll-a loading in the EIO in Fig. 1a. The concentrations of Chlorophyll-a in the SCS and Malacca were higher than the other studied regions. Interestingly, high concentrations of diacids and related compounds (Fig. 3) were consistent with high concentrations of Chlorophyll-a in the marine surface. This consistency indicates that the high abundance of diacids in the SCS and Malacca attributes to the oceanic emissions of BVOCs from high biological area followed by photooxidation, although the influence of anthropogenic emissions associated with the continental outflow cannot be excluded.

4 Conclusions

The dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls in marine aerosols collected over the SCS to EIO from 10 March to 26 April 2015 provided new insight on their origins and formation processes in tropical regions.

The spatial distribution of diacids showed high concentrations in nearshore regions (the SCS, Malacca, SLDP) and relatively low in the remote regions (the EIO-WI and EIO-SL). Their molecular distributions were characterized by the dominance of C₂ diacid. The anthropogenic pollutants under the Asian outflow strongly influenced the concentrations of dicarboxylic acids in the SCS and Malacca whereas the oceanic biogenic emissions and subsequent oxidation influenced the distributions of those species in the EIO-WI and EIO-SL. A close relationship in the temporal variability of C₉, C₆ and total concentration of short-chain diacids (C₂–C₄) implied that the in-situ secondary formation of long-chain (including C₃ and C₄) diacids and related compounds was significant rather than the photochemical breakdown of higher diacids to their lower homologues over the SCS to EIO, despite significant aging of the air masses transported from the terrestrial region, due to enhanced emissions of VOCs from marine biota over the region. This finding was also supported by good linear relationships among C₄, hC₄, and C₃ diacids. The positive correlations obtained between C₂% and C₂/C₄ ratios and negative correlations between C₂% and C₃% as well as C₄% indicated that the formation of C₂ from C₄ diacid *via* C₃ is significant. Further, the enhanced emissions of biogenic unsaturated fatty acids and VOCs from the ocean surface and subsequent photochemical oxidation controlled the high and low levels of diacids and related compounds measured in the SCS/Malacca and EIO-WI/EIO-SL, respectively. These results combined with 5-day backward air mass trajectories and the satellite image of Chlorophyll-a suggest the emission of BVOCs from the biologically active ocean surface and their subsequent oxidation reactions were more significant over the SCS and Malacca than that over the EIO-WI and EIO-SL regions. Finally, SLDP samples were affected by all the sources mentioned above. In addition, terrestrial biogenic and anthropogenic emissions might had played significant role on the loading of diacids and related compounds at SLDP. Further work on the chemical characterization of the marine organic aerosols at a molecular level by high-resolution MS such as Orbitrap and FT-ICR MS are needed to obtain the detailed information on sulfur-containing organics from both biogenic and anthropogenic sources.

Data availability. The dataset for this paper is present in the Tables and Figures and is available upon request from the corresponding author (fupingqing@tju.edu.cn).

Competing interests. The authors declare that they have no conflict of interest.

Author contributions. PF designed this research. PF and WH supported this research. Samples were collected by LW. Laboratory analyses were performed by JY and WZ. The manuscript was written by JY, PF and CMP with consultation from other authors.

Acknowledgements. This work was supported the National Natural Science Foundation of China (Grant Nos. 41977183, 41625014, and 41805118). We sincerely thank the captain and crews of the R/V “Shiyan 1” during the open research cruise NORC2015-10 supported by NSFC Shiptime Sharing Project.

References

Aggarwal, S. G., and Kawamura, K.: Molecular distributions and stable carbon isotopic compositions of dicarboxylic acids and related compounds in aerosols from Sapporo, Japan: Implications for photochemical aging during long-range atmospheric transport, *Journal of Geophysical Research: Atmospheres*, 113, 2008.

Bikkina, S., Kawamura, K., Miyazaki, Y., and Fu, P.: High abundances of oxalic, azelaic, and glyoxylic acids and methylglyoxal in the open ocean with high biological activity: Implication for secondary OA formation from isoprene, *Geophysical Research Letters*, 41, 3649-3657, 10.1002/2014gl059913, 2014.

Bikkina, S., Kawamura, K., and Miyazaki, Y.: Latitudinal distributions of atmospheric dicarboxylic acids, oxocarboxylic acids, and α -dicarbonyls over the western North Pacific: Sources and formation pathways, *Journal of Geophysical Research: Atmospheres*, 120, 5010-5035, 10.1002/2014jd022235, 2015.

Carlton, A. G., Turpin, B. J., Lim, H. J., Altieri, K. E., Seitzinger, S., and Lim, H. J.: Link between isoprene and secondary organic aerosol (SOA): Pyruvic acid oxidation yields low volatility organic acids in clouds, *Geophysical Research Letters*, 33, 272-288, 2006.

Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmospheric Chemistry & Physics Discussions*, 9, 4987-5005, 2009.

Carlton, A. M., J. Turpin, B., Altieri, K., Seitzinger, S., Reff, A., Lim, H.-J., and Ervens, B.: Atmospheric oxalic acid and SOA production from glyoxal: Results of aqueous photooxidation experiments, 7588-7602 pp., 2007.

Cavalli, F., Facchini, M. C., Decesari, S., Mircea, M., and 'Acqua, A. D.: Advances in characterization of size-resolved organic matter in marine aerosol over the North Atlantic, *Journal of Geophysical Research Atmospheres*, 109, 89-92, 2004.

Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655-661, 1987.

Charlson, R. J., SCHWARTZ, S. E., HALES, J. M., CESS, R. D., COAKLEY, J. A., HANSEN, J. E., and HOFMANN, D. J.: Climate Forcing by Anthropogenic Aerosols, *Science*, 255, 423-430, 1992.

Chow, J. C., Watson, J. G., Lu, Z., Lowenthal, D. H., Frazier, C. A., Solomon, P. A., Thuillier, R. H., and Magliano, K.: Descriptive analysis of PM 2.5 and PM 10 at regionally representative locations during SJVAQS/AUSPEX, *Atmospheric Environment*, 30, 2079-2112, 1996.

Claeys, M., Wang, W., Vermeylen, R., Kourchev, I., Chi, X., Farhat, Y., Surratt, J. D., Gómez-González, Y., Sciare, J., and Maenhaut, W.: Chemical characterisation of marine aerosol at Amsterdam Island during the austral summer of 2006–2007, *Journal of Aerosol Science*, 41, 13-22, 2010.

Cong, Z. Y., Kawamura, K., Kang, S. C., and Fu, P. Q.: Penetration of biomass-burning emissions from South Asia through the Himalayas: new insights from atmospheric organic acids, *Sci. Rep.*, 5, 9580; DOI:9510.1038/srep09580, 2015.

Donnelly, T. H., Shergold, J. H., and Southgate, P. N.: Anomalous geochemical signals from phosphatic Middle Cambrian rocks in the southern Georgina Basin, Australia, *Sedimentology*, 35, 549-570, 2010.

Ervens, B., Turpin, B., and Weber, R.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmospheric Chemistry and Physics*, 11, 11069-11102, 2011.

Facchini, M. C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Ceburnis, D., Flanagan, R., and Nilsson, E. D.: Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates, *Geophysical Research Letters*, 35, L17814, 2008.

- Fu, P. Q., Kawamura, K., and Barrie, L. A.: Photochemical and other sources of organic compounds in the Canadian high arctic aerosol pollution during winter-spring, *Environmental Science & Technology*, 43, 286-292, 2009.
- Fu, P. Q., Kawamura, K., and Miura, K.: Molecular characterization of marine organic aerosols collected during a round-the-world cruise, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- 545 Fu, P. Q., Kawamura, K., Usukura, K., and Miura, K.: Dicarboxylic acids, ketocarboxylic acids and glyoxal in the marine aerosols collected during a round-the-world cruise, *Marine Chemistry*, 148, 22-32, 10.1016/j.marchem.2012.11.002, 2013a.
- Fu, P. Q., Kawamura, K., Chen, J., Charrière, B., and Sempéré, R.: Organic molecular composition of marine aerosols over the Arctic Ocean in summer: contributions of primary emission and secondary aerosol formation, *Biogeosciences*, 10, 653-667, 2013b.
- 550 Fu, P. Q., Aggarwal, S. G., Chen, J., Li, J., Sun, Y. L., Wang, Z. F., Chen, H. S., Liao, H., Ding, A. J., Umarji, G. S., Patil, R. S., Chen, Q., and Kawamura, K.: Molecular Markers of Secondary Organic Aerosol in Mumbai, India, *Environmental Science & Technology*, 50, 4659-4667, 10.1021/acs.est.6b00372, 2016.
- Hatakeyama, S., Ohno, M., Weng, J., Takagi, H., and Akimoto, H.: Mechanism for the formation of gaseous and particulate products from ozone-cycloalkene reactions in air, *Environmental Science & Technology*, 21, 52-57, 1987.
- 555 Haywood, and M., J.: Tropospheric Aerosol Climate Forcing in Clear-Sky Satellite Observations over the Oceans, *Science*, 283, 1299-1303, 1999.
- Hegde, P., and Kawamura, K.: Seasonal variations of water-soluble organic carbon, dicarboxylic acids, ketocarboxylic acids, and α -dicarbonyls in Central Himalayan aerosols, *Atmospheric Chemistry and Physics*, 12, 6645-6665, 10.5194/acp-12-6645-2012, 2012.
- 560 Ho, K., Cao, J., Lee, S., Kawamura, K., Zhang, R., Chow, J. C., and Watson, J. G.: Dicarboxylic acids, ketocarboxylic acids, and dicarbonyls in the urban atmosphere of China, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Ho, K., Lee, S., Ho, S. S. H., Kawamura, K., Tachibana, E., Cheng, Y., and Zhu, T.: Dicarboxylic acids, ketocarboxylic acids, α -dicarbonyls, fatty acids, and benzoic acid in urban aerosols collected during the 2006 Campaign of Air Quality Research in Beijing (CAREBeijing-2006), *Journal of geophysical research: atmospheres*, 115, 2010.
- 565 Hoffman, E. J., and Duce, R. A.: Organic carbon in marine atmospheric particulate matter: Concentration and particle size distribution, *Geophysical Research Letters*, 4, 449-452, 2013.
- Hoque, M. M. M., and Kawamura, K.: Longitudinal distributions of dicarboxylic acids, ω -oxoacids, pyruvic acid, α -dicarbonyls, and fatty acids in the marine aerosols from the central Pacific including equatorial upwelling, *Global Biogeochemical Cycles*, 30, 534-548, 10.1002/2015gb005346, 2016.
- 570 Hoque, M. M. M., Kawamura, K., and Uematsu, M.: Spatio-temporal distributions of dicarboxylic acids, ω -oxocarboxylic acids, pyruvic acid, α -dicarbonyls and fatty acids in the marine aerosols from the North and South Pacific, *Atmospheric Research*, 185, 158-168, 10.1016/j.atmosres.2016.10.022, 2017.
- Huang, W. T. K., Ickes, L., Tegen, I., Rinaldi, M., Ceburnis, D., and Lohmann, U.: Global relevance of marine organic aerosol as ice nucleating particles, *Atmospheric Chemistry & Physics*, 18, 2018.
- 575 Iii, D. J. E., and Duce, R. A.: On the global flux of atmospheric sea salt, *Journal of Geophysical Research Oceans*, 93, 14079-14088, 1988.
- Jung, J., and Kawamura, K.: Springtime carbon emission episodes at the Gosan background site revealed by total carbon, stable carbon isotopic composition, and thermal characteristics of carbonaceous particles, *Atmospheric Chemistry and Physics*, 11, 10911-10928, 2011.
- 580 Kanakidou, M., Seinfeld, J., Pandis, S., Barnes, I., Dentener, F., Facchini, M., Dingenen, R. V., Ervens, B., Nenes, A., and Nielsen, C.: Organic aerosol and global climate modelling: a review, *Atmospheric Chemistry and Physics*, 5, 1053-1123, 2005.
- Kang, M. J., Yang, F., Ren, H., Zhao, W. Y., Zhao, Y., Li, L. J., Yan, Y., Zhang, Y. J., Lai, S. C., Zhang, Y. Y., Yang, Y., Wang, Z. F., Sun, Y. L., and Fu, P. Q.: Influence of continental organic aerosols to the marine atmosphere over the East China

585 Sea: Insights from lipids, PAHs and phthalates, *Science of the Total Environment*, 607, 339-350, 10.1016/j.scitotenv.2017.06.214, 2017.

Kawamura, K., and Gagosian, R. B.: Implications of ω -oxocarboxylic acids in the remote marine atmosphere for photo-oxidation of unsaturated fatty acids, *Nature*, 325, 330-332, 1987.

590 Kawamura, K., and Kaplan, I. R.: Motor exhaust as a primary source for dicarboxylic acids in Los Angeles ambient air, *Environ.sci.technol*, 21, 105-110, 1987.

Kawamura, K.: Identification of C2-C10. ω -oxocarboxylic acids, pyruvic acid, and C2-C3. α -dicarbonyls in wet precipitation and aerosol samples by capillary GC and GC/MS, *Analytical Chemistry*, 65, 3505-3511, 1993.

Kawamura, K., and Ikushima, K.: Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere, *Environmental Science & Technology*, 27, 2227-2235, 1993.

595 Kawamura, K., Kasukabe, H., and Barrie, L. A.: Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in arctic aerosols: One year of observations, *Atmospheric Environment*, 30, 1709-1722, 1996a.

Kawamura, K., Seméré, R., Imai, Y., Fujii, Y., and Hayashi, M.: Water soluble dicarboxylic acids and related compounds in Antarctic aerosols, *Journal of Geophysical Research Atmospheres*, 101, 18721-18728, 1996b.

600 Kawamura, K., and Sakaguchi, F.: Molecular distribution of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics, *Journal of Geophysical Research Atmospheres*, 104, 3501-3509, 1999.

Kawamura, K., Kobayashi, M., Tsubonuma, N., Mochida, M., Watanabe, T., and Lee, M.: Organic and inorganic compositions of marine aerosols from East Asia: Seasonal variations of water-soluble dicarboxylic acids, major ions, total carbon and nitrogen, and stable C and N isotopic composition, in: *The Geochemical Society Special Publications*, Elsevier, 243-265, 2004.

605 Kawamura, K., and Yasui, O.: Diurnal changes in the distribution of dicarboxylic acids, ketocarboxylic acids and dicarbonyls in the urban Tokyo atmosphere, *Atmospheric Environment*, 39, 1945-1960, 2005.

Kawamura, K., Kasukabe, H., and Barrie, L. A.: Secondary formation of water-soluble organic acids and α -dicarbonyls and their contributions to total carbon and water-soluble organic carbon: Photochemical aging of organic aerosols in the Arctic spring, *Journal of Geophysical Research: Atmospheres*, 115, 2010.

610 Kawamura, K., Tachibana, E., Okuzawa, K., Aggarwal, S., Kanaya, Y., and Wang, Z.: High abundances of water-soluble dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls in the mountaintop aerosols over the North China Plain during wheat burning season, *Atmospheric chemistry and physics*, 13, 8285-8302, 2013.

615 Kawamura, K., and Bikkina, S.: A review of dicarboxylic acids and related compounds in atmospheric aerosols: Molecular distributions, sources and transformation, *Atmospheric Research*, 170, 140-160, 10.1016/j.atmosres.2015.11.018, 2016.

Kawamura, K., Hoque, M. M. M., Bates, T. S., and Quinn, P. K.: Molecular distributions and isotopic compositions of organic aerosols over the western North Atlantic: Dicarboxylic acids, related compounds, sugars, and secondary organic aerosol tracers, *Organic Geochemistry*, 113, 229-238, 10.1016/j.orggeochem.2017.08.007, 2017.

620 Kumar, A., Sudheer, A., and Sarin, M.: Chemical characteristics of aerosols in MABL of Bay of Bengal and Arabian Sea during spring inter-monsoon: a comparative study, *Journal of earth system science*, 117, 325-332, 2008.

Kundu, S., Kawamura, K., Andreae, T., Hoffer, A., and Andreae, M.: Molecular distributions of dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls in biomass burning aerosols: implications for photochemical production and degradation in smoke layers, *Atmospheric Chemistry and Physics*, 10, 2209-2225, 2010a.

625 Kundu, S., Kawamura, K., and Lee, M.: Seasonal variations of diacids, ketoacids, and α -dicarbonyls in aerosols at Gosan, Jeju Island, South Korea: Implications for sources, formation, and degradation during long-range transport, *Journal of Geophysical Research*, 115, 10.1029/2010jd013973, 2010b.

Kunwar, B., and Kawamura, K.: Seasonal distributions and sources of low molecular weight dicarboxylic acids, ω -oxocarboxylic acids, pyruvic acid, α -dicarbonyls and fatty acids in ambient aerosols from subtropical Okinawa in the western Pacific Rim, *Environmental Chemistry*, 11, 673-689, 2014.

- 630 Kunwar, B., Torii, K., and Kawamura, K.: Springtime influences of Asian outflow and photochemistry on the distributions of diacids, oxoacids and α -dicarbonyls in the aerosols from the western North Pacific Rim, *Tellus B: Chemical and Physical Meteorology*, 69, 10.1080/16000889.2017.1369341, 2017.
- Lau, K.-M., Wu, H.-T., and Yang, S.: Hydrologic processes associated with the first transition of the Asian summer monsoon: A pilot satellite study, *Bulletin of the American Meteorological Society*, 79, 1871-1882, 1998.
- 635 Li, W. J., Xu, L., Liu, X. H., Zhang, J. C., Lin, Y. T., Yao, X. H., Gao, H. W., Zhang, D. Z., Chen, J. M., Wang, W. X., Harrison, R. M., Zhang, X. Y., Shao, L. Y., Fu, P. Q., Nenes, A., and Shi, Z. B.: Air pollution-aerosol interactions produce more bioavailable iron for ocean ecosystems, *Science Advances*, 3, 10.1126/sciadv.1601749, 2017.
- Lim, H.-J., Carlton, A. G., and Turpin, B. J.: Isoprene forms secondary organic aerosol through cloud processing: Model simulations, *Environmental science & technology*, 39, 4441-4446, 2005.
- 640 Liu, K. K., Chao, S. Y., Shaw, P. T., Gong, G. C., Chen, C. C., and Tang, T. Y.: Monsoon-forced chlorophyll distribution and primary production in the South China Sea: observations and a numerical study, *Deep Sea Research Part I Oceanographic Research Papers*, 49, 1387-1412, 2002.
- Matsunaga, S., Kawamura, K., Nakatsuka, T., and Ohkouchi, N.: Preliminary study on laboratory photochemical formation of low molecular weight dicarboxylic acids from unsaturated fatty acid (oleic acid), *Org. Geochem*, 14, 19-25, 1999.
- 645 Miyazaki, Y., Aggarwal, S. G., Singh, K., Gupta, P. K., and Kawamura, K.: Dicarboxylic acids and water-soluble organic carbon in aerosols in New Delhi, India, in winter: Characteristics and formation processes, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Miyazaki, Y., Kawamura, K., and Sawano, M.: Size distributions and chemical characterization of water-soluble organic aerosols over the western North Pacific in summer, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- 650 Mochida, M., Kawabata, A., Kawamura, K., Hatsushika, H., and Yamazaki, K.: Seasonal variation and origins of dicarboxylic acids in the marine atmosphere over the western North Pacific, *Journal of Geophysical Research: Atmospheres*, 108, 2003a.
- Mochida, M., Kawamura, K., Umemoto, N., Kobayashi, M., Matsunaga, S., Lim, H. J., Turpin, B. J., Bates, T. S., and Simoneit, B. R.: Spatial distributions of oxygenated organic compounds (dicarboxylic acids, fatty acids, and levoglucosan) in marine aerosols over the western Pacific and off the coast of East Asia: Continental outflow of organic aerosols during the ACE-Asia campaign, *Journal of Geophysical Research: Atmospheres*, 108, 2003b.
- 655 Narukawa, M., Kawamura, K., Takeuchi, N., and Nakajima, T.: Distribution of dicarboxylic acids and carbon isotopic compositions in aerosols from 1997 Indonesian forest fires, *Geophysical Research Letters*, 26, 3101-3104, 1999.
- Nguyen, T. B., Bateman, A. P., Bones, D. L., Nizkorodov, S. A., Laskin, J., and Laskin, A.: High-resolution mass spectrometry analysis of secondary organic aerosol generated by ozonolysis of isoprene, *Atmospheric Environment*, 44, 1032-1042, 2010.
- 660 O'Dowd, C. D., and De Leeuw, G.: Marine aerosol production: a review of the current knowledge, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 365, 1753-1774, 2007.
- Pavuluri, C., Kawamura, K., Aggarwal, S., and Swaminathan, T.: Characteristics, seasonality and sources of carbonaceous and ionic components in the tropical aerosols from Indian region, *Atmospheric Chemistry and Physics*, 11, 8215-8230, 2011.
- 665 Pavuluri, C. M., Kawamura, K., and Swaminathan, T.: Water-soluble organic carbon, dicarboxylic acids, ketoacids, and α -dicarbonyls in the tropical Indian aerosols, *Journal of Geophysical Research Atmospheres*, 115, 10.1029/2009JD012661, 2010.
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Atmosphere - Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119-2124, 2001.
- 670 Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J. T., Washington, W. M., Fu, Q., Sikka, D. R., and Wild, M.: Atmospheric brown clouds: Impacts on South Asia climate and hydrological cycle, *PNAS*, 102, 5326-5333, 2005.
- Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Fuzzi, S., Ceburnis, D., O'Dowd, C. D., Sciare, J., Burrows, J. P., and Vrekoussis, M.: Evidence of a natural marine source of oxalic acid and a possible link to glyoxal, *Journal of Geophysical Research: Atmospheres*, 116, 2011.

- 675 Rogge, W. F., Mazurek, M. A., Hildemann, L. M., Cass, G. R., and Simoneit, B. R.: Quantification of urban organic aerosols at a molecular level: identification, abundance and seasonal variation, *Atmospheric Environment. Part A. General Topics*, 27, 1309-1330, 1993.
- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R.: Measurement of emissions from air pollution sources. 3. C1-C29 organic compounds from fireplace combustion of wood, *Environmental Science & Technology*, 35, 1716-1728, 2001.
- 680 Schulz, M., Prospero, J. M., Baker, A. R., Dentener, F., Ickes, L., Liss, P. S., Mahowald, N. M., Nickovic, S., Garc a-A-Pando, C. P., and Rodr a-Guez, S.: Atmospheric Transport and Deposition of Mineral Dust to the Ocean: Implications for Research Needs, *Environmental Science & Technology*, 46, 10390-10404, 2012.
- Semp ere, R., and Kawamura, K.: Comparative distributions of dicarboxylic acids and related polar compounds in snow, rain and aerosols from urban atmosphere, *Atmospheric Environment*, 28, 449-459, 1994.
- 685 Semp er e, R., and Kawamura, K.: Trans-hemispheric contribution of C2-C10 α , ω -dicarboxylic acids, and related polar compounds to water-soluble organic carbon in the western Pacific aerosols in relation to photochemical oxidation reactions, *Global Biogeochemical Cycles*, 17, n/a-n/a, 10.1029/2002gb001980, 2003.
- Shaw, S. L., Gantt, B., and Meskhidze, N.: Production and emissions of marine isoprene and monoterpenes: a review, *Advances in Meteorology*, 2010, 2010.
- 690 Simoneit, B. R., Kobayashi, M., Mochida, M., Kawamura, K., Lee, M., Lim, H. J., Turpin, B. J., and Komazaki, Y.: Composition and major sources of organic compounds of aerosol particulate matter sampled during the ACE-Asia campaign, *Journal of Geophysical Research: Atmospheres*, 109, 2004.
- Song, J., Zhao, Y., Zhang, Y., Fu, P., Zheng, L., Yuan, Q., Wang, S., Huang, X., Xu, W., Cao, Z., Gromov, S., and Lai, S.: Influence of biomass burning on atmospheric aerosols over the western South China Sea: Insights from ions, carbonaceous fractions and stable carbon isotope ratios, *Environ Pollut*, 242, 1800-1809, 10.1016/j.envpol.2018.07.088, 2018.
- 695 Sorooshian, A., Varutbangkul, V., Brechtel, F. J., Ervens, B., Feingold, G., Bahreini, R., Murphy, S. M., Holloway, J. S., Atlas, E. L., and Buzorius, G.: Oxalic acid in clear and cloudy atmospheres: Analysis of data from International Consortium for Atmospheric Research on Transport and Transformation 2004, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Sorooshian, A., Lu, M.-L., Brechtel, F. J., Jonsson, H., Feingold, G., Flagan, R. C., and Seinfeld, J. H.: On the source of organic acid aerosol layers above clouds, *Environmental science & technology*, 41, 4647-4654, 2007.
- 700 Sorooshian, A., Padr o, L. T., Nenes, A., Feingold, G., McComiskey, A., Hersey, S. P., Gates, H., Jonsson, H. H., Miller, S. D., and Stephens, G. L.: On the link between ocean biota emissions, aerosol, and maritime clouds: Airborne, ground, and satellite measurements off the coast of California, *Global Biogeochemical Cycles*, 23, 2009.
- Sorooshian, A., Murphy, S. M., Hersey, S., Bahreini, R., Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Constraining the contribution of organic acids and AMSm/z44 to the organic aerosol budget: On the importance of meteorology, aerosol hygroscopicity, and region, *Geophysical Research Letters*, 37, n/a-n/a, 10.1029/2010gl044951, 2010.
- 705 Tan, S. C., Shi, G. Y., Shi, J. H., Gao, H. W., and Yao, X. H.: Correlation of Asian dust with chlorophyll and primary productivity in the coastal seas of China during the period from 1998 to 2008, *Journal of Geophysical Research-Biogeosciences*, 116, 10.1029/2010jg001456, 2011.
- 710 Turpin, B. J., and Huntzicker, J. J.: Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS, *Atmospheric Environment*, 29, 3527-3544, 1995.
- Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Burrows, S. M., Ceburnis, D., and DeMott, P. J.: Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations, *Atmospheric Chemistry and Physics*, 17, 3637-3658, 2017.
- 715 Volkamer, R., San Martini, F., Molina, L. T., Salcedo, D., Jimenez, J. L., and Molina, M. J.: A missing sink for gas-phase glyoxal in Mexico City: Formation of secondary organic aerosol, *Geophysical Research Letters*, 34, 2007.
- Wang, G., Kawamura, K., Cheng, C., Li, J., Cao, J., Zhang, R., Zhang, T., Liu, S., and Zhao, Z.: Molecular distribution and stable carbon isotopic composition of dicarboxylic acids, ketocarboxylic acids, and alpha-dicarbonyls in size-resolved atmospheric particles from Xi'an City, China, *Environ Sci Technol*, 46, 4783-4791, 10.1021/es204322c, 2012.

- 720 Wang, G., Cheng, C., Meng, J., Huang, Y., Li, J., and Ren, Y.: Field observation on secondary organic aerosols during Asian dust storm periods: Formation mechanism of oxalic acid and related compounds on dust surface, *Atmospheric Environment*, 113, 169-176, 10.1016/j.atmosenv.2015.05.013, 2015.
- Wang, H., and Kawamura, K.: Stable carbon isotopic composition of low-molecular-weight dicarboxylic acids and ketoacids in remote marine aerosols, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- 725 Wang, H., Kawamura, K., and Yamazaki, K.: Water-soluble dicarboxylic acids, ketoacids and dicarbonyls in the atmospheric aerosols over the Southern Ocean and western Pacific Ocean, *Journal of Atmospheric Chemistry*, 53, 43-61, 2006.
- Warneck, P.: In-cloud chemistry opens pathway to the formation of oxalic acid in the marine atmosphere, *Atmospheric Environment*, 37, 2423-2427, 2003.
- 730 Yang, L., Ray, M. B., and Liya, E. Y.: Photooxidation of dicarboxylic acids—Part II: Kinetics, intermediates and field observations, *Atmospheric Environment*, 42, 868-880, 2008.
- Yao, T., Wu, F., Ding, L., Sun, J., Zhu, L., Piao, S., Deng, T., Ni, X., Zheng, H., and Ouyang, H.: Multispherical interactions and their effects on the Tibetan Plateau's earth system: a review of the recent researches, *Nat. Sci. Rev.*, 2, 468-488, 2015.
- 735 Zhang, M., Chen, J., Wang, T., Cheng, T., Lin, L., Bhatia, R., and Hanvey, M.: Chemical characterization of aerosols over the Atlantic Ocean and the Pacific Ocean during two cruises in 2007 and 2008, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- Zhao, D., Buchholz, A., Kortner, B., Schlag, P., Rubach, F., Fuchs, H., Kiendler-Scharr, A., Tillmann, R., Wahner, A., and Watne, Å.: Cloud condensation nuclei activity, droplet growth kinetics, and hygroscopicity of biogenic and anthropogenic secondary organic aerosol (SOA), *Atmospheric chemistry and physics*, 16, 1105-1121, 2016a.
- 740 Zhao, W. Y., Kawamura, K., Yue, S. Y., Wei, L. F., Ren, H., Yan, Y., Kang, M. J., Li, L. J., Ren, L. J., Lai, S. C., Li, J., Sun, Y. L., Wang, Z. F., and Fu, P. Q.: Molecular distribution and compound-specific stable carbon isotopic composition of dicarboxylic acids, oxocarboxylic acids and alpha-dicarbonyls in PM_{2.5} from Beijing, China, *Atmospheric Chemistry and Physics*, 18, 2749-2767, 10.5194/acp-18-2749-2018, 2018.
- 745 Zhao, Y., Zhang, Y. Y., Fu, P. Q., Ho, S. S. H., Ho, K. F., Liu, F. B., Zou, S. C., Wang, S., and Lai, S. C.: Non-polar organic compounds in marine aerosols over the northern South China Sea: Influence of continental outflow, *Chemosphere*, 153, 332-339, 10.1016/j.chemosphere.2016.03.069, 2016b.
- Zhuang, G. S., Yi, Z., Duce, R. A., and Brown, P. R.: Link between Iron and Sulfur Cycles Suggested by Detection of Fe(II) in Remote Marine Aerosols, *Nature*, 355, 537-539, 1992.

750

Table 1. Summary of concentrations (ng m^{-3}) of dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls as well as concentrations ($\mu\text{g m}^{-3}$) of EC and OC in TSP samples collected during the NORC2015-10 cruise over the South China Sea to the East Indian Ocean.

Species	SCS (n = 27)		EIO-WI (n = 20)		EIO-SL (n = 28)		Malacca (n = 9)		SLDP (n = 3)	
	Range	Avg±SD	Range	Avg±SD	Range	Avg±SD	Range	Avg±SD	Range	Avg±SD
Dicarboxylic acids										
Oxalic, C ₂	30.3–626	233±143	4.43–70.1	18.2±18.4	5.21–232	58.2±55.3	14.5–373	145±112	179–454	303±140
Malonic, C ₃	2.14–87.9	36±23	0.34–11.4	2.09±2.52	0.76–42.1	8.19±9.95	2.57–74.0	24.2±25.4	25.9–55.4	42.2±15.0
Succinic, C ₄	1.73–50.4	18±14	0.23–6.4	1.35±1.65	0.57–20.6	4.33±4.86	1.18–72.9	18.0±22.7	14.1–78.5	45.2±32.2
Glutaric, C ₅	0.23–8.52	2.60±2.15	bdl–0.79	0.23±0.19	bdl–1.69	0.48±0.46	bdl–9.53	3.49±3.70	2.94–14.6	8.82±5.85
Adipic, C ₆	0.13–4.09	1.11±0.93	0.17–0.83	0.42±0.22	0.03–1.18	0.31±0.29	0.01–3.59	1.28±1.36	1.95–7.17	4.76±2.63
Pimelic, C ₇	bdl–1.88	0.58±0.60	bdl–0.22	0.03±0.07	bdl–0.50	0.12±0.17	bdl–1.30	0.43±0.46	bdl–0.45	0.28±0.25
Suberic, C ₈	bdl	bdl	bdl–0.09	0.004±0.0	bdl–14.3	0.51±2.71	bdl	bdl	bdl	bdl
Azelaic, C ₉	bdl–6.97	1.82±1.57	bdl–2.22	0.45±0.49	0.09–15.1	1.25±2.75	0.32–3.96	1.82±1.29	11.7–140	68.9±65.1
Sebacic, C ₁₀	bdl–2.69	0.28±0.53	bdl–0.82	0.29±0.24	bdl–9.06	0.57±1.68	bdl–1.10	0.58±0.46	0.66–3.43	2.00±1.39
Undecanedioic, C ₁₁	bdl–0.16	0.02±0.05	bdl–0.69	0.16±0.15	bdl–0.38	0.13±0.10	bdl–0.38	0.20±0.15	bdl–1.64	0.62±0.89
Dodecanedioic, C ₁₂	bdl	bdl	bdl–0.35	0.02±0.08	bdl	bdl	bdl–0.29	0.06±0.11	bdl	bdl
Methylmalonic, iC ₄	bdl–2.26	0.95±0.65	bdl–0.29	0.01±0.06	bdl–0.82	0.14±0.23	bdl–3.29	0.93±1.06	1.02–2.47	1.94±0.79
Methylsuccinic, iC ₅	0.30–3.37	1.46±0.98	0.08–0.54	0.35±0.12	0.13–1.18	0.56±0.30	0.21–3.56	1.63±1.28	2.61–10.1	6.99±3.89
Methylglutaric, iC ₆	bdl–0.68	0.05±0.18	bdl–0.18	0.02±0.05	bdl	bdl	bdl–0.59	0.26±0.25	0.67–3.29	1.84±1.33
Maleic, M	0.12–2.70	0.93±0.67	0.07–0.55	0.19±0.11	0.08–1.02	0.28±0.22	0.04–1.42	0.77±0.59	1.58–4.14	2.56±1.38
Fumaric, F	0.17–2.16	0.74±0.48	0.16–1.45	0.77±0.32	0.05–0.82	0.28±0.22	0.08–1.07	0.39±0.33	0.81–2.49	1.49±0.89
Methylmaleic, mM	0.17–2.13	0.82±0.48	0.17–0.76	0.39±0.15	0.06–0.96	0.32±0.21	0.11–1.33	0.73±0.47	2.26–5.49	3.80±1.62
Phthalic, Ph	0.16–6.57	2.29±1.89	bdl–1.25	0.50±0.34	0.06–2.95	0.77±0.59	0.26–3.56	1.64±1.23	8.63–13.2	10.8±2.31
Isophthalic, iPh	bdl–0.60	0.05±0.16	bdl–0.37	0.02±0.08	bdl–0.22	0.02±0.06	bdl–0.49	0.12±0.20	1.08–2.73	1.99±0.84
Terephthalic, tPh	bdl–2.12	0.22±0.57	bdl–1.25	0.13±0.17	bdl–0.38	0.03±0.09	bdl–0.59	0.10±0.21	bdl–1.76	0.59±1.02
Malic, hC ₄	bdl–2.25	0.62±0.53	bdl–0.72	0.05±0.17	bdl–0.81	0.13±0.23	bdl–1.05	0.31±0.35	0.88–1.95	1.45±0.54
Ketomalonic, kC ₃	bdl–5.84	1.76±1.34	bdl–0.83	0.22±0.22	0.05–2.10	0.54±0.47	0.15–2.19	1.00±0.90	1.15–2.75	2.06±0.82
4-Ketopimelic, kC ₇	bdl–7.75	1.85±1.70	bdl–0.63	0.12±0.19	bdl–1.87	0.45±0.56	0.17–9.27	3.59±3.68	1.24–3.34	2.50±1.10
Subtotal	44.6–759	305±186	8.28–96.3	26.1±23.3	8.55–294	77.6±73.1	21.9–501	207±158	259–708	514±231
Oxocarboxylic acids										
Pyruvic, Pyr	0.01–1.70	0.60±0.42	bdl–0.70	0.30±0.17	0.11–0.75	0.33±0.19	bdl–1.23	0.46±0.44	0.83–1.62	1.13±0.42
Glyoxylic, ωC ₂	0.41–9.23	2.68±1.87	bdl–0.75	0.23±0.23	0.09–2.89	0.97±0.86	0.003–4.92	2.05±2.08	3.56–8.65	6.28±2.56
3-Oxopropanoic, ωC ₃	0.14–2.53	0.86±0.56	0.06–0.65	0.27±0.14	bdl–0.75	0.25±0.26	bdl–2.01	0.56±0.61	0.55–2.16	1.13±0.90
4-Oxobutanoic, ωC ₄	bdl–1.09	0.29±0.41	bdl	bdl	bdl–1.42	0.06±0.27	bdl–2.66	0.72±1.02	1.41–9.27	5.10±3.95
5-Oxopentanoic, ωC ₅	bdl–0.09	0.01±0.03	0.05–0.39	0.13±0.08	bdl–0.15	0.04±0.03	bdl–0.15	0.04±0.05	0.07–0.21	0.16±0.07
7-Oxoheptanoic, ωC ₇	bdl–0.23	0.23±0	bdl	bdl	bdl	bdl	bdl	bdl	bdl	bdl
8-Oxo-octanoic, ωC ₈	bdl–3.94	1.37±1.44	bdl	bdl	bdl	bdl	bdl–0.23	0.07±0.10	bdl–16.1	8.47±8.07
9-Oxononanoic, ωC ₉	bdl–1.89	0.69±0.65	bdl–0.29	0.03±0.07	bdl–1.41	0.19±0.36	bdl–0.88	0.32±0.32	0.80–2.38	1.72±0.83
Subtotal	1.48–13.2	6.51±3.99	0.16–2.01	0.96±0.46	0.23–5.06	1.85±1.51	0.29–10.4	4.22±3.75	7.46–40.3	24.0±16.4
α-Dicarbonyls										
Glyoxal, Gly	0.05–7.70	2.28±2.26	0.20–3.20	1.64±0.67	0.11–2.02	0.87±0.40	0.26–0.87	0.52±0.23	2.70–7.04	4.15±2.50
Methylglyoxal, MeGly	0.29–20.0	2.97±3.72	0.88–3.01	1.66±0.54	0.09–7.45	1.22±1.31	0.93–4.01	1.81±0.93	2.31–4.45	3.20±1.11
Subtotal	0.56–20.2	5.25±4.46	1.08–5.29	3.31±1.01	0.95–8.86	2.08±1.47	1.18–4.79	2.34±1.06	5.04–11.5	7.35±3.58
Carbonaceous components										
OC	0.18–2.63	1.26±0.67	0.02–0.66	0.27±0.19	0.06–1.69	0.59±0.45	0.28–4.44	2.02±1.54	3.57–20.9	12.8±8.73
EC	0.05–0.75	0.27±0.15	0.01–0.13	0.05±0.05	0.04–0.91	0.21±0.18	0.07–0.63	0.24±0.18	1.67–7.57	4.24±3.02
WSOC	0.22–2.31	1.05±0.65	0.05–0.58	0.15±0.12	0.06–1.00	0.35±0.25	0.09–2.72	1.15±0.92	1.12–7.44	4.43±3.17



bdl: below detection limit, which is ca. 0.005 ng m^{-3} for the target compounds.

Table 2. The fractions of diacids and related compounds in WSOC and OC in marine aerosols collected during the NORC2015-10 cruise.


Components	Relative Abundance (%)	
	in WSOC	in OC
Oxalic, C ₂	0.61–14.3 (4.78±2.63)	0.28–12.2 (3.52±2.34)
Malonic, C ₃	0.14–2.82 (0.85±0.54)	0.07–2.41 (0.63±0.50)
Succinic, C ₄	0.10–1.94 (0.54±0.37)	0.04–1.86 (0.41±0.34)
ΣC ₂ –C ₄	1.02–22.0 (7.50±4.01)	0.47–18.8 (5.55±3.68)
ΣDiacids	2.00–32.4 (12.0±5.75)	0.92–27.7 (8.84±5.61)
ΣOxocarboxylic acids	0.03–1.22 (0.34±0.20)	0.01–1.93 (0.26±0.26)
Σα-Dicarbonyls	0.02–3.12 (0.60±0.68)	0.02–8.95 (0.47±1.08)

Table 3. Diagnostic mass ratios of selected diacids and related compounds in TSP samples collected during the NORC2015-10 cruise.


Mass Ratio	SCS	EIO-WI	EIO-SL	Malacca	SLDP
	Avg ± SD				
C ₂ /C ₄	16.2±8.14	17.1±10.6	16.1±6.72	15.1±12.4	8.28±3.80
C ₃ /C ₄	2.18±0.95	1.79±0.90	2.05±0.80	1.99±1.14	1.23±0.63
C ₂ /ωC ₂	101±69.8	146±220	69.7±41.2	721±1678	48.1±5.79
C ₂ /Pyr	580±499	70.3±63.7	193±191	294±254	267±71.0
C ₂ /Σ(C ₂ -C ₁₂)	0.80±0.05	0.76±0.07	0.78±0.07	0.76±0.11	0.65±0.12
C ₃ /Σ(C ₂ -C ₁₂)	0.12±0.04	0.09±0.02	0.11±0.03	0.12±0.05	0.09±0.02
C ₄ /Σ(C ₂ -C ₁₂)	0.06±0.02	0.06±0.03	0.06±0.04	0.08±0.06	0.09±0.03
C ₆ /C ₉	0.76±0.93	1.17±0.76	0.49±0.52	0.49±0.39	0.11±0.07
Ph/C ₉	2.02±4.26	1.66±1.33	1.22±1.19	0.90±0.21	0.34±0.34
M/F	1.33±0.52	0.29±0.19	1.30±0.59	2.08±1.72	1.77±0.17
C ₂ /MeGly	129±156	11.3±10.6	57.9±40.8	81.9±55.5	92.1±13.1
C ₂ /Gly	681±1466	14.6±15.2	127±196	295±184	77.4±21.2
Gly/MeGly	0.86±0.67	1.02±0.45	1.43±2.31	0.31±0.12	1.24±0.32




765 **Figure 1.** (a) Cruise tracks used for aerosols collection in the South China Sea and the East Indian Ocean during 10th March
 770 to 26th April 2015. The area where the red line is located is South China Sea (SCS), the yellow line is the East Indian Ocean
 near western Indonesia (EIO-WI), the green line is the East Indian Ocean near Sri Lanka (EIO-SL), the blue line is Malacca,
 and the red solid origins are the collection positions of three samples of Sri Lanka docking point (SLDP). The base map
 (©MATLAB) reflects an average composite of Chl a concentrations in March–April 2015 obtained from the NASA website at
https://modis.gsfc.nasa.gov/data/dataproduct/chlor_a.php. (b) Air mass backward trajectories for the sampling days at arrival
 heights 100 m; the map was produced by the software © Meteoinfo.




775 **Figure 2.** Molecular distributions of dicarboxylic acids and related compounds in TSP samples from (a) SCS, (b) EIO-WI, (c) EIO-SL, (d) Malacca, and (e) SLDP during 10th March to 26th April 2015.




780 **Figure 3.** Daily variations in the concentrations of selected organic acids and water-soluble organic carbon (WSOC) in the TSP aerosols in South China Sea and the East Indian Ocean from 10th March to 26th April 2015 (all parameters are in ng m⁻³ except WSOC is in μg m⁻³ and Chlorophyll a is in mg m⁻³).



785 **Figure 4.** Pie charts showing the percentage contribution of individual diacid to total aliphatic homologous diacids (ΣC_2-C_{12}) in different sea areas aerosols collected during the NORC2015-10 cruise.



790 **Figure 5.** Relative abundances of individual compound class in the total diacids, oxoacids, and α -dicarbonyls in different sampling area aerosols collected during the NORC2015-10 cruise.



795 **Figure 6.** Daily variations in the concentration ratios of (a) C_2/C_4 , (b) C_3/C_4 , (c) $C_2/\Sigma C_2-C_{12}$, (d) $C_2/\omega C_2$, (e) C_6/C_9 , (f) C_6/C_9 , (g) Ph/C_9 , (h) M/F , (i) C_2/Gly , and (j) $C_2/MeGly$ in TSP samples collected during the NORC2015-10 cruise.

5 **Molecular and spatial distributions of dicarboxylic acids,
oxocarboxylic acids and α -dicarbonyls in marine aerosols from
the South China Sea to East Indian Ocean**

Jing Yang^{1,2}, Wanyu Zhao^{2,4}, Lianfang Wei², Qiang Zhang¹, Yue Zhao¹, Wei Hu¹, Libin Wu¹, Xiaodong Li¹, Chandra Mouli Pavuluri¹, Xiaole Pan², Yele Sun², Zifa Wang², Cong-Qiang Liu¹, Kimitaka Kawamura³, and Pingqing Fu¹

10

¹Institute of Surface-Earth System Science, Tianjin University, Tianjin 300072, China

²State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

³Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan

15 ⁴College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China

Correspondence to: Pingqing Fu (fupingqing@tju.edu.cn)

Linear correlations

20 In order to investigate the origins and formation pathways of diacids and related compounds in marine aerosols, we examined the correlations between selected species and also conducted linear regression analyses for mass ratios of the selected chemical species. Moderate correlations were observed for C₆ with C₉ diacids in the EIO-WI and EIO-SL, and significant correlations over the SCS and Malacca regions. Such positive correlations indicate that the C₆ diacid should have also been significantly generated from the biogenic VOCs such as cyclic olefins, in
25 addition to the anthropogenic sources.

Hydroxy succinic acid (malic acid, hC₄), C₄ and C₃ diacids in the SCS correlated well with each other ($p < 0.05$; $R^2 = 0.87$ for hC₄ vs C₃ and $R^2 = 0.78$ for hC₄ vs C₄; Fig. S2a and S2b). The formation of C₃ diacid from C₄ diacid through hC₄ as an intermediate has been proposed by Kawamura and Sakaguchi (1999) and also observed in a laboratory study (Yang et al., 2008). A very good linear relationship of hC₄ diacid with C₄ diacid ($R^2 = 0.85$) and
30 C₂ diacid ($R^2 = 0.78$) has been reported in forest aerosols from central Europe, which have been attributed to *in-situ* photochemical production from biogenic unsaturated fatty acids (Kourtchev et al., 2009). Therefore, the good correlations observed among C₃, C₄ and hC₄ in the SCS implies that they were derived from common but mixed (both biogenic and anthropogenic) sources and the *in-situ* secondary formation might be more intensive, rather than the transformations from C₄ to C₃ via hC₄ diacids. A significant linear correlation was also found between
35 methylmalonic (iC₄) acid with C₃ and C₄ diacids. The branched chain diacids could come from the oxidation of isoprene released from the ocean surface. Such linear correlations also suggest that the diacids and related compounds were photochemical oxidation products of isoprene and biogenic unsaturated fatty acids in the MABL, and also confirm the intensive *in-situ* secondary formation of diacids and related species. A notable feature observed here is the significant ($p < 0.05$) correlations between MeGly, Pyr, and ω C₂ with C₃ and C₄ diacids over Malacca, but not in other regions. Nguyen et al. (2010) found that C₃ and C₄ diacids were formed from isoprene oxidized by
40 ozone based on the laboratory experiment. Furthermore, it has been reported that isoprene emission results in significant enhancement of SOA and O₃ levels at coastal U.S. sites (Gantt et al., 2010). Hence, the loading of C₃ and C₄ diacids also should have been influenced by not only the unsaturated fatty acids but also the isoprene, at least in Malacca region.

45 The C₂/(Σ C₂-C₁₀) ratios showed negative relationships with C₃ and C₄ diacids over the SCS to EIO (Fig. S3a and S3b). This phenomenon again confirms the formation of C₂ diacid from C₃ and C₄ diacids by their photochemical transformations during long-range atmospheric transport. It is also worth noting that the C₄/(Σ C₂-C₁₀) ratios showed a linear relationship with glutaric (C₅) acid in all regions, except the EIO-WI (Fig. S3c). Additionally, a similar variation trend is observed for the concentrations of C₅ and C₉ diacids in all regions, although

50 they were not significant (C_9 ; Fig. S3d). Such positive relations reveal that they all should have been simultaneously produced significantly by in-situ photochemical reactions over the oceanic region, although we do not preclude the photochemical transformations of long-chain diacids to their lower homologues including C_6 - C_4 diacids. Thus, the observed linear correlations among the relative abundances of C_2 , C_3 , C_4 , C_5 , and C_9 infer that the long-chain as well as C_3 and C_4 diacids were mainly derived from biogenic emissions followed by the in-situ secondary reactions
55 whereas the C_2 diacid is mainly produced by the photochemical transformation of their higher over the SCS to EIO. The long-range transported air masses from the continental region that should be enriched with the anthropogenic emissions including oxidant species should have been promoted the oxidation of unsaturated fatty acids and BVOCs, resulting the enhanced levels of diacids and related compounds over the remote oceanic region.


60

Table S1. Correlation matrices of selected diacids and related compounds in TSP collected during the NORC2015-10 cruise.

Correlation	C ₂	C ₃	C ₄	C ₅	C ₆	C ₉	Ph	Pyr	ω C ₂	MeGly
SCS (n=27)										
C ₂	1.00									
C ₃	0.85 **	1.00								
C ₄	0.79 **	0.88 **	1.00							
C ₅	0.86 **	0.73 **	0.87 **	1.00						
C ₆	0.87 **	0.67 **	0.75 **	0.95 **	1.00					
C ₉	0.67 **	0.38	0.36	0.70 **	0.78 **	1.00				
Ph	0.70 **	0.54 **	0.72 **	0.86 **	0.84 **	0.59 **	1.00			
Pyr	0.50 **	0.44 *	0.55 **	0.52 **	0.43 *	0.19	0.52 **	1.00		
ω C ₂	0.88 **	0.71 **	0.68 **	0.80 **	0.79 **	0.68 **	0.69 **	0.66 **	1.00	
MeGly	0.65 **	0.40 *	0.30	0.42*	0.49 *	0.50 **	0.50 **	0.59 **	0.80 **	1.00
EIO-WI (n=20)										
C ₂	1.00									
C ₃	0.90 **	1.00								
C ₄	0.89 **	0.94 **	1.00							
C ₅	0.85 **	0.83 **	0.76 **	1.00						
C ₆	0.26	0.37	0.41	0.41	1.00					
C ₉	0.34	0.36	0.36	0.39	0.51 *	1.00				
Ph	0.11	0.29	0.18	0.51 *	0.67 **	0.40	1.00			
Pyr	0.10	0.13	0.14	0.17	0.04	0.33	0.04	1.00		
ω C ₂	0.68 **	0.51 *	0.53 *	0.71 **	-0.04	0.40	0.12	0.23	1.00	
MeGly	0.11	0.13	0.13	0.00	0.21	0.28	-0.11	0.61 **	-0.12	1.00
EIO-SL (n=28)										
C ₂	1.00									
C ₃	0.89 **	1.00								
C ₄	0.86 **	0.94 **	1.00							
C ₅	0.73 **	0.75 **	0.78 **	1.00						
C ₆	0.41 *	0.31	0.49 **	0.44	1.00					
C ₉	0.31	0.14	0.20	0.31	0.39 *	1.00				
Ph	0.53 **	0.35	0.43 *	0.52 **	0.70 **	0.79 **	1.00			
Pyr	0.40 *	0.20	0.38 *	0.24	0.38 *	0.22	0.27	1.00		
ω C ₂	0.79 **	0.55 **	0.65 **	0.61 **	0.50 **	0.51 **	0.69 **	0.64 **	1.00	
MeGly	0.21	0.21	0.09	0.11	0.14	0.07	0.19	0.32	0.14	1.00
Malacca (n=9)										
C ₂	1.00									
C ₃	0.65	1.00								
C ₄	0.45	0.88 **	1.00							
C ₅	0.69 *	0.76 *	0.87 **	1.00						
C ₆	0.89 **	0.64	0.56	0.85 **	1.00					
C ₉	0.87 **	0.71 *	0.60	0.81 **	0.96 **	1.00				
Ph	0.84 **	0.87 **	0.80 *	0.89 **	0.90 **	0.94 **	1.00			
Pyr	0.55	0.71 *	0.83 **	0.79 *	0.60	0.56	0.71 *	1.00		
ω C ₂	0.80 **	0.75 *	0.80 **	0.97 **	0.91 **	0.86 **	0.91 **	0.84 **	1.00	
MeGly	0.36	0.79 *	0.94 **	0.83 **	0.50	0.52	0.74 *	0.67	0.71 *	1.00


** : At 0.01 level (two-tailed), the correlation is significant.

* : At 0.05 level (two-tailed), the correlation is significant.




65

Figure S1. Chemical structures of selected dicarboxylic acids and other major organic compounds detected in the marine aerosols.



70

Figure S2. Scatterplots for (a) malic versus succinic acids, (b) malonic versus malic acids in aerosols collected over the SCS; (c) relative abundance of oxalic acid in total diacid mass versus C₂/C₄ ratios in aerosols collected over the all sea areas.



75

Figure S3. Scatterplots between the relative abundances of C_2 in their total aliphatic homologues (ΣC_2-C_{10}) with that of (a) malonic acid and (b) succinic acid. Likewise, scatterplots showing the relative abundances of C_5 in ΣC_2-C_{10} with those of (c) C_4 and (d) C_9 diacids in TSP samples collected during the NORC2015-10 cruise.

80

REFERENCES

- Gantt, B., Meskhidze, N., Zhang, Y., and Xu, J.: The effect of marine isoprene emissions on secondary organic aerosol and ozone formation in the coastal United States, *Atmospheric Environment*, 44, 115-121, 2010.
- 85 Kawamura, K., and Sakaguchi, F.: Molecular distribution of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics, *Journal of Geophysical Research Atmospheres*, 104, 3501-3509, 1999.
- Kourtchev, I., Copolovici, L., Claeys, M., and Maenhaut, W.: Characterization of atmospheric aerosols at a forested site in Central Europe, *Environmental science & technology*, 43, 4665-4671, 2009.
- Nguyen, T. B., Bateman, A. P., Bones, D. L., Nizkorodov, S. A., Laskin, J., and Laskin, A.: High-resolution mass spectrometry analysis of secondary organic aerosol generated by ozonolysis of isoprene, *Atmospheric Environment*, 44, 1032-1042, 2010.
- 90 Yang, L., Ray, M. B., and Liya, E. Y.: Photooxidation of dicarboxylic acids—Part II: Kinetics, intermediates and field observations, *Atmospheric Environment*, 42, 868-880, 2008.